

Junctophilin1 and Junctophilin2 in Assembly of Sarcoplasmic Reticulum

Subjects: Physiology

Contributor: Stefano Perni

Contraction of striated muscle is triggered by a massive release of calcium from the sarcoplasmic reticulum (SR) into the cytoplasm. This intracellular calcium release is initiated by membrane depolarization, which is sensed by voltage-gated calcium channels CaV1.1 (in skeletal muscle) and CaV1.2 (in cardiac muscle) in the plasma membrane (PM), which in turn activate the calcium-releasing channel ryanodine receptor (RyR) embedded in the SR membrane. This cross-communication between channels in the PM and in the SR happens at specialized regions, the SR-PM junctions, where these two compartments come in close proximity. Junctophilin1 and Junctophilin2 are responsible for the formation and stabilization of SR-PM junctions in striated muscle and actively participate in the recruitment of the two essential players in intracellular calcium release, CaV and RyR.

Keywords: striated muscle ; ER-PM junctions ; junctophilins ; excitation-contraction coupling

1. Introduction

Striated muscle evolved in early free-living invertebrates to confer locomotion to the individual and allow the search for food and the avoidance of predators or harmful environments. The term striated arises from the typical striated pattern of this tissue when observed in light and electron microscopy and defines both skeletal and cardiac muscle. This striation is the manifestation of the finely organized contractile apparatus in each of the functional contractile units of the muscle, the sarcomeres, which are arranged in series in the longitudinal direction of the muscle fiber. By the 1960s, it was suspected that a chemical activator, later identified as calcium ions (Ca^{2+}), was responsible for muscle contraction, but it was still puzzling how a soluble activator, with a relatively slow diffusion speed, could be responsible for the fast and uniform contraction of a muscle fiber that can be tens of microns thick. Experiments conducted by Andrew Huxley and colleagues on frog, lizard and crab skeletal muscle revealed that the minimal electrical stimulus necessary to achieve local contractions in the muscle fiber was lower in areas of the fiber that fell along specific sections of the sarcomere ^{[1][2]}. Such sections corresponded to regions where particular structures, called triads, were observed in electron microscopy ^[3] (**Figure 1**).

Ultrastructural studies of the triad revealed a tri-partite organization, hence the name triad, in which two enlarged regions of the sarcoplasmic reticulum (SR), called terminal cisternae, sandwich another membrane structure, the T-tubule, which is in direct continuity with the plasma membrane (PM) ^{[4][5]}. The continuity of the T-tubule with the plasma membrane ensures fast transmission of the membrane depolarization into the interior of the fiber, allowing for a uniform contraction. Therefore, the triad in skeletal muscle represents a highly specialized form of ER-PM junction in which the SR membrane and the T-tubule are joined together in close proximity and with considerable strength, since they have to withstand mechanical stress provided by the repetitive contractions and stretching of the muscle fiber, which in some cases can reach extreme levels ^{[6][7]}.

Mammalian ventricular cardiac muscle shows a similar organization to that of skeletal muscle with T-tubules and SR terminal cisternae. However, the specialized ER-PM junctions are arranged as dyads with a T-tubule contacting a single SR terminal cisterna ^[8] rather than triads (**Figure 1**).




Figure 1. Organization of ER-PM junctions in skeletal and cardiac muscle. Thin section electron micrographs illustrating the different organization of ER-PM junctions in skeletal and cardiac muscle. The sarcoplasmic reticulum and the T-tubule are pseudo-colored in yellow and pink, respectively. The gap separating the T-tubule and the SR membrane is pseudo-colored in blue. The T-tubule is absent in cardiac peripheral couplings since the SR is juxtaposed to the plasma membrane at the periphery of the fiber. Images are from Perni et al. ^[9] (triad), Lavorato et al. ^[10] (dyad) and Perni et al. ^[11] (peripheral coupling).

Triads in skeletal muscle and dyads and peripheral couplings in cardiac muscle are the sites at which membrane depolarization is translated into Ca^{2+} release from the SR and, eventually, muscle contraction. This process, called excitation–contraction (EC) coupling, requires the cross-talk between two main players: the ryanodine receptor (RyR), a highly conductive Ca^{2+} channel embedded in the SR membrane and responsible for the rapid release of Ca^{2+} from the SR, and the L-type Ca^{2+} channel in the plasma membrane (T-tubule), which senses membrane depolarization and activates the RyR. The way RyR is activated differs in the cardiac and skeletal muscle systems. In the former, the opening of the cardiac muscle L-Type channel $\text{Ca}_v1.2$ generates a rapid Ca^{2+} influx through the channel from the extracellular environment, causing a rapid increase in Ca^{2+} concentration in the narrow space separating the T-tubule and the SR in the dyad, and inducing the opening of the cardiac RyR isoform, RyR2. This mechanism is defined as calcium-induced calcium release (CICR) ^[12]. In skeletal muscle, the release of Ca^{2+} from the SR is directly triggered by the activation of the skeletal muscle L-type channel, $\text{Ca}_v1.1$. The voltage-induced conformational change in $\text{Ca}_v1.1$ is mechanically transmitted to RyR1 ^[13], causing its opening. This mechanism, which is independent of Ca^{2+} influx through $\text{Ca}_v1.1$, is known as voltage-induced calcium release (VICR).

In addition to the voltage sensor in the membrane (Ca_v) and the Ca^{2+} -releasing channel in the SR (RyR), the voltage-gated Ca^{2+} channels β subunits are also essential for EC coupling ^{[14][15]}. $\text{Ca}_v1.1$ and $\text{Ca}_v1.2$ channels bind to the skeletal $\beta1a$ and cardiac $\beta2a$, respectively, through their alpha interacting domain (AID) located in the intracellular loop connecting transmembrane domains I and II ^[16]. Beta subunits are crucial for facilitating the trafficking of the channel into the plasma membrane and for modulating the channel activity ^{[14][17]}. In skeletal muscle, the adapter protein Stac3 is also required for voltage-induced calcium release ^{[18][19]}. The exact role of Stac3 in skeletal muscle EC coupling is yet to be elucidated. Stac3 facilitates, but is not essential for, the membrane trafficking of $\text{Ca}_v1.1$ ^{[20][21]}; nonetheless, knocking out Stac3 completely abolishes the voltage-induced Ca^{2+} release ^{[18][21]}. The observation that Stac3 binds to the II-III intracellular loop of $\text{Ca}_v1.1$ ^[22], which is critical for the cross-talk between $\text{Ca}_v1.1$ and RyR1 ^[23], suggests that Stac3 might allow or facilitate the mechanical coupling between these two channels.

It appears evident that the association of all the EC coupling essential and accessory proteins and the efficient cross-talk between Ca_v1 in the plasma membrane and RyR in the SR require the accurate formation and organization of ER (SR)-PM junctions, making the proteins responsible for the organization of such junctions also essential for EC coupling.

2. The Junctophilin Family

Junctophilins (JPH1, JPH2, JPH3 and JPH4) were discovered by Takeshima and collaborators in the early 2000s in muscles and neurons ^{[24][25]}. All four isoforms of this family contain a C-terminal transmembrane domain that embeds in

the ER membrane and eight N-terminal domains called membrane occupation and recognition nexuses (MORN), thought to be responsible for the association with the plasma membrane (**Figure 2**).




Figure 2. Schematic representation of junctophilin's domain structure and its arrangement in the ER-PM junction. **(A)** Linear map showing the first (I through IV) and second (VII to VIII) set of MORN domains (in pink) separated by the joining domain (white). The α -helical domain (orange) follows MORN VIII and is separated from the transmembrane (TM, in cyan) by the long divergent domain (green). The numbers underneath the map indicate mutations associated with human cardiomyopathies identified in JPH2 and their relative positions. **(B)** Schematic representation of junctophilin's organization in the ER-PM junctions (adapted from Garbino et al. [26]). The MORN motifs associated with the internal leaflet of the plasma membrane and the C-terminal transmembrane domain embedded in the endo/sarcoplasmic reticulum membrane allow junctophilins to bridge the two membrane systems together. Different domains are color-coded as indicated in (A).

3. Recruitment of Junctional Proteins by Junctophilins1 and 2

3.1. Recruitment of Skeletal Muscle Junctional Proteins

JPH1 and JPH2, and specifically a region in the two junctophilins spanning approximately from the second half of the joining domain to the first half of the putative α -helical domain, co-immunoprecipitate with $\text{Ca}_v1.1$ [27]. A 20-residue sequence in the C-terminal domain of $\text{Ca}_v1.1$ is directly involved in the interactions with junctophilins and in the recruitment of $\text{Ca}_v1.1$ to triads [28]. Additionally, $\text{Ca}_v1.1$ is recruited to junctions formed by JPH2 when the proteins are expressed in non-muscle cell models together with the $\text{Ca}_v1.1$ auxiliary subunit $\beta 1a$ and Stac3 [29]. Overall, this indicates that JPH1 and JPH2 have an active role in recruiting the voltage-gated Ca^{2+} channel to triads by binding directly to the channel; the disruption of this interaction interferes with the assembly of the triad [27][28].

JPH1 co-immunoprecipitates with RyR1, a behavior that has not been observed for JPH2 [30]. Nonetheless, JPH1 KO mice can still perform EC coupling [31][32], suggesting that the presence of RyR1 in junctions does not depend solely on JPH1. Therefore, the recruitment of RyR1 in JPH2-induced junctions might be due to a weak interaction that is not detected in biochemical assays or requires the presence of additional proteins, with $\text{Ca}_v1.1$ being a likely candidate.

3.2. Recruitment of Cardiac Muscle Junctional Proteins

The cardiac L-type Ca^{2+} channel $\text{Ca}_v1.2$ co-immunoprecipitates with JPH2 [33][34], indicating that, as in skeletal muscle, JPH2 likely plays a role in recruiting the voltage-sensor channel in the dyads and peripheral couplings of cardiac muscle. Notably, the same C-terminal sequence identified as the $\text{Ca}_v1.1$ site of interaction with junctophilins, is conserved in $\text{Ca}_v1.2$ [28], suggesting that this sequence might also be involved in $\text{Ca}_v1.2$ –JPH2 interactions.

Differently from what was found with JPH2 and RyR1, co-immunoprecipitation was observed between JPH2 and RyR2 [33][34], indicating a stronger interaction between the two proteins. This interaction is disrupted by the E169K substitution located towards the N-terminal end of the JPH2 joining domain [35] and weakened by the R420Q mutation in RyR2 [36]. A stronger interaction with RyR2 might be required by JPH2 because it is the only junctophilin isoform expressed in cardiac muscle and possibly because cardiac muscle lacks the additional stabilization provided by the mechanical connection between RyR1 and Ca_v1.1 that exists in skeletal muscle [13].

4. Functional Studies on Junctophilins 1 and 2

4.1. Junctophilin 1

JPH1 knock-out mice die within 24 h after birth due to suckling defects leading to undernourishment. The suckling defect is likely due to muscle weakness since the neuronal suckling reflexes are normal in knock-out mice [37]. Functional studies on isolated hindlimb muscle showed abnormal twitch tension and a greater dependency on extracellular calcium in KO mice muscles, suggesting that a significant fraction of RyR1s in the junctional SR are not directly coupled with the Ca_v1.1 channels in the T-tubules and therefore operate via calcium-induced calcium release. Nonetheless, knock-out (KO) mice are still relatively mobile and show skeletal muscle-type EC coupling to a certain degree, indicating that JPH2 can support voltage-induced Ca²⁺ release in the absence of JPH1. From a structural point of view, although no major disorganization of the fiber is noticed at the light microscopy level, evident alterations are noticeable at the ultrastructural level [31][38]. In particular, the skeletal muscle of wt and JPH1 KO mice show a similar development in the embryonic stages until shortly after birth. At this age, wt muscle experiences a significant increase in JPH1 expression, which is temporally correlated with the transition from immature SR-PM junctions, mainly organized in dyads at this stage, into fully formed triads. This transition is absent in JPH1 KO muscle [31][38], suggesting that JPH2 is important in forming the dyads, while JPH1 has a crucial role in the conversion from dyads to triads in the fully mature skeletal muscle. The knocking down of junctophilins using sh-RNA, leads to the impairment of store-operated Ca²⁺ entry (SOCE), altered intracellular calcium release and intracellular calcium stores [39] and to a reduction in RyR1 and Ca_v1.1 co-clustering associated with a decrease in Ca_v1.1 membrane expression [27] both in myotubes and muscle fibers. In both these studies, a shRNA against both JPH1 and JPH2 was used; hence, it was impossible to distinguish each isoform's relative contribution to the resulting phenotype.

4.2. Junctophilin 2

JPH2 knock-out mice die in utero due to cardiac failure. Ultrastructural analyses on embryonic myotubes of KO mice revealed a substantial reduction in the number and extension of peripheral couplings [25]. To avoid the complication related to the early mortality of KO mice, van Oort et al. generated a conditional JPH2 knockdown mice model to assess the effect of JPH2-reduced expression in the mature heart [40]. Inducing JPH2 knockdown led to an increased frequency of heart failure events. At the cellular level, this was explained structurally by T-tubule remodeling and destabilization and disorganization of the dyads, and functionally by Ca_v1.2 and RyR2 uncoupling and the consequent reduction in the efficiency of calcium-induced calcium release. An increase in the frequency of calcium sparks was noticed in knocked down isolated cardiomyocytes, suggesting that JPH2 might also modulate RyR2 by reducing its activity.

A number of point mutations in JPH2 have been discovered in association with hypertrophic cardiomyopathy and atrial fibrillation. The localization of these mutations spans from the N-terminal MORN motifs to the divergent domain at the C-terminus, indicating that multiple regions of JPH2 are involved in supporting cardiac muscle structure and function.

Amino acid substitutions N101R and Y141H in the MORN IV and VI, respectively, and S165F in the joining domain, cause similar phenotypes such as JPH2 mislocalization, reduction in spontaneous Ca²⁺ signaling and increased cell size in HL-1 and H9c2 cell lines [41]. Mutations Y141H and S165F were also tested in skeletal muscle myotubes [42], where they were found to induce myocytes hypertrophy, reduce EC coupling gain and increase intracellular Ca²⁺ concentration. Additionally, Y141H but not S165F pathogenically increased store-operated calcium entry [42]. Mutation E169K, located in the joining domain, causes weaker binding between JPH2 and RyR2 and increased spontaneous Ca²⁺ leakage from the SR in the form of a spontaneous Ca²⁺ release and increased Ca²⁺ sparks in isolated cardiomyocytes from a pseudo-knock-in mouse model [35]. The A405S mutation is located in the putative α -helical region of JPH2. The equivalent mutation introduced in mice (A399S) resulted in cardiomyocytes with an irregular T-tubule pattern but otherwise relatively normal Ca²⁺ signaling with only a moderate increase in sarco-endoplasmic reticulum Ca²⁺ ATPase (SERCA) activity.

5. Post-Transcriptional Regulation of JPH1 and JPH2

Conditions of elevated cytosolic Ca^{2+} concentrations lead to fragmentation of junctophilins 1 and 2 in skeletal and cardiac muscle. Murphy and colleagues [43] determined that exposure to elevated ($\geq 20 \mu\text{M}$) intracellular $[\text{Ca}^{2+}]$ for 60 min led to the almost complete loss of full-length JPH1 and JPH2 in skeletal muscle fibers. This loss is mirrored by a loss of contractile force in skinned skeletal muscle fibers after just one minute of exposure to $40 \mu\text{M}$ Ca^{2+} . The same authors also observed fragmentation of JPH1 after raising the intracellular $[\text{Ca}^{2+}]$ by supraphysiological stimulation of the muscle fiber. Interestingly, the proteolysis of JPH1 temporally matched the autolytic activation of calpain- μ (calpain1). The link between calpain, specifically calpain1, and JPH1 cleavage was recently confirmed by data from Tamminen and colleagues in patients with malignant hyperthermia susceptibility (MHS) and muscle cell lines [44].

After the identification of several putative calpain binding sites also in JPH2 [34][45], the implication of calpain in the proteolytic regulation of JPH2 was verified using calpain inhibitors to rescue the loss of JPH2 in an inducible heart failure mouse model and in mice cardiomyocytes after ischemia/reperfusion [46]. Akin to what was shown by Tamminen and colleagues in skeletal muscle, Guo and collaborators also observed that digestion by calpain1 releases several fragments of JPH2 [34]. Among these fragments, a ~ 75 kDa N-terminal peptide (JPH2-NTP), generated by calpain1 cleavage at residues R565/T566 in the JPH2 divergent region, migrates into the nucleus, where it binds to TATA box regions and interacts with the transcription machinery [45]. Altogether, these results indicate a fine modulation of junctophilin 1 and 2 as a way to regulate intracellular Ca^{2+} homeostasis and possibly reduce EC coupling gain in conditions of excessive intracellular Ca^{2+} concentration.

6. New Insights from Deep Learning Protein Structure Prediction

Remarkable advancements in protein folding prediction were recently achieved by the artificial intelligence software AlphaFold2 [47]. AlphaFold2 is a giant leap forward in the reliability of protein folding prediction compared to similar existing software [48], and it has already been used to predict the structure of nearly the entire human proteome. Based on AlphaFold2 prediction models, junctophilin 1 and 2 show a similar 3D structure, also shared by the neuronal isoforms JPH3 and JPH4 (UniProt protein ID: Q9HDC5, Q9BR39, Q8WXH2, Q96JJ6 for human JPH1, JPH2, JPH3 and JPH4, respectively). The structure of the most ordered domains of junctophilin, specifically the MORN repeats, the α -helical region and the transmembrane domain, are predicted with high confidence by AlphaFold2. In contrast, the joining and divergent domains are likely disordered, at least in the isolated protein, and the structure cannot be predicted with reasonable confidence. According to the prediction model (Figure 3), the MORN domains are arranged in an extended “halfpipe” configuration, with the α -helical domain lying on the convex side of this half-pipe (Figure 3B,C) and establishing interactions with charged residues in the MORN domains (see the zoomed-in region in Figure 3C for an example).




Figure 3. Structure of the MORNs and α -helical domains of human junctophilin1 and junctophilin2. (A) Schematic representation of junctophilin domain as shown in Figure 2; the solid red lines indicate the regions for which the structure is predicted with high fidelity by AlphaFold2 and illustrated in (B,C). (B,C) predicted structures of the MORNs- α -helical domains of junctophilin1 (B) and junctophilin2 (C). The α -helical domain (in red) lies on the convex side of the MORN domains half-pipe structure (in blue) in both junctophilin1 and junctophilin2. β -sheet hairpins forming MORN domains I and VIII (green parentheses) and the position of the N-terminal end (N-t) and C-terminal end (C-t) of the joining domain (in pink), which is absent in this representation, are indicated in (B). The inset in (C) shows some of the residues that form the hydrogen bonds that stabilize the association between the MORN domains and the α -helical domain of junctophilin2.

The structure obtained using AlphaFold2 is substantially different from what was previously predicted using RaptorX software by Gross and colleagues [49]. In the structure described by Gross et al., the α -helical domain extends beyond the MORN domains without interacting with them at all. However, AlphaFold2 software is considered to be more accurate than most (if not all) of the currently existing structure-predicting software, especially for proteins for which no homologous structures exist [50][51], and the reciprocal arrangement of JPH2 MORN motifs and α -helical domain predicted by AlphaFold2 agrees with data from Li and collaborators [52] based on the crystal structure of the protein MORN4. MORN4 contains a series of MORN motifs arranged in a half-pipe configuration followed by a brief α -helical region. The helical region stabilizes the MORN domains by lying over part of the convex side of the half-pipe. The structure solved by Li and colleagues is in many ways very similar to the sequence predicted by AlphaFold2 for JPHs.

Furthermore, in MORN 4, the concave side of the MORN half-pipe structure, containing most of the conserved residues that define the MORN domain, engages in the binding with the α -helical region of myosin3a. It is conceivable that the concave side of the junctophilin MORN motifs could also participate in protein–protein interactions with components of the EC coupling machinery. The particular arrangement of the α -helical domain with respect to the MORN motifs predicted by AlphaFold2 and suggested by the observations of Li and colleagues challenges the classic view of the α -helical domain as the spacer that spans most of the junctional gap (see schematic representation in **Figure 2**) and points to the divergent domain as the region that most likely fulfills this role.

References

1. A. F. Huxley; R. E. Taylor; Local activation of striated muscle fibres. *The Journal of Physiology* **1958**, *144*, 426-441, [10.1113/jphysiol.1958.sp006111](https://doi.org/10.1113/jphysiol.1958.sp006111).
2. A F Huxley; Local activation of striated muscle from the frog and the crab.. *The Journal of Physiology* **1957**, *135*, 269-300, .
3. Keith R. Porter; George E. Palade; STUDIES ON THE ENDOPLASMIC RETICULUM. *Journal of Cell Biology* **1957**, *3*, 269-300, [10.1083/jcb.3.2.269](https://doi.org/10.1083/jcb.3.2.269).
4. C Franzini-Armstrong; FINE STRUCTURE OF SARCOPLASMIC RETICULUM AND TRANVERSE TUBULAR SYSTEM IN MUSCLE FIBERS.. *Federation proceedings* **1964**, *23*, 887-895, .
5. Clara Franzini-Armstrong; STUDIES OF THE TRIAD. *Journal of Cell Biology* **1970**, *47*, 488-499, [10.1083/jcb.47.2.488](https://doi.org/10.1083/jcb.47.2.488).
6. Matthew T. Close; Stefano Perni; Clara Franzini-Armstrong; David Cundall; Highly extensible skeletal muscle in snakes. *Journal of Experimental Biology* **2013**, *217*, 2445-8, [10.1242/jeb.097634](https://doi.org/10.1242/jeb.097634).
7. L. C. Rome; D. A. Syme; S. Hollingworth; S. L. Lindstedt; S. M. Baylor; The whistle and the rattle: the design of sound producing muscles.. *Proceedings of the National Academy of Sciences* **1996**, *93*, 8095-8100, [10.1073/pnas.93.15.8095](https://doi.org/10.1073/pnas.93.15.8095).
8. Clara Franzini-Armstrong; Feliciano Protasi; Venkat Ramesh; Shape, Size, and Distribution of Ca²⁺ Release Units and Couplons in Skeletal and Cardiac Muscles. *Biophysical Journal* **1999**, *77*, 1528-1539, [10.1016/s0006-3495\(99\)77000-1](https://doi.org/10.1016/s0006-3495(99)77000-1).
9. Stefano Perni; Kurt C. Marsden; Matias Escobar; Stephen Hollingworth; Stephen M. Baylor; Clara Franzini-Armstrong; Structural and functional properties of ryanodine receptor type 3 in zebrafish tail muscle. *Journal of General Physiology* **2015**, *145*, 173-184, [10.1085/jgp.201411303](https://doi.org/10.1085/jgp.201411303).
10. Manuela Lavorato; Tai-Qin Huang; Venkat Ramesh Iyer; Stefano Perni; Gerhard Meissner; Clara Franzini-Armstrong; Dyad content is reduced in cardiac myocytes of mice with impaired calmodulin regulation of RyR2. *Journal of Muscle Research and Cell Motility* **2015**, *36*, 205-214, [10.1007/s10974-015-9405-5](https://doi.org/10.1007/s10974-015-9405-5).
11. Stefano Perni; V. Ramesh Iyer; Clara Franzini-Armstrong; Ultrastructure of cardiac muscle in reptiles and birds: optimizing and/or reducing the probability of transmission between calcium release units. *Journal of Muscle Research*

12. A Fabiato; Time and calcium dependence of activation and inactivation of calcium-induced release of calcium from the sarcoplasmic reticulum of a skinned canine cardiac Purkinje cell.. *Journal of General Physiology* **1985**, 85, 247-289, [10.1085/jgp.85.2.247](https://doi.org/10.1085/jgp.85.2.247).
13. Barbara A Block; Toshiaki Imagawa; Kevin Campbell; Clara Franziniarmstrong; Structural evidence for direct interaction between the molecular components of the transverse tubule/sarcoplasmic reticulum junction in skeletal muscle.. *Journal of Cell Biology* **1988**, 107, 2587-2600, [10.1083/jcb.107.6.2587](https://doi.org/10.1083/jcb.107.6.2587).
14. Johann Schredelseker; Anamika Dayal; Thorsten Schwerte; Clara Franzini-Armstrong; Manfred Grabner; Proper Restoration of Excitation-Contraction Coupling in the Dihydropyridine Receptor β 1-null Zebrafish Relaxed Is an Exclusive Function of the β 1a Subunit. *Journal of Biological Chemistry* **2008**, 284, 1242-1251, [10.1074/jbc.m807767200](https://doi.org/10.1074/jbc.m807767200).
15. Petra Weissgerber; Brigitte Held; Wilhelm Bloch; Lars Kaestner; Kenneth R. Chien; Bernd K. Fleischmann; Peter Lipp; Veit Flockerzi; Marc Freichel; Reduced Cardiac L-Type Ca^{2+} Current in $\text{Ca}_v\beta 2^{-/-}$ Embryos Impairs Cardiac Development and Contraction With Secondary Defects in Vascular Maturation. *Circulation Research* **2006**, 99, 749-757, [10.1161/01.res.0000243978.15182.c1](https://doi.org/10.1161/01.res.0000243978.15182.c1).
16. Marion Pragnell; Michel De Waard; Yasuo Mori; Tsutomu Tanabe; Terry P. Snutch; Kevin Campbell; Calcium channel β -subunit binds to a conserved motif in the I-II cytoplasmic linker of the α 1-subunit. *Nature* **1994**, 368, 67-70, [10.1038/368067a0](https://doi.org/10.1038/368067a0).
17. Andy J. Chien; Xiaolan Zhao; Roman E. Shirokov; Tipu S. Puri; Chan Fong Chang; Dandan Sun; Eduardo Rios; M. Marlene Hosey; Roles of a Membrane-localized β Subunit in the Formation and Targeting of Functional L-type Ca^{2+} Channels. *Journal of Biological Chemistry* **1995**, 270, 30036-30044, [10.1074/jbc.270.50.30036](https://doi.org/10.1074/jbc.270.50.30036).
18. Benjamin R. Nelson; Fenfen Wu; Yun Liu; Douglas M. Anderson; John McAnally; Weichun Lin; Stephen Cannon; Rhonda Bassel-Duby; Eric N. Olson; Skeletal muscle-specific T-tubule protein STAC3 mediates voltage-induced Ca^{2+} release and contractility. *Proceedings of the National Academy of Sciences* **2013**, 110, 11881-11886, [10.1073/pnas.1310571110](https://doi.org/10.1073/pnas.1310571110).
19. Eric Horstick; Jeremy Linsley; James J. Dowling; Michael A. Hauser; Kristin K. McDonald; Allison Ashley-Koch; Louis Saint-Amant; Akhila Satish; Wilson W. Cui; Weibin Zhou; et al. Stac3 is a component of the excitation-contraction coupling machinery and mutated in Native American myopathy. *Nature Communications* **2013**, 4, 1-11, [10.1038/ncomms2952](https://doi.org/10.1038/ncomms2952).
20. Alexander Polster; Stefano Perni; Hicham Bichraoui; Kurt G. Beam; Stac adaptor proteins regulate trafficking and function of muscle and neuronal L-type Ca^{2+} channels. *Proceedings of the National Academy of Sciences* **2014**, 112, 602-606, [10.1073/pnas.1423113112](https://doi.org/10.1073/pnas.1423113112).
21. Alexander Polster; Benjamin R. Nelson; Eric N. Olson; Kurt G. Beam; Stac3 has a direct role in skeletal muscle-type excitation-contraction coupling that is disrupted by a myopathy-causing mutation. *Proceedings of the National Academy of Sciences* **2016**, 113, 10986-10991, [10.1073/pnas.1612441113](https://doi.org/10.1073/pnas.1612441113).
22. Alexander Polster; Benjamin R. Nelson; Symeon Papadopoulos; Eric N. Olson; Kurt G. Beam; Stac proteins associate with the critical domain for excitation-contraction coupling in the II-III loop of $\text{Ca}_v1.1$. *Journal of General Physiology* **2018**, 150, 613-624, [10.1085/jgp.201711917](https://doi.org/10.1085/jgp.201711917).
23. Tsutomu Tanabe; Kurt G. Beam; Brett A. Adams; Tetsuhiro Niidome; Shosaku Numa; Regions of the skeletal muscle dihydropyridine receptor critical for excitation-contraction coupling. *Nature* **1990**, 346, 567-569, [10.1038/346567a0](https://doi.org/10.1038/346567a0).
24. Miyuki Nishi; Hiroyuki Sakagami; Shinji Komazaki; Hisatake Kondo; Hiroshi Takeshima; Coexpression of junctophilin type 3 and type 4 in brain. *Molecular Brain Research* **2003**, 118, 102-110, [10.1016/s0169-328x\(03\)00341-3](https://doi.org/10.1016/s0169-328x(03)00341-3).
25. H Takeshima; Junctophilins A Novel Family of Junctional Membrane Complex Proteins. *Molecular Cell* **2000**, 6, 11-22, [10.1016/s1097-2765\(00\)00003-4](https://doi.org/10.1016/s1097-2765(00)00003-4).
26. Alejandro Garbino; Ralph J. Van Oort; Sayali S. Dixit; Andrew P. Landstrom; Michael J. Ackerman; Xander H. T. Wehrens; Molecular evolution of the junctophilin gene family. *Physiological Genomics* **2009**, 37, 175-186, [10.1152/physiolgenomics.00017.2009](https://doi.org/10.1152/physiolgenomics.00017.2009).
27. Lucia Golini; Christophe Chouabe; Christine Berthier; Vincenza Cusimano; Mara Fornaro; Robert Bonvallet; Luca Formoso; Emiliana Giacomello; Vincent Jacquemond; Vincenzo Sorrentino; et al. Junctophilin 1 and 2 Proteins Interact with the L-type Ca^{2+} Channel Dihydropyridine Receptors (DHPs) in Skeletal Muscle. *Journal of Biological Chemistry* **2011**, 286, 43717-43725, [10.1074/jbc.m111.292755](https://doi.org/10.1074/jbc.m111.292755).
28. Tsutomu Nakada; Toshihide Kashihara; Masatoshi Komatsu; Katsuhiko Kojima; Toshikazu Takeshita; Mitsuhiro Yamada; Physical interaction of junctophilin and the $\text{Ca}_v1.1$ C terminus is crucial for skeletal muscle contraction. *Proceedings of the National Academy of Sciences* **2018**, 115, 4507-4512, [10.1073/pnas.1716649115](https://doi.org/10.1073/pnas.1716649115).

29. Stefano Perni; Manuela Lavorato; Kurt G. Beam; De novo reconstitution reveals the proteins required for skeletal muscle voltage-induced Ca²⁺ release. *Proceedings of the National Academy of Sciences* **2017**, *114*, 13822-13827, [10.1073/pnas.1716461115](https://doi.org/10.1073/pnas.1716461115).
30. Phimister, A.J.; Lango, J.; Lee, E.H.; Ernst-Russell, M.A.; Takeshima, H.; Ma, J.; Allen, P.D.; Pessah, I.N. Conformation-dependent stability of junctophilin 1 (JP1) and ryanodine receptor type 1 (RyR1) channel complex is mediated by their hyper-reactive thiols. *J. Biol. Chem.* 2007, *282*, 8667–8677.
31. Koichi Ito; Shinji Komazaki; Kazushige Sasamoto; Morikatsu Yoshida; Miyuki Nishi; Kenji Kitamura; Hiroshi Takeshima; Deficiency of triad junction and contraction in mutant skeletal muscle lacking junctophilin type 1. *Journal of Cell Biology* **2001**, *154*, 1059-1068, [10.1083/jcb.200105040](https://doi.org/10.1083/jcb.200105040).
32. Stefano Perni; Kurt Beam; Junctophilins 1, 2, and 3 all support voltage-induced Ca²⁺ release despite considerable divergence. *Journal of General Physiology* **2022**, *154*, e202113024, [10.1085/jgp.202113024](https://doi.org/10.1085/jgp.202113024).
33. Min Jiang; Mei Zhang; Maureen Howren; Yuhong Wang; Alex Tan; Ravi C. Balijepalli; Jose Huizar; Gea-Ny Tseng; JPH-2 interacts with Cai-handling proteins and ion channels in dyads: Contribution to premature ventricular contraction-induced cardiomyopathy. *Heart Rhythm* **2015**, *13*, 743-752, [10.1016/j.hrthm.2015.10.037](https://doi.org/10.1016/j.hrthm.2015.10.037).
34. Ang Guo; Duane Hall; Caimei Zhang; Tianqing Peng; Jordan Miller; William Kutschke; Chad E. Grueter; Frances L. Johnson; Richard Lin; Long-Sheng Song; et al. Molecular Determinants of Calpain-dependent Cleavage of Junctophilin-2 Protein in Cardiomyocytes. *Journal of Biological Chemistry* **2015**, *290*, 17946-17955, [10.1074/jbc.m115.652396](https://doi.org/10.1074/jbc.m115.652396).
35. David L. Beavers; Wei Wang; Sameer Ather; Niels Voigt; Alejandro Garbino; Sayali S. Dixit; Andrew Landstrom; Na Li; Qionglin Wang; Iacopo Olivotto; et al. Mutation E169K in Junctophilin-2 Causes Atrial Fibrillation Due to Impaired RyR2 Stabilization. *Journal of the American College of Cardiology* **2013**, *62*, 2010-2019, [10.1016/j.jacc.2013.06.052](https://doi.org/10.1016/j.jacc.2013.06.052).
36. Liheng Yin; Alexandra Zahradnikova Jr; Riccardo Rizzetto; Simona Boncompagni; Camille Rabesahala de Meritens; Yadan Zhang; Pierre Joanne; Elena Marques Sule; Yuriana Aguilar-Torres; Miguel Fernandez-Tenorio; et al. Impaired Binding to Junctophilin2 and Nanostructural Alteration in CPVT Mutation. *Circulation Research* **2021**, *129*, e35-e52, [10.1161/circresaha.121.319094](https://doi.org/10.1161/circresaha.121.319094).
37. Ito, K.; Komazaki, S.; Sasamoto, K.; Yoshida, M.; Nishi, M.; Kitamura, K.; Takeshima, H. Deficiency of triad junction and contraction in mutant skeletal muscle lacking junctophilin type 1. *J. Cell Biol.* 2001, *154*, 1059–1067.
38. Shinji Komazaki; Koichi Ito; Hiroshi Takeshima; Hiroaki Nakamura; Deficiency of triad formation in developing skeletal muscle cells lacking junctophilin type 1. *FEBS Letters* **2002**, *524*, 225-229, [10.1016/s0014-5793\(02\)03042-9](https://doi.org/10.1016/s0014-5793(02)03042-9).
39. Yutaka Hirata; Marco Brotto; Noah Weisleder; Yi Chu; Pei-Hui Lin; Xiaoli Zhao; Angela Thornton; Shinji Komazaki; Hiroshi Takeshima; Jianjie Ma; et al. Uncoupling Store-Operated Ca²⁺ Entry and Altered Ca²⁺ Release from Sarcoplasmic Reticulum through Silencing of Junctophilin Genes. *Biophysical Journal* **2006**, *90*, 4418-4427, [10.1529/biophysj.105.076570](https://doi.org/10.1529/biophysj.105.076570).
40. Ralph J. Van Oort; Alejandro Garbino; Wei Wang; Sayali S. Dixit; Andrew Landstrom; Namit Gaur; Angela C. De Almeida; Darlene G. Skapura; Yoram Rudy; Alan R. Burns; et al. Disrupted Junctional Membrane Complexes and Hyperactive Ryanodine Receptors After Acute Junctophilin Knockdown in Mice. *Circulation* **2011**, *123*, 979-988, [10.1161/circulationaha.110.006437](https://doi.org/10.1161/circulationaha.110.006437).
41. Andrew Landstrom; Noah Weisleder; Karin B. Batalden; J. Martijn Bos; David J. Tester; Steve R. Ommen; Xander Wehrens; William C. Claycomb; Jae-Kyun Ko; MoonSun Hwang; et al. Mutations in JPH2-encoded junctophilin-2 associated with hypertrophic cardiomyopathy in humans. *Journal of Molecular and Cellular Cardiology* **2007**, *42*, 1026-1035, [10.1016/j.yjmcc.2007.04.006](https://doi.org/10.1016/j.yjmcc.2007.04.006).
42. Jin Seok Woo; Chung-Hyun Cho; Keon Jin Lee; Do Han Kim; Jianjie Ma; Eun Hui Lee; Hypertrophy in Skeletal Myotubes Induced by Junctophilin-2 Mutant, Y141H, Involves an Increase in Store-operated Ca²⁺ Entry via Orai1*. *Journal of Biological Chemistry* **2012**, *287*, 14336-14348, [10.1074/jbc.m111.304808](https://doi.org/10.1074/jbc.m111.304808).
43. R. M. Murphy; T. L. Dutka; D. Horvath; J. R. Bell; L. M. Delbridge; G. D. Lamb; Ca²⁺-dependent proteolysis of junctophilin-1 and junctophilin-2 in skeletal and cardiac muscle. *The Journal of Physiology* **2012**, *591*, 719-729, [10.1113/jphysiol.2012.243279](https://doi.org/10.1113/jphysiol.2012.243279).
44. Eshwar R. Tammineni; Lourdes Figueroa; Natalia Kraeva; Carlo Manno; Carlos A. Ibarra; Amira Klip; Sheila Riazi; Eduardo Rios; Fragmentation and roles of junctophilin1 in muscle of patients with cytosolic leak of stored calcium. *Journal of General Physiology* **2021**, *154*, 2021ecc32, [10.1085/jgp.2021ecc32](https://doi.org/10.1085/jgp.2021ecc32).
45. Ang Guo; Yihui Wang; Biyi Chen; Yunhao Wang; Jinxiang Yuan; Liyang Zhang; Duane Hall; Jennifer Wu; Yun Shi; Qi Zhu; et al. E-C coupling structural protein junctophilin-2 encodes a stress-adaptive transcription regulator. *Science* **2018**, *362*, science.aan3303, [10.1126/science.aan3303](https://doi.org/10.1126/science.aan3303).

46. Chia-Yen C. Wu; Biyi Chen; Ya-Ping Jiang; Zhiheng Jia; Dwight W. Martin; Shengnan Liu; Emilia Entcheva; Long-Sheng Song; Richard Z. Lin; Calpain-Dependent Cleavage of Junctophilin-2 and T-Tubule Remodeling in a Mouse Model of Reversible Heart Failure. *Journal of the American Heart Association* **2014**, 3, e000527, [10.1161/jaha.113.000527](https://doi.org/10.1161/jaha.113.000527).
47. John Jumper; Richard Evans; Alexander Pritzel; Tim Green; Michael Figurnov; Olaf Ronneberger; Kathryn Tunyasuvunakool; Russ Bates; Augustin Žídek; Anna Potapenko; et al. Highly accurate protein structure prediction with AlphaFold. *Nature* **2021**, 596, 583-589, [10.1038/s41586-021-03819-2](https://doi.org/10.1038/s41586-021-03819-2).
48. Ewen Callaway; 'It will change everything': DeepMind's AI makes gigantic leap in solving protein structures. *Nature* **2020**, 588, 203-204, [10.1038/d41586-020-03348-4](https://doi.org/10.1038/d41586-020-03348-4).
49. Polina Gross; Jaslyn Johnson; Carlos M. Romero; Deborah M. Eaton; Claire Poulet; Jose Sanchez-Alonso; Carla Lucarelli; Jean Ross; Andrew A. Gibb; Joanne F. Garbincius; et al. Interaction of the Joining Region in Junctophilin-2 With the L-Type Ca²⁺ Channel Is Pivotal for Cardiac Dyad Assembly and Intracellular Ca²⁺ Dynamics. *Circulation Research* **2021**, 128, 92-114, [10.1161/circresaha.119.315715](https://doi.org/10.1161/circresaha.119.315715).
50. Andrew W. Senior; Richard Evans; John Jumper; James Kirkpatrick; Laurent Sifre; Tim Green; Chongli Qin; Augustin Žídek; Alexander W. R. Nelson; Alex Bridgland; et al. Improved protein structure prediction using potentials from deep learning. *Nature* **2020**, 577, 706-710, [10.1038/s41586-019-1923-7](https://doi.org/10.1038/s41586-019-1923-7).
51. Hong Su; Wenkai Wang; Zongyang Du; Zhenling Peng; Shang-Hua Gao; Ming-Ming Cheng; Jianyi Yang; Improved Protein Structure Prediction Using a New Multi-Scale Network and Homologous Templates. *Advanced Science* **2021**, 8, 2102592, [10.1002/adv.202102592](https://doi.org/10.1002/adv.202102592).
52. Jianchao Li; Haiyang Liu; Manmeet H. Raval; Jun Wan; Christopher M. Yengo; Wei Liu; Mingjie Zhang; Structure of the MORN4/Myo3a Tail Complex Reveals MORN Repeats as Protein Binding Modules. *Structure* **2019**, 27, 1366-1374.e3, [10.1016/j.str.2019.06.004](https://doi.org/10.1016/j.str.2019.06.004).

Retrieved from <https://encyclopedia.pub/entry/history/show/47907>