

Mycoplasma genitalium Provokes Seminal Inflammation

Subjects: Andrology

Contributor: Stanislav Tjagur

To investigate the prevalence of urethritis-associated STIs (chlamydia, gonorrhoeae, *Mycoplasma genitalium*, trichomoniasis) among infertile males; to analyze the effect of STIs on semen parameters and blood PSA. Case-control study.

Keywords: sexually transmitted infections ; sperm functions ; male ; infertility ; Mycoplasma genitalium

1. Introduction

Infertility is defined as the inability of a couple to become pregnant despite unprotected intercourse for a period of more than twelve months ^[1]. The prevalence rate of infertility varies from 3.5% to 16.7% in more developed countries and from 6.9% to 9.3% in less-developed countries, with an estimated overall median prevalence of 9% ^[2]. In 50% of involuntarily childless couples, a male infertility-associated factor is found usually together with abnormal semen parameters ^[3].

Urogenital infections and inflammation are accepted contributing factors of male infertility in 6.9 to 16% of men ^{[4][5]}. Inflammation and oxidative stress could serve as core mechanisms linking STI with male infertility ^[6]. The impact of sexually transmitted infections (STI) on male fertility is a controversial topic. According to Ochsendorf et al., depending on the local prevalence of STIs and the availability of medical care, the impact of STIs on male urogenital system and fertility (as a consequence of infection) may appear regionally different ^[7]. Another problem is the distinction of the impact of STI agents between infection with pathological consequences and contamination without relevance ^[7]. Inconsistent diagnostic criteria applied to date could also explain the controversy about the role of infection and inflammation in the genital tract as a cause of infertility ^[8]. For example, using detection methods for *Chlamydia trachomatis* other than nucleic acid amplification tests (NAAT) including culture or serological methods is of significant concern. Culture for *C. trachomatis* detects only viable infectious chlamydial elementary bodies and the sensitivity is at most 70–85%. The major drawback of serological tests for *C. trachomatis* is that a positive antibody test will not distinguish a previous from a current infection ^[9].

Ahmadi et al. published a systematic review and meta-analysis on the association of *C. trachomatis* with infertility and clinical manifestations. The authors concluded that urogenital *C. trachomatis* prevalence was significantly higher in the infertile men compared with the fertile men (overall odds ratio 2.2, 95% confidence interval 1.3–3.7) and in symptomatic men compared with asymptomatic men (overall odds ratio 4.9, 95% confidence interval 1.1–21.7). However, the authors guided the attention to the fact that many studies were excluded from the analyses because the detection method was other than NAAT. This may suggest participation bias in the generalization of the meta-analysis results. Secondly, the number of included studies in males in both the fertile-infertile and the symptomatic-asymptomatic groups was fewer than 10 and therefore insufficient for a good meta-analysis and an accurate conclusion ^[10].

Less is known about the association between male infertility and other STIs. There is only limited information about the role of *Trichomonas vaginalis* in male infertility ^[11]. The lack of studies on *Neisseria gonorrhoeae* is likely due to the rarity of this infection in developed countries ^[12]. Little work has been carried out on the possible effects of the newest STI-causing agent *Mycoplasma genitalium* on male infertility, too ^{[13][14]}. To sum up, the association between STIs and male infertility still needs additional investigations, especially concerning *M. genitalium*.

2. Basic Clinical and Semen Parameters of Study Subjects

The basic parameters of study subjects are presented in **Table 1**.

Table 1. Basic and semen parameters of the unselected study and control groups (without exclusion of patients for hormonal, testicular volume, and infertility causal factors).

Parameter	Study Group, Infertile Men (n = 2000)	Control Group, Fertile Men (n = 248)	p-Value
	Median (Range); 25th Centile; 75th Centile	Median (Range); 25th Centile; 75th Centile	
Basic parameters			
Age in years	32.5 (18.2–49.8); 32.3 to 32.8 ^{\$}	31.8 (20.0–50.0); 31.1 to 32.6 ^{\$}	0.081 ^{###}
STI status			
→ Patients without STI	1954 (97.7%)	244 (98.4%)	0.649 ^{##}
→ <i>C. trachomatis</i> and <i>M. genitalium</i> positive patients together	46 (2.3%)	4 (1.6%)	0.649 ^{##}
→ <i>C. trachomatis</i> positive patients	24 (1.2%)	4 (1.6%)	0.541 ^{##}
→ <i>M. genitalium</i> positive patients	22 (1.1%)	0 (0.0%)	0.162 ^{##}
Bitesticular volume, in mL-s	45.0 (0.0–100.0); 44.6 to 45.5 ^{\$,A}	47.3 (23.0–100.0); 46.1 to 48.5 ^{\$,A}	<0.001 ^{###}
→ Left testicle's volume, in mL-s	22.0 (0.0–50.0); 21.8 to 22.3 ^{\$,A}	22.9 (0.0–50.0); 22.2 to 23.6 ^{\$,A}	0.016 ^{###}
→ Right testicle's volume, in mL-s	23.0 (0.0–50.0); 22.8 to 23.3 ^{\$,A}	24.4 (6.0–50.0); 23.7 to 25.1 ^{\$,A}	<0.001 ^{###}
STI in anamnesis	459 (23.0%)	51 (20.6%)	0.422 ^{##}
→ <i>N. gonorrhoeae</i> in anamnesis	95 (4.8%)	7 (2.8%)	0.197 ^{##}
→ <i>C. trachomatis</i> in anamnesis	259 (13.0%)	27 (10.9%)	0.419 ^{##}
→ <i>T. vaginalis</i> in anamnesis	85 (4.2%)	9 (3.6%)	0.739 ^{##}
→ <i>Lues</i> in anamnesis	15 (0.8%)	1 (0.4%)	1 ^{##}
→ HSV in anamnesis	47 (2.4%)	12 (4.8%)	0.032 ^{##}
→ HPV in anamnesis	68 (3.4%)	11 (4.4%)	0.364 ^{##}
Patients with varicocele	467 (23.4%)	61 (24.6%)	0.691 ^{##}
→ Left-sided grade 1 varicocele–	173 (8.7%)	25 (10.1%)	0.668 ^{##}
→ Left-sided grade 2 varicocele	211 (10.6%)	27 (10.9%)	0.777 ^{##}
→ Left-sided grade 3 varicocele	32 (1.6%)	4 (1.6%)	1 ^{##}
→ Right-sided varicocele	5 (0.3%)	0 (0.0%)	1 ^{##}

Parameter	Study Group, Infertile Men (n = 2000)	Control Group, Fertile Men (n = 248)	p-Value
	Median (Range); 25th Centile; 75th Centile	Median (Range); 25th Centile; 75th Centile	
→ Bilateral varicocele	24 (1.2%)	3 (1.2%)	1 ^{##}
→ Operated varicocele	22 (1.1%)	2 (0.8%)	1 ^{##}
Smokers	667 (33.4%)	74 (31.1%)	0.513 ^{##}
Concentration of FSH in serum, in U/L	3.8 (0.1–74.1); 2.6; 5.6 ^B	3.6 (0.6–15.2); 2.6; 5.1	0.065 [#]
Concentration of LH in serum, in U/L	3.9 (0.1–25.9); 2.8; 5.4 ^C	3.6 (0.6–11.7); 2.4; 4.6 ^C	<0.001 [#]
Concentration of total testosterone in serum, in nmol/L	16.0 (0.7–47.8); 12.4; 20.4 ^D	16.6 (6.0–49.3); 12.9; 20.2	0.489 [#]
PSA in serum, in µg/L	0.71 (0.11–8.34); 0.52; 0.99 ^E	0.69 (0.14–7.58); 0.47; 0.96 ^E	0.053 [#]
Height, cm	181.1 (153.0–209.0); 180.8 to 181.4 ^{S, F}	180.8 (167.0–198.0); 180.0 to 181.5 ^{\$}	0.468 ^{###}
Weight, kg	85.1 (47.0–189.0); 75.7; 95.5 ^G	82.0 (52.0–139.0); 74.5; 90.9	0.005 [#]
BMI	26.0 (14.9–76.2); 23.6; 28.7 ^H	24.9 (18.0–42.4); 22.9; 27.8	0.003 [#]
→ Overweighed patients (BMI > 24.9)	812 (42.3%) ^H	85 (34.3%)	0.016 ^{##}
→ Obese patients (BMI > 29.9)	340 (17.7%) ^H	34 (13.7%)	0.129 ^{##}
Waist circumference, cm	93.0 (62.0–152.0); 85.0; 101.0 ^I	90.0 (68.5–124.0); 83.5; 97.1 ^I	<0.001 [#]
Semen parameters			
Abstinence before semen analysis, in days	4.0 (0.0–56.0); 3.0; 4.0 ^J	3.0 (1.0–60.0); 3.0; 5.0	0.238 [#]
Volume of seminal fluid, in mL-s	4.0 (0.0–40.0); 2.9; 5.1 ^J	3.8 (1.2–9.5); 2.9; 5.1	0.361 [#]
Concentration of spermatozoa, in M/mL	50.0 (0.0–487.0); 21.9; 88.0 ^J	79.0 (6.0–355.0); 46.8; 125.0	<0.001 [#]
Total count of spermatozoa, in M	189.0 (0.0–2432.0); 79.2; 341.3 ^J	302.5 (14.4–1657.6); 164.8; 499.3	<0.001 [#]
Spermatozoa with progressive motility, %	45 (0–83); 34; 56 ^K	53 (11–84); 44; 60	<0.001 [#]
Spermatozoa with normal morphology, %	7 (0–53); 4; 11 ^L	10 (0–27); 7; 14	<0.001 [#]
Concentration of round cells in seminal fluid, in M/mL	1.5 (0.0–75.0); 0.6; 3.0 ^M	1.5 (0.0–45.0); 0.5; 3.0	0.231 [#]
Percentage of neutrophils in seminal fluid, %	9.0 (0.0–92.0); 3.0; 18.0 ^M	4.0 (0.0–69.0); 0.0; 12	<0.001 [#]
Concentration of neutrophils in seminal fluid, in M/mL	0.1 (0.0–68.3); 0.0; 0.4 ^M	0.0 (0.0–27.5); 0.0; 0.3	<0.001 [#]
IL-6 in seminal plasma, in ng/L	18.6 (2.0–4928.0); 11.2; 32.9 ^N	19.3 (2.0–486.0); 12.0; 35.0	0.676 [#]

Annotations to this table. Data with non-parametric distribution. ^{\$} The only variables with parametric distribution were age, bitesticular volume, left testicle's volume, right testicle's volume, and height. For these parameters, the mean, range and 95% confidence interval are presented. ^A Testicular volume was not measured for 146 and 3 patients in the study and the control group, respectively; ^B Data not available for 52 patients in the study group only; ^C Data not available for 51 and 1 patients in the study and the control group, respectively; ^D Data not available for 43 patients in the study group only; ^E Data not available for 72 and 16 patients in the study and the control group, respectively; ^F Data not available for 91 patients in the study group only; ^G Data not available for 94 patients in the study group only; ^H Data not available for 80 patients in the study group only; ^I Data not available for 183 and 4 patients in the study and the control group, respectively; ^J Data not available for 4 patients with serious sexual dysfunctions in the study group only; ^K Data not available for 73 patients in the study group only; ^L Data not available for 74 patients in the study group only; ^M Data not available for 5 patients in the study group only; ^N Data not available for 65 patients in the study group only; [#] Mann–Whitney U test; ^{##} Fisher's exact test; ^{###} Unpaired *t*–test.

Among 2000 infertile men, the proportion of primary and secondary infertility was 74.4% (1487 patients) and 25.6% (513 patients), respectively. The median duration of infertility was 1.9 years with the range 1.0–20.0 years.

The infertility group had lower testicular volume, higher body weight, and a larger proportion of overweight patients (BMI > 24.9) compared to the control group. In addition, infertile men had a lower total count and concentration of spermatozoa, lower percentage of spermatozoa with progressive motility, and normal morphology. The concentration and percentage of neutrophils in semen were also higher among infertile men. However, there was no difference in seminal IL-6 levels between groups. Importantly, there was no difference in abstinence time between infertile and fertile groups.

There was a significantly higher blood level of LH among the infertility group compared to the control group. There was also a slight trend for higher FSH and PSA blood levels among infertile men compared to fertile men, albeit statistically insignificant.

3. Prevalence of STI among Study Subjects

In the first part of our study, we aimed to describe the prevalence of STIs among fertile and infertile patients.

The prevalence of STIs was 2.3% in the infertility group and 1.6% in the control group. Among the infertility group patients, there were 22 men with *M. genitalium* and 24 men with *C. trachomatis* that comprise 1.1% and 1.2% prevalence, respectively (**Table 1**). The prevalence of *C. trachomatis* among control group patients was 1.6% while no cases with *M. genitalium* were found. There were no cases with *N. gonorrhoeae*, *T. vaginalis*, or combined infections in neither group. Interestingly, we found two *C. trachomatis* cases with reported acute orchitis in anamnesis that may suggest reinfection or untreated STI condition.

4. Impact of STI on Inflammation in Semen

For the second step, we aimed to describe the association of STIs with inflammation in male genital tract.

We made the following analyses of both infertile and fertile groups together because the total number of STI-positive patients among fertile patients was considerably low (only four *C. trachomatis* cases). The association of *M. genitalium* and *C. trachomatis* with semen neutrophils' concentration and IL-6 concentration is shown in **Figure 1A,B**, respectively. Both markers were statistically higher for *C. trachomatis* and *M. genitalium*-positive patients compared with STI-negative patients.

We also tried to find the best cut-off value to predict *C. trachomatis* and/or *M. genitalium* infection using semen concentration of neutrophils as a descriptive variable. Using different previously proposed ^{[15][16]} cut-off levels for seminal neutrophils' concentrations, the proportion of STI-positive/negative patients changed but not significantly.

We found the best cut-off value for semen neutrophils' at a concentration of 0.28 mill/mL for predicting *C. trachomatis* and/or *M. genitalium* infection with sensitivity 58.0%, specificity 71.3%, positive prognostic value 1.3%, negative prognostic value 95.6%, and area under the curve 0.691. For seminal IL-6, the optimal cut-off value predicting presence or absence of *C. trachomatis* and/or *M. genitalium* infection was 17.5 ng/L with sensitivity 75.5%, specificity 47.8%, positive prognostic value 1.2%, negative prognostic value 96.8%, and area under the curve 0.644.




Figure 1. (A). Distribution of semen leucocyte concentration by sexually transmitted infection status (both unselected fertile and infertile group, without exclusion of patients for hormonal, testicular volume, and infertility causal factors). **(B).** Distribution of seminal plasma IL-6 concentration by sexually transmitted infection status (both fertile and infertile group, without exclusion of patients for hormonal, testicular volume, and infertility causal factors). Abbreviations: STI—sexually transmitted infections. Data for semen leucocyte concentration are not available for five patients. Data for seminal plasma IL-6 are not available for 65 patients. Mann–Whitney U test with Bonferroni correction for two tests (STI negatives vs. Chlamydia positives, STI negatives vs. Genital mycoplasma positives). For subfigure A: A— p -value < 0.001 , B— p -value = 0.007. For subfigure B: A,B— p -value = 0.01. For subfigure B: A,B—y-axis is natural logarithm scale.

5. Impact of STI on Blood PSA and Semen Quality

In the third part of our study, we aimed to reveal the association of particular STI agents with the quality of semen parameters and blood PSA level. The selection of the data was performed to avoid possible confounding before analysis.

To eliminate possible confounding with known factors influencing semen quality and reproductive hormone levels, we excluded 625 male patients from the main study group and 57 male patients from the main control group additionally. The reasons and order of decision to eliminate study subjects additionally were defined according to the following criteria: (I) defined causal factors of male infertility according to Punab et al. [5]: genetic causes, secondary hypogonadism, congenital anomalies: systemic and/or in urogenital tract, serious sexual dysfunctions, oncological diseases, seminal tract obstruction, other testicular factors that include acquired testicular damage (TD) [exposure to high dose radiation in Chernobyl, testis trauma with volume change, mumps orchitis, other orchitis, epididymitis, testicular torsion, hernia operation with ipsilateral TD, epididymal cyst operation with ipsilateral TD, hydrocele operation with ipsilateral TD, other testis operation with ipsilateral TD] and secondary testicular damage [anabolic steroids, medication (salasopyrin, trexan), status diagnosed post kidney transplantation]; (II) bitesticular volume < 30 mL for unexplained reason; (III) hormonal changes: FSH ≥ 8.0 U/L and/or LH > 9.4 U/L and/or FSH < 1.0 U/L and/or LH < 1.0 U/L and/or total testosterone < 10.5 nmol/L).

After performing the abovementioned exclusion, there remained 1345 STI-negative, 14 *C. trachomatis*-positive and 16 *M. genitalium*-positive patients in the selected infertility group and 188 STI-negative and three *C. trachomatis*-positive patients in the selected control group (**Figure 2**).




Figure 2. Description of the study protocol. Abbreviations: STI—sexually transmitted infections.

In this step of the study, we decided not to combine patients from the selected infertility and control groups to avoid possible bias of the results. As the total number of STI-positive patients in the selected control group was low (only three *C. trachomatis*-positive patients), we performed all the following calculations for the selected infertile group's patients only.

After performing the abovementioned data correction, we did not reveal any impact of *M. genitalium* and *C. trachomatis* on blood levels of PSA.

The semen parameters of the selected infertility group's patients according to STI status (**Figure 1**) are presented in **Table 2**.

Generally, STI-positive patients (both *C. trachomatis* and *M. genitalium* cases) had significantly lower total counts of spermatozoa and total counts of spermatozoa with progressive motility compared to STI-negative patients (**Table 2**). The count of spermatozoa with normal morphology was also lower among STI-positive patients, however, the observed difference was over the significance level.

Table 2. Impact of *C. trachomatis*, *M. genitalium*, and their combination on semen parameters (selected infertile group only, after exclusion of patients for hormonal, testicular volume, and infertility causal factors).

Parameter	STI Negative <i>n</i> = 1345	Either <i>C. trachomatis</i> or <i>M. genitalium</i> Positive <i>n</i> = 30	STI Negative vs. Either <i>M. genitalium</i> or <i>M. genitalium</i> Positive, <i>p</i> -Value ^{\$\$}	<i>C. trachomatis</i> Positive <i>n</i> = 14	STI Negative vs. <i>C. trachomatis</i> Positive, <i>p</i> -Value ^{\$\$, ①}	<i>M. genitalium</i> Positive <i>n</i> = 16	STI Negative vs. <i>M. genitalium</i> Positive, <i>p</i> -Value ^{\$\$, ①}
Abstinence before semen analysis, in days	4.0 (0.0–56.0); 3.0; 4.0	4.0 (1.0–7.0); 3.0; 5.0	0.920	3.5 (2.0–7.0); 2.0; 7.0	0.186	4.0 (1.0–7.0); 2.8; 5.0	0.191
Volume of seminal fluid, in mL	4.1 (0.8–40.0); 3.0; 5.2	4.1 (1.8–9.0); 3.4; 5.3	0.812	3.7 (1.8–9.0); 3.6; 4.2	0.188	4.4 (1.8–6.9); 3.0; 5.4	0.159
Concentration of spermatozoa, M/mL	56.0 (0.0–440.0); 30.0; 94.5	47.0 (1.3–210.0); 9.1; 69.8	0.034	49.0 (1.3–110.0); 17.1; 51.5	0.164	36.5 (1.3–210.0); 7.1; 95.0	0.396

Parameter	STI Negative <i>n</i> = 1345	Either <i>C. trachomatis</i> or <i>M. genitalium</i> Positive <i>n</i> = 30	STI Negative vs. Either <i>M. genitalium</i> or <i>M. genitalium</i> Positive, <i>p</i> -Value ^{\$\$}	<i>C. trachomatis</i> Positive <i>n</i> = 14	STI Negative vs. <i>C. trachomatis</i> Positive, <i>p</i> -Value ^{\$\$, ①}	<i>M. genitalium</i> Positive <i>n</i> = 16	STI Negative vs. <i>M. genitalium</i> Positive, <i>p</i> -Value ^{\$\$, ①}
Total count of spermatozoa, in M	226.2 (0.0–2432.0); 110.4; 372.8	151.0 (2.3–1113.0); 39.6; 303.1	0.028	169.5 (10.1–405.0); 55.9; 201.2	0.132	134.4 (2.3–1113.0); 42.0; 444.8	0.380
Spermatozoa with progressive motility, %	47 (0–83); 35; 57 ^A	41 (0–71); 23; 57	0.187	48 (0–71); 39; 58	0.167	34 (3–64); 19; 57	0.088
Total count of spermatozoa with progressive motility, in M	101.8 (0.0–1056.0); 44.3; 186.7 ^A	68.6 (0.0–489.7); 9.3; 156.6	0.034	75.9 (0.0–281.2); 22.8; 113.9	0.356	50.7 (0.3–489.7); 6.4; 202.5	0.194
Spermatozoa with normal morphology, %	8 (0–53); 4; 12 ^B	5 (0–24); 2; 24	0.242	8 (0–22); 3; 14	0.188	4 (0–24); 2; 10	0.184
Total count of spermatozoa with normal morphology, in M	16.4 (0.0–389.1); 5.3; 37.5 ^B	7.9 (0.0–189.2); 0.4; 35.5	0.105	14.3 (0.0–69.2); 1.7; 34.6	0.952	4.8 (0.0–189.2); 0.3; 33.3	0.236
Concentration of round cells in seminal fluid, in M/mL	1.6 (0.0–75.0); 0.6; 3.0	1.8 (0.2–31.0); 0.8; 3.9	0.485	1.2 (0.2–7.2); 0.5; 2.0	0.746	3.0 (0.4–31.0); 1.0; 7.3	0.146
Percentage of neutrophils in seminal fluid, %	8 (0–91); 3; 16	22 (0–63); 11; 39	7.379×10^{-6}	20 (4–48); 13; 30	0.004	25 (0–63); 10; 43	0.002
Concentration of neutrophils in seminal fluid, in M/mL	0.1 (0.0–68.3); 0.0; 0.3	0.2 (0.0–10.70); 0.1; 0.9	0.001	0.2 (0.0–1.4); 0.1; 0.7	0.250	0.4 (0.0–10.7); 0.1; 2.6	0.004
IL-6 in seminal plasma, in ng/L	18.2 (2.0–4928.0); 11.1; 32.3 ^C	30.4 (3.0–1330.0); 17.9; 78.7 ^D	0.002	24.8 (3.0–1330.0); 20.8; 66.2 ^D	0.192	30.6 (9.2–406.0); 17.6; 84.5	0.016

Annotations to this table. Data are presented as median (range), 25th, and 75th percentile. Data are with non-parametric distribution. ^{\$\$} *p*-value is calculated with Mann–Whitney test comparing two groups. ^① Bonferroni correction of *p*-value for two tests. ^A Data accounting progressive motility of spermatozoa are not available for one patient. ^B Data accounting spermatozoa morphology are not available for one patient. ^C Data accounting IL-6 in seminal plasma are not available for 36 patients. ^D Data accounting IL-6 in seminal plasma are not available for one patient.

However, when STI-positive cases were analyzed separately, the impact of *C. trachomatis* on the total count of spermatozoa and progressive motility became slightly over the significance level. Similar tendencies were also seen in the case of *M. genitalium*-positive patients. The spermatozoa count with normal morphology was lower in *M. genitalium*-positive patients though the difference between the groups was slightly over the significance level again.

The impact of both combined *C. trachomatis* and *M. genitalium* on seminal inflammatory parameters (concentration of neutrophils, seminal IL-6) also remained significant in the selected infertility group patients. However, when STI-positive cases were split separately, the impact of *C. trachomatis* on seminal IL-6 and neutrophil concentration disappeared, while remaining for *M. genitalium* (Table 2).

References

1. Zegers-Hochschild, F.; Adamson, G.; de Mouzon, J.; Ishihara, O.; Mansour, R.; Nygren, K.; Sullivan, E.; Vanderpoel, S.; International Committee for Monitoring Assisted Reproductive Technology; World Health Organization. International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary of ART terminology, 2009. *Fertil. Steril.* 2009, 92, 1520–1524.
2. Boivin, J.; Bunting, L.; Collins, J.A.; Nygren, K.G. International estimates of infertility prevalence and treatment-seeking: Potential need and demand for infertility medical care. *Hum. Reprod.* 2007, 22, 1506–1512.
3. Jungwirth, A.; Giwercman, A.; Tournaye, H.; Diemer, T.; Kopa, Z.; Dohle, G.; Krausz, C.G. European Association of Urology Guidelines on Male Infertility: The 2012 Update. *Eur. Urol.* 2012, 62, 324–332.
4. Weidner, W.; Pilatz, A.; Diemer, T.; Schuppe, H.C.; Rusz, A.; Wagenlehner, F. Male urogenital infections: Impact of infection and inflammation on ejaculate parameters. *World J. Urol.* 2013, 31, 717–723.
5. Punab, M.; Poolamets, O.; Paju, P.; Vihljajev, V.; Pomm, K.; Ladva, R.; Korrovits, P.; Laan, M. Causes of male infertility: A 9-year prospective monocentre study on 1737 patients with reduced total sperm counts. *Hum. Reprod.* 2016, 32, 18–31.
6. Dutta, S.; Sengupta, P.; Slama, P.; Roychoudhury, S. Oxidative Stress, Testicular Inflammatory Pathways, and Male Reproduction. *Int. J. Mol. Sci.* 2021, 22, 10043.
7. Ochsendorf, F.R. Sexually transmitted infections: Impact on male fertility. *Andrologia* 2008, 40, 72–75.
8. Schuppe, H.-C.; Pilatz, A.; Hossain, H.; Diemer, T.; Wagenlehner, F.; Weidner, W. Urogenital Infection as a Risk Factor for Male Infertility. *Dtsch. Aerzteblatt Online* 2017, 114, 339–346.
9. Wagenlehner, F.M.; Naber, K.G.; Weidner, W. Chlamydial infections and prostatitis in men. *BJU Int.* 2006, 97, 687–690.
10. Ahmadi, M.H.; Mirsalehian, A.; Bahador, A. Association of Chlamydia trachomatis with infertility and clinical manifestations: A systematic review and meta-analysis of case-control studies. *Infect. Dis.* 2016, 48, 517–523.
11. Mielczarek, E.; Blaszkowska, J. Trichomonas vaginalis: Pathogenicity and potential role in human reproductive failure. *Infection* 2015, 44, 447–458.
12. Fode, M.; Fusco, F.; Lipshultz, L.; Weidner, W. Sexually Transmitted Disease and Male Infertility: A Systematic Review. *Eur. Urol. Focus* 2016, 2, 383–393.
13. Taylor-Robinson, D.; Jensen, J.S. Mycoplasma genitalium: From Chrysalis to Multicolored Butterfly. *Clin. Microbiol. Rev.* 2011, 24, 498–514.
14. Huang, C.; Zhu, H.; Xu, K.; Wang, S.; Fan, L.; Zhu, W.B. Mycoplasma and ureaplasma infection and male infertility: A systematic review and meta-analysis. *Andrology* 2015, 3, 809–816.
15. Punab, M.; Lõivukene, K.; Kermes, K.; Mändar, R. The limit of leucocytospermia from the microbiological view-point. *Andrologia* 2003, 35, 271–278.
16. Gdoura, R.; Kchaou, W.; Znazen, A.; Chakroun, N.; Fourati, M.; Keskes, L.; Hammami, A. Screening for bacterial pathogens in semen samples from infertile men with and without leukocytospermia. *Andrology* 2008, 40, 209–218.

Retrieved from <https://encyclopedia.pub/entry/history/show/41433>