

Bruton's Tyrosine Kinase Signaling in Tumor Microenvironment

Subjects: [Oncology](#)

Contributor: Justin K. Messex , Geou-Yarh Liou

Protein tyrosine kinase Bruton's tyrosine kinase (BTK) is essential for B cell maturation and proliferation.

Dysregulation of BTK signaling in B cells leads to B cell lymphoma. In addition to B cells, BTK is also expressed in other types of immune cells including myeloid derived suppressor cells (MDSC), dendritic cells, mast cells and macrophages, all of which comprise the tumor microenvironment in solid cancers.

Bruton's tyrosine kinase

cell signaling

tumor microenvironment

cancer therapy

solid tumor

1. Introduction

As tumor cells begin to thrive, the microenvironment which they reside in becomes increasingly important to their survival and spread [\[1\]\[2\]\[3\]](#). The microenvironment composed of immune cells, vessels for transportation, structural components, e.g., the extracellular matrix, signaling proteins, and nerve cells provide necessary resources for tumor proliferation. Although the components of the microenvironment may differ slightly depending on the location of the tumor, each microenvironment plays a role in tumor development by utilizing various signaling pathways to control gene expression.

Under times of stress, the microenvironment undergoes various degrees of alterations in an attempt to maintain conditions that are conducive for tumor growth. Modulation of the tumor microenvironment is orchestrated by a vast network of cell-to-cell signaling. For example, activation of the phosphatase and tensin (PTEN) pathway in CD8⁺ T has been highly correlated with a significant decrease in T cell differentiation and enhancing immune suppression in breast cancer patients [\[4\]](#).

B lymphocytes, also known as B cells, originate in the bone marrow. Here, they depend on the signal from a protein tyrosine kinase known as Bruton's tyrosine kinase (BTK) for their maturation [\[5\]](#). Once mature, they become a part of the adaptive immune response and are responsible for the secretion of antibodies. However, aberrant BTK signaling has been recognized in diseases, like X-linked agammaglobulinemia (XLA), as well as various B cell lymphomas [\[6\]](#).

2. Cells of the Tumor Microenvironment

At its core, the tumor microenvironment is supported by the extracellular matrix (ECM). This collage of extracellular molecules, glycoproteins, and collagen provides rigidity to support the tumor cells, as well as remodeling itself to allow expansion of adipose tissue, which is known for its role in colorectal cancer [7][8]. Several factors, including TNF, IL-6, leptin, adiponectin, and visfatin secreted by unregulated adipose tissues and adipocytes, have been shown to enhance cell proliferation, survival, migration, anti-apoptosis and angiogenesis of colorectal cancer [9][10]. Furthermore, the ECM allows multipotent cells like mesenchymal stromal cells (MSCs) to receive signals which allow differentiation into pro-tumor components of the tumor microenvironment [11]. The presence of cytokines, like tumor growth factor β 1 (TGF β 1), has also been shown to induce MSC differentiation into carcinoma-associated fibroblasts (CAF), producing a significant increase in the migratory and invasive properties of prostate cancer cells [11].

Mast cells, another type of cell found in abundance within the tumor microenvironment, contributes to tumor development by causing inflammation at the tumor site, allowing nutrients and blood flow to reach the tumor. Studies have shown that protein kinase D2/3 (PKD 2/3) of the prostate cancer cells increased infiltrating mast cells through turning on transcription factors, including AP-1 and/or nuclear factor kappa light chain enhancer of activated B cell (NF- κ B), to upregulate scf, CCL5, and CCL11 transcripts [12]. Furthermore, blockade of mast cell recruitment by prostate cancer cells through PKD inhibitors diminished tumor volume and growth in an allografted mouse model. Other cells of the tumor microenvironment include cells of the immune system; however, within the tumor microenvironment, these cells are often exploited by tumorous cells and prevented from doing their job.

In lung cancer, for example, CD3⁺ T cells represent the majority of infiltrating immune cells contributing to tumor progression. Furthermore, CD4⁺ helper T cells and CD8⁺ cytotoxic T cells present in lung cancer contribute to the production of immune exhaustion proteins, including cytotoxic T-lymphocyte antigen 4 (CTLA-4), lymphocyte activated gene 3 (LAG3), programmed cell death protein 1 (PD-1), etc., leading to tolerance of tumor antigens and loss of anti-cancer effects in these cells [13]. Increased levels of IL-17, which contribute to an anti-cancer immune response, as well as CTLA-4 positive regulatory T cells, have also been highly associated with the bronchoalveolar lavage of the cancerous lung, suggesting that the anti-immune response was enhanced in the local cancerous microenvironment advocating for lung cancer progression and metastasis [14]. Other studies using Kras^{G12D};p53^{fl/fl} transgenic mice to study the microenvironment of lung cancer, showed a collection of immune cells infiltrating the cancerous area, including B cells, CD4⁺ and CD8⁺ T lymphocytes, macrophages, CD11c⁺ dendritic cells, and Gr1⁺ neutrophils. Depletion of Gr1⁺ neutrophils using the anti-Gr1 neutralizing antibody resulted in an increase of macrophages, as well as CD4⁺ and CD8⁺ T lymphocytes, with a simultaneous decrease in regulatory T cells. In addition, Gr1⁺ depletion also diminished lung cancer growth and re-sensitized lung cancer cells to anti-PD1 treatment allowing activation and proper function of T cells [15]. Other immune cells of the innate response, such as macrophages, are critical for a healthy immune system. However, they are also known to contribute to the tumor microenvironment in a variety of ways and are widely recognized for the vital role they play in tumor progression [16].

Macrophages in their immature non-activated form, indicated as M0, have two major destinations. Upon stimulation by lipopolysaccharide (LPS) or interferon gamma (IFN γ), M0 macrophages polarize into M1 pro-inflammatory

macrophages. On the other hand, M0 macrophages, when exposed to IL-4 or IL-13 cytokines, differentiate into M2 immunosuppressive macrophages and have been known for their role in tumor progression (**Figure 1A**). Tumor associated macrophages (TAMs) possess a M2-like phenotype and are abundant within the tumor microenvironment of various cancers, like colitis-associated carcinoma, contributing to tumor progression [17][18][19].




Figure 1. Macrophage polarization and M2 macrophage-associated signaling pathways that contribute to cancer dissemination. **(A)** Non-activated macrophages (M0 macrophages) are polarized to pro-inflammatory/M1 macrophages which specifically express several molecules, including iNOS, CD80, IFN γ , and TNF, in the presence of LPS and IFN γ . In the presence of IL-4 or IL-13, M0 macrophages are polarized to anti-inflammatory/M2 macrophages which specifically express arginase 1, CCL17, CCL22, IL-4, IL-10, CD163, and CD206, and contributing to tissue repair process and tumor development. **(B)** M2 macrophages potentiate cancer metastasis through activation of ERK and JNK signaling that upregulates cell migration and invasion. Notch signaling inhibits M2 macrophage polarization. Wnt signaling can polarize macrophages to a M2 subtype by activation of c-Myc. Abbreviation: ERK: extracellular signal-regulated kinase; JNK: c-Jun N-terminal kinase. Abbreviation: EMR1: EGF-like module-containing mucin-like hormone receptor-like 1; iNOS: inducible nitric oxide synthase; IFN γ : interferon gamma; TNF: tumor necrosis factor; LPS: lipopolysaccharide; IL-4: interleukin-4; IL-10: interleukin-10.

Mice treated with Azoxymethane (AOM)/Dextran Sodium Sulfate (DSS), used to induce inflammatory colorectal cancer, well recapitulate human colitis-associated cancer of the colon. Blockage of the EGFR signaling pathway present in cancerous cells has been shown to assist in M2 macrophage polarization, reduced tumor growth and showed a reduction in the amount of infiltrating CD206⁺ M2 macrophages in AOM/DSS-induced murine colon cancer mice [20]. This study also showed that the conditioned media of colon cancer cells promoted macrophage polarization to a M2 subtype through an increase in arginase-1, CCL17, CCL22, IL-4, and IL-10. Furthermore, knockout of the EGFR pathway in the colon cancer cells reversed macrophage polarization back to a M1 subtype, suggesting that EGFR signaling of colon cancer cells modulates macrophage polarization and repolarization to promote colon cancer development and progression [20]. In other types of cancers, such as hepatocellular

carcinoma (HCC), TAMs express either CD86⁺ (M1 subtype) or CD206⁺ (M2 subtype). However, using this as a diagnostic tool for HCC is uncertain as one study evaluated 253 patients with HCC and discovered the presence of macrophages was not associated with diagnosis nor prognosis of HCC in clinic. They did, however, determine that increased numbers of CD206⁺ TAMs were linked to tumor aggressiveness, defined by the total tumor numbers, tumor-node-metastasis, poor overall survival, and recurrence, contradicting the results found in breast cancer studies [21].

3. BTK Signaling

BTK is a member of the Tec family kinases, a family of kinases highly associated with regulation of immune functions. However, unlike Tec kinases, which control T helper cell function, BTK is primarily found in B cells and essential for B cell maturation, as well as being known for various other functions when expressed in other types of cells, such as macrophage polarization [22]. Located on the cytoplasmic side of the B cell, the BTK structure contains five interaction domains, including a pleckstrin homology (PH) domain and a kinase domain with enzymatic activity (**Figure 2**). Once antigen binds to the B cell receptor, BCR, the PH domain of BTK interacts with phosphatidylinositol triphosphate (PIP3), a downstream signaling activator generated from phosphoinositide 3-kinase (PI3K). Once bound to PIP3, BTK is phosphorylated by the spleen tyrosine kinase (SYK) within the kinase domain, also known as the Src homology type 1 (SH1) domain, at Y551 [23]. BTK then autophosphorylates itself in the SH3 domain at Y223, allowing full activation of BTK [24][25]. In return, two primary secondary messengers, inositol triphosphate (IP3) and diacylglycerol (DAG), are activated, turning on various processes which regulate cellular homeostasis.




Figure 2. BTK domain diagram. BTK contains five domains, including a PH domain at the amino terminus, a TH domain, a SH3 domain, a SH2 domain and a tyrosine kinase domain at the carboxyl terminus. The PH domain of BTK is able to interact with phosphatidylinositol lipids which allows it to eventually regulate transcription factors through IP3 and DAG. The SH3 domain of BTK mediates protein-protein interactions through proline-rich repeats and also contains a tyrosine at 223 position (Y223), an autophosphorylation site of BTK. The SH2 domain of BTK can interact with phospho-tyrosine. The kinase domain of BTK, also known as the SH1 domain contains a tyrosine at 551 position (Y551) with the activation loop. Once Y551 is phosphorylated by SYK or Src, it results in autophosphorylation of Y223 to fully activate BTK. In addition, cysteine 481 (C481) is the site to which BTK inhibitors covalently bound. Abbreviation: PH: pleckstrin homology; TH: Tec homology; SH: Src homology; IP3: inositol triphosphate; DAG: diacylglycerol.

BTK is considered a non-receptor tyrosine kinase and does not receive direct stimulation from antigen. This is why BTK is able to be activated by multiple different antigens. The short cytoplasmic domain of the BCR associates with the disulfide-linked CD79a/CD79b heterodimers upon stimulation from antigen binding [23]. These heterodimers contain immunoreceptor tyrosine based activation motifs (ITAMS) which are phosphorylated upon antigen binding to create docking sites for SYK, a BTK upstream activator as previously mentioned, ultimately leading to the activation of the BTK signaling cascade. In the absence of BCR stimulation and, consequently, the absence of BTK signaling, B cells undergo a high rate of apoptosis due to a reduction in the expression of the anti-apoptotic protein Bcl-xL [26]. However, stimulation of BTK through the BCR antigen binding ultimately leads to the activation of the Akt pathway, subsequently turning on pro-survival transcription factors, including forkhead box O (FOXO) and NF- κ B [27].

Activation of BTK is also crucial for the function of CXCR4 and CXCR5, G-protein coupled receptors found in B cells which play important roles in B cell entry to lymph nodes [28]. Binding of chemokine to CXCR4 or CXCR5 receptor induces BTK activation, ultimately leading to factors which contribute to B cell homing to the surrounding lymphoid organs. Although these pathways are poorly defined, BTK deficient mice showed significant impairment in adhesion and migration response, suggesting that BTK signaling is critical for proper B cell migration and integrin adhesion [29].

Other receptors, like Toll-like receptors (TLRs), are not only found in B cells but also in myeloid cells, including macrophages, mast cells, etc., and can be utilized for activating the BTK signaling pathway. These receptors recognize conserved regions of bacterial and viral origins, such as the lipopolysaccharide (LPS). Upon TLR binding with antigen, an adaptor protein, like myeloid differentiation primary response (MYD88), interacts with BTK, ultimately activating transcription factors necessary for antibody secretion and immunoglobulin class switching, a critical component to maintaining a healthy immune system [30][31][32][33].

Although BTK is most well-known for its function in B cell maturation and processes necessary for cellular homeostasis, its contribution to tumor progression by activating cells known for their pro-tumor properties is also well documented (**Figure 3**). For example, BTK-activated myeloid derived suppressor cells (MDSC) contain immunosuppressive properties which are critical for evading the immune response. The production of arginase-1,

indoleamine 2, 3-dioxygenase (IDO), nitric oxide (NO), reactive oxygen species (ROS), and suppressive cytokines, such as IL-10 and TGF β , from MDSCs help tumorous cells avoid death from cells of the adaptive immune response. Studies have shown that IFN γ produced by immune cells to combat harmful material is significantly reduced in the presence of MDSC-generated NO, indirectly promoting tumor growth and progression. Furthermore, activation of BTK was detected in MDSCs isolated from tumor-bearing mice, as well as human samples [34].




Figure 3. Activated BTK signaling of the stromal cells promotes cancer cell growth and progression. Activated BTK in myeloid-derived suppressor cells increases several molecules, including arginase-1, ROS, IDO, and suppressive cytokines, which directly and indirectly potentiate growth and development of cancer cells. Activated BTK in macrophages induces Fc γ R expression. Fc γ R⁺ macrophages can directly promote cancer cell proliferation and metastasis or through inhibition of cytotoxic CD8⁺ T cells, which eliminate cancer cells. Activated BTK in the regulatory B cells is able to elevate cancer cell growth and progression by their secreted suppressive cytokines, including IL-10 and IL-35, that suppress cytotoxic CD8⁺ T cells. Abbreviation: MDSC: myeloid-derived suppressor cell; Breg: regulatory B cells; ROS: reactive oxygen species; IDO: Indoleamine 2, 3-dioxygenase; TGF β : transforming growth factor β ; Fc γ R: Fc γ receptor.

Activation of BTK has also been detected in the interaction between Fc γ R-positive macrophages, a type of macrophage with immunosuppressive properties also referred to as M2 macrophages, and B cells in order to

promote pancreatic ductal adenocarcinoma (PDAC) growth and progression. Furthermore, activation of BTK in surrounding macrophages was mediated by PI3Ky protein kinase, resulting in their polarization to FcyR-positive macrophages. Blockade of the signaling pathway using the BTK inhibitor PCI-32765, known as ibrutinib, or PI3Ky inhibitor TG100-115 decreased PDAC growth both in vitro and in vivo (**Table 1**). Additionally, combining these two inhibitors with Gemcitabine in the treatment of PDAC tumors halted macrophage infiltration and increased cytotoxic CD8⁺ T cells numbers inside the tumors [35]. Other studies assessing BTK signaling and PDAC development showed that BTK expression in CD1d^{hi}CD5⁺ regulatory B cells resulted in the production of IL-10 and IL-35, both of which mediate immunosuppressive functions in regulatory B cells. Inhibition of BTK through its specific inhibitor tirabrutinib diminished both regulatory B cell differentiation and their secreted immunosuppressive cytokines. Furthermore, administration of tirabrutinib to the Kras^{G12D} orthotopic mouse model reduced tumor cell growth and infiltrating regulatory B cells with increasing numbers of CD8⁺IFN γ ⁺ T cells, thus also indicating BTK's role in immune function [36].

Table 1. Summary of BTK inhibitors that have been evaluated in cancer studies.

BTK Inhibitor	Study Type	Cancer Type
ibrutinib	in vitro	B cell lymphoma
		chronic lymphocytic leukemia
		pancreatic cancer
	in vivo	chronic lymphocytic leukemia
		pancreatic cancer
		colorectal cancer
in vivo	breast cancer	
acalabrutinib	in vitro	chronic lymphocytic leukemia

BTK Inhibitor	Study Type	Cancer Type	
	in vivo	B cell lymphoma	
		chronic lymphocytic leukemia	
		pancreatic cancer *	
		urothelial cancer *	
tirabrutinib	in vivo	pancreatic cancer	inhibitors possibly other
zanubrutinib	in vivo	mantle cell lymphoma *	

Dendritic cells are responsible for bridging the innate and adaptive immune responses, including eradication of cancer, through T cell mediated immunity. Studies using BTK knockout mice demonstrated BTK's role in regulating dendritic cell development [37]. In addition, it has also been shown that treatment using the BTK inhibitor ibrutinib in murine bone marrow-derived dendritic cells upregulated CD11c in these cells along with MHC-II and CD80 expressions [38]. Stimulation of these ibrutinib-treated dendritic cells with LPS resulted in decreased IL-6, IL-12, and NO production. In addition, these dendritic cells also promoted CD4⁺ T cell proliferation [38]. Altogether, the study suggested that inhibition of BTK in dendritic cells enhanced maturation and activation of dendritic cells and accelerated CD4⁺ T cell growth, which contributes to reshaping the tumor microenvironment during cancer development.

4. BTK Inhibitors and Their Clinic Application for Cancer Therapy

BTK inhibitors, such as ibrutinib, acalabrutinib, and zanubrutinib, are currently being used in clinical trials for hematologic malignancies, including B cell lymphoma, mantle cell lymphoma, and chronic lymphocytic leukemia/small lymphocytic lymphoma, with promising results. Furthermore, the second-generation BTK inhibitor acalabrutinib has also been approved by the FDA for treatment of mantle cell lymphoma, and chronic lymphocytic leukemia/small lymphocytic lymphoma. These inhibitors outcompete ATP by covalently binding to the cysteine 481 residue within the kinase domain to selectively inhibit BTK kinase activity [39][40].

Ibrutinib, also known as PCI-32765, was the first BTK inhibitor approved and tested in clinical trials for patients with B cell lymphomas [41][42][43]. In addition to BTK, ibrutinib also shows significant inhibitory activity against 10 other protein kinases, including BLK, BMX, ITK, TEC, EGFR, ERBB2, and JAK 3, with an IC₅₀ value less than 100 nM [44]. Furthermore, low dosage of ibrutinib, IC₅₀ around 10 nM, showed sufficient blockage of BTK

autophosphorylation, as well as its downstream signaling targets PLC γ and ERK in cultured B cell lymphoma DOHH2 cells [44]. Several other studies demonstrated that, in human primary chronic lymphocytic leukemia (CLL) cells, ibrutinib is able to not only cause apoptosis through activation of caspase 3 but also block CLL cell proliferation via suppression of BCR- and CD40-activated Akt, ERK, and NF- κ B pathways, all of which are downstream targets of BTK [45]. Ibrutinib has also been reported to abolish integrin-mediated adhesion and B cell migration in response to CXCL12 and CXCL13 in CLL cells [45][46][47]. Although ibrutinib has been effective in many cases, CLL patients have encountered drug tolerance with ibrutinib, as well as its off-target effects, resulting in adverse effects like platelet function defect, atrial fibrillation, etc. [48].

Acalabrutinib, also known as ACP-196, is a second-generation BTK inhibitor with a higher potency and stronger selectivity than ibrutinib [49][50]. Unlike ibrutinib, acalabrutinib does not inhibit EGFR, ITK or TEC pathways, kinases which closely mimic BTK, and has a lower IC₅₀ as compared to ibrutinib [51]. In addition, acalabrutinib shows improved pharmacological features, such as rapid oral absorption and a short half-life, with less adverse effects [51]. In vitro studies have shown that acalabrutinib's effectiveness against CLL cells is due to inhibiting activation of ERK, NF- κ B, and Akt [51]. Prior to clinical trials, acalabrutinib has been evaluated in several animal models of B cell non-Hodgkin lymphoma, including canine model, human NSG primary CLL xenografted mouse model, and TCL1 adoptive transfer mouse model, a mouse model which spontaneously develops CLL-like leukemia and has a similar drug response to human CLL cells [52][53][54][55][56]. Acalabrutinib treated TCL1 transgenic mice showed significant suppression of CLL cell growth and subsequently decreased tumor burden by blocking BCR signaling-mediated PLC γ 2 and ERK, proving its in vivo efficacy against CLL [54][55][56]. Furthermore, clinical trials using acalabrutinib for CLL patients indicated acalabrutinib shows less adverse effects in CLL patients than ibrutinib due to its high specificity for BTK and develops resistance far less [54].

As for solid tumors, it has been reported that ibrutinib decreased CD20 mRNA through inhibition of NF- κ B transcription factor activity. In addition, ibrutinib also inhibits the expression of decay-accelerating factor (CD55), leading to the complement protein C3d-mediated opsonization of tumor cells [57]. Furthermore, treatment with ibrutinib in breast cancer xenografted mice decreased tumor growth and metastasis [58]. In addition, these treated mice had more mature dendritic cells and less immature MDSCs as compared to the vehicle-treated mice. Using the primary MDSC isolated from mouse, ibrutinib treatment converted primary murine MDSC to dendritic cells as indicated by the various markers of MDSC and dendritic cells. Treatment using ibrutinib also reduced NO generation and cell migration of MDSC. Furthermore, ibrutinib treatment in tumor-bearing mice significantly decreased splenic MDSC and re-sensitizes tumor to anti-PD-L1 immune checkpoint blockade [34]. Although mouse models and in vitro studies have shown high correlation with BTK signaling/inhibition and its effect on tumor progression, clinical studies have been contradictory. In some cases, focusing on BTK inhibition has proven to be a viable therapy for cancer patients while others have shown BTK inhibitors have dismal effects on patient outcome [59].

A phase 2 trial with 77 pancreatic patients, the dual treatment of the BTK inhibitor acalabrutinib and pembrolizumab (PD-L1 inhibitor) was well tolerated and showed a reduction of MDSC cells in the peripheral blood [60]. However, not all trials have shared the same optimistic view for BTK inhibitors.

Phase 3 clinical trials testing 424 metastatic PDAC patients using ibrutinib plus nab-paclitaxel/gemcitabine as a treatment showed no significant improvement in overall survival nor progression free survival as compared to the current first line treatment, nab-paclitaxel/gemcitabine [61]. Another phase 2 trial with 75 patients for platinum-resistant metastatic urothelial cancer, dual treatment of acalabrutinib and pembrolizumab did not improve the overall survival nor the progressive-free survival of the disease for the platinum-resistant metastatic urothelial cancer as compared to the pembrolizumab treatment alone [62]. The numbers of the peripheral monocytic MDSCs remained same in response to the dual treatment of acalabrutinib and pembrolizumab in comparison to the perbrolizumab alone. Meanwhile, proliferating CD8⁺ T cells increased under the dual treatment.

Although BTK inhibitors like ibrutinib and acalabrutinib are viable treatment options for lymphoma patients, more pre-clinical studies focusing on BTK inhibition within the TME is necessary to completely understand how it contributes to solid tumor development. This will reveal if these FDA approved BTK inhibitors can serve as an adjuvant treatment for solid tumors.

References

1. Catalano, V.; Turdo, A.; di Franco, S.; Dieli, F.; Todaro, M.; Stassi, G. Tumor and Its Microenvironment: A Synergistic Interplay. In *Seminars in Cancer Biology*; Elsevier: London, UK, 2013; Volume 23, pp. 522–532.
2. Ozdemir, B.C.; Pentcheva-Hoang, T.; Carstens, J.L.; Zheng, X.; Wu, C.C.; Simpson, T.R.; Laklai, H.; Sugimoto, H.; Kahlert, C.; Novitskiy, S.V.; et al. Depletion of Carcinoma-Associated Fibroblasts and Fibrosis Induces Immunosuppression and Accelerates Pancreas Cancer with Reduced Survival. *Cancer Cell* 2014, 25, 719–734.
3. Whiteside, T.L. The Tumor Microenvironment and Its Role in Promoting Tumor Growth. *Oncogene* 2008, 27, 5904–5912.
4. Mittendorf, E.A.; Philips, A.V.; Meric-Bernstam, F.; Qiao, N.; Wu, Y.; Harrington, S.; Su, X.; Wang, Y.; Gonzalez-Angulo, A.M.; Akcakanat, A.; et al. PD-L1 Expression in Triple-Negative Breast Cancer. *Cancer Immunol. Res.* 2014, 2, 361–370.
5. Vetrie, D.; Vorechovsky, I.; Sideras, P.; Holland, J.; Davies, A.; Flinter, F.; Hammarstrom, L.; Kinnon, C.; Levinsky, R.; Bobrow, M.; et al. The Gene Involved in X-linked Agammaglobulinaemia is a Member of the SRC Family of Protein-Tyrosine Kinases. *Nature* 1993, 361, 226–233.
6. Tsukada, S.; Saffran, D.C.; Rawlings, D.J.; Parolini, O.; Allen, R.C.; Klisak, I.; Sparkes, R.S.; Kubagawa, H.; Mohandas, T.; Quan, S.; et al. Deficient Expression of a B Cell Cytoplasmic Tyrosine Kinase in Human X-linked Agammaglobulinemia. *Cell* 1993, 72, 279–290.
7. Bauer, J.; Emon, M.A.B.; Staudacher, J.J.; Thomas, A.L.; Zessner-Spitzenberg, J.; Mancinelli, G.; Krett, N.; Saif, M.T.; Jung, B. Increased Stiffness of the Tumor Microenvironment in Colon Cancer

- Stimulates Cancer Associated Fibroblast-Mediated Prometastatic Activin a Signaling. *Sci. Rep.* 2020, 10, 50.
8. Liu, C.; Pei, H.; Tan, F. Matrix Stiffness and Colorectal Cancer. *OncoTargets Ther.* 2020, 13, 2747–2755.
 9. Day, S.D.; Enos, R.T.; McClellan, J.L.; Steiner, J.L.; Velazquez, K.T.; Murphy, E.A. Linking Inflammation to Tumorigenesis in a Mouse Model of High-Fat-Diet-Enhanced Colon Cancer. *Cytokine* 2013, 64, 454–462.
 10. Kern, L.; Mittenbuhler, M.J.; Vesting, A.J.; Ostermann, A.L.; Wunderlich, C.M.; Wunderlich, F.T. Obesity-Induced TNFalpha and IL-6 Signaling: The Missing Link between Obesity and Inflammation-Driven Liver and Colorectal Cancers. *Cancers* 2018, 11, 24.
 11. Barcellos-de-Souza, P.; Comito, G.; Pons-Segura, C.; Taddei, M.L.; Gori, V.; Becherucci, V.; Bambi, F.; Margheri, F.; Laurenzana, A.; Del Rosso, M.; et al. Mesenchymal Stem Cells are Recruited and Activated into Carcinoma-Associated Fibroblasts by Prostate Cancer Microenvironment-Derived TGF-beta1. *Stem Cells* 2016, 34, 2536–2547.
 12. Xu, W.; Qian, J.; Zeng, F.; Li, S.; Guo, W.; Chen, L.; Li, G.; Zhang, Z.; Wang, Q.J.; Deng, F.; et al. Protein Kinase Ds Promote Tumor Angiogenesis through Mast Cell Recruitment and Expression of Angiogenic Factors in Prostate Cancer Microenvironment. *J. Exp. Clin. Cancer Res.* 2019, 38, 114.
 13. Zhou, H.; Liu, T.; Wang, Z. Analysis of Non-Small Cell Lung Cancer Microenvironment Indicates Preponderance of T Cell Exhaustion Marker Expression. *Exp. Cell Res.* 2017, 360, 205–209.
 14. Kwiecien, I.; Stelmaszczyk-Emmel, A.; Polubiec-Kownacka, M.; Dziedzic, D.; Domagala-Kulawik, J. Elevated Regulatory T Cells, Surface and Intracellular CTLA-4 Expression and Interleukin-17 in the Lung Cancer Microenvironment in Humans. *Cancer Immunol. Immunother.* 2017, 66, 161–170.
 15. Faget, J.; Groeneveld, S.; Boivin, G.; Sankar, M.; Zangger, N.; Garcia, M.; Guex, N.; Zlobec, I.; Steiner, L.; Piersigilli, A.; et al. Neutrophils and Snail Orchestrate the Establishment of a Pro-tumor Microenvironment in Lung Cancer. *Cell Rep.* 2017, 21, 3190–3204.
 16. Chen, P.C.; Cheng, H.C.; Wang, J.; Wang, S.W.; Tai, H.C.; Lin, C.W.; Tang, C.H. Prostate Cancer-Derived CCN3 Induces M2 Macrophage Infiltration and Contributes to Angiogenesis in Prostate Cancer Microenvironment. *Oncotarget* 2014, 5, 1595–1608.
 17. Mantovani, A.; Germano, G.; Marchesi, F.; Locatelli, M.; Biswas, S.K. Cancer-Promoting Tumor-Associated Macrophages: New Vistas and Open Questions. *Eur. J. Immunol.* 2011, 41, 2522–2525.
 18. Mantovani, A.; Schioppa, T.; Porta, C.; Allavena, P.; Sica, A. Role of Tumor-Associated Macrophages in Tumor Progression and Invasion. *Cancer Metastasis Rev.* 2006, 25, 315–322.

19. Puig-Kroger, A.; Sierra-Filardi, E.; Dominguez-Soto, A.; Samaniego, R.; Corcuera, M.T.; Gomez-Aguado, F.; Ratnam, M.; Sanchez-Mateos, P.; Corbi, A.L. Folate Receptor Beta is Expressed by Tumor-Associated Macrophages and Constitutes a Marker for M2 Anti-Inflammatory/Regulatory Macrophages. *Cancer Res.* 2009, 69, 9395–9403.
20. De Robertis, M.; Massi, E.; Poeta, M.L.; Carotti, S.; Morini, S.; Cecchetelli, L.; Signori, E.; Fazio, V.M. The AOM/DSS murine model for the study of colon carcinogenesis: From pathways to diagnosis and therapy studies. *J. Carcinog.* 2011, 10, 9.
21. Dong, P.; Ma, L.; Liu, L.; Zhao, G.; Zhang, S.; Dong, L.; Xue, R.; Chen, S. CD86(+)/CD206(+), Diametrically Polarized Tumor-Associated Macrophages, Predict Hepatocellular Carcinoma Patient Prognosis. *Int. J. Mol. Sci.* 2016, 17, 320.
22. Gabhann, J.; Hams, E.; Smith, S.; Wynne, C.; Byrne, J.C.; Brennan, K.; Spence, S.; Kissenpfennig, A.; Johnston, J.A.; Fallon, P.G.; et al. Btk Regulates Macrophage Polarization in Response to Lipopolysaccharide. *PLoS ONE* 2014, 9, e85834.
23. Singh, S.P.; Dammeijer, F.; Hendriks, R.W. Role of Bruton's Tyrosine Kinase in B Cells and Malignancies. *Mol. Cancer* 2018, 17, 57.
24. Park, H.; Wahl, M.I.; Afar, D.E.; Turck, C.W.; Rawlings, D.J.; Tam, C.; Scharenberg, A.M.; Kinet, J.P.; Witte, O.N. Regulation of Btk Function by a Major Autophosphorylation Site within the SH3 Domain. *Immunity* 1996, 4, 515–525.
25. Rawlings, D.J.; Scharenberg, A.M.; Park, H.; Wahl, M.I.; Lin, S.; Kato, R.M.; Fluckiger, A.-C.; Witte, O.N.; Kinet, J.-P. Activation of BTK by a Phosphorylation Mechanism Initiated by SRC Family Kinases. *Science* 1996, 271, 822–825.
26. Solvason, N.; Wu, W.W.; Kabra, N.; Lund-Johansen, F.; Roncarolo, M.G.; Behrens, T.W.; Grillot, D.A.; Nunez, G.; Lees, E.; Howard, M.; et al. Transgene Expression of bcl-xL Permits Anti-Immunoglobulin (Ig)-Induced Proliferation in Xid B Cells. *J. Exp. Med.* 1998, 187, 1081–1091.
27. Craxton, A.; Jiang, A.; Kurosaki, T.; Clark, E.A. Syk and Bruton's Tyrosine Kinase are Required for B Cell Antigen Receptor-Mediated Activation of the Kinase Akt. *J. Biol. Chem.* 1999, 274, 30644–30650.
28. Okada, T.; Ngo, V.N.; Ekland, E.H.; Forster, R.; Lipp, M.; Littman, D.R.; Cyster, J.G. Chemokine Requirements for B Cell Entry to Lymph Nodes and Peyer's Patches. *J. Exp. Med.* 2002, 196, 65–75.
29. De Gorter, D.J.; Beuling, E.A.; Kersseboom, R.; Middendorp, S.; van Gils, J.M.; Hendriks, R.W.; Pals, S.T.; Spaargaren, M. Bruton's Tyrosine Kinase and Phospholipase Cgamma2 Mediate Chemokine-Controlled B Cell Migration and Homing. *Immunity* 2007, 26, 93–104.
30. Alugupalli, K.R.; Akira, S.; Lien, E.; Leong, J.M. MyD88-and Bruton's Tyrosine Kinase-Mediated Signals are Essential for T Cell-Independent Pathogen-Specific IgM Responses. *J. Immunol.*

2007, 178, 3740–3749.

31. Mitsuiki, N.; Yang, X.; Bartol, S.J.; Grosserichter-Wagener, C.; Kosaka, Y.; Takada, H.; Imai, K.; Kanegane, H.; Mizutani, S.; van der Burg, M.; et al. Mutations in Bruton's Tyrosine Kinase Impair IgA Responses. *Int. J. Hematol.* 2015, 101, 305–313.
32. Ng, Y.S.; Wardemann, H.; Chelnis, J.; Cunningham-Rundles, C.; Meffre, E. Bruton's Tyrosine Kinase is Essential for Human B Cell Tolerance. *J. Exp. Med.* 2004, 200, 927–934.
33. Rajaiya, J.; Hatfield, M.; Nixon, J.C.; Rawlings, D.J.; Webb, C.F. Bruton's Tyrosine Kinase Regulates Immunoglobulin Promoter Activation in Association with the Transcription Factor Bright. *Mol. Cell Biol.* 2005, 25, 2073–2084.
34. Mundy-Bosse, B.L.; Lesinski, G.B.; Jaime-Ramirez, A.C.; Benninger, K.; Khan, M.; Kuppusamy, P.; Guenterberg, K.; Kondadasula, S.V.; Chaudhury, A.R.; La Perle, K.M.; et al. Myeloid-Derived Suppressor Cell Inhibition of the IFN Response in Tumor-Bearing Mice. *Cancer Res.* 2011, 71, 5101–5110.
35. Gunderson, A.J.; Kaneda, M.M.; Tsujikawa, T.; Nguyen, A.V.; Affara, N.I.; Ruffell, B.; Gorjestani, S.; Liudahl, S.M.; Truitt, M.; Olson, P.; et al. Bruton Tyrosine Kinase-Dependent Immune Cell Cross-talk Drives Pancreas Cancer. *Cancer Discov.* 2016, 6, 270–285.
36. Das, S.; Bar-Sagi, D. BTK Signaling Drives CD1d(hi)CD5(+) Regulatory B-Cell Differentiation to Promote Pancreatic Carcinogenesis. *Oncogene* 2019, 38, 3316–3324.
37. Kawakami, Y.; Inagaki, N.; Salek-Ardakani, S.; Kitaura, J.; Tanaka, H.; Nagao, K.; Kawakami, Y.; Xiao, W.; Nagai, H.; Croft, M.; et al. Regulation of Dendritic Cell Maturation and Function by Bruton's Tyrosine Kinase via IL-10 and Stat3. *Proc. Natl. Acad. Sci. USA* 2006, 103, 153–158.
38. Natarajan, G.; Oghumu, S.; Terrazas, C.; Varikuti, S.; Byrd, J.C.; Satoskar, A.R. A Tec Kinase BTK Inhibitor Ibrutinib Promotes Maturation and Activation of Dendritic Cells. *Oncoimmunology* 2016, 5, e1151592.
39. Johnson, A.R.; Kohli, P.B.; Katewa, A.; Gogol, E.; Belmont, L.D.; Choy, R.; Penuel, E.; Burton, L.; Eigenbrot, C.; Yu, C.; et al. Battling Btk Mutants With Noncovalent Inhibitors That Overcome Cys481 and Thr474 Mutations. *ACS Chem. Biol.* 2016, 11, 2897–2907.
40. Wu, J.; Liu, C.; Tsui, S.T.; Liu, D. Second-Generation Inhibitors of Bruton Tyrosine Kinase. *J. Hematol. Oncol.* 2016, 9, 80.
41. Brown, J.R. Ibrutinib (PCI-32765), the First BTK (Bruton's Tyrosine Kinase) Inhibitor in Clinical Trials. *Curr. Hematol. Malig. Rep.* 2013, 8, 1–6.
42. Burger, J.A.; Buggy, J.J. Bruton Tyrosine Kinase Inhibitor Ibrutinib (PCI-32765). *Leuk. Lymphoma* 2013, 54, 2385–2391.

43. Sivina, M.; Kreitman, R.J.; Arons, E.; Ravandi, F.; Burger, J.A. The Bruton Tyrosine Kinase Inhibitor Ibrutinib (PCI-32765) Blocks Hairy Cell Leukaemia Survival, Proliferation and B Cell Receptor Signalling: A New Therapeutic Approach. *Br. J. Haematol.* 2014, 166, 177–188.
44. Honigberg, L.A.; Smith, A.M.; Sirisawad, M.; Verner, E.; Loury, D.; Chang, B.; Li, S.; Pan, Z.; Thamm, D.H.; Miller, R.A.; et al. The Bruton Tyrosine Kinase Inhibitor PCI-32765 Blocks B-Cell Activation and is Efficacious in Models of Autoimmune Disease and B-Cell Malignancy. *Proc. Natl. Acad. Sci. USA* 2010, 107, 13075–13080.
45. Herman, S.E.; Gordon, A.L.; Hertlein, E.; Ramanunni, A.; Zhang, X.; Jaglowski, S.; Flynn, J.; Jones, J.; Blum, K.A.; Buggy, J.J.; et al. Bruton Tyrosine Kinase Represents a Promising Therapeutic Target for Treatment of Chronic Lymphocytic Leukemia and is Effectively Targeted by PCI-32765. *Blood* 2011, 117, 6287–6296.
46. De Rooij, M.F.; Kuil, A.; Geest, C.R.; Eldering, E.; Chang, B.Y.; Buggy, J.J.; Pals, S.T.; Spaargaren, M. The Clinically Active BTK Inhibitor PCI-32765 Targets B-Cell Receptor-and Chemokine-Controlled Adhesion and Migration in Chronic Lymphocytic Leukemia. *Blood* 2012, 119, 2590–2594.
47. Ponader, S.; Chen, S.S.; Buggy, J.J.; Balakrishnan, K.; Gandhi, V.; Wierda, W.G.; Keating, M.J.; O'Brien, S.; Chiorazzi, N.; Burger, J.A.; et al. The Bruton Tyrosine Kinase Inhibitor PCI-32765 Thwarts Chronic Lymphocytic Leukemia Cell Survival and Tissue Homing in Vitro and in Vivo. *Blood* 2012, 119, 1182–1189.
48. Thorp, B.C.; Badoux, X. Atrial Fibrillation as a Complication of Ibrutinib Therapy: Clinical Features and Challenges of Management. *Leuk. Lymphoma.* 2018, 59, 311–320.
49. Kriegsmann, K.; Kriegsmann, M.; Witzens-Harig, M. Acalabrutinib, A Second-Generation Bruton's Tyrosine Kinase Inhibitor. *Recent Results Cancer Res.* 2018, 212, 285–294.
50. Wu, J.; Zhang, M.; Liu, D. Acalabrutinib (ACP-196): A Selective Second-Generation BTK Inhibitor. *J. Hematol. Oncol.* 2016, 9, 21.
51. Byrd, J.C.; Harrington, B.; O'Brien, S.; Jones, J.A.; Schuh, A.; Devereux, S.; Chaves, J.; Wierda, W.G.; Awan, F.T.; Brown, J.R.; et al. Acalabrutinib (ACP-196) in Relapsed Chronic Lymphocytic Leukemia. *N. Engl. J. Med.* 2016, 374, 323–332.
52. Heather, B.K.H.; Gardner, L.; Raquel, I.; Ahmed, H.; Allard, K.; Bart, V.L.; Cheryl, A.; London, J.C.B.; Amy, J.J.; William, C.K. ACP-196: A Second Generation Btk Inhibitor Demonstrates Biological Activity in a Canine Model of B-Cell Non-Hodgkin Lymphoma. In Proceedings of the 105th Annual Meeting of the American Association for Cancer Research, San Diego, CA, USA, 5–9 April 2014; Volume 74.
53. Niemann, C.U.; Montraveta, A.; Herman, S.E.M.; Ingallinera, T.; Barf, T.; Colomer, D.; Wiestner, A. The Novel Bruton's Tyrosine Kinase Inhibitor ACP-196 Shows in Vivo Efficacy Against Human

- Chronic Lymphocytic Leukemia Cells Xenografted to the NSG Mouse Model. *Cancer Res.* 2014, 74, 2624.
54. Herman, S.E.; Sun, X.; McAuley, E.M.; Hsieh, M.M.; Pittaluga, S.; Raffeld, M.; Liu, D.; Keyvanfar, K.; Chapman, C.M.; Chen, J.; et al. Modeling Tumor-Host Interactions of Chronic Lymphocytic Leukemia in Xenografted Mice to Study Tumor Biology and Evaluate Targeted Therapy. *Leukemia* 2013, 27, 2311–2321.
55. Herman, S.E.M.; Montraveta, A.; Niemann, C.U.; Mora-Jensen, H.; Gulrajani, M.; Krantz, F.; Mantel, R.; Smith, L.L.; McClanahan, F.; Harrington, B.K.; et al. The Bruton Tyrosine Kinase (BTK) Inhibitor Acalabrutinib Demonstrates Potent On-Target Effects and Efficacy in Two Mouse Models of Chronic Lymphocytic Leukemia. *Clin. Cancer Res.* 2017, 23, 2831–2841.
56. Johnson, A.J.; Lucas, D.M.; Muthusamy, N.; Smith, L.L.; Edwards, R.B.; De Lay, M.D.; Croce, C.M.; Grever, M.R.; Byrd, J.C. Characterization of the TCL-1 Transgenic Mouse as a Preclinical Drug Development Tool for Human Chronic Lymphocytic Leukemia. *Blood* 2006, 108, 1334–1338.
57. Skarzynski, M.; Niemann, C.U.; Lee, Y.S.; Martyr, S.; Maric, I.; Salem, D.; Stetler-Stevenson, M.; Marti, G.E.; Calvo, K.R.; Yuan, C.; et al. Interactions between Ibrutinib and Anti-CD20 Antibodies: Competing Effects on the Outcome of Combination Therapy. *Clin. Cancer Res.* 2016, 22, 86–95.
58. Varikuti, S.; Singh, B.; Volpedo, G.; Ahirwar, D.K.; Jha, B.K.; Saljoughian, N.; Viana, A.G.; Verma, C.; Hamza, O.; Halsey, G.; et al. Ibrutinib Treatment Inhibits Breast Cancer Progression and Metastasis by Inducing Conversion of Myeloid-Derived Suppressor Cells to Dendritic Cells. *Br. J. Cancer* 2020, 122, 1005–1013.
59. Hong, D.; Rasco, D.; Veeder, M.; Luke, J.J.; Chandler, J.; Balmanoukian, A.; George, T.J.; Munster, P.; Berlin, J.D.; Gutierrez, M.; et al. A Phase 1b/2 Study of the Bruton Tyrosine Kinase Inhibitor Ibrutinib and the PD-L1 Inhibitor Durvalumab in Patients with Pretreated Solid Tumors. *Oncology* 2019, 97, 102–111.
60. Overman, M.; Javle, M.; Davis, R.E.; Vats, P.; Kumar-Sinha, C.; Xiao, L.; Mettu, N.B.; Parra, E.R.; Benson, A.B.; Lopez, C.D.; et al. Randomized Phase II Study of the Bruton Tyrosine Kinase Inhibitor Acalabrutinib, Alone or with Pembrolizumab in Patients with Advanced Pancreatic Cancer. *J. Immunother. Cancer* 2020, 8.
61. Tempero, M.; Oh, D.Y.; Tabernero, J.; Reni, M.; Van Cutsem, E.; Hendifar, A.; Waldschmidt, D.T.; Starling, N.; Bachet, J.B.; Chang, H.M.; et al. Ibrutinib in Combination with Nab-Paclitaxel and Gemcitabine for First-Line Treatment of Patients with Metastatic Pancreatic Adenocarcinoma: Phase III RESOLVE Study. *Ann. Oncol.* 2021, 32, 600–608.
62. Zhang, T.; Harrison, M.R.; O'Donnell, P.H.; Alva, A.S.; Hahn, N.M.; Appleman, L.J.; Cetnar, J.; Burke, J.M.; Fleming, M.T.; Milowsky, M.I.; et al. A Randomized Phase 2 Trial of Pembrolizumab Versus Pembrolizumab and Acalabrutinib in Patients with Platinum-Resistant Metastatic Urothelial Cancer. *Cancer* 2020, 126, 4485–4497.

Retrieved from <https://encyclopedia.pub/entry/history/show/73371>