

# Resistance Training on Blood Pressure

Subjects: Nursing

Contributor: Iván Chulvi-Medrano, Andres Loaiza-Betancur

The aim was to systematically review and meta-analyze the current evidence for the effects of resistance training (RT) on blood pressure (BP) as the main outcome and body mass index (BMI) in children and adolescents. Limited evidence suggests that RT has no adverse effects on BP and may positively affect BP in youths. More high-quality studies are needed to clarify the association between RT and BP in light of body composition changes throughout childhood and adolescence.

Keywords: resistance training ; hypertension ; blood pressure ; youth

---

## 1. Introduction

The treatment for hypertension is usually pharmacological and has shown to be effective in 50% of adult patients <sup>[1]</sup>. However, in younger populations pharmacological treatment should be reserved for those who present with persistent elevated blood pressure (BP) despite lifestyle modification <sup>[2]</sup>. Therefore, it is reasonable to investigate non-pharmacologic treatments for youth and to emphasize preventative strategies including regular physical activity. Resistance training (RT) has been suggested as an effective non-pharmacological treatment for the prevention and management of high BP in adults <sup>[3][4]</sup>, yet little is known about the effects of RT on BP in children and adolescents (6–18 years of age) <sup>[3]</sup>.

Research evidence has found that cardiovascular disease has its roots in childhood, with some reports of endothelial damage occurring early in life <sup>[5]</sup>. The prevalence of diagnosed primary pediatric hypertension is increasing <sup>[6][7]</sup>. Primary pediatric hypertension is the cardiovascular condition whereby systolic or diastolic BP values are > 95th percentile for boys and girls up to 12 years of age and > 130/80 mmHg for youth older than 13 years of age <sup>[6]</sup>. Primary pediatric hypertension (as early as 7 years of age) has been associated with pathophysiological changes that tracks into later stages <sup>[6][8]</sup>. Moreover, the prevalence of obesity is increasing among youth and it has been identified as a risk factor for elevated BP <sup>[9][10]</sup>. Thus, the prevention and management of obesity early in life should be a primary consideration for reducing the prevalence of pediatric hypertension <sup>[9]</sup>. Of note, data from diverse populations indicate that childhood BP is associated with BP later in life <sup>[11]</sup>. Therefore, early treatment and management are needed since accelerated weight gain in youth may increase the risk of elevated BP later in life <sup>[12]</sup>. Juonala et al. reported that overweight or obesity early in life was predictive of many comorbidities and found that youth who were overweight or obese but who became nonobese as adults had a cardiovascular risk profile that was similar to those who were never obese <sup>[13]</sup>. Therefore, maintenance of normal body weight in children and adolescents may prevent the clustering of hypertension and other cardiovascular disease risk factors in adulthood <sup>[11]</sup>. Body mass index (BMI) is the most commonly used surrogate measure of adiposity and screening tool for cardio-metabolic risk <sup>[5]</sup>.

Along with weight maintenance, physical activity can improve BP levels in adults independently of pharmacological treatment <sup>[14]</sup>. A clinical report demonstrated a decrease in BP values of  $-5/8$  mmHg in hypertensive adults following aerobic training <sup>[15]</sup>. Traditionally, research and clinical efforts have focused on aerobic training as a means of BP management. Recently, RT has gained attention as an important modulator of BP. Regular participation in RT has been found to reduce BP by  $-4$  mmHg and  $-5$  mmHg in hypertensive adults who performed dynamic and isometric RT, respectively <sup>[15]</sup>.

In addition to increasing muscular strength, muscular power, and local muscular endurance, RT in youth has shown to produce many health benefits including improvements in cardiovascular fitness, body composition, bone mineral density, blood lipid profiles, insulin sensitivity, injury resistance, and mental health <sup>[16][17][18][19][20][21][22][23][24]</sup>. By definition, resistance training is a specialized method of conditioning that involves the use of different modes of training with a wide range of resistive loads including body weight exercises and free weights (barbells and dumbbells) <sup>[16]</sup>. Although the potential health benefits of RT in youth have been widely studied, there is limited understanding about the effects of RT on BP in children and adolescents. Several systematic reviews and meta-analyses have examined the positive effects of RT on BP values in adults. However, no previous systematic review has quantitatively examined the association between RT

on BP and BMI in youth. Given this research gap, a systematic review was conducted to examine the literature regarding the effects of youth RT on systolic and diastolic BP. In addition, a meta-analysis of selected studies was conducted to quantitatively evaluate the effects of RT on systolic and diastolic BP, and BMI, in children and adolescents. Given the potential health-related benefits of RT in adults, we hypothesized that RT would also produce beneficial effects on BP and BMI values in youth.

## 2. Discussion

The aim of this systematic review and meta-analysis was to quantify the effect of RT on the values of SBP, DBP and BMI in youth. To the best of our knowledge, this is the first systematic review with a subsequent meta-analysis that investigates the effects of RT on BP values in children and adolescents. While other studies have investigated the role of physical activity on cardiometabolic health in youth [25][26], no previous reports have examined the influence of RT in this population. As shown in previous research [3][27], RT has been found to offer observable health-related benefits in adults. Thus, we hypothesized that RT would have positive effects on BP and BMI in youth.

Our main findings are that RT resulted in non significant reductions in SBP ( $-1.09$  mmHg;  $P = 0.32$ ) and DBP ( $-0.93$  mmHg;  $P = 0.10$ ) and statistically significant reductions in BMI ( $-0.43$  kg/m<sup>2</sup>;  $P = 0.03$ ) in youth. Although the research reports in this review failed to show statistical significance in terms of the ability of RT to lower systolic and diastolic BP, several factors need to be considered. These factors include the design of the RT protocols (i.e., training intensity, volume, frequency and duration) as well as the health status (all were normotensive) and the training status of the participants. Conflicting findings from several studies are likely due to differences in outcomes measures, study designs and study populations. Regarding the RCTs examined in our review, researchers used different RT protocols. While three studies performed RT with bodyweight exercises [28][29][30], one used sandbags and dumbbells [31], the others used weight machines [32][33][34][35]. Notable, there was wide variation in the prescription of RT variables including intensity, volume, frequency or duration. For instance, some protocols proposed two weekly sessions of high RT (12 RM) [31] while others trained 4 days per week with a moderate to high intensity (8 RM) [35]. Further, two studies added nutritional guidelines along with the RT protocol [28][33]. Interpretation and comparison of results would be more accurate with similar RT protocols and with subgroup analysis (i.e., obese and normal body weight; hypertensive and normotensive). There were also differences in the configuration of the control groups among studies that could have impacted the outcomes. For example, two studies did not advise participants about extra physical activity at school or in community based programs [29][35]. In one study that included adolescents who were obese, the control group consisted of adolescents who were not obese [32]. This aforementioned report showed moderate and substantial heterogeneity values in DBP and SBP, respectively (37% and 67%). The heterogeneous values found in this study could help explain why no statistically significant changes were found in SBP and DBP values following RT [32].

RT is an evidence-based preventative exercise intervention strategy that can promote health and well-being through the life course [27][36]. The benefits of progressive RT on muscular strength, muscular power, and local muscular endurance of children and adolescents is well described in several meta-analysis [37][38][39]. Moreover, RT has shown to produce many health-related benefits including improvements in cardiovascular fitness, body composition, bone mineral density, blood lipid profiles, insulin sensitivity, injury resistance and mental health improvements [16][18][19][20][21][22][23][24][40]. Longitudinal studies have confirmed the inverse relationship between low levels of strength early in life and risk of cardiovascular disease later in life [36][41][42][43]. Therefore, it seems plausible that RT could lower BP concurrent with improvements in other health markers. Some studies have speculated that the reduction in BP following RT in youth might be due to an increase in skeletal muscle mass which, in turn, may lead to a myocardial relaxation [34], diastolic filling peak velocity at the mitral septal annulus [34], an improvement in autonomic modulation [32] and/or an enhanced endothelial function [31]. In obese children, functional and structural cardiac abnormalities (i.e., increased left ventricle and left atrium dimensions, diastolic and systolic left ventricle, and right ventricle dysfunction) have been described in comparison to normo-weight children [44][45]. In this sense, BP mechanisms might be different. Further studies are needed in order to clarify the hypothetical link between RT and BP improvements in youth.

Our findings show a statistically significant improvement in BMI ( $P = 0.03$ ) after an RT intervention. It has been established that exercise interventions can alter body composition (e.g., increase fat free mass) while BMI can remain the same or in some cases increase due to the increase in muscle mass [46]. Indeed, some studies demonstrated no change of BMI following RT despite the remarkable benefits on other health parameters such as endothelial function [47][48]. Therefore, BMI values may underestimate the effectiveness of RT interventions with respect to cardiovascular disease risk [49].

This study has several limitations that should be acknowledged; (1) the lack of systematic quantification of the RT intensity, volume or exercise selection; (2) BP was not a primary outcome in many of the studies included in the analysis; (3) heterogeneity in the outcome measurement procedures; (4) most of the RCTs analyzed did not adequately perform or report random sequence generation, allocation concealment and blinding of outcome assessment and; (5) moderate and substantial values of heterogeneity on SBP and DBP were found.

Although limited research has examined the effects of RT on BP in youth, our results suggest that RT does not have an adverse effect on the BP of children and adolescents and may be beneficial in lowering BP and improving BMI in this young population. Unfortunately, our findings do not allow for a recommendation on a specific dose of RT for effectively managing BP in youth. Nevertheless, a technique-driven and progressive RT program including multijoint exercises that involve the large muscle groups should be considered in the design of youth physical activity programs [16]. Further research is needed to effectively examine the "dose response" (e.g., intensity, volume, frequency) of youth RT interventions while exploring novel modes of RT like low intensity isometric handgrip exercise [50][51].

### 3. Conclusions

The present shows that there is limited data to determine the effects of RT on BP values in youth, while significant improvements in BMI have been demonstrated. Although the studies show a tendency towards reducing systolic and diastolic BP, the heterogeneity of the RT intensity, volume, frequency or duration make the interpretation of results difficult. Mechanisms by which RT may induce favorable adaptations in BP in youth are speculative. More high-quality studies are needed to clarify the association between RT and BP in youth with and without clinical conditions.

---

### References

1. Hajjar, I.; Kotchen, T.A. Trends in Prevalence , Awareness , in the United States , 1988-2000. *J. Am. Med. Assoc.* 2003, 290, 199–206, doi:10.1161/HYPERTENSIONAHA.117.09801.
2. Lande, M.B.; Kupferman, J.C. Pediatric hypertension: The Year in Review. *Clin. Pediatr. (Phila)*. 2014, 39, 57–60, doi:10.1177/0009922813499968.
3. Cornelissen, V.; Smart, N.A. Exercise Training for Blood Pressure: A Systematic Review and Meta-analysis. *J. Am. Heart Assoc.* 2013, doi:10.1161/JAHA.112.004473.
4. Chulvi, I.; Sanchis, J.; Tortosa, J.; Cortell, J.M. Exercise for hypertension. *Fit. Med.* 2016, doi:10.5772/65035.
5. Barlow, S.E. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: summary report. *Pediatrics* 2007, 120 Suppl, doi:10.1542/peds.2007-2329C.
6. Taylor-Zapata, P.; Baker-Smith, C.M.; Burckart, G.; Daniels, S.R.; Flynn, J.T.; Giacoia, G.; Green, D.; Kelly, A.S.; Khurana, M.; Li, J.S.; et al. Research gaps in primary pediatric hypertension. *Pediatrics* 2019, 143, doi:10.1542/peds.2018-3517.
7. Weaver, D.J. Pediatric hypertension: Review of updated guidelines. *Pediatr. Rev.* 2019, 40, 354–358, doi:10.1542/pir.2018-0014.
8. Raitakari, O.; Juonala, M.; Nnema, T.R.; Kerblom, H.; Viikari, J. Cardiovascular risk factors in childhood as predictors of carotid artery intima-media thickness in adulthood. *Atheroscler. Suppl.* 2003, 4, 264, doi:10.1016/s1567-5688(03)91137-8.
9. McMurray, R.G.; Ondrak, K.S. Cardiometabolic Risk Factors in Children: The Importance of Physical Activity. *Am. J. Lifestyle Med.* 2013, 7, 292–303, doi:10.1177/1559827613481429.
10. Falkner, B.; Daniels, S.R. Summary of the fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004, 44, 387–388, doi:10.1161/01.HYP.0000143545.54637.af.
11. Chen, X.; Wang, Y.; Chen, X.; Wang, Y. Tracking of Blood Pressure From Childhood to Adulthood A Systematic Review and Meta— Regression Analysis. *Circulation* 2008, 3171–3180, doi:10.1161/CIRCULATIONAHA.107.730366.
12. Law, C.M.; Shiell, A.W.; Newsome, C.A.; Syddall, H.E.; Shinebourne, E.A.; Fayers, P.M.; Martyn, C.N.; De Swiet, M. Fetal, infant, and childhood growth and adult blood pressure: A longitudinal study from birth to 22 years of age. *Circulation* 2002, 105, 1088–1092, doi:10.1161/hc0902.104677.
13. Juonala, M.; Magnussen, C.G.; Berenson, G.S.; Venn, A.; Burns, T.L.; Sabin, M.A.; Srinivasan, S.R.; Daniels, S.R.; Davis, P.H.; Chen, W.; et al. Childhood adiposity, adult adiposity, and cardiovascular risk factors. *N. Engl. J. Med.* 2011, 365, 1876–1885, doi:10.1056/NEJMoa1010112.

14. Williams, B.; Mancia, G.; Spiering, W.; Rosei, E.A.; Azizi, M.; Burnier, M.; Clement, D.; Coca, A.; De Simone, G.; Dominiczak, A.; et al. 2018 practice guidelines for the management of arterial hypertension of the European society of cardiology and the European society of hypertension ESC/ESH task force for the management of arterial hypertension. *J. Hypertens.* 2018, 36, 2284–2309, doi:10.1097/HJH.0000000000001961.
15. Arnett, D.K.; Blumenthal, R.S.; Albert, M.A.; Buroker, A.B.; Goldberger, Z.D.; Hahn, E.J.; Himmelfarb, C.D.; Khera, A.; Lloyd-Jones, D.; McEvoy, J.W.; et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019, 140, e596–e646, doi:10.1161/CIR.0000000000000678.
16. Stricker, P.R.; Faigenbaum, A.D.; McCambridge, T.M. Resistance training for children and adolescents. *Pediatrics* 2020, 145, doi:10.1542/peds.2020-1011.
17. Faigenbaum, A.D.; Lloyd, R.S.; Myer, G.D. Youth resistance training: Past practices, new perspectives, and future directions. *Pediatr. Exerc. Sci.* 2013, 25, 591–604, doi:10.1123/pes.25.4.591.
18. Myer, G.; Faigenbaum, A.; Chu, D.; Falkel, J.; Ford, K.; Best, T.; Hewett, T. Integrative Training for Children and Adolescents: Techniques and Practices for Reducing Sports-Related Injuries and Enhancing Athletic Performance. *Phys. Sportsmed.* 2011, 39, 74–84, doi:10.3810/psm.2011.02.1854.
19. Lloyd, R.S.; Faigenbaum, A.D.; Stone, M.H.; Oliver, J.L.; Jeffreys, I.; Moody, J.A.; Brewer, C.; Pierce, K.C.; McCambridge, T.M.; Howard, R.; et al. Position statement on youth resistance training: The 2014 International Consensus. *Br. J. Sports Med.* 2014, 48, 498–505, doi:10.1136/bjsports-2013-092952.
20. Hind, K.; Burrows, M. Weight-bearing exercise and bone mineral accrual in children and adolescents: A review of controlled trials. *Bone* 2007, 40, 14–27, doi:10.1016/j.bone.2006.07.006.
21. Bernardoni, B.; Thein-Nissenbaum, J.; Fast, J.; Day, M.; Li, Q.; Wang, S.; Scerpella, T. A school-based resistance intervention improves skeletal growth in adolescent females. *Osteoporos. Int.* 2014, 25, 1025–1032, doi:10.1007/s00198-013-2535-y.
22. Ishikawa, S.; Kim, Y.; Kang, M.; Morgan, D.W. Effects of weight-bearing exercise on bone health in girls: A meta-analysis. *Sport. Med.* 2013, 43, 875–892, doi:10.1007/s40279-013-0060-y.
23. Lauersen, J.B.; Andersen, T.E.; Andersen, L.B. Strength training as superior, dose-dependent and safe prevention of acute and overuse sports injuries: A systematic review, qualitative analysis and meta-analysis. *Br. J. Sports Med.* 2018, 52, 1557–1563, doi:10.1136/bjsports-2018-099078.
24. Collins, H.; Booth, J.N.; Duncan, A.; Fawkner, S.; Niven, A. The Effect of Resistance Training Interventions on 'The Self' in Youth: a Systematic Review and Meta-analysis. *Sport. Med. - Open* 2019, 5, doi:10.1186/s40798-019-0205-0.
25. Martínez-Vizcaino, V.; Pozuelo-Carrascosa, D.P.; García-Prieto, J.C.; Cavero-Redondo, I.; Solera-Martínez, M.; Garrido-Miguel, M.; Díez-Fernández, A.; Ruiz-Hermosa, A.; Sánchez-López, M. Effectiveness of a school-based physical activity intervention on adiposity, fitness and blood pressure: MOVI-KIDS study. *Br. J. Sports Med.* 2020, 54, 279–285, doi:10.1136/bjsports-2018-099655.
26. Pozuelo-Carrascosa, D.P.; Cavero-Redondo, I.; Herraiz-Adillo, A.; Díez-Fernandez, A.; Sanchez-Lopez, M.; Martinez-Vizcaino, V. School-based exercise programs and cardiometabolic risk factors: A meta-analysis. *Pediatrics* 2018, 142, doi:10.1542/peds.2018-1033.
27. Westcott, W.L. Resistance training is medicine: effects of strength training on health. *Curr. Sports Med. Rep.* 2012, 11, 209–216, doi:10.1249/JSR.0b013e31825dabb8.
28. Horner, K.; Barinas-Mitchell, E.; DeGroff, C.; Kuk, J.L.; Drant, S.; Lee, S. et al. Effect of Aerobic versus Resistance Exercise on Pulse Wave Velocity, Intima Media Thickness and Left Ventricular Mass in Obese Adolescents. *Pediatr. Exerc. Sci.* 2015, 27, 494–502, doi:10.1123/pes.2015-0067.
29. Kelly, L.A.; Loza, A.; Lin, X.; Schroeder, E.T.; Hughes, A.; Kirk, A.; Knowles, A.-M. The effect of a home-based strength training program on type 2 diabetes risk in obese Latino boys. *J. Pediatr. Endocrinol. Metab.* 2015, 28, 315–322, doi:10.1515/jpem-2014-0470.
30. Larsen, M.N.; Nielsen, C.M.; Madsen, M.; Manniche, V.; Hansen, L.; Bangsbo, J.; Krstrup, P.; Hansen, P.R.; et al. Cardiovascular adaptations after 10 months of intense school-based physical training for 8- to 10-year-old children. *Scand. J. Med. Sci. Sports* 2018, 28, 33–41, doi:10.1111/sms.13253.
31. Yu, C.C.-W.; McManus, A.M.; So, H.-K.; Chook, P.; Au, C.-T.; Li, A.M.; Kam, J.T.-C.; So, R.C.-H.; Lam, C.W.-K.; Chan, I.H.-S.; et al. Effects of resistance training on cardiovascular health in non-obese active adolescents. *World J. Clin. Pediatr.* 2016, 5, 293–300, doi:10.5409/wjcp.v5.i3.293.
32. Farinatti, P.; Neto, M.; Dias, I.; Cunha, F.A.; Bouskela, E.; Kraemer-Aguiar, L.G. Short-Term Resistance Training Attenuates Cardiac Autonomic Dysfunction in Obese Adolescents. *Pediatr. Exerc. Sci.* 2016, 28, 374–380,

33. Lau, P.W.C.; Yu, C.W.; Lee, A.; Sung, R.Y.T. The physiological and psychological effects of resistance training on Chinese obese adolescents. *J. Exerc. Sci. Fit.* 2004, 2, 115–120.
34. Naylor, L.H.; Watts, K.; Sharpe, J.A.; Jones, T.W.; Davis, E.A.N.N.; Thompson, A.; George, K.; Ramsay, J.M.; Driscoll, G.O.; Green, D.J. Resistance Training and Diastolic Myocardial Tissue Velocities in Obese Children. *Med. Sci. Sport. Exerc.* 2008, 2027–2032, doi:10.1249/MSS.0b013e318182a9e0.
35. Sigal, R.J.; Alberga, A.S.; Goldfield, G.S.; Prud'homme, D.; Hadjiyannakis, S.; Gougeon, R.; Phillips, P.; Tulloch, H.; Malcolm, J.; Doucette, S.; et al. Effects of aerobic training, resistance training, or both on percentage body fat and cardiometabolic risk markers in obese adolescents: The healthy eating aerobic and resistance training in youth randomized clinical trial. *JAMA Pediatr.* 2014, 168, 1006–1014, doi:10.1001/jamapediatrics.2014.1392.
36. García-Hermoso, A.; Ramírez-Campillo, R.; Izquierdo, M. Is Muscular Fitness Associated with Future Health Benefits in Children and Adolescents? A Systematic Review and Meta-Analysis of Longitudinal Studies. *Sport. Med.* 2019, 49, 1079–1094, doi:10.1007/s40279-019-01098-6.
37. Behringer, M.; Vom Heede, A.; Yue, Z.; Mester, J. Effects of resistance training in children and adolescents: A meta-analysis. *Pediatrics* 2010, 126, doi:10.1542/peds.2010-0445.
38. Lesinski, M.; Prieske, O.; Granacher, U. Effects and dose-response relationships of resistance training on physical performance in youth athletes: A systematic review and meta-analysis. *Br. J. Sports Med.* 2016, 50, 781–795, doi:10.1136/bjsports-2015-095497.
39. Harries, S.K.; Lubans, D.R.; Callister, R. Resistance training to improve power and sports performance in adolescent athletes: A systematic review and meta-analysis. *J. Sci. Med. Sport* 2012, 15, 532–540, doi:10.1016/j.jsams.2012.02.005.
40. Faigenbaum, A.D.; Kraemer-Aguiar, L.G.; Blimkie, C.J.R.; Jeffreys, I.; Micheli, L.J.; Nitka, M.; Rowland, T.W. Youth Resistance Training: Updated Position Statement Paper From the National Strength and Conditioning Association. *J. Strength Cond. Res.* 2009, 23, 60–79.
41. Ortega, F.B.; Ruiz, J.R.; Castillo, M.J.; Sjöström, M. Physical fitness in childhood and adolescence: A powerful marker of health. *Int. J. Obes.* 2008, 32, 1–11, doi:10.1038/sj.ijo.0803774.
42. Grontved, A.; Ried-Larsen, M.; Moller, N.C.; Kristensen, P.L.; Froberg, K.; Brage, S.; Andersen, L.B. Muscle strength in youth and cardiovascular risk in young adulthood (the European Youth Heart Study). *Br. J. Sports Med.* 2015, 49, 90–94, doi:10.1136/bjsports-2012-091907.
43. Cohen, D.D.; Gómez-Arbeláez, D.; Camacho, P.A.; Pinzon, S.; Hormiga, C.; Trejos-Suarez, J.; Duperly, J.; Lopez-Jaramillo, P. Low muscle strength is associated with metabolic risk factors in Colombian children: The ACFIES study. *PLoS One* 2014, 9, e93150, doi:10.1371/journal.pone.0093150.
44. Alkholi, U.M.; Ahmed, I.A.; Karam, N.A.; Ali, Y.F.; Yosry, A. Assessment of left ventricular mass index could predict metabolic syndrome in obese children. *J. Saudi Hear. Assoc.* 2016, 28, 159–166, doi:10.1016/j.jsha.2015.06.002.
45. Korkmaz, O.; Gursu, H.A.; Karagun, B.S. Comparison of echocardiographic findings with laboratory parameters in obese children. *Cardiol. Young* 2016, 26, 1060–1065, doi:10.1017/S1047951115001729.
46. Watts, K.; Jones, T.W.; Davis, E.A.; Green, D. Exercise training in obese children and adolescents: Current concepts. *Sport. Med.* 2005, 35, 375–392, doi:10.2165/00007256-200535050-00002.
47. Ferguson, M.A.; Gutin, B.; Owens, S.; Barbeau, P.; Tracy, R.P.; Litaker, M. Effects of physical training and its cessation on the hemostatic system of obese children. *Am. J. Clin. Nutr.* 1999, 69, 1130–1134, doi:10.1093/ajcn/69.6.1130.
48. Watts, K.; Beye, P.; Sifarakas, A.; O'Driscoll, G.; Jones, T.W.; Davis, E.A.; Green, D.J. Effects of exercise training on vascular function in obese children. *J. Pediatr.* 2004, 144, 620–625, doi:10.1016/j.jpeds.2004.02.027.
49. Cote, A.T.; Harris, K.C.; Panagiotopoulos, C.; Sandor, G.G.S.; Devlin, A.M. Childhood Obesity and Cardiovascular Dysfunction. *J. Am. Coll. Cardiol.* 2013, 62, doi:10.1016/j.jacc.2013.07.042.
50. Loaiza-Betancur, A.F.; Pérez Bedoya, E.; Montoya Dávila, J.; Chulvi-Medrano, I. Effect of Isometric Resistance Training on Blood Pressure Values in a Group of Normotensive Participants: A Systematic Review and Meta-analysis. *Sports Health* 2020, 12, 256–262, doi:10.1177/1941738120908070.
51. Loaiza-Betancur, A.F.; Chulvi-Medrano, I. Is Low-Intensity Isometric Handgrip Exercise an Efficient Alternative in Lifestyle Blood Pressure Management? A Systematic Review. *Sports Health* 2020, 12, 470–477, doi:10.1177/1941738120943882.

