

Medicinal Significance of 1,2,3-Thiadiazoles Derivatives

Subjects: **Medical Informatics**


Contributor: Ali Irfan

The 1,2,3-thiadiazole moiety occupies a significant and prominent position among privileged heterocyclic templates in the field of medicine, pharmacology and pharmaceuticals due to its broad spectrum of biological activities.


[1](#)[2](#)[3-thiadiazole](#)[synthetic strategies](#)[biological activities](#)[antiviral](#)[anticancer](#)[antifungal](#)[insecticidal](#)

1. Introduction


The fascinating aromatic 1,2,3-thiadiazole is a structurally active pharmacophore and great interest for researchers due to its versatile and wide array of biological activities in the field of medicine, pharmacology and pharmaceuticals. Thiadiazoles occur naturally in four different isomeric forms, having one sulfur and two nitrogen atoms with hydrogen binding domain as presented in **Figure 1** [\[1\]](#)[\[2\]](#)[\[3\]](#)[\[4\]](#).




1,2,3-Thiadiazole, **1**



1,2,4-Thiadiazole, **2**



1,2,5-Thiadiazole, **3**



1,3,4-Thiadiazole, **4**

Figure 1. Structures of different isomeric forms of thiadiazole (1–4).

Thiadiazole scaffold emerges as an interesting structural motif due to the presence of different heteroatoms in its structure and its promising therapeutic activities in the treatment and cure of several diseases. Different substitutions on the thiadiazole nucleus improve the structure–activity relationship (SAR) of thiadiazole hybrid structures to enhance its therapeutic efficacy against a wide variety of pathogens [1][5]. Thiadiazole scaffolds exhibit a wide range of biological activities including antimicrobial [6][7][8][9][10][11], carbonic anhydrase inhibitors [12][13][14], antifungal [15][16][17][18][19], antibacterial [20][21][22][23][24][25], systemic acquired resistance [26][27], insecticidal activity [28][29], antiviral [30][31][32][33], antioxidant [34][35], anticonvulsant [36][37][38][39][40][41][42], antihypertensive activity [43][44], anticancer and antitumor [45][46][47][48][49][50][51], analgesic [52][53], anti-inflammatory [54][55][56][57][58], plant activator [59][60], antidiabetic [61], antileishmanial activity [62][63][64][65] and antituberculosis [66][67][68][69][70][71]. The different marketed drugs of thiadiazole moiety (5–14) are given below [26][72][73][74][75][76] (Figure 2).




Figure 2. Commercial drugs based on different thiadiazole scaffolds (5–14).

The bioactive 1,2,3-thiadiazole analogues display excellent and outstanding new profiles of medicinal, pharmacological and pharmaceutical activities [1][77]. In this review article, the most recent and relevant synthetic approaches and pharmacological activities of 1,2,3-thiadiazole analogues are discussed and can be classified into antiviral, insecticidal, antifungal and anticancer activities.

2. Biological Activities of 1,2,3-Thiadiazole Derivatives

The main objective of the present section is the search and presentation of the 1,2,3-thiadiazole based most potent therapeutic agents for a variety of biological and pharmacological applications with little or no side effects. In this section, 1,2,3-thiadiazole derivatives have been divided into classes according to their biological activity.

2.1. Antiviral Agents

Pawar et al. described antiviral drugs as such bioactive compounds which are used for the treatment of viral infections [78]. Rossignol et al. reported that most of the antiviral agents are specifically targeted oriented while some other antivirals have a broad spectrum against a vast variety of viral infections [79]. Some antiviral drugs such

as ribavirin (**89**; **Figure 3**) are being used against both RNA and DNA viruses [80][81][82], efavirenz (**90**; **Figure 3**) used for the treatment of HIV/AIDS [83][84][85] and arbidol (**91**; **Figure 3**) was used against influenza viruses [86][87].





Figure 3. Structures of antiviral drugs **89–91**.

Zhan et al. prepared 1,2,3-thiadiazole thioacetanilide by adopting a multi-step methodology and exhibited their anti-HIV (Human Immunodeficiency Virus) activity against MT-4 cells. Among all synthesized derivatives, the thioacetanilide based 1,2,3-thiadiazole scaffold (**92**; **Figure 4**) showed remarkably the best anti-HIV activity in terms of EC_{50} value $0.059 \pm 0.02 \mu\text{M}$, $CC_{50} > 283.25 \mu\text{M}$, $SI > 4883 \mu\text{M}$ compared to the reference compounds such as NVP (nevirapine) DLV (delaviridine), EFV (Efavirenz), AZT (azidothymidine) and VRX-480773. SAR displayed that the significant enhancement in the antiviral efficacy of compound **92** was due to the introduction of the nitro group on the phenyl ring. Anilide scaffold having NO_2 and halogen-substituted aryl ring at o-position took great part to enhance the anti-HIV potential, however, the case was the difference for fluorine atom which decreased activity at a certain level [88].



92

Figure 4. Thioacetanilide based 1,2,3-thiadiazole scaffold **92** exhibiting maximum anti-HIV potential.

Zhan et al. prepared potent 1,2,3-thiadiazole derivatives against HIV-1. The new 1,2,3-thiadiazole (**93**; **Figure 5**) proved as the most active anti HIV-1 agent (EC_{50} value $0.0364 \pm 0.0038 \mu\text{M}$, $CC_{50} > 240.08 \mu\text{M}$ and $SI > 6460 \mu\text{M}$) against MT-4 cells among the synthesized derivatives in comparison with the reference compounds NVP (EC_{50} $0.208 \mu\text{M}$) and DLV (EC_{50} $0.320 \mu\text{M}$) EFV (EC_{50} $0.00440 \mu\text{M}$) and AZT (EC_{50} $0.0151 \mu\text{M}$). Phenyl ring substituted with 2,4-Br₂ group significantly increased the antiviral potential of compound **93** than all other substituents on the phenyl ring. SAR studies indicated decreasing order of antiviral strength, i.e., 2,4-Br₂ > 2,4-Cl₂ > 2,4-F₂ [89].




Figure 5. Anti-HIV activity of 5-(2,4-dibromophenyl)-1,2,3-thiadiazole **93**.

Dong et al. prepared potent antiviral piperidine-based thiadiazole derivatives, and the SAR study showed that chlorine atom substituted compounds exhibited good antiviral activity and the substitution on phenyl ring affects inhibitory action. 1,2,3-thiadiazole (**94**; **Figure 6**) displayed excellent potency with IC_{50} 3.59 $\mu\text{g/mL}$ as compared to lamivudine (IC_{50} value 14.8 $\mu\text{g/mL}$). It was observed that compounds having *p*-substituents exhibited more cytotoxic potency than the *o*-substituted derivatives. Moreover, lowering of the selective index with higher anti-HBV potencies of the synthesized compounds was observed when comparing the results with lamivudine. Compound **94** proved to be the most active one as compared to the other derivatives [90].




Figure 6. Antiviral activity of 1,2,3-thiadiazole **94**.

Synthesis and screening of antiviral potency of tetrazole-based 1,2,3-thiadiazoles structural against tobacco mosaic virus (TMV, anti-TMV) were evaluated [91]. The studies indicated that some obtained compounds (**95–98**; **Figure 7**) have higher anti-TMV activity (33.75–48.73%) than ribavirin at 100mg/mL concentration (33.23%) and comparable potency to ribavirin at 100 µg/mL. Structure **97** displayed the best protection effect among the synthesized derivatives as well as from the ribavirin—reference drug. Anti-TMV activity of synthesized scaffolds and reference drugs decreases in order of **98** > ninamycin > **97** > **95** > **96** > ribavirin while the protection effect decreased in order of ninamycin > **97** > **96** > **95** > ribavirin > **98**. Compound **98** display a higher inhibition potential, i.e., 48.73% as compared to ninamycin and ribavirin. The SAR study showed that the **2-fluorophenyl substituent** derivative **97** displayed **the best protection effect** among the synthesized congeners as well as from the ribavirin reference drug. The protection effect decreased in order of **ninamycin** > **2-fluorophenyl** > **cyclopropyl** > **isopropyl** > **ribavirin** > **4-ethylphenyl**. Overall the scaffold **97** proved to be a promising therapeutic agent, which has a better protection effect as well as the best anti-TMV activity [91].




Figure 7. Antiviral activity of terazole derivatives **95–98**.

Mao et al. reported the synthetic protocol to afford substituted methyl carbohydrazide-based 1,2,3-thiadiazoles and determined the anti-TMV activity. The thiophene containing carbohydrazide 1,2,3-thiadiazole (**99**; **Figure 8**) showed potent direct anti-TMV activity with 58.72% and 61.03% induction potencies at 50 $\mu\text{g/mL}$ when compared with reference drugs, ninamycin and tiadinil. Both the standard reference drugs, ninamycin and tiadinil, exhibited anti-TMV activity (54.93% and 7.94%) and induction activity (18.58% and 59.25%) respectively in vivo at 50 $\mu\text{g/mL}$ concentration. The scaffold **99** showed remarkably and significant higher antiviral potential than the standard reference drugs. The excellent anti-TMV activity of analogue **99** was due to the presence of thiophene moiety as compare to all the other tested compounds. The structural motif **99** showed significantly higher induction activity (61.03%) more than the induction activities of reference drugs, ninamycin and tiadinil (18.58%) and (59.25%), respectively. So the derivative **99** proved to be a plant elicitor and anti-TMV agent simultaneously [92].




Figure 8. Antiviral 1,2,3-thiadiazole scaffold **99**.

Another example of compounds with anti-TMV activity are structures **100** and **101** with a 1,3,4-oxadiazole ring (**Figure 9**) [93]. They showed potent curative, inactivation and protection effects against TMV. The derivative **100** displayed curative rate 54.1%, inactivation rate 90.3% and protection rate 52.8% while the compound **101** showed 47.1, 85.5 and 46.4% curative, inactivation and protection rates, respectively. The ningnanmycin was used as a standard reference drug, exhibiting a 56.1% curative rate, 92.5% inactivation rate and 59.3% protection rate. The incorporation of 1,3,4-oxadiazole moiety in the skeleton of 1,2,3-thiadiazoles significantly increased the anti-TMV activity [93].




Figure 9. 1,2,3-thiadiazoles **100** and **101** with anti-TMV activity.

Other promising anti-TMV agents **102** and **103** were presented by Zheng et al. (**Figure 10**) [94]. The substituted 1,2,3-thiadiazole-4-carboxamide scaffold **102** exhibited the best antiviral curative activity of 60 and 47%, at 500 and 100 µg/mL concentrations, while the standard tiadinil showed curative activity 58 and 46% at 500 and 100 µg/mL concentrations, respectively. Compound **103** showed its potent protective effect 76 and 71% at 500 and 100 µg/mL concentrations in comparison with the standard drug tiadinil, which displayed protective 75 and 57% at 500 and 100 µg/mL. The structure **103** marked a higher protective effect than the tiadinil and **102** at 100 µg/mL concentration. The enhanced protective potential of **103** was dose-dependent and due to the presence of hydroxyphenyl moiety. The fluorophenyl, chlorophenyl and hydroxyphenyl functionalities contributed to better efficacy that led to a future investigation to develop promising anti-TMV agents [94].




Figure 10. 1,2,3-Thiadiazole scaffolds **102** and **103** with anti-TMV activity.

1,2,3-Thiadiazole acetanilide hybrid structures prepared by Zhan et al. were screened for their antiviral activity against HIV-1 [95]. In the group of achieved derivatives structure (**104**; **Figure 11**) exhibited moderate antiviral activity (EC_{50} value 0.95 ± 0.33 µM) against HIV in comparison with the reference standard drugs such as: NVP (EC_{50} value 0.208 µM), DLV (EC_{50} value 0.32 µM), EFV (EC_{50} value 0.00440 µM) and zidovudine (EC_{50} value 0.0151 µM). The scaffold **104** displayed remarkable high and excellent anti-HIV-1 activity than all standards. The reason for the highest anti-HIV-1 activity of structural motif **104** among the synthesized congeners was due to the 4-acetyl substituted anilide phenyl ring of 1,2,3-thiadiazole [95].




Figure 11. Antiviral activity of 4-(3,4-dichlorophenyl)-1,2,3-thiadiazole scaffold **104**.

2.2. Anticancer Agents

Cancer has drawn the attention of today's medical science all over the world because it is the lethal, notably complex, most prominent and serious threat to human health. It forms the lump of uncontrolled growth cells called tumors or neoplasm tumor cells. The neoplasm malignant are diversified heterogeneous cells which have properties of rapid proliferation and capability to invade or spread to other parts of the body through the bloodstream and lymphatic system. Most researchers have devoted their extensive research to developing effective anticancer agents, along with the employment and application of integrated surgical procedures, radiation therapy and chemotherapy [96][97]. Great advances and many discoveries in the development of a large quantity of anticancer therapeutics such as carboplatin (**105**) cisplatin (**106**) [98][99][100], anastrozole (**107**) [101][102] and doxorubicin (**108**) (Figure 12), etc., have been made over the past 60 years [103][104][105][106].




Figure 12. Clinical anticancer drugs **105–108**.

Noteworthy are the 4,5-diaryl-1,2,3-thiadiazole derivatives obtained by Wu et al. due to their method of synthesis and anticancer properties [107], and results pointed out that compounds **109** and **110** are the lead compounds by exhibiting favorable activity (**Figure 13**). Growth inhibition rate of tumor cells (obtained from mice S180) by derivative **109** at a dose of 40 mg/kg administrated for 5 days reached 81.0%, while compound **110** displayed a tumor suppression efficacy of 64.2% (at 100 mg/kg of dose administered to mice for five days). These effects were compared with the standard reference CA-4 (combretastatin A-4), which showed a 64.2% antitumor effect at a dose of 40 mg/kg used for 4 days in a mouse model. The SAR study indicated that the reason of the highest antitumor activity of structural motif **109** than reference drug CA-4 and derivative **110** was because of OMe group at C-4 and H at C-3 of 5-phenyl ring of 1,2,3-thiadiazole, while the replacement of hydrogen at C-3 with the nitro group dropped the antitumor activity of derivative **110**. The scaffold **109** could be a promising candidate due to its lower cytotoxicity and excellent antitumor effect [107].




Figure 13. Antitumor effect of 1,2,3-thiadiazoles **109** and **110**.

New 1,2,3-thiadiazole derivatives (**111** and **112**; **Figure 14**) obtained by Hosny et al. via Hurd–Mori cyclization were tested against MCF-7 tumor cells. The IC_{50} values of compounds **111** and **112** (IC_{50} = 12.8 μ g/mL and 8.1 μ g/mL, respectively) were comparable to doxorubicin as the standard anticancer drug with IC_{50} = 3.13 μ g/mL. The derivative **111** showed the highest anti-breast cancer activity than the scaffold **112** and reference drug doxorubicin [108].




Figure 14. Anticancer activity of 2-thioamide-1,2,3-thiadiazole **111** and **112**.

Cikotiene et al. reported the synthetic protocol to achieve the substituted 4,5-diaryl-1,2,3-thiadiazoles as Hsp90 chaperone protein inhibitors. This protein plays a significant part in developing tumor cells, so its inhibition is the main target of many anticancer drugs. The Hsp90 chaperone stabilizes a number of proteins that are essential for the growth of tumors, due to which researchers all over the world are interested in the development of Hsp90 inhibitors which would be investigated as anticancer agents. In the group of tested derivatives, structure **113** deserves special attention as an inhibitor of cancer cells with GI_{50} value of $0.69\ \mu\text{M}$ for U2OS (osteosarcoma) cells and $0.70\ \mu\text{M}$ for HeLa (cervical carcinoma) cells (**Figure 15**) [109]. Isothermal titration calorimetry was used to determine the binding affinity of compound (113) to Hsp90F and Hsp90N. The derivative (**39**) acted as a binder to both Hsp90N and Hsp90F with the observed K_d of about $42\ \text{nM}$ and $37\ \text{nM}$, while for the reference compound 17-AAG [17-(allylamino)-17-demethoxygeldanamycin] K_d was $200\ \text{nM}$ and $240\ \text{nM}$, respectively [109].




Figure 15. Anticancer activity of 1,2,3-thiadiazole derivative **113** against U2OS and HeLa cell lines.

Synthesized by Cui et al., dehydroepiandrosterone derivatives of thiadiazoles were investigated against T47D and HAF cell lines for their antitumor activity applying the sulforhodamine B (SRB) assay. In this group of compounds, derivative **114** (**Figure 16**) focuses particular attention, due to its strong antitumor and antimetastatic activities, in vitro and in vivo. Its IC_{50} value for T47D cells was $0.058 \pm 0.016\ \mu\text{M}$ and for HAF cells $IC_{50} = 21.1 \pm 5.06\ \mu\text{M}$. Moreover, the selectivity of compound **114** ($SI = 364$) was 214 folds better than adriamycin (positive control) (ADM; $SI = 1.7$). The SAR study showed that the introduction of 1,2,3-thiadiazole and D-proline chemical entities in

derivative **114** significantly enhanced antiproliferative activity than the parent DHEA. The SI of scaffold **114** was 214 folds better than SI of ADM standard drugs. The derivative **114** could be used as the promising and novel class of anticancer agent due to its low cytotoxicity, high selectivity index and excellent antitumor and antiproliferative activities (Figure 16) [110].




Figure 16. 1,2,3-Thiadiazole dehydroepiandrosterone derivative **114** as new antitumor agents.

2.3. Insecticidal Agents

Insecticides are the designed substances of either chemical or biological origin that used to regulate the insect behaviour, control insects, cause death, dysfunction and moving away [111][112]. Some of the insecticidal agents, such as tebufenozide (**115**) act as molting hormones primarily against caterpillar pests [113]; imidacloprid (**116**) is the most well-known broad spectrum insecticide, which used as an insect neurotoxin [114][115][116], and neuroactive pymetrozine (**117**) is a novel class of pyridineazomethine insecticide used for the control of aphids and whiteflies in crop fields (Figure 17) [117].




Figure 17. Known insecticidal agents **115–117**.

An interesting example of 1,2,3-thiadiazole derivatives with insecticidal properties are two groups of *N*-*tert*-butyl-*N,N'*-diacylhydrazines obtained by Wang et al. [29] and tested their activity against *Plutella xylostella* L. and *Culex pipiens pallens*. Derivative **118** (Figure 18) displayed remarkable and significantly the highest insecticidal potential (79% mortality at 200 µg/mL) against *Plutella xylostella* L., while **119** exhibited 68% mortality against the same

insects at concentration 200 µg/mL. Insecticidal activity of reference agent, i.e., tebufenozide, was 40%. The scaffold **118** displayed excellent and the highest insecticidal potency, 79% more than the insecticidal potencies of reference drug tebufenozide (40%) and derivative **119** (68%). The EWD group chloro increased the insecticidal activity of scaffold **118**, while ED group methyl decreased the insecticidal potency of scaffold **119** [29].




Figure 18. Insecticidal activity of *N*-tert-butyl-*N'*-diacylhydrazines **118** and **119**.

Zhang et al. designed the (*E*)- β -farnesene based carboxamides of thiadiazoles and checked their aphicidal behavior against *Myzus persicae* [118]. The three compound **120–122** (**Figure 19**) exhibited LC₅₀ values of 33.4 µg/mL, 50.2 µg/mL and 61.8 µg/mL, respectively. The 1,2,3-thiadiazole carboxamide analogues showed significantly higher aphicidal activity than (*E*)- β -farnesene, but lesser insecticidal activity than pymetrozine insecticide (LC₅₀ = 7.1 µg/mL). It was observed that fluoro or difluoro groups on phenyl moiety enhanced the aphicidal potential significantly in novel chemical entities of E β f 1,2,3-thiadiazole **120** and **121** while the methyl substitution led to lower aphicidal activity of scaffold **122** [118].




Figure 19. Insecticide E β f 1,2,3-thiadiazole carboxamides **120–122**.

Pyrazole oxime derivatives, as a new Fenpyroximate analogue bearing a thiadiazole moiety, were synthesized by Dai et al. and then their insecticidal, acaricidal and cytotoxic activities were examined. The best insecticidal properties possessed oximes **123** and **124** (**Figure 20**) against *Aphis craccivora* at 100 µg/mL with 90% of the mortality rate. The structure–activity relationship study revealed that due to the insertion of F- and Me-groups at *p*-position of phenoxy moiety attached to the pyrazole oxime **123** and **124** displayed good insecticidal activities but slightly lower than commercial drug imidacloprid [119].




Figure 20. Ethers of pyrazole oximes **123** and **124** with insecticidal properties.

Another interesting example of insecticidal compounds are hybrids of substituted triazole and 1,2,3-thiadiazole scaffold that were investigated for their activity against *Aphis laburni* and TMV by Li et al. [120]. Preliminary bioassays indicated compounds **125** and **126** (Figure 21) as lead compounds by depicting enhanced activity against *Aphis laburni* at 100mg/mL (mortality $\geq 95\%$) while the derivative **127** represent the least active compound (due to CF_3 substituted triazole ring) that displayed only 25.25% mortality. The structure–activity study indicated that scaffolds with aromatic moieties attached to the carbonyl carbon of triazole ring have higher insecticidal activity than scaffolds with alkyl moieties [120].




Figure 21. Substituted 4-methyl-1,2,3-thiadiazole derivatives with insecticidal properties against *Aphis laburni*.

2.4. Amoebicidal Agents

A parasitic disease caused by a protozoa *Entamoeba histolytica* is known as amoebiasis. *E. histolytica* has infected 40 to 50 million people worldwide and led to the development of amoebic colitis or extraintestinal abscess that has resulted in 100,000 deaths annually [121][122][123]. The tissue amoebiasis is treated by different drugs such as metronidazole, chloroquine, tinidazole, dehydroemetine and nitazoxanide, while the diloxanide furoate and iodoquinoline drugs are used to treat luminal infections (Figure 22) [124][125][126][127][128].



Figure 22. Amoebicidal commercial drugs **128–130**.

Hayat et al. reported the synthetic route to achieve substituted 1,2,3-thiadiazole scaffolds from the cyclization of quinoline-based hydrazone. All substituted 1,2,3-thiadiazole hybrids and quinoline-based hydrazones were evaluated for antiamoebic activity against HM1 and IMSS strains of *E. histolytica*. The scaffold 4-bromo phenyl-1,2,3-thiadiazole (**131**, **Figure 23**) and furan based 1,2,3-thiadiazole (**132**, **Figure 23**) exhibited potent antiamoebic activity ($IC_{50} = 0.24 \mu M$ and $IC_{50} = 0.23 \mu M$, respectively) when compared with metronidazole ($IC_{50} = 1.80 \mu M$). The parent intermediate quinoline-based hydrazone displayed better antiamoebic potential than the targeted derivatives of 1,2,3-thiadiazoles. The SAR study revealed that the significant loss of antiamoebic potential in scaffolds **131** and **132** was due to the absence of quinoline acetohydrazone moiety in final thiadiazoles [129].




Figure 23. The most active antiamoebic 1,2,3-thiadiazoles **131** and **132**.

2.5. Plant Activator Agents

Plant activators are synthetic or natural chemicals entities that protect plants from different pathogens. Among them, pyrimidin-type plant activator 2 (PPA2 **133**; **Figure 24**) significantly enhanced plant defense system against bacterial infections unlikely traditional commercially available plant activators (INA **134**, BABA **135**, BHT **7**; **Figure 24**). Plant activators not only showed their therapeutic potential against wide variety of pathogens but also increased growth of plants, rate of photosynthesis and overall yield of crops [130][131][132][133][134].




Figure 24. Structures of commercially available plant activators **133**–**135**.

Du et al. reported a synthetic pathway for the preparation of carboxylate derivatives of thiadiazoles as potential plant activators against seven plant diseases [135]. The 1,2,3-thiadiazole derivatives **136** and **137** (Figure 25) proved to be good plant activators and displayed excellent inhibition activities against diseases such as *M. melonis*, (90% for **136** and 69% for **137**), *C. cassicola* (77% for **136** and 52% for **137**), *P. syringae* pv. *Lachrymans* (42% for **136** and 42% for **137**) and *P. infestans* (81% for **136** and 67% for **137**) when results were compared with commercial BTH plant activator. The structural motifs **136** and **137** displayed significantly better therapeutic efficacy than BHT commercial drug, but the scaffold **136** had better plant activating potency than scaffold **137** [136].




Figure 25. Potential new plant activator with thieno [2,3-*d*]-1,2,3-thiadiazoles **136** and **137**.

2.6. Fungicidal Agents

The chemical agents that have the potential to control and eradicate fungal infections are termed as fungicides or antifungal drugs [137]. The commercial antifungal agents are chlorothalonil (**65**) which is a non-systemic fungicide [138][139], fluconazole (**66**) [140] and ketoconazole (**67**) (Figure 26) [141].




Figure 26. Structures of commercial antifungal drugs **138–140**.

As the first example, it is worth bring up oxadiazoles containing thiadiazole derivatives [\[142\]](#). In this group, two oxadiazole derivatives **141** and **142** (**Figure 27**) displayed the best antifungal activity against *Puccinia triticina* at a concentration of 500 µg/mL. Both structures showed a remarkable high growth inhibition activity (98 and 83%, respectively) which was slighter lower than the standard drug chlorothalonil with 100% fungal growth inhibition activity. The remarkable high and excellent antifungal activity of scaffold **141** could lead to the development of novel oxadiazole-based 1,2,3-thiadiazole fungicides [\[142\]](#).




Figure 27. Novel potent 1,2,3-thiadiazole fungicides **141** and **142**.

Another example of compounds with good antifungal activities are new structures designed by Sun et al., which are 1,2,4-triazole derivatives substituted with a 1,2,3-thiadiazole ring [\[143\]](#). One of the synthesized compounds **143** (**Figure 28**), exhibited remarkably high fungicidal 93.19% against *Corynespora cassicola*, while reference compounds iprodione, validamycin and topsin-M possessed 78.20, 53.52 and 78.75% antifungal activity, respectively. The authors declared that compound containing phenyl group on triazole ring **143** exhibited higher activity, however, the results were different in the case of *Pseudoperonospora cubensis* as compound **144** containing cyclopropyl group displayed higher activity than that of **143**. The most active compound with regard to *Pseudoperonospora cubensis* was **144**, which inhibited the growth of this pathogen in 81.62%. However, SAR analysis did not clear the results against *pseudomonas syringae* pv. *Lachrymans*. Both the scaffolds showed excellent and the highest antifungal potential than all three commercial reference drugs that will lead to the development of promising antifungal agents [\[143\]](#).




Figure 28. Antifungal activity of triazole moiety containing 1,2,3-thiadiazolew **143** and **144**.

Less antifungal activity was observed in the derivative **145** (Figure 29) reported by Tan et al. [144]. Its inhibition rates to *Phytophthora infestans* (Mont) de Bary, *Alternaria solani*, *Gibberella sanbinetti*, *Physalospora piricola* Nose, *Botrytis cinerea*, *Phytophthora capsici* Leonian, *Cercospora arachidicola*, *Rhizoctonia solanii* and *Fusarium oxysporum* reached 32.3, 17.4, 32.3, 25.8, 43.5%, 36.1, 15.8, 30.3 and 26.9% at 50 µg/mL respectively. The results indicated that the structural motif **145** proved to be broad-spectrum fungicide because it exhibited significant antifungal activity against different fungal strains [144].




Figure 29. Piperidine based 1,2,3-Thiadiazole **145** with moderate antifungal activity.

Another group reported the fungicidal properties of 1,2,3-thiadiazole derivatives against *Alternaria solani* (AS), *Botrytis cinerea* (BC), *Cercospora arachidicola* (CA), *Cercospora beticola* (CB), *Colletotrichum lagenarium* (CL), *Fusarium oxysporum* (FO), *Gibberella zeae* (GZ), *Macrophoma kuwatsukai* (MK), *Phytophthora infestans*(Mont) de Bary (PI), *Physalospora piricola* (PP), *Pellicularia sasakii* (Shirai) (PS), *Puccinia triticina* Eriks (PT) and *Rhizoctonia solani* Kuhn (RS) [145]. The most active occurred to be structure **146** (**Figure 30**) showed potent fungicidal inhibitory activity 84.8, 83.9, 83.3, 78.9, 75.0, 71.4, 70.7, 66.7, 64.7 and 63.6% against PS, PP, RS, PT, FO, CB, PI, AS, CL and CA, respectively. On the other hand, the propyl containing fused 1,2,4-triazolo[1,3,4]thiadiazole **147** (**Figure 30**) exhibited the best fungicidal inhibition activity 93.0% for FO, 84.9% for BC, 77.8% for PS, 75.8% for PI, 75.0% for GZ, 62.1% for CA, 50.0% for CL, 40.5% for PP, 34.4% for AS and 33.3% for CB. The results of the antifungal study showed that fused 1,2,4-triazolo[1,3,4]thiadiazole led to the development of wide-spectrum antifungal agents [145].




Figure 30. 1,2,3-thiadiazoles with broad antifungal activity.

Wang et al. reported a synthetic approach to new series of organotin 4-methyl-1,2,3-thiadiazole-5-carboxylates and benzo[1,2,3]thiadiazole-7-carboxylates [146] and antifungal properties against *P. piricola* and *Gibberella zeae*. The triethyltin-based 1,2,3-thiadiazole carboxylate analogue **148** (**Figure 31**) displayed antifungal efficacy EC₅₀ = 0.12 µg/mL and 0.16 µg/mL against *P. piricola*, and *Gibberella zeae*, respectively. The overall results of the present study indicated the antifungal effectiveness of organotin-based benzo-1,2,3-thiadiazole or 1,2,3-thiadiazoles analogues proved to be broad-spectrum fungicides [146].




Figure 31. Antifungal benzo[1,2,3]thiadiazole-7-carboxylate derivative **148**.

Antifungal properties have also been determined for carboxamide derivatives of thiadiazoles obtained by Zuo et al. utilized one-pot four-component Ugi reaction strategy under green synthetic conditions [147]. All the synthesized analogues were investigated against eleven fungal strains to develop novel fungicidal candidates. The 1,2,3-thiadiazole containing carboxamide moiety **149** (Figure 32) displayed broad-spectrum fungicidal inhibition activities against CA 71%, AS 100%, GZ 62%, PP 85%, PG 14%, PS, 74%, CL 95%, PI 88% and RS 97%. Similarly good antifungal activity was observed for scaffold **150** that exhibited fungicidal inhibition against various fungal strains such as: CB (*Cercospora beticola*) 67%, CA 79%, AS 44%, GZ 34%, PP 53%, PG 57%, PS 16%, CL 43%, RS 51% and PI 57%. The scaffold **149** displayed maximum fungicidal potential 100, 97 and 95% against fungal strains AS, RS and CL, respectively, while the zero percent antifungal activity was shown against CB and FO fungal strains, just as scaffold **150** displayed the best antifungal potential 67% against CB fungal strains and 0% against FO [147].




Figure 32. The most active 4-methyl-1,2,3-thiadiazole-5-carboxamides.

References

1. Irfan, A.; Batool, F.; Ahmad, S.; Ullah, R.; Sultan, A.; Sattar, R.; Nisar, B.; Rubab, L. Recent trends in the synthesis of 1,2,3-thiadiazoles. *Phosphorus Sulfur Silicon Relat. Elem.* 2019, 194, 1098–1115.
2. Singh, A.K.; Parthsarthy, R.; Kshitiz, J.; Mishra, G. Synthesis, characterization and antibacterial activity of 1,3,4-thiadiazole derivatives. *Int. J. Sci. Innov. Discov.* 2011, 3, 353–361.
3. Siddiqui, N.; Ahuja, P.; Ahsan, W.; Pandeya, S.N.; Alam, S.M. Thiadiazoles: Progress report on biological activities. *J. Chem. Pharm. Res.* 2009, 1, 19–30.
4. Jalhan, S.; Jindal, A.; Gupta, A.; Hemraj, H. Synthesis, biological activities and chemistry of thiadiazole derivatives and schiff bases. *Asian J. Pharm. Clin. Res.* 2012, 3, 199–208.
5. Srivastava, S.; Prasad, K.R.; Saini, R. Thiadiazole: A brief review. *WJPPS* 2014, 3, 1198–1212.
6. Paulrasu, K.; Duraikannu, A.; Palrasu, M.; Shanmugasundaram, A.; Kuppusamy, M.; Thirunavukkarasu, B. Synthesis of 4-methyl-N'-(3-alkyl-2r,6cdiaryl)piperidin-4-ylidene)-1,2,3-thiadiazole-5-carbohydrazides with antioxidant, antitumor and antimicrobial activities. *Org. Biomol. Chem.* 2014, 12, 911–921.
7. Kanakaraju, S.; Suresh, L. Design, synthesis, in vitro antimicrobial and cytotoxic evaluation of novel 1,2,3-selena/thiadiazolyltetrazole derivatives. *RSC Adv.* 2015, 5, 29325–29334.
8. Dogan, H.N.; Duran, A.; Rollas, S.; Sener, G.; Uysal, M.K.; Gulen, D. Synthesis of new 2,5-disubstituted-1,3,4-thiadiazoles and preliminary evaluation of anticonvulsant and antimicrobial activities. *Bioorg. Med. Chem.* 2002, 10, 2893–2898.

9. Hussain, S.; Sharma, J.; Amir, M. Synthesis and Antimicrobial activities of 1,2,4-triazole and 1,3,4-thiadiazole derivatives of 5-amino-2-hydroxybenzoic acid. *E J. Chem.* 2008, 5, 963–968.
10. Demirbas, A.; Sahin, D.; Demirbas, N.; Karaoglu, A.S. Synthesis of some new 1,3,4-thiadiazol-2-ylmethyl-1,2,4-triazole derivatives and investigation of their antimicrobial activities. *Eur. J. Med. Chem.* 2009, 44, 2896–2903.
11. Farshori, N.N.; Banday, M.R.; Ahmad, A.; Khan, A.U.; Rauf, A. Synthesis, characterization and in vitro antimicrobial activities of 5-alkenyl/hydroxyalkenyl-2-phenylamine-1,3,4-oxadiazoles and thiadiazoles. *Bioorg. Med. Chem. Lett.* 2010, 20, 1933–1938.
12. Almajan, G.L.; Innocenti, A.; Puccetti, L.; Manole, G.; Barbuceanu, S.; Saramet, I.; Scozzafava, A.; Supuran, C.T. Carbonic anhydrase inhibitors. Inhibition of the cytosolic and tumor-associated carbonic anhydrase isozymes I, II, and IX with a series of 1,3,4-thiadiazole- and 1,2,4-triazole-thiols. *Bioorg. Med. Chem. Lett.* 2005, 15, 2347–2352.
13. Abdel-Hamid, M.K.; Abdel-Hafez, A.A.; El-Koussi, N.A.; Mahfouz, N.M.; Innocenti, A.; Supuran, C.T. Design, synthesis, and docking studies of new 1,3,4-thiadiazole-2-thione derivatives with carbonic anhydrase inhibitory activity. *Bioorg. Med. Chem.* 2007, 15, 6975–6984.
14. Dudutien, V.; Baranauskien, L.; Matulis, D. Benzimidazo[1,2-c][1,2,3]thiadiazole-7-sulfonamides as inhibitors of carbonic anhydrase. *Bioorg. Med. Chem. Lett.* 2007, 17, 3335–3338.
15. Jalilian, A.R.; Sattari, S.; Bineshmarvasti, M.; Daneshtalab, M.; Shafiee, A. Synthesis and in vitro antifungal and cytotoxicity evaluation of substituted 4,5-dihydronaphtho[1,2-d][1,2,3]thia(orselena)diazoles. *Il Farmaco* 2003, 58, 63–68.
16. Chu, C.H.; Hui, X.P.; Xu, P.F.; Zhang, Z.Y.; Li, Z.C.; Liao, R.A. Synthesis and antifungal activities of ω -(5-aryl amino-1,3,4-thiadiazol-2-thio)- ω -(1H-1,2,4-triazol-1-yl)acetophenones. *Indian J. Chem.* 2002, 41B, 2436–2438.
17. Moawad, E.B.; Yousif, M.Y.; Metwally, M.A. Synthesis of certain heteroaryl-fused pyrimidines and pyridines and seleno- and thia-diazoles with naphthyl substituent as potential antifungal agents. *Pharmazie* 1989, 44, 820–822.
18. Yadav, L.S.; Zaidi, M.G.H.; Singh, B.N. Synthesis and fungicidal evaluation of some 1,3,4-thiadiazoles against *P. oryzae* and *R. solani*. *Asian J. Chem.* 2003, 15, 1805–1807.
19. Zou, X.-J.; Lai, L.H.; Jin, G.Y.; Zhang, Z.X. Synthesis, fungicidal activity, and 3D-QSAR of pyridazinone-substituted 1,3,4-oxadiazoles and 1,3,4-thiadiazoles. *J. Agric. Food. Chem.* 2002, 50, 3757–3760.
20. Balasankar, T.; Gopalakrishnan, M.; Nagarajan, S. Synthesis and antibacterial activity of some 5-(4-biphenyl)-7-aryl[3,4-d][1,2,3]-benzothiadiazoles. *Eur. J. Med. Chem.* 2005, 40, 728–731.

21. Padmavathi, V.; Mahesh, K.; Mohan, N.V.A.; Mohan, N.; Padmaja, A. Synthesis and bioassay of oxazolyl/thiazolyl selenadiazoles, thiadiazoles and diazaphospholes. *Chem. Pharm. Bull.* 2009, 57, 561–566.
22. Taha, S.S.; Ahmad, A.A.; Mawlood, S.I.; Ali, N.O. Synthesis of some series of 2-amino-1,3,4-thiadiazole derivatives with their pathogenic bacterial activity. *J. Raparin Univ.* 2017, 4, 63–78.
23. Othman, A.A.; Kihel, M.; Amara, S. 1,3,4-Oxadiazole, 1,3,4-thiadiazole and 1,2,4-triazole derivatives as potential antibacterial agents. *Arab. J. Chem.* 2014, 12, 1660–1675.
24. Paruch, K.; Popiołek, Ł.; Biernasiuk, A.; Berecka-Rycerz, A.; Malm, A.; Gumieniczek, A.; Wujec, M. Novel derivatives of 4-methyl-1,2,3-thiadiazole-5-carboxylic acid hydrazide: Synthesis, lipophilicity, and in vitro antimicrobial activity screening. *Appl. Sci.* 2021, 11, 1180.
25. Camoutsis, C.; Geronikaki, A.; Ciric, A.; Sokovic, M.; Zoumpoulakis, P.; Zervou, M. Sulfonamide-1,2,4-thiadiazole derivatives as antifungal and antibacterial agents: Synthesis, biological evaluation, lipophilicity and conformational studies. *Chem. Pharm. Bull.* 2010, 58, 160–167.
26. Yasuda, M.; Kusajima, M.; Nakajima, M.; Akutsu, K.; Kudo, T.; Yoshida, S.; Nakashita, H. Thiadiazole carboxylic acid moiety of tiadinil, SV-03, induces systemic acquired resistance in tobacco without salicylic acid accumulation. *J. Pest. Sci.* 2006, 31, 329–334.
27. Xu, Y.F.; Zhao, Z.J.; Qian, X.H.; Qian, Z.; Tian, W.; Zhong, J. Novel, unnatural benzo-1,2,3-thiadiazole-7-carboxylate elicitors of taxoid biosynthesis. *J. Agric. Food Chem.* 2006, 54, 8793–8798.
28. Nombela, G.; Pascual, S.; Aviles, M.; Guillard, E.; Muñizet, M. Benzothiadiazole induces local resistance to *Bemisia tabaci* (Hemiptera:Aleyrodidae) in tomato plants. *J. Econ. Entomol.* 2005, 98, 2266–2271.
29. Wang, H.; Yang, Z.; Fan, Z.; Wu, Q.; Zhang, Y.; Mi, N.; Wang, S.; Zhang, Z.; Song, H.; Liu, F. Synthesis and insecticidal activity of N-tert-314 butyl-N,N0-diacylhydrazines containing 1,2,3-thiadiazoles. *J. Agric. Food Chem.* 2011, 59, 628–634.
30. Bloom, J.D.; DiGrandi, M.J.; Dushin, R.G.; Curran, K.J.; Ross, A.A.; Norton, E.B.; Terefenko, E.; Jones, T.R.; Feld, B.; Lang, S.A. Thiourea inhibitors of herpes viruses. Part 1: Bis-(aryl)thiourea inhibitors of CMV. *Bioorg. Med. Chem. Lett.* 2003, 13, 2929–2932.
31. Bloom, J.D.; Dushin, R.G.; Curran, K.J.; Donahue, F.; Norton, E.B.; Terefenko, E.; Jones, T.R.; Ross, A.A.; Feld, B.; Lang, S.A.; et al. Thiourea inhibitors of herpes viruses. Part 2: N-Benzyl-N0-arylthiourea inhibitors of CMV. *Bioorg. Med. Chem. Lett.* 2004, 14, 3401–3406.
32. Pannecouque, C.; Szafarowicz, B.; Volkova, N.; Bakulev, V.; Dehaen, W.; Mely, Y.; Daelemans, D. N,N'-Bis(1,2,3-thiadiazol-5-yl)benzene-1,2-diamine targets the HIV-1 retroviral nucleocapsid zinc fingers. *Antimicrob. Agents Chemother.* 2010, 54, 1461–1468.

33. Tatar, E.; Kucukguzel, S.G.; Karakus, S.; Clercq, E.D.; Andrei, G.; Snoeck, R.; Pannecouque, C.; Okullu, S.O.; Ünübol, N.; Kocagöz, T.; et al. Synthesis and biological evaluation of some new 1,3,4-thiadiazole and 1,2,4-triazole derivatives from L-methionine as antituberculosis and antiviral agents. *Marmara Pharm. J.* 2015, 19, 88–102.
34. Azaam, M.M.; Kenawy, E.; El-Din, A.S.B.; Khamis, A.A.; El-Magdd, M.A. Antioxidant and Anticancer Activities of α -Aminophosphonates Containing Thiadiazole. *J. Saudi Chem. Soc.* 2018, 22, 34–41.
35. Cressier, D.; Prouillac, C.; Hernandez, P.; Amourette, C.; Diserbo, M.; Lion, C.; Rima, G. Synthesis, antioxidant properties and radioprotective effects of new benzothiazoles and thiadiazoles. *Bioorg. Med. Chem.* 2009, 17, 5275–5284.
36. Kaur, H.; Kumar, S.; Vishwakarma, P.; Sharma, M.; Saxena, K.K.; Kumar, A. Synthesis and antipsychotic and anticonvulsant activity of some new substituted oxa/thiadiazolyl azetidinonyl/thiazolidinonyl carbazoles. *Eur. J. Med. Chem.* 2010, 45, 2777–2783.
37. Siddiqui, N.; Ahuja, P.; Ahsan, W.; Pandeya, S.N.; Alam, S.M. Synthesis of 3-arylamino-4-aryl-5-(N-arylthiocarbonylimino)-4,5-dihydro-1,2,4-thiadiazoles as anticonvulsant agents. *Indian J. Heterocycl. Chem.* 2004, 14, 159–160.
38. Siddiqui, N.; Arshad, M.F.; Khan, S.A.; Ahsan, W. Synthesis, anticonvulsant and neurotoxicity screening of 1-(substituted phenyl)-3-[(5-substituted phenyl)-1,3,4-thiadiazole-2-yl]-2-thioxodihydro pyrimidine-4,6 (1H, 5H)-diones. *J. Pharm. Res.* 2008, 7, 122–125.
39. Siddiqui, N.; Ahsan, W. Synthesis, anticonvulsant and toxicity screening of thiazolyl-thiadiazole derivatives. *Med. Chem. Res.* 2011, 20, 261–268.
40. Mullick, P.; Khan, S.A.; Verma, S.; Alam, O. Thiadiazole derivatives as potential anticonvulsant agents. *B Korean Chem. Soc.* 2011, 32, 1011–1016.
41. Chapleo, C.B.; Myers, M.; Myers, P.L.; Saville, J.F.; Smith, A.C.B.; Stillings, M.R.; Tulloch, J.F.; Walter, D.S.; Welbourn, A.P. Substituted 1,3,4-thiadiazoles with anticonvulsant activity. 1-Hydrazines. *J. Med. Chem.* 1986, 29, 2273–2280.
42. Rajak, H.; Behera, C.K.; Pawar, R.S.; Singour, P.K.; Kharya, M.D. Synthesis and anticonvulsant evaluation of some novel 2,5-disubstituted 1,3,4-thiadiazoles: Pharmacophore model studies. *Acta Pol. Pharm.* 2010, 67, 503–510.
43. Samel, A.B.; Pai, N.R. Synthesis of novel aryloxypropanoylthiadiazoles as potential antihypertensive agents. *Chin. Chem. Soc. TAIP* 2010, 57, 1327–1330.
44. Turner, S.; Myers, M.; Gadie, B.; Nelson, A.J.; Pape, R.; Saville, J.F.; Doxey, J.C.; Berridge, T.L. Antihypertensive thiadiazoles 1. Synthesis of some 2-aryl-5-hydrazino-1,3,4- thiadiazole with vasodilator activity. *J. Med. Chem.* 1988, 31, 902–906.

45. Wei, M.-X.; Feng, L.; Li, X.-Q.; Zhou, X.-Z.; Shao, Z.-H. Synthesis of new chiral 2,5-disubstituted 1,3,4-thiadiazoles possessing γ -butenolide moiety and preliminary evaluation of in vitro anticancer activity. *Eur. J. Med. Chem.* 2009, 44, 3340–3344.
46. Tripathy, R.; Ghose, A.; Singh, J.; Bacon, E.R.; Angeles, T.S.; Yang, S.X.; Albom, M.S.; Aimone, L.D.; Herman, J.L.; Mallamo, J.P. 1,2,3-Thiadiazole substituted pyrazolones as potent KDR/VEGFR-2 kinase inhibitors. *Bioorg. Med. Chem. Lett.* 2007, 17, 1793–1798.
47. Ibrahim, D.A. Synthesis and biological evaluation of 3,6-disubstituted [1,2,4]triazolo[3,4-b][1,3,4]thiadiazole derivatives as a novel class of potential anti-tumor agents. *Eur. J. Med. Chem.* 2009, 44, 776–2781.
48. Padmavathi, V.; Reddy, G.S.; Padmaja, A.; Kondaiah, P.P.; Shazia, A. Synthesis, antimicrobial and cytotoxic activities of 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazoles. *Eur. J. Med. Chem.* 2009, 44, 2106–2112.
49. Karakuü, S.; Çoruh, U.; Barlas-Durgun, B.; Vázquez-López, E.M.; Özbaü-Turan, S.; Akbuüa, J.; Rollas, S. Synthesis and cytotoxic activity of some 1,2,4-triazoline-3-thione and 2,5-disubstituted-1,3,4-thiadiazole derivatives. *Marmara Pharm. J.* 2010, 14, 84–90.
50. Holla, B.S.; Poorjary, K.N.; Rao, B.S.; Shivananda, M.K. New bis-aminomercaptotriazoles and bis-triazolothiadiazoles as possible anti-cancer agents. *Eur. J. Med. Chem.* 2002, 37, 511–517.
51. Upadhyay, P.K.; Mishra, P. Synthesis, antimicrobial and anticancer activities of 5-(4-substituted phenyl)-1,3,4-thiadiazole-2-amines. *Rasayan J. Chem.* 2017, 10, 254–262.
52. Schenone, S.; Brullo, C.; Bruno, O.; Bondavalli, F.; Ranise, A.; Filippelli, W.; Rinaldi, B.; Capuano, A.; Falcone, G. New 1,3,4-thiadiazole derivatives endowed with analgesic and anti-inflammatory activities. *Bioorg. Med. Chem.* 2006, 14, 1698–1705.
53. Rostom, S.A.F.; El-Ashmawy, I.M.; el Razik, H.A.A.; Badr, M.H.; Ashour, H.M.A. Design and synthesis of some thiazolyl and thiadiazolyl derivatives of antipyrine as potential non-acidic anti-inflammatory, analgesic and antimicrobial agents. *Bioorg. Med. Chem.* 2009, 17, 882–895.
54. Labanauskas, L.; Kalcas, V.; Udrenaite, E.; Gaidelis, P.; Brukstus, A.; Dauksas, A. Synthesis of 3-(3,4-dimethoxyphenyl)-1H-1,2,4-triazole-5-thiol and 2-amino-5-(3,4-dimethoxyphenyl)-1,3,4-thiadiazole derivatives exhibiting anti-inflammatory activity. *Pharmazie* 2001, 56, 617–619.
55. Kadi, A.A.; El-Brolossy, N.R.; Al-Deeb, O.A.; Habib, E.E.; Ibrahim, T.M.; El-Emam, A.A. Synthesis, antimicrobial, and anti-inflammatory activities of novel 2-(1-adamantyl)-5-substituted-1,3,4-oxadiazoles and 2-(1-adamantylamino)-5-substituted-1,3,4-thiadiazoles. *Eur. J. Med. Chem.* 2007, 42, 235–242.
56. Kadi, A.A.; Al-Abdullah, E.S.; Shehata, I.A.; Habib, E.E.; Ibrahim, T.M.; El-Emam, A.A. Synthesis, antimicrobial and anti-inflammatory activities of novel 5-(1-adamantyl)-1,3,4-thiadiazole derivatives. *Eur. J. Med. Chem.* 2010, 45, 5006–5011.

57. Bekhit, A.A.; Ashour, H.M.A.; Ghany, Y.S.A.; Bekhit, A.E.A.; Baraka, A. Synthesis and biological evaluation of some thiazolyl and thiadiazolyl derivatives of 1H-pyrazole as anti-inflammatory antimicrobial agents. *Eur. J. Med. Chem.* 2008, 43, 456–463.
58. Kamal, M.D.; Gawad, H.A.; Rageb, E.A.; Ellithey, M.; Mohamed, H.A. Synthesis, anticonvulsant, and anti-inflammatory evaluation of some new benzotriazole and benzofuran-based heterocycles. *Bioorg. Med. Chem.* 2006, 14, 3672–3680.
59. Du, Q.S.; Zhu, W.P.; Zhao, Z.J.; Qian, X.H.; Xu, Y.F. Novel benzo-1,2,3-thiadiazole-7-carboxylate derivatives as plant activators and the development of their agricultural applications. *J. Agric. Food Chem.* 2012, 60, 346–353.
60. Stanetty, P.; Kremslehner, M.; Veollenkle, H. A new type of plant activator: Synthesis of thieno[2,3-d][1,2,3]-thiadiazole-6-carboxylic acid derivatives via Hurd-Mori cyclization. *J. Chem. Soc. Perkin Trans.* 1998, 1, 853–856.
61. Pattan, S.R.; Kekare, P.; Dighe, N.S.; Nirmal, S.A.; Musmade, D.S.; Parjane, S.K.; Daithankar, A.V. Synthesis and biological evaluation of some 1,3,4-thiadiazoles. *J. Chem. Pharm. Res.* 2009, 1, 191–198.
62. Ram, V.J.; Goel, A.; Kandpal, M. Tetraazaacenaphthene, tetraazaphenalene and 1,3,4-thiadiazole derivatives as potential leishmanicides. *Bioorg. Med. Chem. Lett.* 1997, 7, 651–656.
63. Silva, E.F.; Canto-Cavalheiro, M.M.; Braz, V.R.; Cysne-Finkelstein, L.; Leon, L.L.; Echevarria, A. Synthesis, and biological evaluation of new 1,3,4-thiadiazolium-2-phenylaminenderivatives against *Leishmania amazonensis* promastigotes and amastigotes. *Eur. J. Med. Chem.* 2002, 37, 979–984.
64. Behrouzi-Fardmoghadam, M.; Poorrajab, F.; Ardestani, S.K.; Emami, S.; Shafiee, A.; Foroumadi, A. Synthesis and in vitro anti-leishmanial activity of 1-[5-(5-nitrofuranyl)-1,3,4-thiadiazol-2-yl]- and 1-[5-(5-nitrothiophen-2-yl)-1,3,4-thiadiazol-2-yl]-4-aryl piperazines. *Bioorg. Med. Chem.* 2008, 16, 4509–4515.
65. Foroumadi, A.; Emami, S.; Pournourmohammadi, S.; Kharazmi, A.; Shafiee, A. Synthesis and in vitro leishmanicidal activity of 2-(1-methyl-5-nitro-1H-imidazol-2-yl)-5-substituted-1,3,4-thiadiazole derivatives. *Eur. J. Med. Chem.* 2005, 40, 1346–1350.
66. Oruc, E.E.; Rollas, S.; Kandermirli, S.; Shvets, N.; Dimoglo, A.S. 1,3,4-thiadiazole derivatives. Synthesis, structure elucidation, and structure-antituberculosis activity relationship investigation. *J. Med. Chem.* 2004, 47, 6760–6767.
67. Karakus, S.; Rollas, S. Synthesis and antituberculosis activity of new N-phenyl-N'-[4-(5-alkyl/aryl-amino-1,3,4-thiadiazole-2-yl)phenyl]thioureas. *Farmaco* 2002, 57, 577–581.
68. Solak, N.; Rollas, S. Synthesis and antituberculosis activity of 2-(aryl/alkyl-amino)-5-(4-aminophenyl)-1,3,4-thiadiazoles and their Schiff bases. *ARKIVOC* 2006, 12, 173–181.

69. Patole, J.; Shingnapurkar, D.; Padhye, S.; Ratledge, C. Schiff base conjugates of p-aminosalicylic acid as antimycobacterial agents. *Bioorg. Med. Chem. Lett.* 2006, 16, 1514–1517.
70. Foroumadi, A.; Kargar, Z.; Sakhteman, A.; Sharifzadeh, Z.; Mohammadi, R.F.; Kazemi, M.; Shafiee, A. Synthesis and antimycobacterial activity of some alkyl [5-(nitroaryl)-1,3,4-thiadiazol-2-ylthio]propionates. *Bioorg. Med. Chem. Lett.* 2006, 16, 1164–1167.
71. Tahghighi, A.; Babalouei, F. Thiadiazoles: The appropriate pharmacological scaffolds with leishmanicidal and antimalarial activities: A review. *Iran. J. Basic Med. Sci.* 2017, 20, 613–622.
72. Gudala, S.; Ambati, S.R.; Patel, J.L.; Vedula, R.R.; Penta, S. An Efficient Synthesis of Pyrazolyl-1,2,3-thiadiazoles via Hurd-Mori Reaction. *J. Heterocycl. Chem.* 2019, 56, 2163–2169.
73. Jain, K.A.; Sharma, S.; Vaidya, A.; Ravichandran, V.; Agrawal, R.K. 1,3,4-Thiadiazole and its Derivatives: A Review on Recent Progress in Biological Activities. *Chem. Biol. Drug Des.* 2013, 81, 557–576.
74. Supuran, C.T.; Scozzafava, A. Carbonic anhydrase inhibitors. *Curr. Med. Chem. Immunol. Endocrinol. Metab. Agents* 2001, 1, 61–97.
75. Georgeta, S.; Oana, S.; Eugenia, S.; Santa, B. 2-Amino-1,3,4-thiadiazole as a potential scaffold for promising antimicrobial agents. *Drug Des. Devel. Ther.* 2018, 12, 1545–1566.
76. Iizawa, Y.; Okonogi, K.; Hayashi, R.; Iwahi, T.; Yamazaki, T.; Imada, A. Therapeutic effect of cefozopran (SCE-2787), a new parenteral cephalosporin, against experimental infections in mice. *Antimicrob. Agents Chemother.* 1993, 37, 100–105.
77. Sharma, B.; Verma, A.; Prajapati, S.; Sharma, K.U. Synthetic methods, chemistry, and the anticonvulsant activity of thiadiazoles. *Int. J. Med. Chem.* 2013, 2013.
78. Pawar, P.; Chawla, A.; Sharma, P.; Arora, S. Novel drug delivery approaches on antiviral and antiretroviral agents. *J. Adv. Pharm. Technol. Res.* 2012, 3, 147–159.
79. Rossignol, J.F. Nitazoxanide: A first-in-class broad-spectrum antiviral agent. *Antivir. Res.* 2014, 110, 94–103.
80. Raymund, R.; Razonable, M.D. Antiviral Drugs for Viruses Other Than Human Immunodeficiency Virus. *Mayo Clin. Proc.* 2011, 86, 1009–1026.
81. Nystrom, K.; Waldenstrom, J.; Tang, K.-W.; Lagging, M. Ribavirin: Pharmacology, multiple modes of action and possible future perspectives. *Future Virol.* 2019, 14, 153–160.
82. Sidwell, R.W.; Huffman, J.H.; Khare, G.P.; Allen, L.B.; Witkowski, J.T.; Robins, R.K. Broad-spectrum antiviral activity of virazole: 1-beta-D-ribofuranosyl-1,2,4-triazole-3-carboxamide. *J. Sci.* 1972, 177, 705–706.

83. Rakhmanina, N.Y.; Vanda, V.J.N. Efavirenz in the therapy of HIV infection. *Expert Opin. Drug Metab. Toxicol.* 2009, 6, 95–103.
84. Larru, B.; Eby, J.; Lowenthal, E.D. Antiretroviral treatment in HIV-1 infected pediatric patients: Focus on efavirenz. *Pediatric Health Med. Ther.* 2014, 2014, 29–42.
85. Kryst, J.; Kawalec, P.; Pilc, A. Efavirenz-Based Regimens in Antiretroviral-Naive HIV-Infected Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *PLoS ONE* 2015, 10, e0124279.
86. Leneva, I.A.; Russell, R.J.; Boriskin, Y.S.; Hay, A.J. Characteristics of arbidol-resistant mutants of influenza virus: Implications for the mechanism of anti-influenza action of arbidol. *Antivir. Res.* 2009, 81, 132–140.
87. Liu, Q.; Xiong, H.; Lu, L.; Liu, Y.; Luo, F.; Hou, W.; Yang, Z. Antiviral and anti-inflammatory activity of arbidol hydrochloride in influenza A (H1N1) virus infection. *Acta Pharmacol. Sin.* 2013, 34, 1075–1083.
88. Zhan, P.; Liu, X.; Cao, Y.; Wang, Y.; Pannecouque, C.; Clercq, E.D. 1,2,3-thiadiazole thioacetanilides as a novel class of potent HIV-1 non-nucleoside reverse transcriptase inhibitors. *Bioorg. Med. Chem. Lett.* 2008, 18, 5368–5371.
89. Zhan, P.; Liu, X.; Li, Z.; Fang, Z.; Li, Z.; Wang, D.; Pannecouque, C.; Clercq, E.D. Novel 1,2,3-thiadiazole derivatives As HIV-1 NNRTIs with improved potency: Synthesis and preliminary SAR studies. *Bioorg. Med. Chem. Lett.* 2009, 17, 5920–5927.
90. Dong, W.-L.; Liu, Z.-X.; Liu, X.-H.; Li, Z.-M.; Zhao, W.-G. Synthesis and antiviral activity of new acrylamide derivatives containing 1,2,3-thiadiazole as inhibitors of hepatitis B virus replication. *Eur. J. Med. Chem.* 2010, 45, 1919–1926.
91. Wang, S.-X.; Fang, Z.; Fan, Z.J.; Wang, D.; Li, Y.-D.; Li, X.-T.; Hua, X.-W.; Huang, Y.; Kalinina, A.T.; Bakulev, A.V.; et al. Synthesis of tetrazole containing 1,2,3-thiadiazole derivatives via U-4CR and their anti-TMV activity. *Chin. Chem. Lett.* 2013, 24, 889–892.
92. Mao, W.T.; Zhao, H.; Fan, Z.J.; Ji, X.T.; Hua, X.W.; Kalinina, T.; Yury, Y.M.; Vasiliy, A.B. Synthesis and bioactivity of N-tert-butyl-N'-acyl-5-methyl-1,2,3-thiadiazole-4-carbohydrazides. *Chin. Chem. Lett.* 2012, 23, 1233–1236.
93. Xu, W.-M.; Li, S.-Z.; He, M.; Yang, S.; Li, X.-Y.; Li, P. Synthesis and bioactivities of novel thioether/sulfone derivatives containing 1,2,3-thiadiazole and 1,3,4-oxadiazole/thiadiazole moiety. *Bioorg. Med. Chem. Lett.* 2013, 23, 5821–5824.
94. Zheng, Q.; Mi, N.; Fan, Z.; Zuo, X.; Zhang, H.; Wang, H.; Yang, Z. 5-Methyl-1,2,3-thiadiazoles Synthesized via Ugi Reaction and Their Fungicidal and Antiviral Activities. *J. Agric. Food Chem.* 2010, 58, 7846–7855.

95. Zhan, P.; Liu, X.Y.; Li, Z.Y.; Fang, Z.-J.; Pannecouque, C.; Clercq, E.D. Synthesis and biological evaluation of a new series of 2-[[4-(3,4-dichlorophenyl)-1,2,3-thiadiazol-5-yl] sulfanyl]acetanilides as HIV-1 inhibitors. *Chem. Biodivers.* 2010, 7, 1717–1727.
96. Grasso, C.S.; Wu, Y.M.; Robinson, D.R.; Cao, X.; Dhanasekaran, S.M.; Khan, A.P.; Quist, M.J.; Jing, X.J.; Lonigro, R.J.; Brenner, J.C.; et al. The mutational landscape of lethal castration-resistant prostate cancer. *Nature* 2012, 487, 239–243.
97. Irfan, A.; Batool, F.; Naqvi, S.A.Z.; Islam, A.; Osman, S.M.; Nocentini, A.; Alissa, S.A.; Supuran, C.T. Benzothiazole derivatives as anticancer agents. *J. Enzym. Inhib. Med. Chem.* 2020, 35, 265–279.
98. Pattabiraman, V.R.; Bode, J.W. Rethinking amide bond synthesis. *Nature* 2011, 480, 471–479.
99. He, Z.C.; Ling, G.L.; Yi, Z.Y.; Fei, Z.F.; Zhou, W.G.; Lei, J.; Xia, G.R. Review on supermolecules as chemical drugs. *Sci. China Ser. B* 2009, 52, 415–458.
100. Dasari, S.; Tchounwou, P.B. Cisplatin in cancer therapy: Molecular mechanisms of action. *Eur. J. Pharmacol.* 2014, 740, 364–378.
101. Gangadhara, S.; Bertelli, G. Long-term efficacy and safety of anastrozole for adjuvant treatment of early breast cancer in postmenopausal women. *Ther. Clin. Risk Manag.* 2009, 5, 291–300.
102. Barros-Oliveira, M.C.; Costa-Silva, D.R.; Andrade, D.B.; Borges, U.S.; Tavares, C.B.; Borges, R.S.; Silva, J.M.; Silva, B.B. Use of anastrozole in the chemoprevention and treatment of breast cancer: A literature review. *Rev. Assoc. Med. Bras.* 2017, 63, 371–378.
103. Tacar, O.; Sriamornsak, P.; Dass, C.R. Doxorubicin: An update on anticancer molecular action, toxicity and novel drug delivery systems. *J. Pharm. Pharmacol.* 2013, 65, 157–170.
104. Pereyra, C.E.; Dantas, R.F.; Ferreira, S.B.; Gomes, L.P.; Silva, F.P. The diverse mechanisms and anticancer potential of naphthoquinones. *Cancer Cell Int.* 2019, 19, 207.
105. Thorn, C.F.; Oshiro, C.; Marsh, S.; Hernandez, B.T.; McLeod, H.; Klein, T.E.; Altman, R.B. Doxorubicin pathways: Pharmacodynamics and adverse effects. *Pharm. Genom.* 2011, 21, 440–446.
106. Al-Malky, S.H.; Damanhour, Z.A.; Aama, J.Y.A.; Qahtani, A.A.A.; Ramadan, W.S.; AlKreathy, H.M.; Harthi, S.E.A.; Osman, A.M.M. Diltiazem potentiation of doxorubicin cytotoxicity and cellular uptake in human breast cancer cells. *Breast Cancer Manag.* 2019, 8, BMT31.
107. Wu, M.; Sun, Q.; Yang, C.; Chen, D.; Ding, J.; Chen, Y.; Lin, L.; Xie, Y. Synthesis and activity of Combretastatin A-4 analogues: 1,2,3-thiadiazoles as potent antitumor agents. *Bioorg. Med. Chem. Lett.* 2007, 17, 869–873.
108. Hosny, M.A.; El-Sayed, T.H.; El-Sawi, E.A. A New Type of Synthesis of 1,2,3-thiadiazole and 1,2,3-diazaphosphole derivatives Via-Hurd-Mori cyclization. *J. Chem.* 2012, 9, 1276–1287.

109. Cikotiene, I.; Kazlauskas, E.; Matulienė, J.; Michailovienė, V.; Torresan, J.; Jachno, J.; Matuliset, D. 5-Aryl-4-(5-substituted-2,4-dihydroxyphenyl)-1,2,3-thiadiazoles as inhibitors of Hsp90 chaperone. *Bioorg. Med. Chem. Lett.* 2009, 19, 1089–1092.
110. Cui, H.W.; Peng, S.; Gu, X.Z.; Chen, H.; He, Y.; Gao, W.; Lv, F.; Wang, J.-H.; Wang, Y.; Xie, J.; et al. Synthesis and biological evaluation of D-ring fused 1,2,3-thiadiazole dehydro epiandro sterone derivatives as antitumor agents. *Eur. J. Med. Chem.* 2016, 111, 126–137.
111. Oberemok, V.V.; Laikova, K.V.; Gninenko, Y.I.; Zaitsev, A.S.; Nyadar, P.M.; Adeyemi, T.A. A short history of insecticides. *J. Plant Prot. Res.* 2015, 55, 221–226.
112. Xu, Z.; Shi, L.; Jiang, D.; Cheng, J.; Shao, X.; Li, Z. Azobenzene Modified Imidacloprid Derivatives as Photoswitchable Insecticides: Steering Molecular Activity in a Controllable Manner. *Sci. Rep.* 2015, 5, 13962.
113. Carlson, G.R. Tebufenozide: A Novel Caterpillar Control Agent with Unusually High Target Selectivity. *Green Chemical Syntheses and Processes. ACS Symp. Ser.* 2000, 767, 8–17.
114. Zhang, A.; Kayser, H.; Maeinfisch, P. Insect nicotinic acetylcholine receptor: Conserved neonicotinoid specificity of imidacloprid binding site. *J. Neurochem.* 2002, 75, 1294–1303.
115. Kumar, A.; Verma, A.; Kumar, A. Accidental human poisoning with a neonicotinoid insecticide, imidacloprid: A rare case report from rural India with a brief review of literature. *Egypt. J. Forensic Sci.* 2013, 3, 123–126.
116. Mundhe, S.A.; Birajdar, S.V.; Chavan, S.S.; Pawar, N.R. Imidacloprid Poisoning: An Emerging Cause of Potentially Fatal Poisoning. *Indian J. Crit. Care Med.* 2017, 21, 786–788.
117. Ausborn, J.; Wolf, H.; Mader, W.; Kayser, H. The insecticide pymetrozine selectively affects chordotonal mechanoreceptors. *J. Exp. Biol.* 2005, 28, 4451–4466.
118. Zhang, J.P.; Qin, Y.G.; Dong, Y.W.; Song, D.-L.; Duan, H.-X.; Yang, X.-L. Synthesis and biological activities of (E)-B-farnesene analogues 4 containing 1,2,3-thiadiazole. *Chin. Chem. Lett.* 2016, 28, 372–376.
119. Dai, H.; Ge, S.; Li, G.; Chen, J.; Shi, Y.; Ye, L.; Ling, Y. Synthesis and bioactivities of novel pyrazole oxime derivatives containing a 1,2,3-thiadiazole moiety. *Bioorg. Med. Chem. Lett.* 2016, 28, 4504–4507.
120. Li, Y.-D.; Mao, W.-T.; Fan, Z.-J.; Li, J.-J.; Fang, Z.; Ji, X.-T.; Hua, X.-W.; Zong, G.-N.; Li, F.-Y.; Liu, C.-L.; et al. Synthesis and biological evaluation of novel 1,2,4-triazole containing 1,2,3-thiadiazole derivatives. *Chin. Chem. Lett.* 2013, 24, 1134–1136.
121. Bercu, T.E.; Petri, W.A.; Behm, J.W. Amebic colitis: New insights into pathogenesis and treatment. *Curr. Gastroenterol. Rep.* 2007, 9, 429–433.

122. Choudhuri, G.; Rangan, M. Amebic infection in humans. *Indian J. Gastroenterol.* 2012, 31, 153–162.
123. Shirley, D.T.; Farr, L.; Watanabe, K.; Moonah, S. A review of the global burden, new diagnostics, and current therapeutics for amebiasis. *Open Forum Infect. Dis.* 2018, 5, ofy161.
124. Farrar, J.; Hotez, P.; Junghanss, T.; Kang, G.; Laloo, D.; White, N. *Manson's Tropical Diseases*; Elsevier Health Sciences: Amsterdam, The Netherlands, 2013; pp. 664–671. ISBN 9780702053061.
125. Lofmark, S.; Edlund, C.; Nord, C.E. Metronidazole Is Still the Drug of Choice for Treatment of Anaerobic Infections. *Clin. Infect. Dis.* 2010, 50, S16–S23.
126. Steinitz, H. The Treatment of Chronic Amoebic Infection with Metronidazole (Flagyl). *Digestion* 1972, 6, 75–82.
127. Gonzales, M.L.M.; Dans, L.F.; Martinez, E.G. Antiamoebic drugs for treating amoebic colitis. *Cochrane Database Syst. Rev.* 2009, 2, CD006085.
128. Ghaskadbi, S.; Vaidya, V.G. In vivo antimutagenic effect of ascorbic acid against mutagenicity of the common antiamebic drug diiodohydroxyquinoline. *Mutat. Res.* 1989, 222, 219–222.
129. Hayat, F.; Salahuddin, A.; Zargan, J.; Azam, A. Synthesis, characterization, antiamoebic activity and cytotoxicity of novel 2-(quinolin-8-yloxy) acetohydrazones and their cyclized products (1,2,3-thiadiazole and 1,2,3-selenadiazole derivatives). *Eur. J. Med. Chem.* 2010, 45, 6127–6134.
130. Aranega-Bou, P.O.; Leyva, M.; Finiti, I.; Garcia-Agustin, P.; Gonzalez-Bosch, C. Priming of plant resistance by natural compounds. Hexanoic acid as a model. *Front. Plant Sci.* 2014, 5, 488.
131. Li, J.; Long, T.; Sun, T.-J.; Lu, Y.; Yin, J.; Yang, Y.-B.; Dai, G.Y.; Zhu, X.Y.; Yao, N. Pyrimidin-Like Plant Activator Stimulates Plant Disease Resistance and Promotes the Synthesis of Primary Metabolites. *Int. J. Mol. Sci.* 2020, 21, 2705.
132. Conrath, U.; Beckers, G.J.M.; Langenbach, C.J.G.; Jaskiewicz, M.R. Priming for enhanced defense. *Annu. Rev. Phytopathol.* 2015, 53, 97–119.
133. Cavalcanti, F.R.; Resendeb, M.L.V.; Limac, J.P.M.S.; Silveirac, J.A.G.; Oliveira, J.T.A. Activities of antioxidant enzymes and photosynthetic responses in tomato pretreated by plant activators and inoculated by *Xanthomonas vesicatoria*. *Int. J. Plant Pathol.* 2006, 68, 198–208.
134. Martinez-Medina, A.; Flors, V.; Heil, M.; Mauch-Mani, B.; Pieterse, C.M.J.; Pozo, M.J.; Ton, J.; Dam, N.M.V. Recognizing Plant Defense Priming. *Trends Plant Sci.* 2016, 21, 818–822.
135. Zhang, L.; Sun, B.; Liu, Q.; Mo, F. Addition of Diazo Compounds ipso-C-H Bond to Carbon Disulfide: Synthesis of 1,2,3-Thiadiazoles under Mild Conditions. *J. Org. Chem.* 2018, 83, 4275–4278.

136. Du, Q.-S.; Shi, Y.-X.; Li, P.-F.; Zhao, Z.-J.; Zhu, W.-P.; Qian, X.-H.; Li, B.-J.; Xu, Y.-F. Novel plant activators with thieno[2,3-d]-1,2,3-thiadiazole-6-carboxylate scaffold: Synthesis and bioactivity. *Chin. Chem. Lett.* 2013, 24, 967–969.
137. Gupta, P.K.; Agarwal, M. Toxicity of fungicides. In *Veterinary Toxicology: Basic and Clinical Principles*, 2nd ed.; Gupta, R.C., Ed.; Elsevier: Amsterdam, The Netherland, 2012; pp. 653–670.
138. Ramudu, A.C.; Mohiddin, G.J.; Srinivasulu, M.; Madakka, M.; Rangaswamy, V. Impact of Fungicides Chlorothalonil and Propiconazole on Microbial Activities in Groundnut (*Arachis hypogaea* L.) Soils. *ISRN Microbiol.* 2011, 2011.
139. Bacmaga, T.M.; Wyszowska, J.; Kucharski, J. The influence of chlorothalonil on the activity of soil microorganisms and enzymes. *Ecotoxicology* 2018, 27, 1188–1202.
140. Martin, M.V. The use of fluconazole and itraconazole in the treatment of *Candida albicans* infections: A review. *J. Antimicrob.* 1999, 44, 429–437.
141. Greer, D.; Jolly, H.W. Topical ketoconazole treatment of cutaneous candidiasis. *J. Am. Acad. Dermatol.* 1988, 18, 748–750.
142. Fan, Z.; Shi, Z.; Zhang, H.; Liu, X.; Bao, L.; Ma, L.; Zuo, X.; Zheng, Q.; Mi, N. Synthesis and biological activity evaluation of 1,2,3-thiadiazole derivatives as potential elicitors with highly systemic acquired resistance. *J. Agric. Food Chem.* 2009, 57, 4279–4286.
143. Sun, N.-B.; Fu, J.Q.; Weng, J.-Q.; Jin, J.-Z.; Tan, C.-X.; Liu, X.-H. Microwave Assisted Synthesis, Antifungal Activity and DFTTheoretical Study of Some Novel 1,2,4-Triazole Derivatives Containing the 1,2,3-Thiadiazole Moiety. *Molecules* 2013, 18, 12725–12739.
144. Tan, C.-X.; Weng, J.-Q.; Liu, Z.-X.; Liu, X.-H.; Zhao, W.-G. Synthesis, Crystal Structure, and Fungicidal Activity of a Novel 1,2,3-Thiadiazole Compound. *Phosphorus Sulfur Silicon Relat. Elem.* 2012, 187, 990–996.
145. Fan, Z.; Yang, Z.; Zhang, H.; Mi, N.; Wang, H.; Cai, F.; Zuo, X.; Zheng, Q.; Song, H. Synthesis, crystal structure, and biological activity of 4-methyl-1,2,3-thiadiazole-containing 1,2,4-triazolo[3,4-b][1,3,4]thiadiazoles. *J. Agric. Food Chem.* 2010, 58, 2630–2636.
146. Wang, Z.-H.; Guo, Y.-Z.; Zhang, J.; Ma, L.; Song, H.-B.; Fan, Z.-J. Synthesis and biological activity of organotin 4-Methyl-1,2,3-thiadiazole-5-carboxylates and benzo[1,2,3]thiadiazole-7-carboxylates. *J. Agric. Food Chem.* 2010, 58, 2715–2719.
147. Zuo, X.; Mi, N.; Fan, Z.; Zheng, Q.; Zhang, H.; Wang, H.; Yang, Z. Synthesis of 4-methyl-1,2,3-thiadiazole derivatives via Ugi reaction and their biological Activities. *J. Agric. Food Chem.* 2010, 58, 2755–2762.

Retrieved from <https://encyclopedia.pub/entry/history/show/28664>