

Global Solar Radiation

Subjects: **Meteorology & Atmospheric Sciences**

Contributor: Jianhui Bai

An empirical model of global solar irradiance (EMGSI) under all sky conditions was developed by using solar radiation and meteorological parameters at Sodankylä. The calculated hourly global solar irradiance is in agreement with that observed at the ground during 2008–2011 and at the top of the atmosphere (TOA). Using this empirical model, the albedos at the TOA and the surface were calculated, and they are in agreement with the satellite-retrieved values.

absorbing and scattering factors

energy

aerosol optical depth

1. Introduction

The surface temperatures in the Arctic increase much faster than the global average, a phenomenon known as Arctic Amplification. The most significant increase occurs in the winter [\[1\]](#)[\[2\]](#)[\[3\]](#). The Arctic warming is related to the imbalance of energy budget at the ground and the top of the atmosphere (TOA), sea-ice–albedo feedback, lapse-rate feedback, temperature inversion, enhanced greenhouse effect [\[1\]](#) and the reasons for the climate change in the Arctic region are still unclear.

The Sun provides important energy to the atmospheric gases, liquids and particles (GLPs) and initiates their changes in the three phases through chemical and photochemical reactions (CPRs). Therefore, solar radiation transfers in and interacts with the atmospheric GLPs. The attenuation in the atmosphere and reflections at the ground and the TOA are necessary to be investigated; these physical and chemical processes control/interact with the atmospheric movement through short- and long-wave radiation in different ways [\[4\]](#)[\[5\]](#)[\[6\]](#)[\[7\]](#)[\[8\]](#)[\[9\]](#)[\[10\]](#)[\[11\]](#)[\[12\]](#), and should be studied for further understanding the regional climate and climate change in the Arctic.

The popular radiative transfer models and empirical models are used to estimate global, direct and diffuse solar radiation at the ground and briefly described in References [\[12\]](#)[\[13\]](#)[\[14\]](#)[\[15\]](#)[\[16\]](#)[\[17\]](#)[\[18\]](#)[\[19\]](#)[\[20\]](#). To thoroughly investigate the Sun–atmosphere–Earth system, variations of solar radiation at the surface and the TOA, the loss of solar radiation in the atmosphere, the albedos at the surface and the TOA, and the interactions between solar radiation and meteorological variables are required.

2. Global Solar Irradiance during 2000–2018

During 2000–2018, annual averages of observed global and diffuse solar irradiance ($n= 92,941$) were 0.59 and 0.29 MJ m^{-2} , thus, the direct horizontal irradiance was a little larger than the diffuse irradiance at the surface,

average of S/G was 0.64, and averages of surface air temperature and relative humidity were 0.76 °C (ranged from -39.90 to 31.50 °C) and 80.31%, respectively.

Applying the empirical model of global solar irradiance, hourly global irradiance was calculated for Sodankylä during 1 January 2000 to 31 December, 2018. The input data were observed hourly global and diffuse solar irradiance for S/G and water vapor pressure at the ground. Global solar irradiance in December was too low and not considered in the later analysis, and n decreased to 69,171. The calculated and observed hourly global solar irradiance varied similarly, and calculated values were larger than the observed by 18.11% for the average in 2000–2018. It is reasonable that the empirical model represents the global irradiance and its relationships with absorbing and scattering processes at relative clean atmospheric conditions (i.e., relative low S/G at 0.50 in 2008–2011 in the model development). The calculated and observed monthly global irradiance, diffuse irradiance and S/G are shown in Figure 1. Global solar irradiance in June or July was higher over clean atmospheric conditions, e.g., 2001 to 2003, 2007 and 2011 at low S/G (<0.60), but lower over high GLP loading, e.g., 2008, 2009 and 2017 at high S/G (>0.60). Scattering factor S/G was lower during April to September and higher in October to March.




Figure 1. Monthly global solar irradiance calculated and observed (G), observed diffuse irradiance (S) and scattering factor (S/G) at Sodankylä.

During 19 years, the monthly averages of the calculated global irradiance decreased by 0.01% and diffuse irradiance increased by 0.11% per year, respectively. They were associated with the increases of S/G by 0.14% and water vapor by 0.06%. Air temperature and relative humidity increased by 0.83% (corresponding to 0.01 °C) and 0.23% for monthly average, respectively. On the average, the annual air temperature increased about 2.09 °C during 2000–2018, also indicating the Arctic warming [21].

Long term interannual variations were also investigated during 2000–2018 (Figure 2 and Figure 3). The calculated annual global solar irradiance decreased by 0.92% and diffuse irradiance increased by 1.28% per year, respectively. Both were associated with the increases of S/G by 1.73% and water vapor by 0.43% per year. Annual air temperature increased by 4.64% (corresponding to 0.07 °C), and relative humidity increased by 0.23% per year, respectively. The annual air temperature showed a higher correlation with water vapor ($R = 0.79$) than with relative humidity ($R = 0.20$). A good correlation was determined in annual water vapor pressure (E) and air temperature (T) and relative humidity (RH) as $E = 0.293 \times T + 0.024 \times RH + 3.994$ ($R = 0.828$, $n = 19$, $\alpha = 0.001$).




Figure 2. Annual global solar irradiance calculated and observed (G), observed diffuse irradiance (S) and scattering factor (S/G) at Sodankylä.




Figure 3. Annual air temperature (T), relative humidity (RH) and water vapor pressure (E) at Sodankylä.

Based on above as well as the following integrated results, the increase of air temperature is evidently associated with the increased GLPs, as well as water vapor. If the output of the Sun remains, solar global radiation at the ground would decrease and air temperature increase with the increase of atmospheric substances in the future.

3. The Losses of Global Solar Irradiance in the Atmosphere during 2000–2018

The global solar irradiance losses caused by absorbing and scattering GLPs (G_{LA} and G_{LS}) were calculated by using $A_1(1-e^{-kWm} \times \cos(Z))$ and $A_2(1-e^{-S/G})$, respectively. The total loss G_L is $G_{LA} + G_{LS}$. The monthly losses due to absorption G_{LA} dominated the total loss G_L and showed clear seasonal variations, lower in April–September and

higher in October–March, whereas G_{LS} did not have evident seasonal variation and most peaks appeared in October–February. During January 2000 to December, 2018, (1) monthly G_{LA} decreased slightly by 0.003%, associated with the increase of water vapor by 0.06%; (2) monthly G_{LS} increased by 0.06%, associated with the increase of S/G by 0.14%; and (3) monthly G_L increased by 0.02% (Figure 4).




Figure 4. Monthly losses of global solar irradiance caused by absorbing and scattering substances (G_{LA} , G_{LS}) and total loss ($G_L = G_{LA} + G_{LS}$) at Sodankylä.

For the annual losses of global solar irradiance during 2000–2018, G_{LA} decreased by 0.02% per year, associating with the increase of E by 0.43%; G_{LS} increased by 0.72% per year, associating with the increase of S/G by 1.73%; annual G_L increased by 0.24% per year.

The contributions of monthly mean absorbing and scattering losses (R_{LA} and R_{LS}) to monthly mean total loss were 63.33% (ranged from 53.95 to 71.99%) and 36.67% (28.01–46.05%), respectively, during January 2000 to December 2018 (Figure 5), corresponding to the monthly averages of water vapor at 6.48 hPa (0.57–14.33) and S/G at 0.55 (0.30–0.94). Generally, R_{LA} was lower in April–September and higher in October–March, whereas R_{LS} varied inversely compared to R_{LA} (i.e., most peaks appeared in October–February). The corresponding annual mean absorbing and scattering losses (R_{LA} and R_{LS}) to annual mean total loss were 63.32% (60.09–65.48%) and 36.68% (34.52–39.04%), respectively.




Figure 5. Contributions (R_{LA} , R_{LS}) of monthly absorbing and scattering losses to monthly total loss at Sodankylä.

During 2000–2018 at Sodankylä, the annual mean monthly loss of G_{LA} , G_{LS} and G_L were 1.94 (1.57–2.41), 1.17 (0.82–1.53) and 3.11 (2.50–3.90) MJ m⁻², corresponding to 539.82, 323.86 and 863.68 W m⁻², respectively.

References

1. Cao, Y.F.; Liang, S.L. Recent advances in driving mechanisms of the Arctic amplification: A review. *Chin. Sci. Bull.* 2018, 63, 2757–2771. (In Chinese)
2. Cohen, J.; Screen, J.A.; Furtado, J.; Barlow, M.; Whittleston, D.; Coumou, D.; Francis, J.A.; Dethloff, K.; Entekhabi, D.; Overland, J.E.; et al. Recent Arctic amplification and extreme mid-latitude weather. *Nat. Geosci.* 2014, 7, 627–637.
3. Liu, S.Y.; Wang, B.Y.; Xie, Y.X.; Hu, S.H.; Wang, Z.Y.; Yue, L.Y. The Variation Characteristics of Temperature in Barrow Alaska during 1925–2018. *Clim. Chang. Res. Lett.* 2019, 8, 769–774.
4. Solanki, S.K. Solar variability and climate change: Is there a link. *Astron. Geophys.* 2002, 43, 5.
5. Elminir, H.K. Relative influence of weather conditions and air pollutants on solar radiation—Part 2: Modification of solar radiation over urban and rural sites. *Meteorol. Atmos. Phys.* 2007, 96, 257–264.
6. Lean, J.; Rind, D. Climate forcing by changing solar radiation. *J. Clim.* 1998, 11, 3069–3094.
7. Andreae, M.O.; Ramanathan, V. Climate's Dark Forcings. *Sciences* 2013, 340, 280–281.
8. García, R.D.; Cuevas, E.; García, O.E.; Cachorro, V.E.; Pallé, P.; Bustos, J.J.; Romero-Campos, P.M.; de Frutos, A.M. Reconstruction of global solar radiation time series from 1933 to 2013 at the Izaña Atmospheric Observatory. *Atmos. Meas. Tech.* 2014, 7, 3139–3150.
9. Rosenfeld, D.; Sherwood, S.; Wood, R.; Donner, L. Climate Effects of Aerosol-Cloud Interactions. *Sciences* 2014, 343, 379–380.
10. Williamson, C.E.; Zepp, R.G.; Lucas, R.M.; Madronich, S.; Austin, A.T.; Ballaré, C.L.; Norval, M.; Sulzberger, B.; Bais, A.F.; McKenzie, R.L.; et al. Solar ultraviolet radiation in a changing climate. *Nat. Clim. Chang.* 2014, 4, 434–441.
11. Calabró, E.; Magazù, S. Correlation between Increases of the Annual Global Solar Radiation and the Ground Albedo Solar Radiation due to Desertification—A Possible Factor Contributing to Climatic Change. *Climate* 2016, 4, 64.
12. Bai, J.H.; Zong, X.M. Global solar radiation transfer and its loss in the atmosphere. *Appl. Sci.* 2021, 11, 2651.
13. Glover, J.; McCulloch, J.S.G. The empirical relation between solar radiation and hours of sunshine. *Q. J. R. Meteorol. Soc.* 1958, 84, 172–175.

14. Zhang, J.; Zhao, L.; Deng, S.; Xu, W.; Zhang, Y. A critical review of the models used to estimate solar radiation. *Renew. Sustain. Energy Rev.* 2017, 70, 314–329.
15. Gueymard, C.A. Critical analysis and performance assessment of clear sky solar irradiance models using theoretical and measured data. *Sol. Energy* 1993, 51, 121–138.
16. Gueymard, C.A. Clear-sky solar irradiance predictions for large-scale applications using 18 radiative models: Improved validation methodology and detailed performance analysis. *Sol. Energy* 2012, 86, 2145–2169.
17. Badescu, V.; Gueymard, C.A.; Cheval, S.; Oprea, C.; Baciu, M.; Dumitrescu, A.; Iacobescu, F.; Milos, I.; Rada, C. Accuracy analysis for fifty-four clear-sky solar radiation models using routine hourly global irradiance measurements in Romania. *Renew. Energy* 2013, 55, 85–103.
18. Bayrakc, H.C.; Demircan, C.; Keceba, A. The development of empirical models for estimating global solar radiation on horizontal surface: A case study. *Renew. Sustain. Energy Rev.* 2018, 81, 2771–2782.
19. Antonanzas-Torres, F.; Urraca, R.; Polo, J.; Perpinan-Lamigueiro, O.; Escobar, R. Clear sky solar irradiance models: A review of seventy models. *Renew. Sustain. Energy Rev.* 2019, 107, 374–387.
20. Zang, H.; Cheng, L.; Ding, T.; Cheung, K.W.; Wang, M.; Wei, Z.; Sun, G. Estimation and validation of daily global solar radiation by day of the year-based models for different climates in China. *Renew. Energy* 2019, 135, 984–1003.
21. Post, E.; Alley, R.B.; Christensen, T.R.; Macias-Fauria, M.; Forbes, B.; Gooseff, M.N.; Iler, A.; Kerby, J.T.; Laidre, K.L.; Mann, M.E.; et al. The polar regions in a 2 °C warmer world. *Sci. Adv.* 2019, 5, eaaw9883.

Retrieved from <https://www.encyclopedia.pub/entry/history/show/38933>