

# General Anesthesia

Subjects: Anesthesiology

Contributor: Alessandro De Cassai

Vertebral lumbar surgery can be performed under both general anesthesia (GA) and spinal anesthesia. A clear benefit from spinal anesthesia (SA) remains unproven.

Keywords: general anesthesia ; spinal anesthesia

---

## 1. Introduction

Vertebral lumbar surgery can be performed under both general anesthesia (GA) and spinal anesthesia (SA). Each has possible advantages and complications in the perioperative period <sup>[1]</sup>. In particular, SA does not require airway device placement for intraoperative sedation and analgesia; however, it could be associated with patient discomfort and intraoperative patient movements <sup>[2]</sup>. Moreover, fear of neuraxial damage caused by either local anesthetic toxicity or direct damage with an associated prolonged hospital length of stay (LOS) may discourage its use <sup>[3]</sup>.

According to a systematic review and meta-analysis published in 2016 <sup>[4]</sup>, patients undergoing lumbar spine surgery under SA required less analgesia in post-anesthesia care units and had less nausea and vomiting (PONV) in the first postoperative day, but no difference in intraoperative hypotension, bradycardia, blood loss, and surgical time was reported.

Given the above, a clear benefit from SA during lumbar spine surgery remains unproven. Furthermore, relevant clinical outcomes remain unexplored.

## 2. Study Selection and Data Retrieval

Bibliographic search results are shown in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) diagram (Figure 1). Notably, three RCTs did not report quantitative data. Two papers were excluded because, despite our best efforts, we were not able to retrieve the full text <sup>[5][6]</sup>, in one case <sup>[7]</sup> the paper did not contain any variable of interest and the authors were not able to provide any missing information.

**Figure 1.** The PRISMA flowchart.

Eleven studies counting a total of 896 patients entered the quantitative and qualitative analysis [8][9][10][11][12][13][14][15][16][17][18]. All controversies were solved by discussion and the third reviewer was not required.

We asked all the corresponding authors for missing data, and five of them replied to our query. Only two of them, nevertheless, provided part of the missing data required [10][11][12][13][14][15][16][17][18].

Additional records were identified by checking the reference lists of included studies.

### 3. Study Characteristics

Among the 896 patients, half (449; 50.12%) underwent GA, while the remainder underwent SA (447; 49.88%). The characteristics of included studies are shown in Table 1 and Table S2.

**Table 1.** Study characteristics.

Study	N (%)	Inclusion Criteria			Protocols			Surgery	Time to Pain Assessment
		Age	ASA-PS	BMI (Kg/m <sup>2</sup> )	SA	GA	PO Therapy		
Attari (2011) [8]	72 (46%)	18–60	I–II	NR	3.0–3.2 mL Hyperbaric Bupivacaine 0.5% + 25 mcg Fentanyl	- Induction: propofol, lidocaine, fentanyl, atracurium Maintenance: isoflurane 1,2%, N <sub>2</sub> O 50%	Pethidine 0.4 mg/kg on VAS (rescue pethidine 0.2 mg/kg).	Laminectomy, Discectomy	NR
Baenziger (2020) [15]	100 (46%)	Adult	I–III	NR	3.0–4.0 mL Hyperbaric Bupivacaine 0.5% + 25 mcg Fentanyl	- Induction: propofol, fentanyl, atracurium Maintenance: Propofol TCI, Remifentanyl TCI	NR	Laminectomy, Discectomy	3 h
Chowdhury (2010) [9]	80 (38%)	Adult	I–II	NR	2.5–2.8 mL Hyperbaric Bupivacaine 0.5% + 12.5 mcg Fentanyl	- Induction: propofol, fentanyl, rocuronium Maintenance: halothane 0.8%, N <sub>2</sub> O 60%	Pethidine 2 mg/kg six hourly and on request.	Discectomy	6 h

Hussain (2015) <sup>[10]</sup>	60 (50%)	20–50	I–II	NR	2 mL Bupivacaine 0.75%	- Induction: propofol, atracurium - Maintenance: sevoflurane 1.5–2%, nalbuphine	NR	Micro-discectomy	Peak at 6 h
Jellish (1996) <sup>[11]</sup>	122 (46%)	Adult	I–III	NR	1.5 mL Hyperbaric Bupivacaine 0.75%	- Induction: thiopental, fentanyl, vecuronium - Maintenance: isoflurane, N <sub>2</sub> O 70%	PACU: morphine 2 mg IV ward: meperidine 25–50 mg IV or 50–100 mg intramuscularly.	Laminectomy, Discectomy	Peak
Kahveci (2014) <sup>[12]</sup>	80 (38%)	≥18	I–II	≤25	3 mL Hyperbaric Bupivacaine 0.5%	- Induction: propofol, fentanyl, atracurium - Maintenance: sevoflurane 1.5–2%, atracurium	Pethedine 25 mg IV on VAS.	Single-level spinal surgery	NR
Kara (2011) <sup>[15]</sup>	60 (45%)	Adult	I–II	NR	2 mL Levobupivacaine 0.5%	- Induction: propofol, fentanyl, rocuronium - Maintenance: desflurane 6%, N <sub>2</sub> O 40–60%	Morphine 2 mg on VAS.	Discectomy	Peak
Kilic (2019) <sup>[17]</sup>	111 (45%)	18–65	I–III	NR	3 mL Hyperbaric Bupivacaine 0.5%	- Induction: propofol, fentanyl, rocuronium - Maintenance: sevoflurane 1.5–2%, remifentanyl	NR	Micro-discectomy	3 h

Sadrolsadat (2009) <sup>[13]</sup>	100 (-)	Adult	I–III	NR	4 mL Bupivacaine 0.5%	- Induction:  propofol,  fentanyl, atracurium  - Maintenance: propofol, alfentanil, atracurium	Pethedine 25 mg IV on VAS (lock 30 min in PACU and 4 h in ward).	Laminectomy	NR
Vural (2014) <sup>[14]</sup>	66 (-)	23– 74	ND	NR	4 mL Hyperbaric Bupivacaine 0.5%	- Induction:  thiopental,  fentanyl, rocuronium  - Maintenance: desflurane 5–6%, N <sub>2</sub> O 40– 60%,fentanyl	NR	Disc herniation surgery	6 h
Yildirim Güçlü (2014) <sup>[18]</sup>	56 (-)	18– 60	I–II	≤35	3 mL Hyperbaric Bupivacaine 0.5%	- Induction:  thiopental,  fentanyl, vecuronium  - Maintenance: desflurane 4–5%, N <sub>2</sub> O 50%, remifentanil	Pethidine 0.5 mg/kg on VAS (Rescue pethidine 0.2 mg/kg).	Micro- discectomy	NR

ASA-PS: ASA Physical Status, F: females, GA: General Anesthesia, SA: Spinal Anesthesia, PACU: Post-Anesthesia Care Unit, PO: Post-Operative, NR: Not Reported.

There were concerns of bias in ten studies, where one study was evaluated at high risk of bias <sup>[17]</sup> (Figure 2).

**Figure 2.** Summary of bias evaluated by the Risk of Bias 2 Tool. R: Bias arising from the randomization process, D: Bias due to deviations from intended interventions, Mi: Bias due to missing outcome data, Me: Bias in measurement of the outcome, S: Bias in the selection of the reported result, and O: Overall risk of bias. Green "+": Low risk of bias, Yellow "?": Some concerns, Red "-": High risk of bias.

## References

1. Parker, M.J.; Unwin, S.C.; Handoll, H.H.; Griffiths, R. General versus spinal/epidural anaesthesia for surgery for hip fractures in adults. *Cochrane Database Syst. Rev.* 2000, 4, CD000521.
2. Mergeay, M.; Verster, A.; Van Aken, D.; Vercauteren, M. Regional versus general anesthesia for spine surgery. A comprehensive review. *Acta Anaesthesiol. Belg.* 2015, 66, 1–9.
3. Zorrilla-Vaca, A.; Healy, R.J.; Mirski, M.A. A Comparison of Regional versus General Anesthesia for Lumbar Spine Surgery: A Meta-Analysis of Randomized Studies. *Neurosurg. Anesthesiol.* 2017, 29, 415–425.
4. Meng, T.; Zhong, Z.; Meng, L. Impact of spinal anaesthesia vs. general anaesthesia on peri-operative outcome in lumbar spine surgery: A systematic review and meta-analysis of randomised, controlled trials. *Anaesthesia* 2017, 72, 391–401.
5. Dagher, C.; Naccache, N.; Narchi, P.; Hage, P.; Antakly, M.C. Regional anesthesia for lumbar microdiscectomy. *J. Liban.* 2002, 50, 206–210.
6. Khan, M.W.; Ibrahim, N.O.; Iqbal, M.; Khan, A.A.; Abbasi, S.M. A prospective randomized trial comparing the spinal and general anesthesia in lumbar disc surgery: A study of 44 cases. *Forum. Mon.* 2012, 23, 61–64.
7. Pinar, H.U.; Kaşdoğan, Z.E.A.; Başaran, B.; Çöven, İ.; Karaca, Ö.; Doğan, R. The effect of spinal versus general anesthesia on intraocular pressure in lumbar disc surgery in the prone position: A randomized, controlled clinical trial. *Clin. Anesth.* 2018, 46, 54–58.
8. Attari, M.A.; Mirhosseini, S.A.; Honarmand, A.; Safavi, M.R. Spinal anesthesia versus general anesthesia for elective lumbar spine surgery: A randomized clinical trial. *Res. Med. Sci.* 2011, 16, 524–529.
9. Chowdhury, M.S.; Roy, S.S.; Rahman, M.M.; Hossain, M.M.; Alim, S. Comparative study in prolapse lumbar intervertebral disc (PLID) surgery by spinal vs general anaesthesia. *J. Bangladesh. S. Anesth.* 2010, 23, 47–50.
10. Hussain, Z.; Ghaffar, A.; Mushtaq, M.J.; Qasmi, S.A. Can spinal anaesthesia be a routine for single level lumbar discectomy? *Pak. Armed Forces Med. J.* 2015, 65, 397–401.
11. Jellish, W.S.; Thalji, Z.; Stevenson, K.; Shea, J. A prospective randomized study comparing short-and intermediate-term perioperative outcome variables after spinal or general anesthesia for lumbar disk and laminectomy surgery. *Anesth. Analg.* 1996, 83, 559–564.
12. Kahveci, K.; Doger, C.; Ornek, D.; Gokcinar, D.; Aydemir, S.; Ozay, R. Perioperative outcome and cost-effectiveness of spinal versus general anesthesia for lumbar spine surgery. *Neurol. Neurochir. Pol.* 2014, 48, 167–173.
13. Sadrolsadat, S.H.; Mahdavi, A.R.; Moharari, R.S.; Khajavi, M.R.; Khashayar, P.; Najafi, A.; Amirjamshidi, A. A prospective randomized trial comparing the technique of spinal and general anesthesia for lumbar disk surgery: A study of 100 cases. *Surg. Neurol.* 2009, 71, 60–65.
14. Vural, C.Y. Comparison of patient satisfaction and cost in spinal and general anesthesia for lumbar disc surgery. *Turk. Neurosurg.* 2014, 24, 380–384.
15. Kara, I.; Celik, J.B.; Bahar, O.C.; Apillioğullari, S.; Karabagli, H. Comparison of spinal and general anesthesia in lumbar disc surgery. *J. Neurol. Sci.* 2011, 28, 487–496.
16. Baenziger, B.; Nadi, N.; Doerig, R.; Proemmel, P.; Gahl, B.; Hodel, D.; Hausmann, O.N. Regional Versus General Anesthesia: Effect of Anesthetic Techniques on Clinical Outcome in Lumbar Spine Surgery: A Prospective Randomized Controlled Trial. *J. Neurosurg. Anesthesiol.* 2020, 32, 29–35.
17. Kilic, E.T.; Naderi, S. Effects of Anesthesia Protocol on Perioperative Outcomes and Costs of Lumbar Microdiscectomies. *Turk. Neurosurg.* 2019, 29, 843–850.
18. Yilidrim Güçlü, Ç.; Keçik, Y.; Yörükoğlu, D.; Attar, A. Neuroendocrine and Hemodynamic Effects of General Anesthesia and Spinal Anesthesia for Minimally Invasive Lumbar Disc Surgery: A Randomized Trial. *Neurol. Sci.* 2014, 3, 586–595.