Bell Peppers (Capsicum annum L.)

Subjects: Cell Biology Contributor: Zuamí Villagrán

The bell pepper is highly consumed worldwide due to its exotic colors (green, red, yellow, orange, and purple), flavor, and nutritional value. However, after processing bell pepper products, waste remains (seeds, peel, stem, and leaves), representing desirable raw material to obtain phytochemical compounds. They contain diverse bioactive compounds with interesting biological activities (in vivo and in vitro) and applications.

food revalorization

bioactive compounds

biological activities

health benefits

1. Introduction

Despite the nutritional contribution of food (regardless of its plant or animal origin), much of it will not be consumed by humans or animals and will be discarded as waste ^[1]. Thus, the prevention of food loss and waste promotes a favorable impact on the environment and food security of the world population, contributing to economic development ^[2]. Besides prevention, the revalorization of these food wastes is a technologically viable strategy, using them as bioactive ingredients to generate new, potentially functional or nutraceutical products ^[3]. Worldwide, food losses and waste mainly occur in fresh fruits and vegetables (>40%), and are mainly associated with poor handling and storage during post-harvest ^{[2][3][4]}, which can be a raw source of bioactive compounds.

The Capsicum genus belongs to the Solanaceae family, Solanoicleae subfamily, Solaneae tribe. Chili (*Capsicum*), along with corn, beans, and squash, is one of the oldest cultivated plants in America ^[4]. There are five commercially cultivated species of chili (*C*. *chinense*, *C*. *annuum*, *C*. *pubescens*, *C*. *baccatum*, *and C*. *frutescens*) and around 25 wild and semi-cultivated species ^[5]. Peppers (*C*. *annuum L*.) are classified as hot or sweet; they are grown in subtropical climate regions throughout the world, including Mexico ^[6].

The production of bell peppers has increased considerably in recent years; however, the annual losses of this crop are estimated to be 40% ^[8]. Bell peppers can be of different colors (red, green, orange, and yellow) depending on their ripening stages and capacity to synthesize chlorophylls or carotenoids. Besides having an exotic flavor, bell peppers are an important source of vitamins (provitamin A, E, and C) and various bioactive compounds (phenolic compounds and carotenoids) that are beneficial for the health of consumers ^[9]. Additionally, scientific evidence shows that bioactive compounds extracted from bell peppers have anti-inflammatory, antidiabetic, antimicrobial, and immunomodulatory effects, among others ^[10][11][12].

2. Description of Bell Peppers

Capsicum annuum is an annual or biennial herbaceous plant. Its lifecycle comprises four phases: seedling, vegetation, flowering and fruiting ^[13]. The bell pepper fruit is large and fleshy, quadrangular, of variable size (7–16 cm long/6–11 cm width) and weight (from 100 to 500 g). The consumers appreciate them for their exotic colors (green, red, yellow, and orange), flavor, and texture. Furthermore, they are usually consumed fresh, but are also frequently used to enhance food dishes or other food products ^{[14][15]}. Moreover, they can be processed in commercial products, such as sauces, puree, and powders, among others ^[16]. On the other hand, bell pepper fruits could be a good source of human nutrition and health by providing energy and bioactive compounds ^[17], as discussed below.

According to the literature, bell peppers have high levels of water and carbohydrates with a low protein and fat content, which makes them a low-calorie food; they also have an adequate content of dietary fiber to be considered a food that is high in fiber, which has important implications for the health and nutrition levels of consumers. Additionally, bell peppers contain some nutritionally important compounds such as vitamins (B, A, D, C, E, and K) and minerals (potassium, sodium, magnesium, calcium, and phosphorus) ^[18]. In this sense, the frequent consumption of bell peppers provides essential nutrients for human health ^{[19][20][21]}. For example, fresh bell pepper consumption (100 g) provides the total ascorbic acid recommended daily intake ^[22]. On the other hand, the nutritional content of bell peppers depends directly on the color of the fruit, growing conditions, and postharvest processing, among other factors ^[23].

Additionally, the total pepper production has increased significantly (25%) in recent years (from 2006 to 2016) ^[24], and it is one of the most commercially cultivated vegetable crops worldwide ^[25]. For example, in Mexico, the total production of bell peppers was 676,216 tons during 2019 ^[13]. Therefore, the recovery of bell pepper phytochemicals offers a viable strategy to obtain bioactive compounds, which could be used as natural ingredients for the food and pharmaceutical industries, as an alternative to replacing synthetic compounds and also in the revalorization of a plant's wastes and losses ^[26].

3. Phytochemicals Present in Bell Peppers

In general, the concentration of polyphenol compounds in bell peppers varies with variety and color, ranging from 5.59 to 52.65 mg of gallic acid equivalent per gram of edible portion ^{[18][19][20]}. The main phenolic compounds found in green bell peppers include myricetin (658 μ g/g), pyrogallol (572 μ g/g), chlorogenic acid (290 μ g/g), catechol (279 μ g/g), protocatechuic acid (116 μ g/g), caffeic acid (108 μ g/g), ellagic acid (106 μ g/g), and gallic acid (89 μ g/g) ^[19]^{[23][27]}. In red bell pepper, pyrogallol (757 μ g/g), P-OH benzoic acid (395 μ g/g), myricetin (244 μ g/g), chlorogenic acid (221 μ g/g), ellagic acid (172 μ g/g), gallic acid (115 μ g/g), and benzoic acid (111 μ g/g) ^{[19][22][23][27]} were found; while in yellow bell pepper, pyrogallol (2175 μ g/g), catechol (225 μ g/g), ellagic acid, benzoic acid (173 μ g/g), myricetin (151 μ g/g), ellagic acid (144 μ g/g), chlorogenic acid (136 μ g/g), and P-OH benzoic acid (123 μ g/g) ^{[19][23]} were found. Furthermore, it has been reported that orange bell pepper contains gallic acid (900 μ g/g), chlorogenic acid (117 μ g/g), myricetin (100 μ g/g), resveratrol (89 μ g/g), caffeic (38 μ g/g), ferulic (13.45 μ g/g) and p-coumaric acids (13.45 μ g/g) ^{[25][28][27]}. Additionally, the presence of gallic acid has been reported in purple (1200 μ g/g) and

dark violet (1150 μ g/g) bell peppers ^[27]. These compounds exert potent antioxidant activity against reactive oxygen species and reactive nitrogen species ^[29].

Evidence indicates that bell pepper fruits are rich in phenolic compounds and flavonoids that may improve the human health status ^{[18][19]}; moreover, there is an association between uptake diets rich in phytochemicals and the risk reduction of chronic non-communicable diseases such as diabetes, osteoporosis, and cancer ^[26]. In most cases, the biological effects of polyphenols and flavonoids have been attributable to their antioxidant capacity, which can mitigate oxidative stress ^[23]. Thus, the regular consumption of bell peppers can improve human health and prevent degenerative diseases ^{[30][31]}.

In general, the concentration of carotenoids in bell pepper depends on their color ^[22] and ripening state ^[32], where the highest concentration was reported in red bell pepper (7137-8800 µg/g), followed by orange (5292 µg/g), yellow (2236.3–2834 μ g/g), and green (1219–1513.5 μ g/g) peppers [19][22][23][33]. The main carotenoids reported in green peppers include neoxanthin (190 µg/g), chlorophyll (150 µg/g), lutein (76 µg/g), zeaxanthin (35 µg/g), and capsanthin (16 µg/g) [18][20][34][32], while, in red bell peppers, the most reported carotenoids were 5.6,-epoxide capsanthin (513 μg/g), lycopene (322 μg/g), capsanthin (178 μg/g), cucurbitaxanthin (81 μg/g), and Zeaxanthin (70 $\mu q/q$) [18][23][34][32][35][36]. In yellow bell peppers, the most common were lutein (115 $\mu q/q$), chlorophyll (61 $\mu q/q$), zeaxanthin (48 μ g/g), capsanthin (45 μ g/g), and α -carotene (21 μ g/g) [18][20][37][32][38]. Moreover, the main carotenoids reported in orange bell peppers were zeaxanthin (191 μg/g), β-carotene (56 μg/g), lutein (45 μg/g), capsanthin (45 μ g/g), and β -cryptoxanthin (19 μ g/g) [18][23][37][32][38]. Carotenoids are excellent antioxidant compounds with several human health benefits; their consumption may prevent coronary heart diseases and some types of cancer (gastrointestinal, lung, prostate, and breast), and can reduce the risk of age-related macular degeneration, as well as having a beneficial effect on cognitive function ^[35]. However, specific carotenoids may provide specific health benefits; for example, α - carotene, β -carotene, and β - cryptoxanthin are provitamin A compounds ^[39]. Moreover, β-carotene has positive effects on cognitive functions, while lutein and zeaxanthin provide eye protection ^[39]. Furthermore, lycopene exhibited potent antioxidant activity and may reduce cholesterol in animals [39]. According to these data, the regular consumption of bell pepper fruits may improve human health.

Additionally, Dias-Games et al. ^[40] isolated a Hevein-like peptide from the leaves of bell pepper crops. These compounds showed antibacterial and antifungal activities and exhibited great potential for biotechnological use ^[41].

4. Biological Activities of Bell Pepper Extracts

As discussed in the preceding sections, bell pepper fruits, seeds, and leaves contain bioactive compounds (phenolic, flavonoid, carotenoids, tocopherol, and pectic polysaccharides), which are associated with different biological activities for diverse applications, as described below.

The antioxidant activity of any fruit or vegetable is determined by different bioactive compounds, showing different action mechanisms to inhibit radicals. Therefore, more than one method should be used to clarify the mode of

action of each extract or compound from bell peppers. In this context, the antioxidant properties of bell peppers are an important parameter for establishing their health functionality.

Park et al. ^[42] investigated the antioxidant activity of methanolic extracts from four different-colored bell peppers by different methods (ABTS, DPPH, and SOD-like activity). They informed that extracts from orange bell peppers showed the highest antioxidant activity by ABTS test (880 µmol TE/g) compared to the red (800 µmol TE/g), yellow (790 µmol TE/g), and green (630 µmol TE/g) bell peppers; in contrast, the green bell pepper exhibited the highest antioxidant activity by DPPH assay (1153 µg/mL) compared to red (882 µg/mL), yellow (811 µg/mL), and orange (694 µg/mL) bell peppers. Furthermore, the green bell pepper showed the highest SOD-like activity (IC 50 = 1472 µg/mL) compared to the yellow (IC 50 = 1676 µg/mL), red (IC 50 = 1826 µg/mL), and orange (IC 50 = 1893 µg/mL) bell peppers. The authors argued that differences in results in colored bell peppers are attributable to the mode of action of each bioactive compound present in the extract and their ability to reduce or inhibit oxidative stress, which is dependent on the color of the fruit. Additionally, the extracts from the four bell peppers exhibited a protective effect against H 2O 2- and HNE-induced DNA damage at low concentrations (1 µg/mL), preventing cellular damage at physiological levels. These compounds could form a reversible complex that may act as a desmutagenic molecule or interceptor, inhibiting genotoxicity ^[42].

In an other hand, it has been reported that the extracts from bell pepper fruits or leaves exhibited antibacterial and antifungal activities, which could be potentially used for food and pharmaceutical applications [^[1][2][3][53][54][55][56][57] [58]]. Also, it has been studied the effects of bell pepper extracts on diverse pathologies, showing ^{[43][44][45][46][47][48]}it ´s anti-hyperglycemic, cytotoxic, and neuroprotective effects (Table 1)

Pathology	Bell Pepper Color	Source	Bioactive Extracts or Compounds	Dose	Model Assay	Effect	Ref.
Diabetes	Green	Fruit juice	Whole juice/ethanol extracts	50 mg/mL	α-glucosidase inhibitory activity	Extract exhibited α- glucosidase inhibitory effects.	[<u>49</u>]
	Green	Fruit juice	Ethanol extracts	100 µg/mL	growth of 3T3-L1 preadipocytes	Extract increased the survive rate of preadipocyte cells	[49]
	Green	Fruit juice	Ethanol extracts	100 μg/mL	3T3-L1 differentiation into adipocytes induced	Extracts promote the 3T3-L1 cells	[<u>49]</u>

Table 1 Effects of bell pepper extracts on diverse pathologies

Pathology	Bell Pepper Color	Source	Bioactive Extracts or Compounds	Dose	Model Assay	Effect	Ref.
			-			differentiation rate	
	NI	Fruit juice	Fruit juice	100 mL/twice a day	Randomized controlled study in humans	Fruit juice reduces post- prandial blood glucose and blood pressure.	[50]
	Red	Fruit pulp	Ethyl acetate extracts	20 µL	In vitro in HeLa cells	Extract inhibited the protein islet amyloid polypeptide.	[<u>51</u>]
	Red	Fruit pulp	Extract mixed with virgin olive oil	2 to 8 mL/kg body weight	In vivo animal assay in adult male rats	The mixture inhibited amylase and α-glucosidase activity.	[<u>52</u>]
	Green Red Yellow	NI	methanol extract	NI	NI	Extracts had α- glucosidase- inhibitory effects.	[<u>53]</u>
Cancer	NI	Powdered	Aqueous extracts	10% <i>v\w</i>	animal assay with <i>Drosophila</i> larval (SMART assay)	Aqueous extracts showed antimutagenic activity.	[<u>54</u>]
	Red	Fruit pulp	Methanol extract	125 μg/mL	In vitro in NIH3T3 and A549 cells	The extract exhibited strong cytotoxicity in A549 cells.	[55]
	Yellow	Fruit pulp	Methanol extract	125 µg/mL	In vitro in NIH3T3 and A549 cells	Selective cytotoxic activity against A549 cells.	[<u>55</u>]
	Green	Fruit pulp	Pectic polysaccharides	150 mg/kg	In vivo animal model in Ehrlich tumor- bearing mice	Significantly reduced tumor growth.	[<u>56</u>]

Pathology	Bell Pepper Color	Source	Bioactive Extracts or Compounds	Dose	Model Assay	Effect	Ref.
	Green	Fruit pulp	Pectic polysaccharides	0.1 mg/mL	In vitro in lineages of human mammary cancer cells (MCF-7, MDA-MB-231, and MDA-MB-436)	Selective cytotoxic activity against MCF- 7, MDA-MB- 231, and MDA-MB-436 cell lines.	[<u>56</u>]
	Green Yellow Red	Fruit pulp	Polyphenol mixtures	1.2 mg/L	In vitro in human gastric adenocarcinoma cells, A549 human lung carcinoma cells, and HeLa human cervical carcinoma cells	Extracts showed cytotoxic effects against all cancer cell lines in a dose- dependent response.	[<u>57</u>]
Alzheimer's disease	NI	Powdered	Aqueous extracts / E- capsiate, Z- capsiate, dihydrocapsiate and nor- dihydrocapsiate,	1–10 g/L	In vitro Peptides agregation test	Bell pepper extracts were able to inhibit b- secretase activity and aggregation of Ab1–40 peptides	[<u>58</u>]

NI: no information.

5. Concluding Remarks

Sustainability in food production and consumption benefits all food chain and the environment. Efficient food production and supply chains lead to less food loss is only part of sustainability. The other part is the use through the revalorization of waste and food losses. The valorization of food waste for edible purposes is an area of opportunity for researchers due to the significant content of biactive compounds present, which can be reused in numerous products to which they will add value and thereby reducing the generation of waste food.

The revalorization of waste from fruits and vegetables, particularly bell pepper, mainly focuses on the use of bioactive compounds whose presence depends on the variety, color, growing conditions, degree of maturity at harvest and post-harvest handling of the fruit; in addition to the source (seeds or leaves). In general, the bioactive compounds reported in bell pepper are phenolic compounds, flavonoids, carotenoids, tocopherol and pectic polysaccharides, which have antioxidant, antimicrobial, immunomodulatory activity. Besides having positive effects

in the treatment of diseases such as diabetes mellitus, cancer and Alzheimer's. Also, bell pepper extracts can be used as food additives (preservatives and colorants).

In this context, the revalorization of food waste is an area of opportunity for research and technological innovation with beneficial effects for the population, the economy and the environment. However, most studies have been carried out in vitro, so in vivo investigations are also necessary to demonstrate the effectiveness of the aforementioned compounds, in addition to guaranteeing their safety and understanding their mechanisms of action.

References

- 1. Viaggi, D. The Bioeconomy: Delivering Sustainable Green Growth; CABI Publishing: Boston, MA, USA, 2018.
- 2. Malorgio, G.; Marangon, F. Agricultural business economics: The challenge of sustainability. Agric. Food Econ. 2021, 9, 6.
- 3. Brunori, G. Biomass, Biovalue and Sustainability: Some Thoughts on the Definition of the Bioeconomy. EuroChoices 2013, 12, 48–52.
- Sawatdeenarunat, C.; Nguyen, D.; Surendra, K.C.; Shrestha, S.; Rajendran, K.; Oechsner, H.; Xie, L.; Khanal, S.K. Anaerobic biorefinery: Current status, challenges, and opportunities. Bioresour. Technol. 2016, 215, 304–313.
- Budzianowski, W.M. High-value low-volume bioproducts coupled to bioenergies with potential to enhance business development of sustainable biorefineries. Renew. Sustain. Energy Rev. 2017, 70, 793–804.
- 6. Mohan, S.V.; Butti, S.K.; Amulya, K.; Dahiya, S.; Modestra, J.A. Waste Biorefinery: A New Paradigm for a Sustainable Bioelectro Economy. Trends Biotechnol. 2016, 34, 852–855.
- Smith, P.; Calvin, K.; Nkem, J.; Campbell, D.; Cherubini, F.; Grassi, G.; Korotkov, V.; Le Hoang, A.; Lwasa, S.; McElwee, P.; et al. Which practices co-deliver food security, climate change mitigation and adaptation, and combat land degradation and desertification? Glob. Chang. Biol. 2020, 26, 1532–1575.
- Scoma, A.; Rebecchi, S.; Bertin, L.; Fava, F. High impact biowastes from South European agroindustries as feedstock for second-generation biorefineries. Crit. Rev. Biotechnol. 2016, 36, 175– 189.
- European Commission Closing the Loop—An EU Action Plan for the Circular Economy COM/2015/0614 Final—European Environment Agency. Available online: https://www.eea.europa.eu/policy-documents/com-2015-0614-final (accessed on 26 July 2021).

- 10. Samtiya, M.; Aluko, R.E.; Dhewa, T.; Moreno-Rojas, J.M. Potential health benefits of plant foodderived bioactive components: An overview. Foods 2021, 10, 839.
- Sagar, N.A.; Pareek, S.; Sharma, S.; Yahia, E.M.; Lobo, M.G. Fruit and vegetable waste: Bioactive compounds, their extraction, and possible utilization. Compr. Rev. Food Sci. Food Saf. 2018, 17, 512–531.
- Coman, V.; Teleky, B.E.; Mitrea, L.; Martău, G.A.; Szabo, K.; Călinoiu, L.F.; Vodnar, D.C. Bioactive potential of fruit and vegetable wastes. In Advances in Food and Nutrition Research; Academic Press: Cambridge, MA, USA, 2020; Volume 91, pp. 157–225. ISBN 9780128204702.
- Servicio Nacional de Inspección y Certificación de Semillas Chile (Capsicum spp.). Available online: https://www.gob.mx/snics/acciones-y-programas/chile-capsicum-spp (accessed on 17 July 2021).
- Padilha, H.K.M.; Pereira, E.D.S.; Munhoz, P.C.; Vizzotto, M.; Valgas, R.A.; Barbieri, R.L. Genetic variability for synthesis of bioactive compounds in peppers (Capsicum annuum) from Brazil. Food Sci. Technol. 2015, 35, 516–523.
- 15. Ribes-Moya, A.M.; Raigón, M.D.; Moreno-Peris, E.; Fita, A.; Rodríguez-Burruezo, A. Response to organic cultivation of heirloom Capsicum peppers: Variation in the level of bioactive compounds and effect of ripening. PLoS ONE 2018, 13, e0207888.
- 16. Šeregelj, V.; Tumbas Šaponjac, V.; Lević, S.; Kalušević, A.; Ćetković, G.; Čanadanović-Brunet, J.; Nedović, V.; Stajčić, S.; Vulić, J.; Vidaković, A. Application of encapsulated natural bioactive compounds from red pepper waste in yogurt. J. Microencapsul. 2019, 36, 704–714.
- Dagadkhair, R.A.; Shinde, D.B.; Shelke, S.A.; Kale, K.B. Vegetables & Its Health Benefits, 1st ed.; Dagadkhair, R.A., Shinde, D.B., Shelke, S.A., Kale, K.B., Eds.; Taurean Publications: Nehru Place, India, 2020; ISBN 9788194488880.
- Thuphairo, K.; Sornchan, P.; Suttisansanee, U. Bioactive compounds, antioxidant activity and inhibition of key enzymes relevant to Alzheimer's disease from sweet pepper (Capsicum annuum) extracts. Prev. Nutr. Food Sci. 2019, 24, 327–337.
- 19. Abdalla, M.U.E.; Taher, M.; Sanad, M.I.; Tadros, L.K. Chemical properties, phenolic profiles and antioxidant activities of pepper fruits. J. Agric. Chem. Biotechnol. 2019, 10, 133–140.
- 20. Kaur, R.; Kaur, K. Preservation of sweet pepper purees: Effect on chemical, bioactive and microbial quality. J. Food Sci. Technol. 2021, 58, 3655–3660.
- Rahim, R.; Mat, I. Phytochemical contents of Capsicum frutescens (Chili Padi), Capsicum annum (Chili Pepper) and Capsicum annum (Bell Peper) aqueous extracts. Int. Conf. Biol. Life Sci. 2012, 40, 164–167.

- 22. Zhuang, Y.; Chen, L.; Sun, L.; Cao, J. Bioactive characteristics and antioxidant activities of nine peppers. J. Funct. Foods 2012, 4, 331–338.
- Blanco-Ríos, A.K.; Medina-Juárez, L.Á.; González-Aguilar, G.A.; Gámez-Meza, N. Antioxidant activity of the phenolic and oily fractions of different sweet bell peppers. J. Mex. Chem. Soc. 2013, 57, 137–143.
- 24. FAO. World Food and Agriculture—Statistical Yearbook 2020; FAO: Rome, Italy, 2020.
- 25. FAO. Save Food: Global Initiative on Food Loss and Waste Reduction; Food and Agriculture Organization United Nations: Rome, Italy, 2016; pp. 01–02.
- Baenas, N.; Belović, M.; Ilic, N.; Moreno, D.A.; García-Viguera, C. Industrial use of pepper (Capsicum annum L.) derived products: Technological benefits and biological advantages. Food Chem. 2019, 274, 872–885.
- 27. Ghasemnezhad, M.; Sherafati, M.; Payvast, G.A. Variation in phenolic compounds, ascorbic acid and antioxidant activity of five coloured bell pepper (Capsicum annum) fruits at two different harvest times. J. Funct. Foods 2011, 3, 44–49.
- 28. Almadhoun, H.R. Bell pepper Classification using Deep Learning. IJAER 2021, 5, 75–79.
- 29. Guevara, L.; Domínguez-Anaya, M.Á.; Ortigosa, A.; González-Gordo, S.; Díaz, C.; Vicente, F.; Corpas, F.J.; Pérez Del Palacio, J.; Palma, J.M. Identification of compounds with potential therapeutic uses from sweet pepper (Capsicum annuum L.) fruits and their modulation by nitric oxide (no). Int. J. Mol. Sci. 2021, 22, 4476.
- Oboh, G.; Ademiluyi, A.O.; Faloye, Y.M. Effect of combination on the antioxidant and inhibitory properties of tropical pepper varieties against α-amylase and α-gflucosidase activities in vitro. J. Med. Food 2011, 14, 1152–1158.
- González-García, Y.; Cárdenas-álvarez, C.; Cadenas-Pliego, G.; Benavides-Mendoza, A.; Cabrera-De-la-fuente, M.; Sandoval-Rangel, A.; Valdés-Reyna, J.; Juárez-Maldonado, A. Effect of three nanoparticles (Se, Si and Cu) on the bioactive compounds of bell pepper fruits under saline stress. Plants 2021, 10, 217.
- 32. Howard, L.R.; Talcott, S.T.; Brenes, C.H.; Villalon, B. Changes in phytochemical and antioxidant activity of selected pepper cultivars (Capsicum species) as influenced by maturity. J. Agric. Food Chem. 2000, 48, 1713–1720.
- González-Saucedo, A.; Barrera-Necha, L.L.; Ventura-Aguilar, R.I.; Correa-Pacheco, Z.N.; Bautista-Baños, S.; Hernández-López, M. Extension of the postharvest quality of bell pepper by applying nanostructured coatings of chitosan with Byrsonima crassifolia extract (L.) Kunth. Postharvest Biol. Technol. 2019, 149, 74–82.

- 34. Marín, A.; Gil, M.I.; Flores, P.; Hellín, P.; Selma, M.V. Microbial quality and bioactive constituents of sweet peppers from sustainable production systems. J. Agric. Food Chem. 2008, 56, 11334–11341.
- 35. Chávez-Mendoza, C.; Sánchez, E.; Carvajal-Millán, E.; Munoz-Márquez, E.; Guevara-Aguilar, A. Characterization of the nutraceutical quality and antioxidant activity in bell pepper in response to grafting. Molecules 2013, 18, 15689–15703.
- Navarro, J.M.; Flores, P.; Garrido, C.; Martinez, V. Changes in the contents of antioxidant compounds in pepper fruits at different ripening stages, as affected by salinity. Food Chem. 2006, 96, 66–73.
- Hallmann, E.; Marszałek, K.; Lipowski, J.; Jasińska, U.; Kazimierczak, R.; Średnicka-Tober, D.; Rembiałkowska, E. Polyphenols and carotenoids in pickled bell pepper from organic and conventional production. Food Chem. 2019, 278, 254–260.
- Guil-Guerrero, J.L.; Martínez-Guirado, C.; Del Mar Rebolloso-Fuentes, M.; Carrique-Pérez, A. Nutrient composition and antioxidant activity of 10 pepper (Capsicum annuun) varieties. Eur. Food Res. Technol. 2006, 224, 1–9.
- 39. Eggersdorfer, M.; Wyss, A. Carotenoids in human nutrition and health. Arch. Biochem. Biophys. 2018, 652, 18–26.
- Games, P.D.; da Silva, E.Q.G.; de Oliveira Barbosa, M.; Almeida-Souza, H.O.; Fontes, P.P.; de Magalhães, M.J., Jr.; Pereira, P.R.G.; Prates, M.V.; Franco, G.R.; Faria-Campos, A.; et al. Computer aided identification of a Hevein-like antimicrobial peptide of bell pepper leaves for biotechnological use. BMC Genom. 2016, 17, 999.
- Games, P.D.; Koscky-Paier, C.R.; Almeida-Souza, H.O.; Barbosa, M.O.; Antunes, P.W.P.; Carrijo, L.C.; Pereira, P.R.G.; Baracat-Pereira, M.C. In vitro anti-bacterial and anti-fungal activities of hydrophilic plant defence compounds obtained from the leaves of bell pepper (Capsicum annuum L.). J. Hortic. Sci. Biotechnol. 2013, 88, 551–558.
- Park, J.H.; Jeon, G.I.; Kim, J.M.; Park, E. Antioxidant activity and antiproliferative action of methanol extracts of 4 different colored bell peppers (Capsicum annuum L.). Food Sci. Biotechnol. 2012, 21, 543–550.
- Dorantes, L.; Colmenero, R.; Hernandez, H.; Mota, L.; Jaramillo, M.E.; Fernandez, E.; Solano, C. Inhibition of growth of some foodborne pathogenic bacteria by Capsicum annum extracts. Int. J. Food Microbiol. 2000, 57, 125–128.
- 44. Hu, X.; Saravanakumar, K.; Jin, T.; Wang, M.H. Effects of yellow and red bell pepper (paprika) extracts on pathogenic microorganisms, cancerous cells and inhibition of survivin. J. Food Sci. Technol. 2021, 58, 1499–1510.

- 45. López-Muñoz, N.R.; Romero-Bastidas, M.; Arce-Amézquita, P.M.; Hernández-Rubio, J.S. Antifungal activity of antioxidants derived from four cultivars of Capsicum spp. against phytopathogenic fungi. Ecosistemas Recur. Agropecu. 2019, 6, 487–498.
- da Silva França, K.R.; Paiva, Y.F.; de Azevedo, P.T.M.; da Nóbrega, L.P.; da Silva, E.V.; Cardoso, T.A.L. Extratos de pimentão vermelho (Capsicum annuum) sobre Colletotrichum gloeosporioides in vitro. Rev. Verde Agroecol. Desenvolv. Sustentável 2019, 14, 382–388.
- 47. Careaga, M.; Fernández, E.; Dorantes, L.; Mota, L.; Jaramillo, M.E.; Hernandez-Sanchez, H. Antibacterial activity of Capsicum extract against Salmonella typhimurium and Pseudomonas aeruginosa inoculated in raw beef meat. Int. J. Food Microbiol. 2003, 83, 331–335.
- 48. Aljaloud, S.O.; Gyawali, R.; Reddy, M.R.; Ibrahim, S.A. Antibacterial activity of red bell pepper against Escherichia coli O157:H7 in ground beef. Internet J. Food Saf. 2012, 14, 44–47.
- Park, J.H.; Kim, R.Y.; Park, E. Antidiabetic activity of fruits and vegetables commonly consumed in Korea: Inhibitory potential against α-glucosidase and insulin-like action in vitro. Food Sci. Biotechnol. 2012, 21, 1187–1193.
- 50. Nagasukeerthi, P.; Mooventhan, A.; Manjunath, N.K. Short-term effect of add on bell pepper (Capsicum annuum var. grossum) juice with integrated approach of yoga therapy on blood glucose levels and cardiovascular functions in patients with type 2 diabetes mellitus: A randomized controlled study. Complement. Ther. Med. 2017, 34, 42–45.
- 51. Fuentes, A.L.; Hennessy, K.; Pascual, J.; Pepe, N.; Wang, I.; Santiago, A.; Chaggan, C.; Martinez, J.; Rivera, E.; Cota, P.; et al. Identification of plant extracts that inhibit the formation of diabeteslinked IAPP amyloid. J. Herb. Med. 2016, 6, 37–41.
- 52. Al-Obaidi, W.M.L. Study of synergistic inhibitory effectiveness of mixed of sweet bell pepper extract and virgin olive oil in α-amylase and α-glucosidase activity in serum of male rats infected with experimental diabetes. Tikrit J. Pure Sci. 2015, 20, 90–96.
- 53. Shukla, S.; Kumar, D.A.; Anusha, S.V.; Tiwari, A.K. Antihyperglucolipidaemic and anticarbonyl stress properties in green, yellow and red sweet bell peppers (Capsicum annuum L.). Nat. Prod. Res. 2016, 30, 583–589.
- 54. El Hamss, R.; Idaomar, M.; Alonso-Moraga, A.; Muñoz Serrano, A. Antimutagenic properties of bell and black peppers. Food Chem. Toxicol. 2003, 41, 41–47.
- 55. Hu, X.; Saravanakumar, K.; Jin, T.; Wang, M.H. Effects of yellow and red bell pepper (paprika) extracts on pathogenic microorganisms, cancerous cells and inhibition of survivin. J. Food Sci. Technol. 2021, 58, 1499–1510.
- 56. Adami, E.R.; Corso, C.R.; Turin-Oliveira, N.M.; Galindo, C.M.; Milani, L.; Stipp, M.C.; do Nascimento, G.E.; Chequin, A.; da Silva, L.M.; de Andrade, S.F.; et al. Antineoplastic effect of

pectic polysaccharides from green sweet pepper (Capsicum annuum) on mammary tumor cells in vivo and in vitro. Carbohydr. Polym. 2018, 201, 280–292.

- 57. Jeong, W.Y.; Jin, J.S.; Cho, Y.A.; Lee, J.H.; Park, S.; Jeong, S.W.; Kim, Y.H.; Lim, C.S.; Abd El-Aty, A.M.; Kim, G.S.; et al. Determination of polyphenols in three Capsicum annuum L. (bell pepper) varieties using high-performance liquid chromatographytandem mass spectrometry: Their contribution to overall antioxidant and anticancer activity. J. Sep. Sci. 2011, 34, 2967–2974.
- 58. Ogunruku, O.O.; Oboh, G.; Passamonti, S.; Tramer, F.; Boligon, A.A. Capsicum annuum var. grossum (Bell Pepper) inhibits β -secretase activity and β -amyloid1-40 aggregation. J. Med. Food 2017, 20, 124–130.

Retrieved from https://encyclopedia.pub/entry/history/show/36147