

Vegetation Optical Depth

Subjects: **Geography**

Contributor: Frédéric Frappart , Jean-Pierre Wigneron , Xiaojun Li , Xiangzhuo Liu , Amen Al-Yaari , Lei Fan , Mengjia Wang , Christophe Moisy , Erwan Le Masson , Zacharie Aoulad Lafkih , Clément Vallé , Bertrand Ygorra , Nicolas Baghdadi

The vegetation optical depth (VOD) parameterizes the extinction (attenuation & scattering) effects due to the vegetation affecting the microwave radiations propagating through the vegetation canopy. It is retrieved from both passive and active microwave remotely sensed observations. It is related almost linearly to the vegetation water content and indirectly to vegetation water status and biomass. It has been retrieved from several satellites operating at different frequencies and provides a long-term record of the vegetation dynamics over land surfaces since 1978. VOD is complementary to classically used vegetation indices derived from multi-spectral images but it is less affected by sun illumination and atmospheric effects than optical vegetation indices.

vegetation optical depth (VOD)

brightness temperature (TB)

backscattering

radiative transfer

vegetation

above ground biomass (AGB)

vegetation water content (VWC)

1. Introduction

Vegetation is a key element in the energy, water and carbon balances over the land surfaces and is strongly impacted by climate change and anthropogenic effects. Remotely sensed observations are commonly used for the monitoring of vegetation dynamics and its temporal changes from regional to global scales. Among the different indices derived from Earth observation satellites to study the vegetation, the vegetation optical depth (VOD), which is related to the intensity of extinction effects within the vegetation canopy layer in the microwave domain and which can be derived from both passive and active microwave observations, is increasingly used for monitoring a wide range of ecological vegetation variables. Based on different frequency bands used to derive VOD, from L- to Ka-bands, these variables include, among others, the vegetation water content/status and the above ground biomass for the top-layer canopy (at high frequencies) and the whole canopy (at lower frequencies as L-band)

2. VOD retrievals from active and passive microwave observations

VOD can be retrieved from both passive and active microwave data ^[1]. Radiometers measure the thermal radiation emitted by the Earth's surface at a given frequency. It is given in terms of brightness temperature (T_B) representing the temperature of an equivalent blackbody emitting the same radiation ^[2]. Emission of land surfaces (i.e., a rough soil covered with vegetation) in the microwave electromagnetic (EM) domain can be simulated using a radiative transfer (RT) model ^[3]. The brightness temperature measured by the sensor onboard a satellite platform is the linear combination of effective temperatures T_{soil} and T_{veg} of soil and vegetation, and the downwelling cold sky T_{sky}

weighted by EM dependent emissivities. Experimental studies have revealed a good linear relationship between VOD and vegetation water content (VWC, kg/m²) over crops fields (VOD ~ b. VWC, where b is depending on vegetation type and structure, growth stage, water status, etc.) [4][5]. 2-Parameter retrievals of soil moisture and VOD from multi-angular L-band observations has been first demonstrated from in situ data sets [6][7]. Following studies based on the L-MEB model (L-band microwave emission of the biosphere)[7], were extended to simulated [8] and actual SMOS observations [9]. Multi-orbit and/or multi-angular retrievals of VOD, and SM or other vegetation or soil parameters (soil roughness, effective scattering albedo) have also been developed [8][10][25].

Radars measure the power of the electromagnetic wave (EM) backscattered by the Earth's surface (P_r). The reflecting properties of the surface are expressed through the backscattering coefficient (σ^0) which is function of the emitted and reflected powers (P_e and P_r respectively), the characteristics of the antenna and of the illuminated area. Numerous models were developed to simulate the radar backscattering of land surfaces in the active domain, VOD is one of the variable used to describe the effects of vegetation in the backscattering of a soil-vegetation media. The Water Cloud Model (WCM) expressed the total backscattering of the land surface as the sum of the direct backscattering of the vegetation, the backscattering of the soil twice attenuated by the vegetation cover and the multiple scattering from soil-vegetation interactions.

3. Advantages and Drawbacks of VOD

Compared to vegetation indices derived from reflectance acquired by sensors operating in the visible and the infra-red domains, VOD is not affected by the presence of clouds and less sensitive to the presence of water in the atmosphere. Thus, the different microwave sensors are likely to provide reliable information related to VWC on both the green and non-green vegetation components [11], at a quasi-daily temporal resolution, at different frequencies and over several decades [12][13][14]. VOD gives access to complementary information to NDVI or LAI on the vegetation dynamics. For instance, significant time-lags were observed between LAI and LPRM (AMSR-E) VOD in the Amazon rainforest [15]. It was observed that the growth of the biomass canopy and the phenological activity exhibit maximum time-lags in areas characterized by shorter dry seasons in the west of the basin. This time-lag progressively decreases along a west-to-east longitudinal wetness gradient as the length of the dry season increases indicating an adaptation of forest to water and light availability [16]. In a recent study, a time-lag of three to six months between LAI and SMOS-IC L-VOD was determined in the savannah/woodland region of Miombo, in southern Africa. The latter one exhibits a peak at the end of the rainy season and in beginning of the dry season that was attributed to tree growth during this period and was confirmed by in situ observations [17]. X-band VOD is strongly correlated to leaf water potential [18]. As VWC is related to leaf water potential (Ψ_L), the differences between VOD estimates at mid—day and midnight was used to derive an isohydricity parameter at ecosystem-scale [19]. As isohydricity is an indicator of the stomatal and xylem flow regulation, which is an important factor in drought-induced reduction of the biomass and plant mortality that cannot be derived from land cover type, VOD is a good candidate for analyzing the spatio-temporal responses of vegetation to water stress. VOD presents saturation for higher biomass values than NDVI, the saturation occurring for lower biomass values as the frequency increases [20]. L-band VOD from SMOS VOD was reported to reach saturation for biomass around 400 Mg ha⁻¹ in equatorial forests of South America and Africa [21][22]. In spite of differences in spatio-temporal patterns between

VODs derived from different sensor types, and at various frequencies and, for different inversion approaches (see for instance [Figure 1](#)), very few inter-comparison studies have been achieved yet. A first study based on one year (2003) of AMSR-E T_B acquired at C- (6.925 GHz), X- (10.65 GHz), and K -(18.7 GHz) bands showed a greater sensitivity of microwave data acquired at higher frequencies to the photosynthetically active component of vegetation (crown and leaves), whereas, T_B acquired at lower frequencies are more sensitive to the woody component of the vegetation. These results were confirmed when comparing L- (SMAP MT-DCA data), X- and C- (AMSR-2 LPRM version 5 data) bands VOD to AGB over tropical forests [\[23\]](#). Comparisons between different L-band VOD from SMOS (i.e., SMOS Level 2, SMOS Level 3 and SMOS-IC v105) and three different AGB datasets, as well as three height NDVI and EVI also exhibited very good agreement between the different parameters. Among the L-VOD products, SMOS-IC obtained the higher correlations and lower dispersion against all the evaluated VOD datasets [\[24\]](#).

Figure 1. (a) Average Vegetation Optical Depth (VOD) over Africa from SMOS-IC v105 for 2016 (left) and corresponding zonal average (right); (b) Same for AMSR-2 from Land Parameter Retrieval Model (LPRM) v05; (c) Same for Advanced Scatterometer (ASCAT) from TU Wien.

A large intercomparison was recently made and highlighted the large complementarity between VODs from different sensors and at different frequencies for monitoring vegetation [\[25\]](#). High frequency VODs are better correlated to optical indices (in time and space) and low frequency VODs are better correlated to biomass; but this general rule has many exceptions depending on sensors and products

The major drawbacks of the current VOD products from radiometers and scatterometers are their sensitivity to RFI, the loss of accuracy in presence of water bodies in the sensor footprint, and their coarse spatial resolutions.

Concerning the important issue of the RFI, preprocessing approaches are increasingly improving to filter out the data impacted by these perturbations. For example, in the version 2 of the SMOS-IC data, available at [26], a special attention was devoted to the prefiltering step allowing to improve the quality of the VOD retrievals compared to v105 [25]. To tackle with the loss of accuracy of VOD in presence of open water, the solution currently used is to filter out pixels containing a percentage of water higher than a threshold. Corrections are difficult as they would require very accurate and Near Real Time (NRT) maps of the water bodies. But inundation fractions are retrieved from SSM/I [27][28] and SMOS [29] for instance. A possible solution would be to retrieve simultaneously the surface water fraction, SM and VOD combining information from several frequencies, and/or polarizations. Improvement in the VOD spatial resolution already started with the availability of Level 1c SMAP T_B at a spatial resolution of 9 km instead of 36 km owing to the use of a Backus-Gilbert approach [30]. But no similar product has been yet developed for SMOS. A promising technique, based on the combination of successive acquisitions and the application of constraints based on the expected climatology-based range of variations of the inverted parameters, and the Sobolev-norm regularization, was recently applied on T_B from SMAP over the Contiguous United States (CONUS). This approach, denominated “combined constrained multi-channel algorithm” (C-CMCA), allowed simultaneous retrievals of SM and VOD at spatial resolutions from 1 to 5 km decreasing the RMSE values on VOD estimates by 81% and 7% respectively compared with non-constrained and single source techniques, respectively [31]. Other techniques were also implemented to other space-borne observations. Using pan-sharpening technique on T_B acquired by AMSR-E at Ka- (higher resolution) and C- (lower resolution) band, high resolution SM and VOD were obtained from LPRM at the spatial resolution of 10 km instead of 25 km [32].

4. Conclusions

VOD retrieved from both passive and active microwave remotely sensed observations has been providing a long-term record of the vegetation dynamics over land surfaces since 1978. It has been shown, in many studies that VOD offers complementary information to the commonly used vegetation indices derived from multi-spectral images as NDVI, LAI or fPAR and presents several advantages compared to these indices. It is much less affected by the presence of water in the atmosphere and is a proxy of the vegetation water content and indirectly of the biomass of the whole canopy even in the tropical forests (at low frequencies as L-band). The various frequencies available and the different sensing modes of the space-borne sensors give access to a wide range of vegetation variables as essential as the water status of the top-canopy layer at high frequencies and the biomass at low frequencies. Yet, the synergy between observations acquired for different frequencies and sensing modes (active and passive) need to be explored further to better understand the functioning of the vegetation in a context of both climate change and increased anthropogenic pressure on the vegetation ecosystems. The launch of satellite missions such as CIMR and Biomass in the years to come will ensure the continuity in the monitoring of land surfaces using passive microwave sensors and offer new information based on the use of multi-frequency bands (for CIMR) of low frequency P-band (for Biomass), respectively.

References

1. Wigneron J P; Jackson T. J. et al.; Modelling the passive microwave signature from land surfaces: A review of recent results and application to the L-band SMOS & SMAP soil moisture retrieval algorithms. *Remote Sensing of Environment* **2017**, *192*, 238-262, 10.1016/j.rse.2017.01.024.
2. Njoku, E.G.; Entekhabi, D. Passive microwave remote sensing of soil moisture. *J. Hydrol.* 1996, *184*, 101–129.
3. Mo, T.; Choudhury, B.J.; Schmugge, T.J.; Wang, J.R.; Jackson, T.J. A model for microwave emission from vegetation-covered fields. *J. Geophys. Res.* 1982, *87*, 11229–11237.
4. T.J. Jackson; T.J. Schmugge; Vegetation effects on the microwave emission of soils. *Remote Sensing of Environment* **1991**, *36*, 203-212, 10.1016/0034-4257(91)90057-d.
5. Wigneron J-P; Pardé M. et al.; Characterizing the Dependence of Vegetation Model Parameters on Crop Structure, Incidence Angle, and Polarization at L-Band. *IEEE Transactions on Geoscience and Remote Sensing* **2004**, *42*, 416-425, 10.1109/tgrs.2003.817976.
6. Wigneron J-P; Chanzy A. et al.; A simple algorithm to retrieve soil moisture and vegetation biomass using passive microwave measurements over crop fields. *Remote Sensing of Environment* **1995**, *51*, 331-341, 10.1016/0034-4257(94)00081-w.
7. Wigneron J-P; Kerr Y. et al.; L-band Microwave Emission of the Biosphere (L-MEB) Model: Description and calibration against experimental data sets over crop fields. *Remote Sensing of Environment* **2007**, *107*, 639-655, 10.1016/j.rse.2006.10.014.
8. J.-P Wigneron; Philippe Waldteufel; A. Chanzy; J.-C. Calvet; Y. Kerr; Two-Dimensional Microwave Interferometer Retrieval Capabilities over Land Surfaces (SMOS Mission). *Remote Sensing of Environment* **2000**, *73*, 270-282, 10.1016/s0034-4257(00)00103-6.
9. Fernandez-Moran R.; Al-Yaari A. A. et al.; SMOS-IC: An Alternative SMOS Soil Moisture and Vegetation Optical Depth Product. *Remote Sensing* **2017**, *9*, 457, 10.3390/rs9050457.
10. Alexandra G. Konings; Maria Piles; Narendra Das; Dara Entekhabi; L-band vegetation optical depth and effective scattering albedo estimation from SMAP. *Remote Sensing of Environment* **2017**, *198*, 460-470, 10.1016/j.rse.2017.06.037.
11. Shi, J.; Jackson, T.; Tao, J.; Du, J.; Bindlish, R.; Lu, L.; Chen, K.S. Microwave vegetation indices for short vegetation covers from satellite passive microwave sensor AMSR-E. *Remote Sens. Environ.* 2008, *112*, 4285–4300.
12. Liu, Y.Y.; van Dijk, A.I.J.M.; McCabe, M.F.; Evans, J.P.; de Jeu, R.A.M. Global vegetation biomass change (1988–2008) and attribution to environmental and human drivers. *Glob. Ecol. Biogeogr.* 2013, *22*, 692–705.
13. Jones, M.O.; Kimball, J.S.; Jones, L.A.; McDonald, K.C. Satellite passive microwave detection of North America start of season. *Remote Sens. Environ.* 2012, *123*, 324–333.

14. Guan, K.; Wood, E.F.; Medvigy, D.; Kimball, J.; Pan, M.; Caylor, K.K.; Sheffield, J.; Xu, X.; Jones, M.O. Terrestrial hydrological controls on land surface phenology of African savannas and woodlands. *J. Geophys. Res. G Biogeosci.* 2014, 119, 1652–1669.
15. Jones, M.O.; Kimball, J.S.; Jones, L.A. Satellite microwave detection of boreal forest recovery from the extreme 2004 wildfires in Alaska and Canada. *Glob. Chang. Biol.* 2013, 19, 3111–3122.
16. Jones, M.O.; Kimball, J.S.; Nemani, R.R. Asynchronous Amazon forest canopy phenology indicates adaptation to both water and light availability. *Environ. Res. Lett.* 2014, 9, 124021.
17. Tian, F.; Wigneron, J.-P.P.; Ciais, P.; Chave, J.; Ogée, J.; Peñuelas, J.; Ræbild, A.; Domec, J.-C.C.; Tong, X.; Brandt, M.; et al. Coupling of ecosystem-scale plant water storage and leaf phenology observed by satellite. *Nat. Ecol. Evol.* 2018, 2, 1428–1435.
18. Momen, M.; Wood, J.D.; Novick, K.A.; Pangle, R.; Pockman, W.T.; McDowell, N.G.; Konings, A.G. Interacting Effects of Leaf Water Potential and Biomass on Vegetation Optical Depth. *J. Geophys. Res. Biogeosci.* 2017, 122, 3031–3046.
19. Konings, A.G.; Gentine, P. Global variations in ecosystem-scale isohydricity. *Glob. Chang. Biol.* 2017, 23, 891–905.
20. Jones, M.O.; Jones, L.A.; Kimball, J.S.; McDonald, K.C. Satellite passive microwave remote sensing for monitoring global land surface phenology. *Remote Sens. Environ.* 2011, 115, 1102–1114.
21. Brandt M; Wigneron J-P; Chave J et al.; Satellite passive microwaves reveal recent climate-induced carbon losses in African drylands. *Nature Ecology & Evolution* **2018**, 2, 827-835, 10.1038/s41559-018-0530-6.
22. Fan L.; Wigneron J-P; Ciais P. et al.; Satellite-observed pantropical carbon dynamics. *Nature Plants* **2019**, 5, 944-951, 10.1038/s41477-019-0478-9.
23. Chaparro, D.; Duveiller, G.; Piles, M.; Cescatti, A.; Vall-llossera, M.; Camps, A.; Entekhabi, D. Sensitivity of L-band vegetation optical depth to carbon stocks in tropical forests: A comparison to higher frequencies and optical indices. *Remote Sens. Environ.* 2019, 232, 111303.
24. Rodríguez-Fernández, N.J.; Mialon, A.; Mermoz, S.; Bouvet, A.; Richaume, P.; Al Bitar, A.; Al-Yaari, A.; Brandt, M.; Kaminski, T.; Le Toan, T.; et al. An evaluation of SMOS L-band vegetation optical depth (L-VOD) data sets: High sensitivity of L-VOD to above-ground biomass in Africa. *Biogeosciences* 2018, 15, 4627–4645.
25. Li X.; Wigneron J-P et al.; Global-scale assessment and inter-comparison of recently developed / reprocessed microwave satellite vegetation optical depth products. *Remote Sensing of Environment* **2020**, in press, 0, 10.1016/j.rse.2020.112208.

26. INRAE BORDEAUX Soil Moisture and VOD PRODUCTS—Soil Moisture and Vegetation Products. Available online: <https://ib.remote-sensing.inrae.fr/> (accessed on 8 September 2020).
27. Prigent, C.; Papa, F.; Aires, F.; Rossow, W.B.; Matthews, E. Global inundation dynamics inferred from multiple satellite observations, 1993–2000. *J. Geophys. Res. Atmos.* 2007, 112.
28. Prigent, C.; Papa, F.; Aires, F.; Jimenez, C.; Rossow, W.B.; Matthews, E. Changes in land surface water dynamics since the 1990s and relation to population pressure. *Geophys. Res. Lett.* 2012, 39.
29. Parrens, M.; Bitar, A.A.; Frappart, F.; Papa, F.; Calmant, S.; Crétaux, J.-F.; Wigneron, J.-P.; Kerr, Y. Mapping dynamic water fraction under the tropical rain forests of the Amazonian basin from SMOS brightness temperatures. *Water* 2017, 9, 350.
30. Chan, S.K.; Bindlish, R.; O’Neill, P.; Jackson, T.; Njoku, E.; Dunbar, S.; Chaubell, J.; Piepmeier, J.; Yueh, S.; Entekhabi, D.; et al. Development and assessment of the SMAP enhanced passive soil moisture product. *Remote Sens. Environ.* 2018, 204, 931–941.
31. Gao, L.; Sadeghi, M.; Ebtehaj, A. Microwave retrievals of soil moisture and vegetation optical depth with improved resolution using a combined constrained inversion algorithm: Application for SMAP satellite. *Remote Sens. Environ.* 2020, 239, 111662.
32. Gevaert, A.I.; Parinussa, R.M.; Renzullo, L.J.; van Dijk, A.I.J.M.; de Jeu, R.A.M. Spatio-temporal evaluation of resolution enhancement for passive microwave soil moisture and vegetation optical depth. *Int. J. Appl. Earth Obs. Geoinf.* 2016, 45, 235–244.

Retrieved from <https://encyclopedia.pub/entry/history/show/10239>