

MDA5 Deficiency

Subjects: Genetics & Heredity

Contributor: Camila Xu

MDA5 deficiency is a disorder of the immune system (immunodeficiency) that leads to recurrent, severe infections of the lungs and airways (respiratory tract) beginning in infancy.

Keywords: genetic conditions

1. Introduction

These MDA5 deficiency infections are most frequently caused by rhinovirus (the virus that causes the common cold). Respiratory syncytial virus (RSV) and the influenza (flu) virus may also cause recurrent infections in affected individuals. While infection by these viruses is common in all children, it usually causes mild symptoms and lasts only a short time before being cleared by a healthy immune system. In contrast, individuals with MDA5 deficiency frequently require hospitalization due to the severity of the symptoms caused by the infection. Repeated infections can contribute to chronic lung disease.

Infections usually become less frequent with age in people with MDA5 deficiency, as the body's immune system matures and develops other mechanisms for fighting viruses.

2. Frequency

MDA5 deficiency is likely a rare disorder. Its prevalence is unknown.

3. Causes

MDA5 deficiency is caused by mutations in the *IFIH1* gene, which provides instructions for making the MDA5 protein. These mutations lead to production of an altered MDA5 protein that cannot function, resulting in a shortage (deficiency) of MDA5 activity.

The MDA5 protein plays an important role in innate immunity, the body's early, nonspecific response to foreign invaders (pathogens) such as viruses and bacteria. In particular, the protein recognizes a molecule called double-stranded RNA (a chemical cousin of DNA), which certain viruses, including rhinovirus, RSV, and the flu virus, have as their genetic material or produce when they infect cells and copy (replicate) themselves. (Another subset of viruses has DNA as their genetic material.) When the MDA5 protein recognizes pieces of viral RNA inside the cell, it helps turn on the production of immune system proteins called interferons. Interferons stimulate the immune system to fight infections.

Deficiency of MDA5 protein activity reduces interferon production in response to RNA-containing viruses. A lack of the important early immune response stimulated by interferons leads to severe viral infections in infants with MDA5 deficiency.

3.1. The gene associated with MDA5 deficiency

- *IFIH1*

4. Inheritance

The inheritance pattern of MDA5 deficiency is unclear. In some cases, the condition seems to follow an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition. In other cases, it appears that the condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder.

5. Other Names for This Condition

- IFIH1

deficiency

References

1. Asgari S, Schlapbach LJ, Anchisi S, Hammer C, Bartha I, Junier T, Mottet-Osman G, Posfay-Barbe KM, Longchamp D, Stocker M, Cordey S, Kaiser L, Riedel T, Kenna T, Long D, Schibler A, Telenti A, Tapparel C, McLaren PJ, Garcin D, Fellay J. Severe viral respiratory infections in children with IFIH1 loss-of-function mutations. *Proc Natl Acad Sci U S A*. 2017 Aug 1;114(31):8342-8347. doi:10.1073/pnas.1704259114.
 2. Berke IC, Yu X, Modis Y, Egelman EH. MDA5 assembles into a polar helical filament on dsRNA. *Proc Natl Acad Sci U S A*. 2012 Nov 6;109(45):18437-41. doi:10.1073/pnas.1212186109.
 3. Lamborn IT, Jing H, Zhang Y, Drutman SB, Abbott JK, Munir S, Bade S, Murdock HM, Santos CP, Brock LG, Masutani E, Fordjour EY, McElwee JJ, Hughes JD, Nichols DP, Belkadi A, Oler AJ, Happel CS, Matthews HF, Abel L, Collins PL, Subbarao K, Gelfand EW, Ciancanelli MJ, Casanova JL, Su HC. Recurrent rhinovirus infections in a child with inherited MDA5 deficiency. *J Exp Med*. 2017 Jul 3;214(7):1949-1972. doi: 10.1084/jem.20161759.
 4. Loo YM, Gale M Jr. Immune signaling by RIG-I-like receptors. *Immunity*. 2011 May 27;34(5):680-92. doi: 10.1016/j.immuni.2011.05.003. Review.
 5. Zaki M, Thoenes M, Kawalia A, Nürnberg P, Kaiser R, Heller R, Bolz HJ. Recurrent and Prolonged Infections in a Child with a Homozygous IFIH1 Nonsense Mutation. *Front Genet*. 2017 Sep 22;8:130. doi: 10.3389/fgene.2017.00130.
-

Retrieved from <https://encyclopedia.pub/entry/history/show/11700>