

Tubular Aggregate Myopathy

Subjects: Genetics

Submitted by:  Bruce

Ren

(This entry belongs to Entry Collection "[MedlinePlus](#)")

Definition

Tubular aggregate myopathy is a disorder that primarily affects the skeletal muscles, which are muscles the body uses for movement.

1. Introduction

Tubular aggregate myopathy is a disorder that primarily affects the skeletal muscles, which are muscles the body uses for movement. This disorder causes muscle pain, cramping, or weakness that begins in childhood and worsens over time. The muscles of the lower limbs are most often affected, although the upper limbs can also be involved. Affected individuals can have difficulty running, climbing stairs, or getting up from a squatting position. The weakness may also lead to an unusual walking style (gait). Some people with this condition develop joint deformities (contractures) in the arms and legs.

Skeletal muscles are normally made up of two types of fibers, called type I and type II fibers, in approximately equal quantities. Type I fibers, also called slow twitch fibers, are used for long, sustained activity, such as walking long distances. Type II fibers, also known as fast twitch fibers, are used for short bursts of strength, which are needed for activities such as running up stairs or sprinting. In people with tubular aggregate myopathy, type II fibers waste away (atrophy), so affected individuals have mostly type I fibers. In addition, proteins build up abnormally in both type I and type II fibers, forming clumps of tube-like structures called tubular aggregates. Tubular aggregates can occur in other muscle disorders, but they are the primary muscle cell abnormality in tubular aggregate myopathy.

2. Frequency

Tubular aggregate myopathy is a rare disorder. Its prevalence is unknown.

3. Causes

Tubular aggregate myopathy can be caused by mutations in the *STIM1* gene. The protein produced from this gene is involved in controlling the entry of positively charged calcium atoms (calcium ions) into cells. The *STIM1* protein recognizes when calcium ion levels are low and stimulates the flow of ions into the cell through special channels in the cell membrane called calcium-release activated calcium (CRAC) channels. In muscle cells, the activation of CRAC channels by *STIM1* is thought to help replenish calcium stores in a structure called the sarcoplasmic reticulum. *STIM1* may also be involved in the release of calcium ions from the sarcoplasmic reticulum. This release of ions stimulates muscle tensing (contraction).

The *STIM1* gene mutations involved in tubular aggregate myopathy lead to production of a *STIM1* protein that is constantly turned on (constitutively active), which means it continually stimulates calcium ion entry through CRAC channels regardless of ion levels. It is unknown how constitutively active *STIM1* leads to the muscle weakness characteristic of tubular aggregate myopathy. Evidence suggests that the tubular aggregates are composed of proteins that are normally part of the sarcoplasmic reticulum. Although the mechanism is unknown, some researchers speculate that the aggregates are the result of uncontrolled calcium levels in muscle cells, possibly due to abnormal *STIM1* activity.

Mutations in other genes, some of which have not been identified, are also thought to cause some cases of tubular aggregate myopathy.

3.1 The gene associated with Tubular aggregate myopathy

- STIM1

4. Inheritance

Most cases of tubular aggregate myopathy, including those caused by *STIM1* gene mutations, are inherited in an autosomal dominant pattern. Autosomal dominant inheritance means one copy of the altered gene in each cell is sufficient to cause the disorder. In some cases, the mutation is passed through generations in a family. Other cases result from new mutations in the gene and occur in people with no history of the disorder in their family.

Rarely, tubular aggregate myopathy is inherited in an autosomal recessive pattern, which means both copies of a gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition. Researchers are still working to determine which gene or genes are associated with autosomal recessive tubular aggregate myopathy.

5. Other Names for This Condition

- myopathy with tubular aggregates
- TAM

References

1. Böhm J, Chevessier F, Maues De Paula A, Koch C, Attarian S, Feger C, Hantaï D, Laforêt P, Ghorab K, Vallat JM, Fardeau M, Figarella-Branger D, Pouget J, Romero NB, Koch M, Ebel C, Levy N, Krahn M, Eymard B, Bartoli M, Laporte J. Constitutive activation of the calcium sensor STIM1 causes tubular-aggregate myopathy. *Am J Hum Genet.* 2013 Feb 7;92(2):271-8. doi: 10.1016/j.ajhg.2012.12.007.
2. Chevessier F, Bauché-Godard S, Leroy JP, Koenig J, Paturneau-Jouas M, Eymard B, Hantaï D, Verdière-Sahuqué M. The origin of tubular aggregates in human myopathies. *J Pathol.* 2005 Nov;207(3):313-23.
3. Endo Y, Noguchi S, Hara Y, Hayashi YK, Motomura K, Miyatake S, Murakami N, Tanaka S, Yamashita S, Kizu R, Bamba M, Goto Y, Matsumoto N, Nonaka I, Nishino I. Dominant mutations in ORAI1 cause tubular aggregate myopathy with hypocalcemia via constitutive activation of store-operated Ca²⁺ channels. *Hum Mol Genet.* 2015 Feb 1;24(3):637-48. doi: 10.1093/hmg/ddu477.
4. Hedberg C, Niceta M, Fattori F, Lindvall B, Ciolfi A, D'Amico A, Tasca G, Petrini S, Tulinius M, Tartaglia M, Oldfors A, Bertini E. Childhood onset tubular aggregate myopathy associated with de novo STIM1 mutations. *J Neurol.* 2014 May;261(5):870-6. doi: 10.1007/s00415-014-7287-x.
5. Lee KJ, Hyun C, Woo JS, Park CS, Kim DH, Lee EH. Stromal interaction molecule 1 (STIM1) regulates sarcoplasmic/endoplasmic reticulum Ca²⁺-ATPase 1a (SERCA1a) in skeletal muscle. *Pflugers Arch.* 2014 May;466(5):987-1001.
6. Lee KJ, Woo JS, Hwang JH, Hyun C, Cho CH, Kim DH, Lee EH. STIM1 negatively regulates Ca²⁺ release from the sarcoplasmic reticulum in skeletal myotubes. *Biochem J.* 2013 Jul 15;453(2):187-200. doi: 10.1042/BJ20130178.
7. Nesin V, Wiley G, Kousi M, Ong EC, Lehmann T, Nicholl DJ, Suri M, Shahrizaila N, Katsanis N, Gaffney PM, Wierenga KJ, Tsiokas L. Activating mutations in STIM1 and ORAI1 cause overlapping syndromes of tubular myopathy and congenital myosis. *Proc Natl Acad Sci U S A.* 2014 Mar 18;111(11):4197-202. doi:10.1073/pnas.1312520111.

Keywords

genetic conditions