Combination Strategies of Different Antimicrobials

Subjects: Biotechnology & Applied Microbiology Contributor: Nagaraj Basavegowda , Kwang-Hyun Baek

Despite the discovery and development of an array of antimicrobial agents, multidrug resistance poses a major threat to public health and progressively increases mortality. The use of antimicrobial agents in combination can produce synergistic effects if each drug invades a different target or signaling pathway with a different mechanism of action. Therefore, drug combinations can achieve a higher probability and selectivity of therapeutic responses than single drugs.

multidrug resistance

bacterial infections

synergistic effects

antimicrobial agents

1. Introduction

The rapid emergence and spread of multidrug-resistant (MDR) bacteria has become a serious global public health threat ^[1]. Long-term exposure and increased use and abuse of antibiotics could result in bacterial tolerance, which renders them less effective or even ineffective, and the mechanism includes changing the targets of antibiotics 2. MDR species are not only restricted to hospitals or healthcare environments; they are also found in humans, animals, plants, food, water, soil, and air. Moreover, they can be passed from person to person and between animals and persons. Antibiotic resistance is observed in various extracellular, intracellular, pathogenic, and nonpathogenic bacterial species. Among Gram-positive MDR species, Staphylococcus aureus, Streptococcus pneumoniae, Enterococcus faecium, and Enterococcus faecalis are the most common. Among Gram-negative strains, Pseudomonas aeruginosa, Escherichia coli, Klebsiella pneumoniae, and Acinetobacter baumannii are the most common MDR species ^[3]. However, methicillin-resistant Staphylococcus aureus causes pneumonia, diseases [<u>4</u>] bacteremia. soft tissue infections. and other fatal Similarly. multiple antibioticresistant Acinetobacter spp. and Klebsiella spp. are the most commonly reported.

Recent studies suggest that biofilm-associated infections account for more than 65% of all infections, and antibiotics lack effectiveness against biofilm-associated bacteria ^[5]. Biofilms can shield bacteria from host defenses, disinfectants, antibiotics, and many antimicrobial agents. This leads to a reduced bacterial growth rate, decreased metabolic activity, and promotion of tolerance to antibiotics ^[6]. Moreover, the excessive use of antibiotics is often not tolerated by the host organism, whereas lower doses are ineffective. In addition, conventional antibiotics support antibiotic resistance in viable bacteria ^[7]. Pathogens growing in biofilms exhibit both adaptive resistance to all antimicrobial agents and the host immune system by 10- to 1000-fold compared to their free-living, planktonic counterparts ^[8]. Hence, there is urgent need to search for alternative, novel, efficient antimicrobial agents and more targeted treatment strategies to overcome antibiotic resistance. An alternative

strategy currently in practice or under trials includes using different antimicrobial agents in combination to produce synergistic effects ^[9]. Combination therapy is an attractive and optional treatment because it represents potential adjuvant targets of non-overlapping signaling pathways and decreases the risk of developing cross-resistance ^[10].

Many plants have been used as sources of natural products to maintain good health, especially antimicrobial compounds ^[11]. Plants have evolved many alternative strategies against pathogens, which involve various phytochemicals, secondary metabolites, and other chemical compounds ^[12]. Bioactive compounds derived from plants, such as alkaloids, phenols, flavonoids, tannins, peptides, and other medicinally important compounds, are responsible for their antimicrobial ability against MDR pathogens ^[13]. Combining two or more plant extracts or their phytochemical components produces mutual antimicrobial enhancement, an unlimited pool of compounds, and the expansion or strengthening of their effects when combined as a multidrug ^[14]. Combinations of different drugs elicit several advantages over their use as individual moieties, including enhancing the effectiveness of other antimicrobial agents, reduction in dosage, fewer side effects, better synergistic effect, attack of multiple target sites, reduced risk, and exhibition of potent and rapid antibacterial effects against MDR pathogens ^[15]. The pharmacological effects of these combinations could be initiated by multiple mechanisms of action of herbal–herbal interactions.

Similarly, combining plant extracts or active phytochemicals with antibiotics improves their efficacy against resistant bacterial pathogens ^[16]. Synergism due to this combination helps minimize the minimum inhibitory concentrations (MICs) of these agents and reduces the economic cost and sensory impact ^[17]. Another strategic approach to combat MDR bacteria involves using essential oils (EOs) combined with conventional antibiotics or plant-derived phytochemicals. EOs have been widely used for their unique flavors; fragrances; and antibacterial, antioxidant, antifungal, anti-inflammatory, and anticarcinogenic properties ^[18]. Combining two or more EOs or their components or interactions between EOs and their components with antibiotics is a promising alternative strategy to increase their additive and synergistic antimicrobial effects. EOs and antibiotics, in combination, produce stronger bacterial inhibition compared to when they are individually administered because they target different pathways to create multifaceted effects against powerful bacterial defenses, consequently needing a decreased dose of each component ^[19]. The synergism between EOs and antibiotics may be attributed partly to the EO-induced permeabilization of the cell membrane, resulting in the immediate transport of antibiotics into the interior of the cell ^[20].

Antimicrobial nanomaterials represent another strategic approach to fighting MDR bacteria in clinical practice. Metal and metal oxide-based nanoparticles (NPs) have been widely investigated over the last decade, owing to their favorable chemical, physical, magnetic, electrical, thermal, optical, and biological properties ^[21]. Consequently, nanomaterials have emerged as new tools to combat deadly bacterial infections due to their specific features, such as size, shape, morphology, stability, and surface charge ^[22]. A combination of EOs and nanomaterials might establish functional materials with modified surfaces, improved inhibitory effects, and the ability to bind target microorganisms to achieve maximum synergistic performance ^[23]. Thus, combining nanomaterials with either EOs or plant extracts may improve their interaction with the bacterial cell membrane, thereby inducing the disruption, damage, and killing of bacteria ^[22].

2. Antibacterial Activities of Plant-Derived Compounds

Although different kinds of synthetic antimicrobial agents have been introduced to the market in many countries, natural medicine from plants might effectively treat certain diseases such as diarrhea, cold, labor pain, and dental diseases. Globally, approximately 60,000 plant species are used for medicinal purposes, of which approximately 28,000 are well-documented, and only 3000 are estimated to be traded internationally ^[24]. As a result, the search for herbal medicines with relevant biological activity has gained additional value as they are associated with fewer side effects and are much cheaper and affordable ^[25]. Plants usually produce two types of metabolites, primary and secondary, that can be found in extracts of their flowers, roots, leaves, bulbs, seeds, and bark (**Figure 1**). Primary metabolites are crucial for plant growth and development, whereas secondary metabolites are involved in plant defense, physiology, and environmental communication ^[26]. Secondary metabolites include many specialized and active compounds derived from primary metabolites. These compounds show promising results in controlling the development of resistance against bacterial pathogens, including MDR bacteria, and combating other bacterial infections. Plant secondary metabolites are classified into three categories on the basis of their biosynthetic origins: terpenoids, phenolics, and alkaloids ^[27].



Figure 1. The extracts of plant organs, namely, the root, bark, bulbs, leaf, flower, and seed, may encompass distinctive phytochemicals with antimicrobial properties.

2.1. Terpenoids

Terpenes are an extensive and diverse group of naturally occurring, highly enriched compounds of secondary plant metabolites. On the basis of the number of their isoprene structures or units, they are classified as monoterpenes, diterpenes, triterpenes, tetraterpenes, or sesquiterpenes. Terpenes are also called isoprenoids, and their derivatives that contain additional elements, such as oxygen, are usually termed terpenoids. Monoterpenes are the smallest terpenes, comprising two isoprene units. Monoterpenes contain volatile compounds found in EOs extracted from different flowers, fruits, and leaves and are commonly used in fragrances and aromatherapy. The antimicrobial properties of these compounds have been studied for two decades, and several studies have reported that thymol, carvacrol, eugenol, and menthol exhibit significant activity against many pathogens ^[28]. Geraniol and thymol have shown the most activity against Enterobacter species and S. aureus and E. coli, respectively ^{[29][30]}. Diterpenes are naturally occurring chemical compounds that contain active groups such as vitamin A. Phytol is an acyclic diterpene alcohol that acts as an antitumor, cytotoxic, and anti-inflammatory agent. Diterpenes also inhibit the growth of Staphylococcus aureus, Pseudomonas aeruginosa, Vibrio cholerae, and Candida spp. [31]. Triterpenes contain six isoprene units derived from mevalonic acid and have been shown to inhibit the growth of Mycobacterium tuberculosis. The combination of rifampicin and oleanolic acid has shown synergistic antibacterial effects against some pathogens ^[32]. Tetraterpenes are also known as carotenoids because beta-carotene is a yellow pigment in carrots. Similarly, yellow, orange, and red organic pigments are produced by plants, and these substances have effective antifungal and antibacterial properties [33]. Sesquiterpenes are the most diverse group of terpenoids, consisting of three units of isoprene with a lower vapor pressure than monoterpenes because of their high molecular weight. Farnesol, a natural sesquiterpene, demonstrated antibacterial activity against *S. aureus* and *S. epidermidis* [34].

2.2. Phenolics

Phenols are the simplest bioactive phytochemicals. They are monomeric components of polyphenols and acids with a single substituted phenolic ring and are typically found in plant tissues such as melanin and lignin. The components catechol, orcinol, tarragon, pyrogallol, phloroglucinol, pyrocatechol, resorcinol, and thyme are effective against viruses, bacteria, and fungi ^[35]. Both catechol and pyrogallol are hydroxylated phenolic compounds that are toxic against microorganisms; catechol has two hydroxyl groups, while pyrogallol has three. The microbial toxicity of phenolic compounds depends mainly on the number and position of their hydroxyl groups, as hydroxylation increases toxicity ^[36]. The presence of two or more hydroxyl groups located at ortho, para, or meta positions to each other is the key factor for their antimicrobial activity. The presence of a hydroxyl group at the meta position of thymol makes it a more effective antibacterial agent than carvacrol, which has a similar structure, whose hydroxyl group is in the ortho position.

2.3. Alkaloids

Alkaloids are cyclic-nitrogen-containing organic compounds that have various chemical structures. More than 18,000 alkaloids have been discovered and studied phytochemically from different sources. Alkaloids are grouped

into several classes, as natural, semi-synthetic, or synthetic, on the basis of their heterocyclic ring systems and biosynthetic precursors ^[37]. Alkaloids have various pharmacological activities, including antitumor, antihyperglycemic, anti-allergic, antidiabetic, antihyperlipidemic, and antibacterial. Piperine, berberine, quinolone, reserpine, sanguinarine, tomatidine, chanoclavine, conessine, and squalamine are the most important alkaloids with potent antibacterial activity. Piperine isolated from *Piper nigrum* and *Piper longum* inhibited the growth of mutant *S. aureus* when co-administered with ciprofloxacin ^[38].

3. Antimicrobial Efficacy of EOs

EOs are aromatic, lipophilic, and complex mixtures of volatile secondary metabolites that are mainly obtained from different parts of plants, such as leaves, herbs, flowers, buds, fruits, twigs, wood, bark, roots, and seeds ^[39]. EOs are extracted using hydrodistillation or steam distillation. EOs are lighter than water, with a strong flavor and odor reminiscent of their plant origin. The chemical composition of EOs is highly complex, with the main components being flavonoids, flavones, flavonols, phenols, polyphenols, tannins, alkaloids, quinones, coumarins, terpenoids, polypeptides, and lectins ^[22]. These compounds show potential pharmacological activities such as hepatoprotective, anti-inflammatory, antioxidant, anticancer, antiseptic, insecticidal, anti-parasitic, anti-allergic, antiviral, and antimicrobial properties ^[40]. Essential-oil-based products are in high demand in aromatherapy; as flavor-enhancers in food, beverages, cosmetics, perfumes, soaps, plastics, and resins; and in the pharmaceutical industries ^[41]. EOs have more than 50 components; however, only two or three of them are the major components present in high proportions. The other minor components are present in low amounts.

The amount of the different components of EOs varies with the parts and species of plants, as they are chemically derived from compounds and their derivatives ^[42]. The major constituents of EOs are terpenes and terpenoids, while other important compounds include aromatic and aliphatic constituents. The volatile components of EOs include a variety of chemicals such as alcohols (such as menthol, borneol, nerol, and linalool), acids (such as geranic acid and benzoic acid), aldehydes (such as citral), esters (such as linalyl acetate, citronellyl acetate, and menthyl), ketones (such as carvone, camphor, and pulegone), hydrocarbons (such as α -pinene, α -terpinene, myrcene, camphene, and p-cimene), ketones (such as carvone), pulegone, and carvone), phenols (such as carvacrol and thymol), lactones (such as bergapten), and peroxides (such as ascaridole), all of which play major roles in the composition of EOs (**Figure 2**) ^[43]. EOs show strong antibacterial activity against various pathogenic bacteria, including MDR pathogens, by penetrating the membrane of bacterial cells and disrupting their cellular structure. The antibacterial effectiveness of EOs differs across plant species and target bacteria depending on their cell wall structure (Gram-positive or Gram-negative). The association of some major constituents of EOs, such as eugenol, thymol, carvacrol, carvone, p-cymene, terpinene-4-ol, and cinnamic aldehyde, which easily penetrate and split in the lipid membrane, could disrupt the cell membrane; prevent cellular respiration; and lead to the loss of cell membrane integrity, removal of cellular contents, and finally cell death ^[44].



Figure 2. Schematic representation of some important EOs containing various components that have been screened for their antimicrobial properties.

EOs of *Thymus serrulatus* and *Thymus schimperi* were shown to possess strong antibacterial activity against *Lactobacillus* and *S. mutans*, with higher contents of thymol and carvacrol compounds reported to be the cause of this inhibition ^[45]. Similarly, EOs have been isolated from various parts of *Eugenia caryophylata*, such as the buds, leaves, and stems, with the main components being eugenol, β -caryophyllene, and eugenyl acetate. These EOs are effective against *S. aureus*, *E. coli*, *B. subtilis*, and *S. typhimurium* ^[46]. Likewise, tea tree EO has been reported to cause changes in the membrane permeability and mycelial morphology of *Monilinia fructicola* ^[47]. Furthermore, a recent study of lavender EO against *A. hydrophila*, *A. caviae*, *A. dhakensis*, *C. freundii*, *P. mirabilis*, and *S. enterica* showed the presence of the major compounds linalool and linalyl acetate ^[48]. In another study, winter savory EO exhibited the strongest inhibitory effect against clinical oral isolates of *Candida* spp., with thymol as the major compound ^[49]. It has been reported that among the commercially available EOs, such as anise, cinnamon, clove, cumin, laurel, Mexican lime, and Mexican oregano, oregano EO has the highest antibacterial activity against *S. typhimurium* and *E. coli*, with thymol as its major compound ^[50].

4. Antimicrobial Nanomaterials

With the emergence of bacterial resistance and biofilm-associated infections, clinical research is needed to develop novel, effective, long-term antibacterial and biofilm-preventative agents. Metals have been extensively studied

among the most promising novel antimicrobial agents ^[51]. Recently, metal-based nanomaterials have become the most extensively and rapidly emerging materials in the field of medicine. Different types of metallic NPs have demonstrated strong antibacterial activity in many recent studies ^[52]. Generally, NPs have fascinating characteristics, such as a high surface area-to-volume ratio, size, shape, and surface activity, and exhibit superior electrical, catalytic, and optical properties. Due to their unique properties, NPs have a more well-developed surface than their microscale counterparts, affecting their antimicrobial efficiency and effectiveness ^[52]. Similar to antibiotics, metals can selectively inhibit metabolic pathways by interacting with bactericidal activity and ultimately kill MDR bacteria ^[52]; however, cells deviate from metal transport systems and metalloproteins ^[54]. Hence, NPs showed noticeable antimicrobial activity against both Gram-negative and Gram-positive pathogens such as *E. faecalis*, *B. subtilis*, S. *epidermidis*, multidrug-resistant *S. aureus*, and *E. coli* strains.

Metal NPs such as Ag, Au, Cu, Zn, Ti, Ga, Al, and Pt ^[55] and metal oxide NPs such as CuO, MgO, ZnO, TiO₂, NiO₂, SiO₂, and Fe₃O₄ are known to display various antimicrobial properties, which have been known and applied for decades ^[56]. In addition, graphene oxide (GO) and carbon nanotubes (CNTs), such as single-walled carbon nanotubes (SWCNTs) and multi-walled carbon nanotubes (MWCNTs), are also excellent candidates due to their antimicrobial activities (**Figure 3**). Recently, metal–organic frameworks (MOF) and metal sulfide nanomaterials, such as FeS-, Ags-, ZnS-, and CdS-MOFs and Zn-, Cu-, and Mn-based MOFs, have also been proven to have antibacterial activities ^[57]. Multimetallic NPs, particularly NPs formed by at least two metals, such as bimetallic, trimetallic, and quadrametallic NPs, display rich optical, electronic, and magnetic properties. The properties of multimetallic NPs, including size, shape, surface area, and zeta potential, enhance their interaction with bacterial cell membranes. They could disrupt cell membranes, produce reactive oxygen species (ROS), damage the DNA, induce protein dysfunction, and may be potentiated by the host immune system ^[23].



Figure 3. Schematic representation of different nanomaterials that possess antimicrobial activity.

Metallic biopolymer-based nanocomposite systems are well-known candidates as antimicrobial nanomaterials. In particular, cationic chitosan-based NPs bind to anionic cell membranes, resulting in alterations to the cell membrane, leakage of intracellular compounds, and eventually cell death ^[58]. Antimicrobial peptides (AMPs) have attracted great interest because of their high biocompatibility and low probability of inducing bacterial resistance. AMP-conjugated nanomaterials can hinder the growth of pathogens and kill bacteria on the basis of the inherent action of typical combination strategies ^[59].

AgNPs are considered the most common antimicrobial agents that can destroy a wide range of Gram-negative and Gram-positive bacteria. Ag ions combine with disulfide or sulfhydryl groups of enzymes, disrupt normal metabolic processes, and ultimately lead to cell death ^[60]. The bactericidal efficacy of Au NPs might have a greater chance of penetrating the bacterial cell wall by generating holes, leading to increased permeability and higher oxidative stress within the cytoplasm. Similarly, ZnO NPs displayed vigorous antimicrobial activity by releasing Zn²⁺ ions and generating ROS, owing to their electrostatic interaction and internalization. In contrast, smaller ZnO NPs increased the interaction and abrasiveness of the bacterial cell wall ^[61]. Cu and CuO NPs showed excellent antimicrobial

activity against different strains of bacteria by releasing Cu^{2+} ions and stimulating ROS production ^[62]. Various studies have revealed the visible-light-induced antibacterial properties of Fe-, Cu-, Ni-, and Ag-doped TiO₂ NPs against *E. coli* and *S. aureus* ^{[63][64]}; however, TiO₂ NPs adversely affect human cells and tissues, so their use remains limited. SiO₂ NPs, especially mesoporous NPs, have attracted considerable attention because their properties, such as size, matrix, and surface functions, which can be tuned to improve their interaction with and penetration of biofilm-producing bacteria ^[65].

Compared to monometallic NPs, multimetallic NPs, such as bi-, tri-, and quadrametallic NPs, have gained great importance and interest due to their unique physical, chemical, electrical, optical, and catalytic properties and applications in different fields ^[66]. Multimetallic NPs can be altered or tuned by controlling their structure, morphology, and chemical composition to achieve strong synergistic interactions and performance ^[67]. When at least two metals are formed as NPs, combinatorial approaches, such as structural changes, deduction of the lattice parameters, and total electronic charge shift improvements, are expected ^[68]. Recently, bimetallic Ag/Cu and Cu/Zn ^[69], and trimetallic Cu/Cr/Ni ^[70] and CuO/NiO/ZnO ^[71] NPs have exhibited remarkably improved antimicrobial performance compared to monometallic NPs.

Core–shell quantum dots (CSQDs) are a new type of fluorescent antibacterial nanomaterial with unique physical and chemical properties. Owing to their high electron transfer, CSQDs produce a large number of free electrons and holes that accumulate ROS inside the cell, inhibiting their respiration and replication ^[72]. CSQDs exert antimicrobial effects by destroying cell walls, binding with genetic material, and inhibiting energy production.

Additionally, the antibacterial properties of graphene involve both chemical and physical modes of action. The chemical action is associated with oxidative stress generated by charge transfer and ROS, while the physical action is induced by the direct contact of graphene with bacterial membranes ^[73]. Similarly, CNTs are more effective and cost-efficient, exhibiting strong antimicrobial properties owing to their remarkable structure. This mechanism is based on the interaction of CNTs with microorganisms and the disruption of their metabolic processes, cellular membranes, and morphology.

Dendrimers are macromolecules with highly branched tree-like dendritic structures, narrow sizes, relatively large molecular masses, and well-defined globular structures ^[74]. Dendrimers peripherally cationic and highly water soluble due to numerous peripheral hydrophilic groups compatible with water ^[75]. Dendrimers can incorporate biologically active agents in the interior or periphery; therefore, they serve as carriers of biologically active agents [^{76]}. Antimicrobial polymers or their composites can prevent or suppress the growth of microbes on their surfaces or in the environment. Positively charged polymer surface groups are attracted to negatively charged cell membranes, leading to cell membrane damage and cell death ^[77].

5. Synergistic Antimicrobial Activity of Plant Extracts, EOs, and Nanomaterials

A synergistic effect is a process in which chemical substances or biological structures interact or combine to create an effect greater than the sum of the effects of the individual components. Synergy is the concept wherein the performance of two or more antimicrobial agents is combined and the effects of such mixtures are greater than those of the separate or individual components, enhancing solubility. In recent years, the synergistic combination of different antimicrobials and plant extracts has been considered a unique strategy to increase the spectrum of antimicrobial activity of these substances and prevent the development of resistant strains [78]. Plant extracts, EOs, and nanomaterials are commonly used as antimicrobial agents for the treatment of many infectious diseases. However, these antimicrobials are not very effective against acute infections because they lack a standardized and clinically applicable pharmaceutical form. Consequently, various antibiotics have been discovered as synthetic antimicrobials; however, these drugs are highly toxic and have poor tolerability, so bacteria develop resistance against them. Hence, a possible approach to improve and enhance antibacterial activity is to use combinations of different antimicrobials. These combinatorial approaches can be used alone or with other antimicrobials against a wide range of pathogens [79]. Compared to the individual substances or components, multicomponent antimicrobials display increased antimicrobial activity; therefore, other molecules present in the antimicrobial agents could control the function of the main components and improve their synergistic effects. Moreover, combining different antimicrobials offers many advantages, including a reduction in dosage, fewer side effects, decreased toxicity, extensive antibacterial action, and the ability to attack multiple target sites with increased efficacy [80]. Some combinations of antimicrobial agents, such as plant extracts, EOs, antibiotics, and NPs, are summarized in Table 1.

Table 1	Antimicrobial	activity of	combinations	of plant	extract,	EOs,	antibiotics,	and NPs	against	different
				pathoge	ens.					

Antimicrobial Agents	Combinations	Pathogens	Ref.
EOs/EOs	Melaleuca alternifolia/Cupressus sempervirens	E. coli	[<u>79</u>]
EOs/antibiotics	Eucalyptus globulus/oxacillin	S. aureus	[<u>79</u>]
EOs/NPs	Lemongrass/chitosan NP	E. coli, S. aureus	[<u>81</u>]
Plant extract/antibiotics	Salvadora persica/amoxicillin	P. gingivalis, T. forsythia	[<u>82</u>]
Plant extract/EOs	Origanum vulgare/carvacrol	S. aureus	[<u>83</u>]
Plant extract/NPs	Vatica diospyroides/Ag NPs	S. aureus, B. subtilis	[<u>84</u>]
NPs/antibiotics	AgNPs/fluconazole	S. aureus, E. coli	[<u>85</u>]
β-Lactam/β-lactamase inhibitor	amoxicillin/potassium clavulanate	S. aureus	[<u>86</u>]

Combinations of antimicrobial agents, such as EOs/EOs, plant extract/plant extract, and NPs/NPs, already have confirmed antimicrobial activities ^[87]. Ncube et al. evaluated the bulb and leaf extracts of three medicinal plants, independently and in combination, against *S. aureus*. Their results showed the strongest synergistic effect compared with the effects observed with individual extracts ^[12]. Similarly, a combination of *Bulbine frutescens* and *Vernonia lasiopus* plant extracts showed improved antimicrobial activity against *E. coli* ^[88]. Obuekwe et al. found the largest zones of inhibition against *S. aureus* using a combination of *Ocimum gratissimum* and *Ficus exasperate*, and *Bryophyllum pinnatum* and *Ocimum gratissimum* against *E. coli* ^[89]. Recently, EO–EO associations showed a synergistic effect against vancomycin-resistant enterococci (VRE), methicillin-resistant *S. aureus* (MRSA), and extended-spectrum β-lactamase (ESBL)-producing *Escherichia coli* ^[79]. A mixture of *R. abyssinicus* and *D. penninervium* EOs showed strong synergistic effects against MRSA and *P. aeruginosa* ^[90].

The synergistic antibacterial activities of cumin, cardamom, and dill weed EOs against *C. coli* and *C. jejuni* have been reported ^[91]. In a previous study, a combination of cinnamon and clove EOs showed synergistic antibacterial activity against foodborne *S. aureus*, *L. monocytogenes*, *S. typhimurium*, and *P. aeruginosa* ^[92]. Garza-Cervantes et al. examined the synergistic antimicrobial activities of silver in combination with other transition metals (Zn, Co, Cd, Ni, and Cu). Their results exhibited synergism since the antimicrobial effects of the combinations against *E. coli* and *B. subtilis* increased up to eightfold when compared to the individual metals ^[93]. Similarly, β-lactam is the most common bactericidal agent recommended for the treatment of several infectious diseases. However, the increasing emergence of β-lactam resistance due to β-lactamase enzyme production is one of the most serious public health threats. Hence, current clinical trials suggest the use of proper combinations of β-lactam and β-lactamase inhibitors ^[94]. β-Lactam inhibitors are associated with β-lactam antibiotics because they are hydrolyzed by β-lactamases, and their main objective is to protect the associated antibiotics. β-Lactam inhibitors prevent the hydrolytic action of β-lactam antibiotics by binding to the active site of β-lactamase enzymes ^[95].

References

- Pang, X.; Xiao, Q.; Cheng, Y.; Ren, E.; Lian, L.; Zhang, Y.; Gao, H.; Wang, X.; Leung, W.; Chen, X. Bacteria-responsive nanoliposomes as smart sonotheranostics for multidrug resistant bacterial infections. ACS Nano 2019, 13, 2427–2438.
- 2. Baym, M.; Stone, L.K.; Kishony, R. Multidrug evolutionary strategies to reverse antibiotic resistance. Science 2016, 351, aad3292.
- Sun, Y.; Ye, J.; Hou, Y.; Chen, H.; Cao, J.; Zhou, T. Predation efficacy of Bdellovibrio bacteriovorus on multidrug-resistant clinical pathogens and their corresponding biofilms. Jpn. J. Infect. Dis. 2017, 70, 485–489.
- 4. Dong, P.; Mohammad, H.; Hui, J.; Leanse, L.G.; Li, J.; Liang, L.; Dai, T.; Seleem, M.N.; Cheng, J. Photolysis of Staphyloxanthin in methicillin-resistant Staphylococcus aureus potentiates killing by

reactive oxygen species. Adv. Sci. 2019, 6, 1900030.

- 5. Jacqueline, C.; Caillon, J. Impact of bacterial biofilm on the treatment of prosthetic joint infections. J. Antimicrob. Chemother. 2014, 69, i37–i40.
- Dincer, S.; Uslu, F.M.; Delik, A. Antibiotic resistance in biofilm. In Bacterial Biofilms; IntechOpen: London, UK, 2020; Available online: https://www.intechopen.com/chapters/72109 (accessed on 30 July 2022).
- 7. Beyth, N.; Houri-Haddad, Y.; Domb, A.; Khan, W.; Hazan, R. Alternative Antimicrobial Approach: Nano-Antimicrobial Materials. Evid.-Based Complement. Altern. Med. 2015, 2015, 246012.
- 8. Dostert, M.; Belanger, C.R.; Hancock, R.E.W. Design and assessment of anti-biofilm peptides: Steps toward clinical application. J. Innate Immun. 2019, 11, 193–204.
- 9. Kaur, I. Novel strategies to combat antimicrobial resistance. J. Infect. Dis. Ther. 2016, 4, 292.
- Bozic, I.; Reiter, J.G.; Allen, B.; Antal, T.; Chatterjee, K.; Shah, P.; Moon, Y.S.; Yaqubie, A.; Kelly, N.; Le, D.T. Evolutionary dynamics of cancer in response to targeted combination therapy. Elife 2013, 2, e00747.
- 11. Nascimento, G.G.F.; Locatelli, J.; Freitas, P.C.; Silva, G.L. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. Braz. J. Microbiol. 2000, 31, 247–256.
- 12. Ncube, B.; Finnie, J.F.; Van Staden, J. In vitro antimicrobial synergism within plant extract combinations from three South African medicinal bulbs. J. Ethnopharmacol. 2012, 139, 81–89.
- 13. Aggarwal, B.; Lamba, H.S.; Ajeet, P.S. Various pharmacological aspects of Cocos nucifera—A review. Am. J. Pharmacol. Sci. 2017, 5, 25–30.
- Olajuyigbe, O.O.; Afolayan, A.J. Evaluation of combination effects of ethanolic extract of Ziziphus mucronata Willd. subsp. mucronata Willd. and antibiotics against clinically important bacteria. Sci. World J. 2013, 2013, 769594.
- León-Buitimea, A.; Garza-Cárdenas, C.R.; Garza-Cervantes, J.A.; Lerma-Escalera, J.A.; Morones-Ramírez, J.R. The demand for new antibiotics: Antimicrobial peptides, nanoparticles, and combinatorial therapies as future strategies in antibacterial agent design. Front. Microbiol. 2020, 11, 1669.
- Cheesman, M.J.; Ilanko, A.; Blonk, B.; Cock, I.E. Developing new antimicrobial therapies: Are synergistic combinations of plant extracts/compounds with conventional antibiotics the solution? Pharmacogn. Rev. 2017, 11, 57.
- Reda, F.M.; El-Zawahry, Y.A.; Omar, A.R. Synergistic effect of combined antibiotic and methanol extract of Eucalyptus camaldulensis leaf against Staphylococcus aureus and Pseudomonas aeruginosa. Int. J. Appl. Sci. Biotechnol. 2017, 5, 486–497.

- 18. Basavegowda, N.; Baek, K.-H. Synergistic Antioxidant and Antibacterial Advantages of Essential Oils for Food Packaging Applications. Biomolecules 2021, 11, 1267.
- 19. Horváth, G.; Bencsik, T.; Ács, K.; Kocsis, B. Sensitivity of ESBL-Producing Gram-Negative Bacteria to Essential Oils, Plant Extracts, and Their Isolated Compounds; Academic Press: Amsterdam, The Netherlands, 2016; pp. 239–269.
- Aleksic, V.; Mimica-Dukic, N.; Simin, N.; Nedeljkovic, N.S.; Knezevic, P. Synergistic effect of Myrtus communis L. essential oils and conventional antibiotics against multi-drug resistant Acinetobacter baumannii wound isolates. Phytomedicine 2014, 21, 1666–1674.
- 21. Dikshit, P.K.; Kumar, J.; Das, A.K.; Sadhu, S.; Sharma, S.; Singh, S.; Gupta, P.K.; Kim, B.S. Green synthesis of metallic nanoparticles: Applications and limitations. Catalysts 2021, 11, 902.
- 22. Basavegowda, N.; Patra, J.K.; Baek, K.-H. Essential oils and mono/Bi/tri-metallic nanocomposites as alternative sources of antimicrobial agents to combat multidrug-resistant pathogenic microorganisms: An overview. Molecules 2020, 25, 1058.
- 23. Basavegowda, N.; Baek, K.-H. Multimetallic nanoparticles as alternative antimicrobial agents: Challenges and perspectives. Molecules 2021, 26, 912.
- 24. Jenkins, M.; Timoshyna, A.; Cornthwaite, M. Wild at Home: Exploring the global harvest, trade and use of wild plant ingredients. Retrieved Jan. 2018, 26, 2019.
- 25. Manzo, L.M.; Moussa, I.; Ikhiri, K.; Yu, L. Toxicity studies of Acacia nilotica (L.): A review of the published scientific literature. J. Herbmed Pharmacol. 2019, 8, 163–172.
- 26. Pott, D.M.; Osorio, S.; Vallarino, J.G. From central to specialized metabolism: An overview of some secondary compounds derived from the primary metabolism for their role in conferring nutritional and organoleptic characteristics to fruit. Front. Plant Sci. 2019, 10, 835.
- 27. Eljounaidi, K.; Lichman, B.R. Nature's chemists: The discovery and engineering of phytochemical biosynthesis. Front. Chem. 2020, 8, 1041.
- Kusakizako, T.; Miyauchi, H.; Ishitani, R.; Nureki, O. Structural biology of the multidrug and toxic compound extrusion superfamily transporters. Biochim. Biophys. Acta BBA-Biomembr. 2020, 1862, 183154.
- 29. Dreier, J.; Ruggerone, P. Interaction of antibacterial compounds with RND efflux pumps in Pseudomonas aeruginosa. Front. Microbiol. 2015, 6, 660.
- 30. Zacchino, S.A.; Butassi, E.; Cordisco, E.; Svetaz, L.A. Hybrid combinations containing natural products and antimicrobial drugs that interfere with bacterial and fungal biofilms. Phytomedicine 2017, 37, 14–26.
- 31. Batista, O.; Duarte, A.; Nascimento, J.; Simões, M.F.; de la Torre, M.C.; Rodríguez, B. Structure and antimicrobial activity of diterpenes from the roots of Plectranthus hereroensis. J. Nat. Prod.

1994, 57, 858-861.

- 32. Cox-Georgian, D.; Ramadoss, N.; Dona, C.; Basu, C. Therapeutic and medicinal uses of terpenes. In Medicinal Plants; Springer: Berlin/Heidelberg, Germany, 2019; pp. 333–359.
- 33. Pan, N.; Sun, G. Functional Textiles for Improved Performance, Protection and Health; Elsevier: Amsterdam, The Netherlands, 2011; ISBN 0857092871.
- 34. Gomes, F.I.A.; Teixeira, P.; Azeredo, J.; Oliveira, R. Effect of farnesol on planktonic and biofilm cells of Staphylococcus epidermidis. Curr. Microbiol. 2009, 59, 118–122.
- 35. Gyawali, R.; Ibrahim, S.A. Natural products as antimicrobial agents. Food Control 2014, 46, 412–429.
- 36. Cowan, M.M. Plant products as antimicrobial agents. Clin. Microbiol. Rev. 1999, 12, 564–582.
- 37. Thawabteh, A.; Juma, S.; Bader, M.; Karaman, D.; Scrano, L.; Bufo, S.A.; Karaman, R. The biological activity of natural alkaloids against herbivores, cancerous cells and pathogens. Toxins 2019, 11, 656.
- 38. Khan, I.A.; Mirza, Z.M.; Kumar, A.; Verma, V.; Qazi, G.N. Piperine, a phytochemical potentiator of ciprofloxacin against Staphylococcus aureus. Antimicrob. Agents Chemother. 2006, 50, 810–812.
- Stephane, F.F.Y.; Jules, B.K.J. Terpenoids as important bioactive constituents of essential oils, essential oils. In Essential Oils-Bioactive Compounds, New Perspectives and Applications; IntechOpen: London, UK, 2020.
- Korinek, M.; Handoussa, H.; Tsai, Y.-H.; Chen, Y.-Y.; Chen, M.-H.; Chiou, Z.-W.; Fang, Y.; Chang, F.-R.; Yen, C.-H.; Hsieh, C.-F. Anti-inflammatory and antimicrobial volatile oils: Fennel and cumin inhibit neutrophilic inflammation via regulating calcium and MAPKs. Front. Pharmacol. 2021, 12, 674095.
- 41. Mahanta, B.P.; Bora, P.K.; Kemprai, P.; Borah, G.; Lal, M.; Haldar, S. Thermolabile essential oils, aromas and flavours: Degradation pathways, effect of thermal processing and alteration of sensory quality. Food Res. Int. 2021, 145, 110404.
- 42. Chouhan, S.; Sharma, K.; Guleria, S. Antimicrobial activity of some essential oils—Present status and future perspectives. Medicines 2017, 4, 58.
- 43. Swamy, M.K.; Akhtar, M.S.; Sinniah, U.R. Antimicrobial properties of plant essential oils against human pathogens and their mode of action: An updated review. Evid.-Based Complement. Altern. Med. 2016, 2016, 3012462.
- 44. Teixeira, B.; Marques, A.; Ramos, C.; Neng, N.R.; Nogueira, J.M.F.; Saraiva, J.A.; Nunes, M.L. Chemical composition and antibacterial and antioxidant properties of commercial essential oils. Ind. Crops Prod. 2013, 43, 587–595.

- 45. Damtie, D.; Mekonnen, Y. Antibacterial activity of essential oils from Ethiopian thyme (Thymus serrulatus and Thymus schimperi) against tooth decay bacteria. PLoS ONE 2020, 15, e0239775.
- 46. Sohilait, H.J.; Kainama, H.; Nindatu, M. Chemical composition and antibacterial activity of the essential oils from different parts of Eugenia caryophylata, Thunb grown in Amboina Island. Int. J. Org. Chem. 2018, 8, 229–239.
- 47. Xu, Y.; Wei, J.; Wei, Y.; Han, P.; Dai, K.; Zou, X.; Jiang, S.; Xu, F.; Wang, H.; Sun, J. Tea tree oil controls brown rot in peaches by damaging the cell membrane of Monilinia fructicola. Postharvest Biol. Technol. 2021, 175, 111474.
- 48. Hossain, S.; Heo, H.; De Silva, B.C.J.; Wimalasena, S.; Pathirana, H.; Heo, G.-J. Antibacterial activity of essential oil from lavender (Lavandula angustifolia) against pet turtle-borne pathogenic bacteria. Lab. Anim. Res. 2017, 33, 195–201.
- Nikolić, M.; Jovanović, K.K.; Marković, T.; Marković, D.; Gligorijević, N.; Radulović, S.; Soković, M. Chemical composition, antimicrobial, and cytotoxic properties of five Lamiaceae essential oils. Ind. Crops Prod. 2014, 61, 225–232.
- Andrade-Ochoa, S.; Chacón-Vargas, K.F.; Sánchez-Torres, L.E.; Rivera-Chavira, B.E.; Nogueda-Torres, B.; Nevárez-Moorillón, G.V. Differential antimicrobial effect of essential oils and their main components: Insights based on the cell membrane and external structure. Membranes 2021, 11, 405.
- Middaugh, J.; Hamel, R.; Jean-Baptiste, G.; Beriault, R.; Chenier, D.; Appanna, V.D. Aluminum triggers decreased aconitase activity via Fe-S cluster disruption and the overexpression of isocitrate dehydrogenase and isocitrate lyase: A metabolic network mediating cellular survival. J. Biol. Chem. 2005, 280, 3159–3165.
- 52. Macomber, L.; Elsey, S.P.; Hausinger, R.P. Fructose-1, 6-bisphosphate aldolase (class II) is the primary site of nickel toxicity in Escherichia coli. Mol. Microbiol. 2011, 82, 1291–1300.
- 53. Seil, J.T.; Webster, T.J. Antimicrobial applications of nanotechnology: Methods and literature. Int. J. Nanomed. 2012, 7, 2767.
- 54. Lemire, J.A.; Harrison, J.J.; Turner, R.J. Antimicrobial activity of metals: Mechanisms, molecular targets and applications. Nat. Rev. Microbiol. 2013, 11, 371–384.
- 55. Malarkodi, C.; Rajeshkumar, S.; Paulkumar, K.; Vanaja, M.; Gnanajobitha, G.; Annadurai, G. Biosynthesis and antimicrobial activity of semiconductor nanoparticles against oral pathogens. Bioinorg. Chem. Appl. 2014, 2014, 347167.
- 56. Biswal, A.K.; Misra, P.K. Biosynthesis and characterization of silver nanoparticles for prospective application in food packaging and biomedical fields. Mater. Chem. Phys. 2020, 250, 123014.

- 57. Yaqoob, A.A.; Ahmad, H.; Parveen, T.; Ahmad, A.; Oves, M.; Ismail, I.M.I.; Qari, H.A.; Umar, K.; Mohamad Ibrahim, M.N. Recent Advances in Metal Decorated Nanomaterials and Their Various Biological Applications: A Review. Front. Chem. 2020, 8, 1–23.
- 58. Díez-Pascual, A.M. Antibacterial activity of nanomaterials. Nanomaterials 2018, 8, 359.
- Pardhi, D.M.; Karaman, D.Ş.; Timonen, J.; Wu, W.; Zhang, Q.; Satija, S.; Mehta, M.; Charbe, N.; McCarron, P.A.; Tambuwala, M.M. Anti-bacterial activity of inorganic nanomaterials and their antimicrobial peptide conjugates against resistant and non-resistant pathogens. Int. J. Pharm. 2020, 586, 119531.
- 60. Egger, S.; Lehmann, R.P.; Height, M.J.; Loessner, M.J.; Schuppler, M. Antimicrobial properties of a novel silver-silica nanocomposite material. Appl. Environ. Microbiol. 2009, 75, 2973–2976.
- 61. Yuan, P.; Ding, X.; Yang, Y.Y.; Xu, Q. Metal nanoparticles for diagnosis and therapy of bacterial infection. Adv. Healthc. Mater. 2018, 7, 1701392.
- 62. Hsueh, Y.-H.; Tsai, P.-H.; Lin, K.-S. Ph-dependent antimicrobial properties of copper oxide nanoparticles in staphylococcus aureus. Int. J. Mol. Sci. 2017, 18, 793.
- Yadav, H.M.; Kolekar, T.V.; Pawar, S.H.; Kim, J.-S. Enhanced photocatalytic inactivation of bacteria on Fe-containing TiO2 nanoparticles under fluorescent light. J. Mater. Sci. Mater. Med. 2016, 27, 1–9.
- 64. Moongraksathum, B.; Chen, Y.-W. Anatase TiO2 co-doped with silver and ceria for antibacterial application. Catal. Today 2018, 310, 68–74.
- 65. Şen Karaman, D.; Manner, S.; Rosenholm, J.M. Mesoporous silica nanoparticles as diagnostic and therapeutic tools: How can they combat bacterial infection? Ther. Deliv. 2018, 9, 241–244.
- 66. Mishra, K.; Basavegowda, N.; Lee, Y.R. Biosynthesis of Fe, Pd, and Fe-Pd bimetallic nanoparticles and their application as recyclable catalysts for cycloaddition reaction: A comparative approach. Catal. Sci. Technol. 2015, 5, 2612–2621.
- 67. Zhang, J.; Ma, J.; Fan, X.; Peng, W.; Zhang, G.; Zhang, F.; Li, Y. Graphene supported Au-Pd-Fe3O4 alloy trimetallic nanoparticles with peroxidase-like activities as mimic enzyme. Catal. Commun. 2017, 89, 148–151.
- 68. Deepak, F.L.; Mayoral, A.; Arenal, R. Advanced Transmission Electron Microscopy: Applications to Nanomaterials; Springer: Berlin/Heidelberg, Germany, 2015; ISBN 3319151770.
- 69. Merugu, R.; Gothalwal, R.; Deshpande, P.K.; De Mandal, S.; Padala, G.; Chitturi, K.L. Synthesis of Ag/Cu and Cu/Zn bimetallic nanoparticles using toddy palm: Investigations of their antitumor, antioxidant and antibacterial activities. Mater. Today Proc. 2021, 44, 99–105.
- 70. Vaseghi, Z.; Tavakoli, O.; Nematollahzadeh, A. Rapid biosynthesis of novel Cu/Cr/Ni trimetallic oxide nanoparticles with antimicrobial activity. J. Environ. Chem. Eng. 2018, 6, 1898–1911.

- 71. Paul, D.; Mangla, S.; Neogi, S. Antibacterial study of CuO-NiO-ZnO trimetallic oxide nanoparticle. Mater. Lett. 2020, 271, 127740.
- 72. Rajendiran, K.; Zhao, Z.; Pei, D.-S.; Fu, A. Antimicrobial activity and mechanism of functionalized quantum dots. Polymers 2019, 11, 1670.
- 73. Radhi, A.; Mohamad, D.; Rahman, F.S.A.; Abdullah, A.M.; Hasan, H. Mechanism and factors influence of graphene-based nanomaterials antimicrobial activities and application in dentistry. J. Mater. Res. Technol. 2021, 11, 1290–1307.
- 74. Alfei, S.; Schito, A.M.; Zuccari, G. Considerable Improvement of Ursolic Acid Water Solubility by its Encapsulation in Dendrimer Nanoparticles: Design, Synthesis and Physicochemical Characterization. Nanomaterials 2021, 11, 2196.
- 75. Alfei, S.; Brullo, C.; Caviglia, D.; Zuccari, G. Preparation and Physicochemical Characterization of Water-Soluble Pyrazole-Based Nanoparticles by Dendrimer Encapsulation of an Insoluble Bioactive Pyrazole Derivative. Nanomaterials 2021, 11, 2662.
- 76. Jiang, G.; Liu, S.; Yu, T.; Wu, R.; Ren, Y.; van der Mei, H.C.; Liu, J.; Busscher, H.J. PAMAM dendrimers with dual-conjugated vancomycin and Ag-nanoparticles do not induce bacterial resistance and kill vancomycin-resistant Staphylococci. Acta Biomater. 2021, 123, 230–243.
- 77. Alfei, S.; Schito, A.M. Positively charged polymers as promising devices against multidrug resistant gram-negative bacteria: A Review. Polymers 2020, 12, 1195.
- Aiyegoro, O.A.; Afolayan, A.J.; Okoh, A.I. Synergistic interaction of Helichrysum pedunculatum leaf extracts with antibiotics against wound infection associated bacteria. Biol. Res. 2009, 42, 327–338.
- 79. Iseppi, R.; Mariani, M.; Condò, C.; Sabia, C.; Messi, P. Essential oils: A natural weapon against antibiotic-resistant bacteria responsible for nosocomial infections. Antibiotics 2021, 10, 417.
- Bassolé, I.H.N.; Lamien-Meda, A.; Bayala, B.; Tirogo, S.; Franz, C.; Novak, J.; Nebié, R.C.; Dicko, M.H. Composition and antimicrobial activities of Lippia multiflora Moldenke, Mentha x piperita L. and Ocimum basilicum L. essential oils and their major monoterpene alcohols alone and in combination. Molecules 2010, 15, 7825–7839.
- Soltanzadeh, M.; Peighambardoust, S.H.; Ghanbarzadeh, B.; Mohammadi, M.; Lorenzo, J.M. Chitosan nanoparticles encapsulating lemongrass (Cymbopogon commutatus) essential oil: Physicochemical, structural, antimicrobial and in-vitro release properties. Int. J. Biol. Macromol. 2021, 192, 1084–1097.
- 82. Saquib, S.A.; AlQahtani, N.A.; Ahmad, I.; Kader, M.A.; Al Shahrani, S.S.; Asiri, E.A. Evaluation and comparison of antibacterial efficacy of herbal extracts in combination with antibiotics on periodontal pathobionts: An in vitro microbiological study. Antibiotics 2019, 8, 89.

- Bahmani, M.; Taherikalani, M.; Khaksarian, M.; Rafieian-Kopaei, M.; Ashrafi, B.; Nazer, M.; Soroush, S.; Abbasi, N.; Rashidipour, M. The synergistic effect of hydroalcoholic extracts of Origanum vulgare, Hypericum perforatum and their active components carvacrol and hypericin against Staphylococcus aureus. Futur. Sci. OA 2019, 5, FSO371.
- 84. Musimun, C.; Papiernik, D.; Permpoonpattana, P.; Chumkaew, P.; Srisawat, T. Synergy of greensynthesized silver nanoparticles and Vatica diospyroides fruit extract in inhibiting Gram-positive bacteria by inducing membrane and intracellular disruption. J. Exp. Nanosci. 2022, 17, 420–438.
- Aabed, K.; Mohammed, A.E. Synergistic and antagonistic effects of biogenic silver nanoparticles in combination with antibiotics against some pathogenic microbes. Front. Bioeng. Biotechnol. 2021, 9, 652362.
- 86. Lee, N.L.S.; Yuen, K.Y.; Kumana, C.R. β-Lactam antibiotic and β-lactamase inhibitor combinations. JAMA 2001, 285, 386–388.
- Rakholiya, K.D.; Kaneria, M.J.; Chanda, S.V. Medicinal plants as alternative sources of therapeutics against multidrug-resistant pathogenic microorganisms based on their antimicrobial potential and synergistic properties. In Fighting Multidrug Resistance with Herbal Extracts, essential Oils and Their Components; Academic Press: Cambridge, MA, USA, 2013; pp. 165– 179.
- 88. Rachuonyo, H.; Ogola, P.; Arika, W.; Wambani, J.; Gatheri, G. Combined effect of crude leaf extracts of selected medicinal plants against selected enteric bacterial pathogens and Candida albicans. J. Antimicrob Agents 2016, 2, 1212–2472.
- 89. Obuekwe, I.S.; Okoyomo, E.P.; Anka, U.S. Effect of Plant Extract Combinations on Some Bacterial Pathogens. J. Appl. Sci. Environ. Manag. 2020, 24, 627–632.
- Gadisa, E.; Usman, H. Evaluation of Antibacterial Activity of Essential Oils and Their Combination against Multidrug-Resistant Bacteria Isolated from Skin Ulcer. Int. J. Microbiol. 2021, 2021, 6680668.
- Mutlu-Ingok, A.; Tasir, S.; Seven, A.; Akgun, N.; Karbancioglu-Guler, F. Evaluation of the single and combined antibacterial efficiency of essential oils for controlling Campylobacter coli, Campylobacter jejuni, Escherichia coli, Staphylococcus aureus, and mixed cultures. Flavour Fragr. J. 2019, 34, 280–287.
- Purkait, S.; Bhattacharya, A.; Bag, A.; Chattopadhyay, R.R. Synergistic antibacterial, antifungal and antioxidant efficacy of cinnamon and clove essential oils in combination. Arch. Microbiol. 2020, 202, 1439–1448.
- 93. Garza-Cervantes, J.A.; Chávez-Reyes, A.; Castillo, E.C.; García-Rivas, G.; Antonio Ortega-Rivera, O.; Salinas, E.; Ortiz-Martínez, M.; Gómez-Flores, S.L.; Peña-Martínez, J.A.; Pepi-Molina,

A. Synergistic antimicrobial effects of silver/transition-metal combinatorial treatments. Sci. Rep. 2017, 7, 1–16.

- 94. Alfei, S.; Schito, A.M. β-Lactam Antibiotics and β-Lactamase Enzymes Inhibitors, Part 2: Our Limited Resources. Pharmaceuticals 2022, 15, 476.
- 95. Alfei, S.; Zuccari, G. Recommendations to Synthetize Old and New β-Lactamases Inhibitors: A Review to Encourage Further Production. Pharmaceuticals 2022, 15, 384.

Retrieved from https://encyclopedia.pub/entry/history/show/67554