

Management of Andean Lupin Anthracnose

Subjects: [Biology](#)

Contributor: César Falconi , Viviana Yáñez-Mendizábal

The lupin (*Lupinus mutabilis* Sweet) is a legume domesticated and cultivated for more than 4000 years by the pre-Hispanic cultures of the Andean zone. Due to its good taste and protein content, the lupin seed contributes significantly to the food and nutritional security of the Andean population. However, lupin is susceptible to diseases, and of these, anthracnose is the most devastating as it affects the whole crop, including leaves, stems, pods, and seeds. Seed treatment with fungicides reduces transmission from seed to seedling, but it does not eradicate anthracnose. Attention is given to alternative strategies from sowing to harvest to limit this seed borne pathogen as well as to enhance plant resistance and to promote plant growth. For anthracnose management in the field, integrated practices that encompass control of volunteer plants, lupin ontogenetic resistance, and rotation of biocontrol with chemical fungicides at susceptible phenological stages are considered.

Andean lupin

anthracnose

integrated management

plant protection

dry heat

UV-C

UV-B

Bacillus subtilis

biological control

ontogenetic resistance

chemical control

1. Introduction

Lupinus mutabilis Sweet is a Fabaceae used as food since pre-Inca times in the Andean countries. It is characterized by its high nutritional value, atmospheric nitrogen fixation in soil, and adaptation to various climatic conditions with minimal soil and water requirements ^{[1][2]}. In the Andean areas of Peru, Ecuador, and Bolivia, lupin constitutes one of the main sources of food and income for indigenous populations ^[2]. Since the 1960s, researchers have attempted to solve some of the main agronomic problems related to planting, fertilization, crop rotation, and crop management to improve productivity ^[3], postharvest quality, and marketing ^[4]. However, the food industry demands continuous production of high quality and uniform lupin seeds to sustain their processes ^[5]. Diseases, and particularly anthracnose, have a direct negative effect on production and quality of lupin seed ^{[5][6]}.

2. Andean Lupin

2.1. Common Names

Lupinus mutabilis, in the family of legumes, is similar to *L. albus*, known as lupin in Spain; it is also known as bitter lupin. The local names in the Andean region are “chocho” in Colombia, Ecuador, and northern Peru; tarwi or tarhui in the Quechua language in central and south central Peru; tauri in the Aymara language south of Lake Titicaca in

Peru and Bolivia, and chuchus muti in the Quechua area of Cochabamba [2]. The name ullus is used in different places in southern Peru and Bolivia. The name in English is “Andean lupin” or “pearl lupin” [7].

2.2. Origin

At least four thousand years ago, two ancient cultures in Egypt and the Andes were the first to domesticate and use species of Lupin in their diets: *L. luteus* in Egypt and *L. mutabilis* in the Andes. These species were used for similar food purposes. Curiously, both cultures first developed a maceration and washing processes to eliminate the alkaloids before using the seeds as food [8]. Remains of lupin seeds have been found in the tombs of the Nazca culture (100 to 500 BC) on the desert coast of Peru [9]. In the south of Peru, paintings depicting the lupin in ceremonial vessels from the Tiahuanaco culture (500–1000 AD) are an indication of its wide distribution [2].

2.3. Seed Yield Production

Lupin is characterized by low yields in the Andean zone. The national average yields are 0.40, 0.65, and 1.33 t/ha in Ecuador, Bolivia, and Peru, respectively [10][11]. Australia represents a special case, where currently more than 500,000 ha of sweet lupinus (*Lupinus angustifolius*) are cultivated mainly for use in animal feed [12]. The contribution of Australia to world production of lupin has averaged between 74 and 80% and the crop is produced in the grain belt of Western Australia [13]. Average yields in Australia and Western Europe exceed 1.5 t/ha [14].

It might be thought that low yields in the highland Andes are due to adverse conditions for plant growth, such as low temperatures, droughts, or soils deficient in nutrients, but this is not always the case. On experiment stations, as well as on high input farms where appropriate soil fertilization is used, Andean lupin yields often exceed 1.8 t/ha [11][15]. The large gap between achievable yields and current national averages suggests a tremendous potential for improving livelihoods through more profitable and sustainable production of Andean lupin cultivation.

2.4. Nutritive Value and Potential Pharmacological Use

After a debittering process, the Andean lupin grain is used for human consumption [16]. The nutritional composition of the cultivated lupins varies greatly among species. *L. mutabilis* contains the highest level of protein and oil in seeds [17]. Protein content varies from 41 to 51% on a dry weight basis [18][19]. The low proportion of testa (13%) in the seed of *L. mutabilis* could explain the high levels of protein in whole seeds compared with others lupin species [17]. Grain lupin protein compares favorably with other legumes, such as soybeans (Table 1). Lupin is sometimes referred to as “Andean soya” due to it has the potential for improving human nutritional standards in the Andean Highlands [17], and currently is also of high interest in European countries [20][21]. Oils and crude fiber contents are similar to that of soybean (Table 1). Oils in Andean lupin seeds are of high quality and appropriate for human consumption [22]. The fibre in whole seed, when finely ground, proved to be highly digestible in pigs and increased the digestible energy in pigs to higher than expected levels [22]. The specific carbohydrate composition of Andean lupin grain is related to anti-nutritional factors [12], characterized by low levels of starch and high levels of non-starch polysaccharides (NSP) (Table 1). In relation to the composition of amino acids, Andean lupin grain has lower content of the nutritionally essential, sulfur-containing amino acids (methionine and cysteine) [12] and, like

most legumes, is deficient in tryptophan and tyrosine (**Table 1**). Therefore, it is important to consume lupin accompanied by a cereal such as corn, rice, or wheat [23] and even better in combination with quinoa, which results in an ideal food to combat malnutrition [2].

Table 1. Chemical composition of Andean lupin compared with soybean.

	<i>L. mutabilis</i> *	<i>Glycine max</i>
Crude protein (% of DM)	44.3	33.4
Oil (%)	16.5	16.4
Non-starch polysaccharides (%)	28.2	18.0
Crude fiber (% of DM)	7.5	6.45
Essential amino acids (g/16 gN)		
Glutamic acid	13.79	15.5
Aspartic acid	6.85	11.0

Ancestral knowledge was the starting point for a group of researchers who set out to isolate and characterize secondary metabolites in each of the fractions of this plant (leaves, flowers, stems, grains) [24] and validate its use in the pharmaceutical field [2]. Lupin alkaloids show a good antimicrobial effect against bacteria and fungi of clinical interest [25]. Lupin has been used to enrich wheat bread, and lupin-wheat bread has a texture quite similar to standard wheat bread, but the average content of protein, fat, and dietary fiber increased significantly, and its regular intake lowered hydrolytic and glycemic indexes of those who consumed it [26]. At a pharmacological level, frequent consumption of *L. mutabilis* grain resulted in improved metabolic control, decreased blood pressure, and increased HDL-cholesterol in type 2 diabetes-mellitus (T2DM) patients [27].

2.5. Andean Lupin in Livestock Farming Systems

In addition to its nutritional properties, the cultivation of lupin is very beneficial for the productive system of the highlands [3][19]. Its roots fix atmospheric nitrogen and therefore contribute to improved soil fertility. Because of this, lupin is part of the traditional systems of crop rotation of the Northern Altiplano, including: quinoa-lupin-potato, quinoa-lupin-barley, or quinoa-lupin-quinoa [6], without use of other fertilizers. Potato usually yields more and better quality. Lupin incorporates into the soil a nitrogen equivalent of 150 to 300 units, which in economic terms would be like providing 7 to 10 bags of urea per ha, and that is a very important contribution to small farmer incomes [28]. In addition, some soil-borne pathogens reduce their levels in soils due to a non-host lupin crop rotation [13].

3. Methods to Eradicate the Pathogen from Infected Seed as the Primary Means of Pathogen Survival

3.1. Traditional Methods for Seed Disinfection

Infected seed is the primary inoculum for *C. lupini*. Unfortunately, in the Andean zone most lupin growers use their own seed, which is often infected [3][29]. Local farmers themselves collect the lupin seed. Due to the good price in the market, they prefer to sell commercial seed and keep small, wrinkled seeds with small reddish-brown spots for the next sowing [29]. Wrinkled and/or decolorized seed usually means that the seed is infected [30][29]. This practice makes it more likely that the next generation of plants will be anthracnose infected and is one of the main reasons for low productivity of lupin in marginal areas. The farmers need to be more aware of how important it is to use good quality seed and learn about the use of good methods for seed disinfection [29].

3.2. Alternative Methods for Seed Disinfection

Emergence of lupin is notoriously irregular in the Andean countries for several reasons, but mainly because the seed has poor sanitary quality and variable physiological development. The social context is an important factor in lupin anthracnose management in the Andean zone [29]. Farmers generally wait for a significant portion of plants to emerge before making a fungicide treatment, and by this time many plants have been exposed to inoculum. Researchers have treated seed with systemic fungicides prior to planting to solve the problem of early infection [3], but this generally does not protect plants after emergence [31]. The use of alternative methods, alone or in combination with chemicals, can offer a greater spectrum of action and higher efficiency in seed disinfection.

3.2.1. Dry Heat

A valuable option to reduce fungal infections in seeds is dry heat treatment. Studies conducted in Australia have shown that *L. angustifolius*, *L. luteus*, and *L. albus* seed heat treated with 60–80 °C for a two-week period or less reduced lupin anthracnose infection to an undetectable level, and these treatments did not greatly reduce seed germination [32]. In *L. mutabilis*, treatments at 65 °C on infected seed for 24 h reduced disease incidence to undetectable levels. In addition, pathogen transmission from seed to seedlings was reduced by 75% with 8 h dry heat treatment (Table 2) [33].

Table 2. Alternative methods for the management of Andean lupin anthracnose infected seed.

Method	Doses—Exposure Time	Infection reDuction Efficiency	Additional Benefit on Lupin Plant
Dry heat	65 °C—8 h	75%	Promote seedling emergence [33]
UV-C	57.6 kJ/m ²	90%	Enhance physiological and biochemical processes in seedlings [34]
UV-B + thermal radiation	2.83 kJ/m ² around 76 °C—45 min	95%	Seed invigoration, positive response in plant developed, increase activity of antioxidant enzymes [35]

Method	Doses—Exposure Time	Infection reDuction Efficiency	Additional Benefit on Lupin Plant
<i>B. subtilis</i>	Living cells (LC) from 72-h-old cultures, 10 ⁹ CFU/mL or cell-free compounds (CFC)	76 to 100% (LC), 68 to 96 (CFC)	Positive effect on plant physiology, stimulate host defense and growth mechanisms [36][37][38]

germination of *L. luteus* and *L. albus* [32]. In *L. mutabilis*, exposure times of 12 h at 65 °C reduced germination [33], comparable with the results reported by [32], with reductions of seed germination on *L. luteus* after 60 or 70 °C treatment for one week. However, treatments of 65 °C for 8 h on *L. mutabilis* did not greatly affect emergence of seedlings [33], comparable with the results reported by [32] on *L. angustifolius* exposed for 4 days at 65 °C. Development of *L. angustifolius* cultivars resulted from an introgression of a large number of wild and external accessions within a breeding program in which selections were made for seed adapted to weather extremes [39]. In contrast, *L. mutabilis* cultivars were selected based on agronomic traits and adaptation to the cold environments of the tropic highlands. Seed coat thickness of *L. mutabilis* is 50% lower and saturated fatty acids are 60% less than in *L. angustifolius* [40][41]. These facts may confer less resistance of *L. mutabilis* to dry heat in comparison with *L. angustifolius* [33].

Interestingly, emergence of seedlings increased with dry heat treatments at 65 °C for 8 h in four *L. mutabilis* cultivars (Table 2) [33]. This seedling emergence promotion has also been observed for cotton seed dry heat treated for 8 h at 60 °C [42]. Treatment of Andean lupin seed with moderate dry heat exposure time at 65 °C reduces anthracnose infections and promotes seedlings emergence.

2.2.2. UV-C Radiation

One of the natural components of sunlight is ultraviolet radiation (UV), which is divided according to its wavelength into long UV-A (315–390 nm), medium UV-B (280–315 nm), and short UV-C (100–280 nm). Within these subtypes, UV-C radiation is used as an effective biocidal agent for the disinfection of water, surfaces, and the air [43]. Similarly, seed infections can be reduced or eradicated by UV-C treatment. Treatment of anthracnose infected Andean lupin seed reduced infections by 90%, whereas doses of 86.4 kJ/m² reduced infection to undetectable levels [34]. In other pathosystems, the mycelium of *Colletotrichum musae* was inhibited with 45 min of exposure to UV-C radiation, whereas treatments of 2 h reduced the diameter of the lesion in bananas inoculated with the same pathogen [44]. UV-C exposure times shorter than 20 min suppressed microbial decay on mangoes when the fruit received full irradiation [45]. Legumes, grains, and vegetables seeds are suitable for UV-C treatment. Optimal doses vary according to the species and are in the range of 0.025–480 kJ/m² [46]. The direct germicidal effect of UV-C radiation could be the primary factor leading to disease suppression [47]. The control of anthracnose in lupin through the use of UV-C radiation is beneficial.

UV-C radiation may also reduce seed germination. Exposures for 24 h reduced germination of *Acacia ampliceps* seed from 80.1% to 65.2%. However, UV-C exposure of 6 h did not critically reduce germination [44]. In *L. mutabilis*, UV-C exposures for 8 h reduced seed borne anthracnose and had a slight effect on seed germination [34].

UV-C seed treatments enhance some physiological processes in seedlings. Andean lupin seedlings grown from UV-C pretreated seed had equal protein and chlorophyll content compared to those grown from non-infected seed, and higher than seedlings grown from infected seed (**Table 2**) [34]. UV radiation doses stimulate the biosynthesis of UV-absorbing compounds and carotenoids, both of which perform a photoprotective role [48][49]. Red bean seedlings grown from seed pretreated with low doses of UV-C significantly increased protein content [50]. The plant response to environmental stress is given by the breakdown and recycling of proteins, which depends on the levels of proteolytic enzymes [51]. The pretreatment on lupin seed with UV-C for 8 h did not affect total chlorophyll and proteins, and doses of 57.6 and 86.4 kJ/m² resulted in higher seed emergence and greater seedling dry weight. The positive effects were even better than that of fungicide seed treatment and contributed to further seedling establishment [34]. Oxidative stress in plants may be induced by UV radiation, increasing reactive oxygen species [51]. Several UV-absorbing compounds that activate antioxidant defense mechanisms are accumulated by plants to cope with this stress [52]. For instance, a significant increase in peroxidase and catalase activity was found in papaya peel as a result of UV-C treatment compared to initial values [53]. Components of cross-linking cell wall, polymerization of lignin and suberin monomers are associated with peroxidase enzymes and subsequent resistance in other host–pathogen interactions [54]. In the work of Falconi and Yanez [34], Andean lupin seedlings that grew from UV-C treated seed recorded higher enzymatic activity, similar to what occurs in tomato plants treated with azoxystrobin or *Pseudomonas fluorescens*, other treatments that induce plant resistance [55]. As a result of the research carried out by Falconi and Yáñez [34] on the use of UV-C for treatment of *L. mutabilis* seed and its further effect on plant physiology and biochemistry, it is possible to hypothesize that UV-C on lupin seed acts as an inducer of systemic resistance. Low incidence of anthracnose observed in lupin seedlings grown under greenhouse conditions can be associated with changes in enzymatic activities [34].

2.2.3. UV-B + Thermal Radiation

Solar radiation can efficiently reduce the survival of plant pathogens. Several types of fungi, such as *Pseudoperonospora cubensis* [56], *Peronospora tabacina*, *Uromyces phaseoli*, *Alternaria solani* [57], and *Phytophthora infestans* [58] have reduced viability as a result of exposure to solar radiation. The UV spectrum of sunlight causes most of the harmful effects on biological systems. Of the total solar radiation, approximately 8–9% corresponds to ultraviolet radiation. Although UV-B (280–315 nm) represents only 1.5% of the total UV spectrum, is of particular interest because it is the most active form [59]. Conidia of most species of phytopathogenic fungi directly exposed to UV-B radiation for a few hours can lose viability, but sub lethal doses of solar UV-B radiation can also delay conidia germination or reduce their virulence [60]. Differences in spore pigmentation may be associated with the effect of UV radiation on spore viability, as has been reported in other plant pathogenic fungi. For instance, spores of *Peronospora tabacina* were shown to be more sensitive to solar radiation than the more pigmented spores of *Alternaria solani* and *Uromyces phaseoli*, which supports the hypothesis that spore pigmentation affects sensitivity to solar radiation [61]. The effect of UV-B alone was determined [61] by exposing Andean lupin seed lots at ambient temperature under an absorption UV-B film for 60 min. UV-B doses of 3.75 kJ/m² reduced anthracnose infection seed by 60% [35]. Although *Colletotrichum* species can overwinter as mycelia under the seed coat [62], solar radiation reduced its viability [35].

Seeds of some plant species have seed coats that are more resistant to UV-B rays [63]. In this sense, depending on the plant species, low to moderate doses of UV-B may not reduce seed germination, and may even increase it [64], whereas high doses of UV-B radiation can reduce seed germination [65]. In addition, solar radiation generates heat, and this radiated energy can also have an effect on disease [66][67]. However, seed exposure to UV-B radiation can induce a variety of responses in seedlings [68]. A solar oven was designed by Falconi and Yanez [35] to collect thermal radiation. Because it is unknown how protective the Andean lupin seed coat is against the sun's extremely incandescent gases, the oven was covered with a film that partially absorbs UV-B. Thus, only the portion of UV-B that enters the solar oven will generate thermal radiation that will be transmitted to the infected seed (**Figure 1**).




Figure 1. UV-B radiation solar oven schematic representation to treat Andean lupin seed lots infected with anthracnose. Source: [35].

Anthracnose infection in lupin seed was reduced more efficiently in the solar oven where the cumulative UV-B radiation and high temperature were combined, as compared to the individual effect of UV-B radiation at ambient temperature or of dry heat. On sunny days between 11 am and 2 pm, UV-B doses of 2.83 or 3.75 kJ/m² were reached after 45 or 60 min, and temperature fluctuated between 73 and 79 °C in the solar oven; the ambient temperature varied from 19 to 23 °C. The high temperatures in the oven represent harsh conditions for seed borne anthracnose [35]. Anthracnose seed infection was reduced by 95% with UV-B doses of 2.83 kJ/m² in the solar oven (**Table 2**), being more efficient than the isolated dry heat treatment at 65 °C for 60 min or the same UV-B dose of 3.75 kJ/m² at ambient temperature. Viability of anthracnose and its subsequent potential plant infection were reduced with pretreatments of infected seed in the solar oven. Acervuli and mycelium of *C. lupini* living in the inner layer of the testa of lupin seed may be radically affected by active UV-B plus temperature [35].

The pretreatment of seed with UV-B irradiation can induce a variety of responses in seedlings and plants [69], but information about seed invigoration due to the combination of UV-B radiation with high temperature is scarce [69].

Combined UV-B doses and temperature achieved after 75 min in the solar oven decreased viability of *L. mutabilis* seed. However, moderate exposure times of 45 min, effective for seed borne disease control, do not reduce germination percentage [35]. Germination percentages of buckwheat, Chinese cabbage, turnip, cabbage, or parsley seeds were not greatly reduced with UV-B exposure times from 10 to 45 min [64]. New studies could elucidate the combined effect of UV-B and high temperature on seed invigoration and its potential expression into positive responses in the plant.

The synthesis of some heat shock physiological and biochemical processes in plants may be stimulated by pretreatment with UV-B thermal-heat of seed. No oxidative damage of chlorophyll and proteins was detected in Andean lupin seedlings grown from seed pre-treated for 45 min. Conversely, these exposure times enhanced peroxidase activity in seed-lings (Table 2) [35]. Diseases, high UV-B levels, and extreme temperatures can elevate the amounts of reactive oxygen species (ROS), including superoxide radicals and hydrogen peroxide [65]. These molecules are very destructive as they participate in many chemical reactions with lipids, proteins, and nucleic acids resulting, for example, in damage to cell membranes due to lipid peroxidation [66]. However, increased activity of antioxidant enzymes that serve as “sunscreens” may also be elicited as an acclimatization response to UV-B solar radiation [67]. For the elimination of H₂O₂, the antioxidant defense system uses peroxidase as an important component [68][69], purifying and participating in the polymerization of lignin and suberin monomers and other cross-linked cell wall components [70]. Dry heat pretreatment of cotton seed at 80 °C for 8 h enhanced physiological and biochemical process that have a photoprotective role for seedlings [42]. In another study using *Indigofera tinctoria*, peroxidase activity slightly increased with supplemental UV-B radiation for 2 h per day for 2 days [71]. In this sense, the combination of UV-B and the high temperature achieved in the solar oven designed by Falconi and Yanez [35] may have contributed to biochemical changes (Table 2). Future studies could elucidate whether such UV-B treatments of lupin seed enhance molecules implicated in host defense or plant growth development.

2.2.4. Biological Treatment for the Seeds

Biological treatments to control anthracnose in seeds based on the use of microorganisms, such as *Bacillus* spp., have been described as effective. Studies conducted by Yanez et al. [36] demonstrated that several *Bacillus subtilis* strains collected from lupin fields strongly suppressed *C. lupini* on Andean lupin seed [36]. Furthermore, living cells, cell-free compounds, and lipopeptides obtained from these strains significantly reduced anthracnose on seeds (Table 2 and Figure 2A–D) [37][38]. The efficacy of these strains is due to their biological characteristic to produce antifungal lipopeptides that progressively inhibited the mycelial growth and conidial sporulation and killed the fungal structures of *C. lupini* (Figure 2D) [36][37][38].




Figure 2. Antifungal activity of *Bacillus subtilis* against *C. lupini*. Black arrows show pathogen mycelial inhibition by bacterial biomass (A), cells free supernatant (B), and lipopeptides extract (C) compared with untreated control. Microscopic images of pathogen from growth zones (D) showing deformation and death of mycelia from bacterial treatments compared with untreated control. Source: [36][37][38].

Several studies found that the broad antimicrobial effect of *Bacillus* spp. against pre- and postharvest disease is due to production of powerful bioactive compounds. *Bacillus* spp. mainly produce cyclic lipopeptides from three families: fengycins, iturins, and surfactins [36][37][72][73][74][75][76][77][78][79]. For anthracnose, the biological control potential of native *B. subtilis* strains by the production of these lipopeptides has been demonstrated [36][37][38]. Living cells of *B. subtilis*, including vegetative cells and endospores, showed a strong biocide effect (Figure 2A) against *C. lupini*. Cell-free supernatant produced by these cells showed the same biocide effect as living cells (Figure 2B,C). Thin layer chromatography analysis (TLC) of *Bacillus* lipopeptide extracts demonstrated the presence of fengycin, iturin, and surfactin families [36][37][38] (Figure 2B). Furthermore, when TLC analysis was combined with the growth inhibition test in TLC-bioautography, the lipopeptide fractions corresponding to fengycin showed the highest antifungal activity against *C. lupini* when compared with iturins and surfactin fractions (Figure 2B). These results strongly support the role of lipopeptide production as the major antifungal factor used by *Bacillus* spp. [36][37][38] against *C. lupini*, similar to that described for other biocide compounds with antifungal activity against other *Colletotrichum* species in plants [76][77].

The biocontrol ability of native *B. subtilis* was confirmed by antifungal lipopeptide-related genes. PCR analysis, which showed fengycin E, iturin B, and surfactin B gene amplification associated with biocide fractions detected by TLC-bioautography, supported the role fengycins and iturins play as the major antifungal factors of *B. subtilis* native strains [37]. From a physiological and pathological perspective, the antifungal action of fengycins is due to their ability to interact with wall components of the fungal pathogen, such as ergosterol, and to alter its cell packing structure and permeability in a dose-dependent way [78] (Figure 2D). This mechanical damage on cell pathogen structures was reported by [77][78][79], who demonstrated that fengycins produced by *B. subtilis* were able to cause

morphological alterations on a wide range of plant fungal pathogens by formation of a fengycin–ergosterol complex. Thus, the susceptibility of *C. lupini* to *B. subtilis* strains mediated by lipopeptides could be explained by the effect of fengycins on the target membranes, leading to the inhibition of mycelial growth, conidial germination, and subsequent pathogen death [37].

The efficacy of native *B. subtilis* strains to control anthracnose was evident when inoculated on *L. mutabilis* seeds, verifying that the *B. subtilis* strains have high antifungal activity, inhibit mycelial growth and conidia germination, reduce seed infection to undetectable levels, and increase lupin germination percentage (Table 2, Figure 3A) [37] [38]. In addition, rhizobacteria trigger induced systemic resistance (ISR) of plants against pathogens through the production of several bioactive metabolites such as antibiotics, siderophores, and volatile compounds [79][80][81][82] [83][84]. *Bacillus* spp. can elicit systemic resistance by inducing the synthesis of antioxidant defense enzymes [85][86] [87]. Native *B. subtilis* strains also promoted the synthesis of catalase, peroxidase, and superoxide dismutase in Andean lupin seedlings, which are precursor substances of plant resistance (Figure 3B) [38]. *Bacillus* spp. increased synthesis of antioxidant defense enzymes and resulted in induced systemic resistance against other plant pathogens in tomato seedlings [83][84], and suppressed anthracnose disease of chili by the activation of defense-related enzymes [85]. Similarly, *B. subtilis* is capable of impairing disease incidence, promoting seedling growth, and increasing activities of antioxidant enzymes (POD, PPO, PAL) in cucumber plants [86]. Native *B. subtilis* promoted the growth of Andean lupin radicles and plumules in pre-inoculated seed (Table 1, Figure 3A) [37]. Plant hormone biosynthesis by *Bacillus* spp. has been directly related to subsequent growth promotion in different plants [83][84].




Figure 3. Effect of *Bacillus subtilis* treatments (A) in the expression of the *Lupinus mutabilis* growth-/ defense-related genes. (B) Relative quantity expression (RQ) of PIN-1, PIN-3, SOD-2, PR-1, and PR-4 genes induced by *B. subtilis* cells, lipopeptide extract, or sterile PBS as control was normalized against actin gene. Source: [38].

4. Strategies to Suppress Secondary Infection and Pathogen Spread

4.1. Biological Control in the Field

From a commercial point of view, the use of *Bacillus*-based preparations has been widely demonstrated for efficiency controlling multiple field fungal pathogens [84]. However, research for the biocontrol of lupin anthracnose is scarce. Studies conducted in the field with anthracnose natural inoculum pressure showed that all *B. subtilis* treatments reduce the area under disease progress curve (AUDPC) compared with controls that showed large stem lesions, accompanied by necrotic tissue 90 days after sowing (DAS) and abundant sporulation, and in some cases dead plant from 105 to 120 DAS [88]. Lupin anthracnose can crack the stem at the site of infection (Figure 1A); multiple lesions can lead to stem twisting, or to complete stem collapse [89][30]. The AUDPC reduced with sequential applications of antagonists 1×10^9 UFC/mL, every two weeks, compared to the AUDPC of untreated control in susceptible cultivars (Figure 4). In general, treatments with bacterial suspensions result in healthier stems, leaves, and pods, which are agronomic characteristics of lupin linked to production [88]. The significant reduction of AUDPC may be due to *B. subtilis* having a cascade effect on the different components of the disease triangle, that is, on the environment, the host plant, or the pathogen. Therefore, it can affect the onset and progress of the disease and also turn on resistance genes in the host [90]. *B. subtilis* induces disease resistance in the plant [82][84][85], or in turn promotes plant growth, a fact that makes it easier for plants to control pathogenic infections [79][81][86] or suppress diseases based on systemic resistance [82][83][84][86]. *B. subtilis* have also shown to control ginseng anthracnose, *Colletotrichum panacicola*, where leaf lesion diameter was not significantly affected compared with the fungicide treatment [91]. The protection mechanism of *B. subtilis* was associated with reduction of the incidence and with the initial processes of the infection cycle such as spore germination, appressorium formation, and penetration [90].




Figure 4. The area under disease progress curve (AUDPC) of anthracnose caused by *C. lupini* on the Andean lupin susceptible cultivar I.450 Andino. Natural pressure of inoculum represented as the control (Ck) and the AUDPC by effect of *Bacillus subtilis* Ctpx-S1 , Ctpx-S2 , and Ctpx-Z3 . AUDPC calculates scores for lupin anthracnose severity on the 1-to-6 scale [92]. Pooled data from the 2015 and 2019 growing seasons at El Chaupi, Cantón Mejía, Ecuador, each point at disease onset plus 90, 105, and 120 days. Source: [88].

A *B. subtilis* strain selection program generally begins with screening under controlled laboratory or semi-controlled greenhouse conditions; however, it is difficult to predict how the bacteria will respond when released into the natural environment [93]. The native bacterial population on the phylloplane was estimated to confirm its protectant effect. Viable cell concentrations of *B. subtilis* were reduced in approximately 2.0 logarithm after 15 days of the first application on Andean lupin (**Figure 5**) [88] or reduced by 50% in strawberry leaves in the open field after 8 days of application [93]. Environmental stresses such as intense sunlight, dryness, or high temperature could reduce initial colonization [88][90], or rainfall could easily remove the bacterial population from the plant surface in several hours [93]. However, the population of *B. subtilis* will vary to some degree naturally once it has been established in the field. The average of native *B. subtilis* population on the phyllosphere of lupin remains stable at approximately 7.0 LOG after successive two-week-sprays for two months, which suggests that after the population has become stable, it can resist harsh environmental conditions (**Figure 5**) [88].




Figure 5. Population of *Bacillus subtilis* Ctpx-S1 , Ctpx-S2 , and Ctpx-Z3 recovered from the phyllosphere of lupin cultivar I-451 Guaranguito during 2015 and 2019 growing seasons. First spray at disease onset and fortnightly for eight weeks. Colony counted and results expressed as CFU *B. subtilis* per gram of lupin leaves. To improve homogeneity of variances, data of bacterial concentration were log 10-transformed (\log_{10} CFU/g). Each point represents the mean \pm SD of four independent repetitions of 10 g leaves. Source: [88].

The total viable epiphytic population on the phylloplane can be determined by a serial dilution plating method considering that bacteria can colonize stomata, trichomes, vein endings, cell wall junctions, or even be beneath the leaf cuticle [94]. In the work of Falconi et al. [88], lupin stems and leaves sprayed with bacterial antagonists were consistently protected from natural infection of *C. lupini* fungi. Control was evident in a significant reduction in the AUDPC and suggests that native *B. subtilis* are more likely the cause of reduction in anthracnose severity. In addition, plants release organic compounds, such as sugars, organic acids, and growth regulators [95] that can stabilize cell concentration of *B. subtilis* due to nutrient availability [96]. Thus, when bacterial antagonist adhered, invaded, and survived on flowers, they effectively prevented the pathogen from colonizing the flowers [93]. Addition of 0.5% unrefined sugar to *B. subtilis* suspensions provides an initial food supply and increases adherence and colonization of bacteria in cocoa pods, improving monilia pod rot control [96][97]. Other studies show that a single inoculation of *B. pumilus* reduced banana sigatoka, *Mycosphaerella fijiensis*, by 33.6% and delayed progress of the disease by 21–28 days compared with the control [98]. Lower AUDPC values in comparison with the control indicates reduction of the initial inoculum or reduction of pathogen on the plant [98]. By associating prior findings [36][37][38] with the field observations where infections and disease severity were reduced, it can be speculated that biocontrol by native *B. subtilis* resulted not only from damage of cellular structures (hyphae, mycelium, spores) of the pathogen, as occurred in the in vitro studies, but also by induction of systemic resistance (ISR) and promotion of plant growth mechanism (PGM). An innovative strategy for the integrated management of anthracnose in lupin could involve the activation of ISR by *B. subtilis*. Future studies could demonstrate more efficient use of these native strains and perhaps allow the establishment of biocontrol strategies.

4.2. Chemical Control

As there are no resistant varieties of Andean lupin and in the absence of any accurate method of controlling the disease, chemical control has been recommended as the most effective measure to reduce the spread of the disease. Synthetic fungicides reported for control of lupin anthracnose in the Andean zone and around the world are listed in **Table 3**. Seed treatment with fungicides reduces transmission of the pathogen from seed coat to seedling, but it does not provide complete control [99]. The synthetic fungicides generally recommended for controlling anthracnose disease are based on copper compounds, dithiocarbamates, benzimidazole and triazole compounds, manganese ethylenebisdithiocarbamate (Maneb) [99][100][101], and carbendazim [100], although the use of fungicides has been found ineffective under severe disease outbreak [99]. Newer chemicals like strobilurins (e.g., azoxystrobin, pyraclostrobin) have also been used for lupin anthracnose management [102][99][103]. Different classes of fungicides have specific modes of action and also differ in the duration of disease control. Farmers should take prevailing environmental conditions into consideration in their choice of fungicides. Rotation of two or more different classes of fungicides is highly recommended for increasing the chance of better protection against the disease in the fields [101][103][104][105]. Applications of Boscalid + Piraclostrobin or Azoxystrobin + Difeconazole significantly reduced the severity and infection in seed and significantly increased grain production [102].

Table 3. Fungicides and bioproducts reported for the control of lupin anthracnose.

Active Ingredient	Chemical or Biological Group	Target Site *	FRAC Code **	References
Iprodione	dicarboximides	MAP/Histidine-Kinase in osmotic signal transduction	E3/2	[99][100]
Procymidone	dicarboximides	MAP/Histidine-Kinase in osmotic signal transduction	E3/2	[99][100]
Carbendazim	benzimidazoles	β -tubulin assembly	B1/1	[99][100]
Difeconazole	triazoles	C14- demethylase in sterol biosynthesis	G1/3	[99]
Copper	inorganic	Multi-site contact activity	M 01	[99]
Mancozeb, Maneb	dithiocarbamates	Multi-site contact activity	M 03	[99][100][101]
Captan	phthalimides	Multi-site contact activity	M 04	[99]
Folpet	chloronitriles	Multi-site contact activity	M 05	[99]
Clorotalonil	chloronitriles	Multi-site contact activity	M 05	[99][101]
Dithianon	quinones	Multi-site contact activity	M 05	[103][104]
Ciprodinil + Fludioxonil	anilino-pyrimidines + phenylpyrroles	Multi-site contact activity	M 09	[104]
Fenhexamid + Tebuconazole	hydroxyanilides+ triazoles	3-keto reductase, C4- de-methylation + C14- demethylase in sterol biosynthesis	G3/17 + G1/3	[104]
Azoxistrobin	QoI-fungicides	Complex III: cytochrome bc1	C3/11	[99][100][101]
Pyraclostrobin + Boscalid	methoxy-carbamates + pyridine-carboxamides	Complex III: cytochrome bc1 at Qo site + complex II: succinate-dehydrogenase	C3/11 + C2/7	[102]
Azoxystrobin + Difeconazole	QoI-fungicides + triazoles	Complex III: cytochrome bc1 at Qo site + C14- demethylase in sterol biosynthesis	C3/11 + G1/3	[102]
<i>B. subtilis</i>	microbial	Multiple effects antibiosis, membrane disruption by fungicidal lipopeptides, lytic enzymes, induced plant defense	BM 02	[38][84][106]

Effective control through the use of chemical fungicides is possible by timely application during the critical period favorable for the onset of the disease [99][100][101][103][104]. Generally, fungicides should be applied for seed disinfection [99], at young seedling emergence, including early flowering or full bloom, and early pod formation to restrict the entry of the pathogen to the plant system [99][100][101][103][104].

Rotating fungicides with different FRAC codes can help delay development of lupin anthracnose resistance to fungicides. Code numbers on fungicide labels, called “FRAC” groups, can help to develop chemical rotations that delay fungicide-resistance (**Table 4**). The optimal application time is when the *C. lupini* population is small and contained, when weather conditions make the pathogen susceptible to fungicide, or when lupin is at the most susceptible growth stage (**Figure 6**).




Figure 6. Anthracnose susceptible phenological stages of lupin.

B. subtilis for control of lupin anthracnose is also listed in **Table 3**. Application of sus-pensions based on *B. subtilis* [38][88][106] into a rotation with foliar fungicide containing mancozeb, chlorothalonil, or azoxystrobin [102][107][108] at sowing, plant emergence, early flowering, and early pod formation can better prevent transmission of the pathogen to the seedling and suppress anthracnose in the field. Using a biocontrol agent can reduce damage to human health and the environment.

4.3. Control Volunteer Lupins and Alternative Hosts

Lupin production in the tropical Andean highlands varies in the amount planted, which in general follows local rainfall patterns and prevailing prices. The highland tropics are also the home of several alternative hosts for anthracnose, including tamarillo (*Solanum betaceum*), that are geographically grown in neighboring lands with lupin and give the pathogen more opportunities to reproduce and survive [109]. Other studies also showed that *C. lupini* was able to cross-infect other host plant and *Colletotrichum* species from soybean also cross-infected lupin with varying degrees of virulence [110]. The plasticity of the interaction of *Colletotrichum* spp. with different hosts is significant. In addition, several species of wild lupin, such as lupin afelpado (*L. alopecuroides*), lupin rastrero (*L. sarmentosus*), miniature lupin or allpa chocho in Quechua language (*L. microphyllus*), or Quito’s chocho (*L. pubescens*) [111], grow in tropical highland areas and thereby act as a reservoir for pathogen growth during drier periods [112]. *L. cosentinii* can act as another source of infection in Western Australia, where control of naturalized populations is an important component for disease management [13]. In the Andean areas, high levels of inoculum could come from other susceptible cultivated or wild lupins and other alternative hosts in the surrounding areas; consequently, sanitation at the field or farm level is very important for anthracnose management [109].

4.4. Ontogenic Resistance

The ability of whole plants or plant parts to resist or tolerate disease as they age and mature is called ontogenic or age-related resistance. Levels of resistance that develop during aging of plant tissue may greatly reduce eventual disease severity and may even lead to escape from infection or immunity [113]. In lupin, anthracnose resistance is not equally expressed at all developmental stages. An ontogenic model was developed for three lupin species—*L. angustifolius*, *L. luteus*, and *L. albus*—and integrated into a decision support system for anthracnose management. The beginning of flowering and full bloom represent growth stages with high susceptibility [114]. Based on a series of in vivo assays involving Andean lupin genotypes, early cotyledonal stages (2–3 leaves) and the beginning of flowering (9–11 leaves) were susceptible to anthracnose, whereas the vegetative development stage (6–8 leaves) was resistant [92]. Cotyledonal and flowering stages of *L. mutabilis* were confirmed as susceptible to anthracnose through spectroradiometry and unmanned aerial vehicles [115]. This plant response of susceptible genotypes of Andean lupin in Ecuador [3] reflects the current situation of resistant and susceptible cultivars of *L. angustifolius* in Australia [99][116], where epidemics of lupin anthracnose progress more slowly in plots of intermediate-aged plants and dramatically increase during flowering. Some stage-specific resistance genes might not be expressed in early and late lupin developmental stages [92]. Targeting disease susceptibility during the life cycle of the Andean lupin plant will contribute to development of a more viable management option for production in areas of high anthracnose risk in Ecuador and maybe to more efficient lupin anthracnose management around the world. Early growth, beginning of flowering, and early pod formation [92][99][114][115] are the stages when lupin is most susceptible to anthracnose. An integrated ontogenic plan can be used during these major plant susceptibility periods for scheduling application of biological or chemical treatments.

4.5. Integrated Methods at Critical Phenological Stages

No single management technique has been found to efficiently control lupin anthracnose, therefore integration of techniques is essential. The first critical step is to reduce infection in seed (**Figure 5**) by using dry heat [33], UV-C radiation [34], or UV-B radiation [35]. Further protection can be added by combining physic treatments with other techniques, such as biological [36][37][38][83][87] or chemical control [102][99][100][101][103][104]. If one of the physic treatments of seed is used for high rainfall regions, it should be supplemented with foliar sprays to suppress pathogen dispersion in the field and subsequent infections. Application of fungicides on flowering and podding stages reduced incidence and increased yield of susceptible and resistant cultivars grown in high-risk areas [99].

Applications of *B. subtilis* theoretically prevent the establishment of the pathogen, interrupt its development, promote plant growth, and induce acquired resistance in lupin [38]. Because local results obtained under field conditions are consistent [88], bacterial suspensions at 1×10^9 UFC/mL should be applied prior to pathogen infection, at sowing time (**Figure 6**). Success in controlling lupin anthracnose disease can be achieved with a sequential two-week application [88]. Shorter interval applications of *B. subtilis* are often recommended during the rapid growth stages of the plant to obtain a commercially acceptable product [106]. Special attention should be given to cotyledonal, early flowering, full bloom, and pod-filling stages (**Figure 6**) because lupin has been shown to be more anthracnose susceptible at these phenological stages [90][92][114][115].

Other technological components are needed to further reduce the use of synthetic pesticides. For this, it is necessary to develop *B. subtilis* at a semi-industrial or industrial level or to use the pathogen-inhibiting substances extracted from bacteria in rotation plans with chemical products. These rotation plans would include the interaction of *B. subtilis*-based biopesticides with protective and systemic fungicides to determine dose, frequency, and times of application [117]. A low-cost medium for *B. subtilis* lipopeptide production is available [118], and a spray drying formulation [119][120] has been developed to expand survival and reproduction of the antagonist under field environmental conditions. A plan that includes the rotation of biological fungicides with chemicals and their application during the most susceptible plant growth stages will greatly minimize infection in the field and reduce losses at harvest for a more sustainable production of Andean lupin, until resistant varieties are developed.

References

1. Mujica, A.; Moscoso, G. La planta del tarwi. In *Lupinus mutabilis* (Tarwi) Leguminosa Andina Con Gran Potencial Industrial; Zavaleta, A.I., Ed.; Fondo Editorial de la Universidad Nacional Mayor de San Marcos: Lima, Perú, 2018; p. 11. 164p.
2. Tapia, M.E. El Tarwi, Lupino Andino. Tarwi, Tauri, o Chocho (*Lupinus mutabilis* Sweet); Corporación Gráfica Universal SAC: Huaylas, Perú, 2015; p. 115.
3. Peralta, E.; Mazón, N.; Murillo, A.; Rivera, M.; Monar, C. Manual Agrícola de Granos Andinos. Chocho, Quinoa, Amaranto y Ataco. Cultivos, Variedades, Costos de Producción; Manual No. 69; INIAP: Quito, Ecuador, 2008; 71p.
4. Caicedo, C.; Peralta, E.; Villacrés, E.; Rivera, M. Postcosecha y Mercadeo de Chocho (*Lupinus mutabilis* Sweet) en Ecuador; Publicación Miscelánea No. 105; INIAP: Quito, Ecuador, 2001; 50p.
5. Vegas-Niño, R.; Vegas-Pérez, C.; Peña-Suasnabar, C. Potencial tecnológico de la semilla de tarwi. In *Lupinus mutabilis* (Tarwi) Leguminosa Andina Con Gran Potencial Industrial; Zavaleta, A.I., Ed.; Fondo Editorial de la Universidad Nacional Mayor de San Marcos: Lima, Perú, 2018; p. 61.
6. Jacobsen, S.; Mujica, A. El tarwi (*Lupinus Mutabilis* Sweet.) y sus Parientes Silvestres. *Botánica Económica de los Andes Centrales*; Moraes, R., Øllgaard, B., Kvist, L.P., Borchsenius, F., Balslev, H., Eds.; Universidad Mayor de San Andrés: La Paz, Bolivia, 2006; pp. 458–482.
7. León, J. Plantas Alimenticias Andinas; Boletín Técnico # 6; Instituto Interamericano de Ciencias Agrícolas: Zona Andina, Lima, Perú, 1964.
8. Carrillo, E. Revisión del Género *Lupinus* en el Perú. Ph.D. Thesis, Universidad Nacional de San Agustín, Arequipa, Perú, 1956.
9. Antúnez de Mayolo, S. Tarwi in Ancient Perú. In *Proceedings of the First International Lupine Workshop*, Lima-Cusco, Perú, 12–21 April 1982.

10. Vicente, J. El cultivo de Tarwi (*Lupinus mutabilis* Sweet) en el Estado Plurinacional de Bolivia. *Rev. Científica Investig. INFO-INIAF* 2016, 1, 88–100.
11. Mazón, N. El Chocho o tarwi Como Recurso Genético de la Región Andina ; Interaprendizaje—IPDRS: Quito, Ecuador, 29 October 2018; Available online: <https://bit.ly/2rsbfFF> (accessed on 20 January 2022).
12. Abraham, E.M.; Ganopoulos, I.; Madesis, P.; Mavromatis, A.; Mylona, P.; Nianiou-Obeidat, I.; Parissi, Z.; Polidoros, A.; Tani, E.; Vlachostergios, D. The Use of Lupin as a Source of Protein in Animal Feeding: Genomic Tools and Breeding Approaches. *Int. J. Mol. Sci.* 2019, 20, 851.
13. French, R.J. Chapter Lupin: Agronomy. *Ref. Modul. Food Sci.* 2016, 1–9.
14. FAOSTAT Data, Lupin Yield, Australia and Western Europe, Crops and livestock products. Food and Agriculture Organization of the United Nations, 2019. Available online: <https://www.fao.org/faostat/en/#data/QCL> (accessed on 15 February 2022).
15. MINAGRI—Ministerio de Agricultura y Riego. Anuario Estadístico de la Producción Agrícola y Pecuaria 2017; III Censo Nacional Agropecuario; MINAGRI: Quito, Ecuador, 2018.
16. Tello, F. *Lupinus mutabilis* sweet—a potent food source from the Andean region. *Am. J. Clin. Nutr.* 1976, 29, 933.
17. Wallace, C.; Buirchell, B.J.; Tapia, M.E.L.; Lupinus, L. Promoting the Conservation and Use of Underutilized and Neglected Crops. 23; Institute of Plant Genetics and Crop Plant Research, Gatersleben/International Plant Genetic Resources Institute: Rome, Italy, 1998; 104p.
18. Von Baer, E. Protein and oil content and yield of *L. mutabilis* in South America and Western Europe. In Proceedings of the 6th International Lupin Conference, Chilean Lupin Association, Temuco, Chile, 25–30 November 1991; pp. 13–15.
19. Gross, R. El Cultivo y la Utilización del tarwi (*Lupinus mutabilis* Sweet); No. 581.632 F3; FAO: Rome, Italy, 1981.
20. Lucas, M.M.; Stoddard, F.L.; Annicchiarico, P.; Frías, J.; Martínez-Villaluenga, C.; Sussmann, D.; Duranti, M.; Seger, A.; Zander, P.M.; Pueyo, J.J. The future of lupin as a protein crop in Europe. *Front. Plant Sci.* 2015, 6, 705.
21. Bebeli, P.J.; Lazaridi, E.; Chatzigeorgiou, T.; Suso, M.J.; Hein, W.; Alexopoulos, A.A.; Canha, G.; van Haren, R.J.F.; Jóhannsson, M.H.; Mateos, C.; et al. State and Progress of Andean Lupin Cultivation in Europe: A Review. *Agronomy* 2020, 10, 1038.
22. Petterson, D.S.; Sipsas, S.; Mackintosh, J.B. The Chemical Composition and Nutritive Value of Australian Grain Legumes, 2nd ed.; Grain Research and Development Corporation: Canberra, Australia, 1997.

23. Camarena, F. El Cultivo del Tarwi; Programa de leguminosas de Grano; Universidad Nacional Agraria La Molina: Lima, Perú, 2000.
24. Villacrés, E.; Cuadrado, L.; Falconí, F. Los Granos Andinos: Chocho (*Lupinus mutabilis* Sweet), Quinoa (*Chenopodium quinoa* Willd), Amaranto (*Amaranthus caudatus* L.) y Sangorache (*Amaranthus hybridus* L.), fuente de Metabolitos Secundarios y Fibra Dietética; Boletín Técnico No. 165; INIAP- Estación Experimental Santa Catalina: Quito, Ecuador, 2013; 42p.
25. Romeo, F.V.; Fabroni, S.; Ballistreri, G.; Muccilli, S.; Spina, A.; Rapisarda, P. Characterization and Antimicrobial Activity of Alkaloid Extracts from Seeds of Different Genotypes of *Lupinus* spp. *Sustainability* 2018, 10, 788.
26. Villacrés, E.; Cueva, P.; Díaz, M.; Rosell, C.M. Replacing Wheat Flour with Debittered and Fermented Lupin: Effects on Bread's Physical and Nutritional Features. *Plant Foods Hum. Nutr.* 2020, 75, 569–575.
27. Fornasini-Salvador, M.V.; Abril-Ulloa, S.V.; Beltrán-Carreño, J.P.; Elena Villacrés, E.; Cuadrado-Merino, L.; Robalino, F.; Sánchez, R.; Ricaurte-Ortiz, P.S.; Muñoz, E.B.; Benítez-Loza, N.B.; et al. Efficacy of a *Lupinus mutabilis* Sweet snack as complement to conventional type 2 diabetes mellitus treatment. *Nutr. Hosp.* 2019, 36, 905–911.
28. Canahua, A.; Roman, P. Tarwi. Leguminosa andina de gran potencial. *Rev. De Agroecol. LEISA* 2016, 32, 2. Available online: <https://www.servindi.org/actualidad-opinion/27/07/2016/tarwi-leguminosa-andina-de-gran-potencial> (accessed on 13 January 2022).
29. Falconi, C.E. *Lupinus mutabilis* in Ecuador with Special Emphasis on Anthracnose Resistance. Ph.D. Thesis, Wageningen University, Wageningen, The Netherlands, 2012; 150p. Available online: <https://edepot.wur.nl/210228> (accessed on 18 February 2022).
30. Thomas, G.; Jones, R.; Vanstone, V. Diseases of lupin. In *Producing Lupins*, 2nd ed.; White, P., French, B., McLarty, A., Eds.; Department of Agriculture and Food: Bulletin, WA, USA, 2008; pp. 101–120. ISSN 1833-7236.
31. Thomas, G.J.; Sweetingham, M.W. Fungicide seed treatments reduce seed transmission and severity of lupin anthracnose caused by *Colletotrichum gloeosporioides*. *Australas. Plant Pathol.* 2003, 32, 39–46.
32. Thomas, G.J.; Adcock, K.G. Exposure to dry heat reduces anthracnose infection of lupin seed. *Australas. Plant Pathol.* 2004, 33, 537–540.
33. Falconí, C.E.; Yáñez-Mendizábal, V. Dry heat treatment of Andean lupin seed to reduce anthracnose infection. *Crop Prot.* 2016, 89, 178–183.
34. Falconí, C.E.; Yáñez-Mendizábal, V. Efficacy of UV-C radiation to reduce seedborne anthracnose (*Colletotrichum acutatum*) from Andean lupin (*Lupinus mutabilis*). *Plant Pathol.* 2017, 67, 831–838.

35. Falconí, C.E.; Yáñez-Mendizábal, V. Solar UV-B radiation limits anthracnose infection in seed and promote biochemical and physiological response in lupin plants. *Plant Pathol.* 2019, 68, 1635–1644.
36. Yáñez-Mendizábal, V.; Falconí, C.E.; Grijalva, A.C. *Bacillus* spp. evaluation to control anthracnose infection on Andean lupin seed (*Lupinus mutabilis* Sweet). *Phytopathology* 2015, 105, S4–S153.
37. Yáñez-Mendizábal, V.; Falconí, C.E. Efficacy of spp. to biocontrol of anthracnose and enhance plant growth on Andean lupin seeds by lipopeptide production. *Biol. Control* 2018, 122, 67–75.
38. Yáñez-Mendizábal, V.; Falconí, C.E. *Bacillus subtilis* CtpxS2-1 induces systemic resistance against anthracnose in Andean lupin by lipopeptide production. *Biotechnol. Lett.* 2021, 43, 719–728.
39. Cowling, W.A. *Pedigrees and Characters of Narrow-Leafed Lupin Cultivars Released in Australia from 1967 to 1998*; Lupin Breeding Bulletin 4365; Agriculture Western Australia: Perth, Australia, 1999.
40. Sweetingham, M.; Clements, J.; Buirchell, B.; Sipsas, S.; Thomas, G.; Quealy, J.; Jones, R.; Francis, C.; Smith, C.G. Preliminary breeding and development of Andean lupin for Australian agriculture. In *Where Old and New World Lupins Meet, Proceedings of the 11th International Lupin Conference, Guadalajara, Jalisco, Mexico, 4–9 May 2005*; Van Santen, E., Hill, G.D., Eds.; International Lupin Association: Canterbury, New Zealand, 2005.
41. Gulisano, A.; Alves, S.; Neves Martins, J.; Trindade, L.M. Genetics and Breeding of *Lupinus mutabilis*: An Emerging Protein Crop. *Front. Plant Sci.* 2019, 10, 1–13.
42. Basra, S.M.A.; Ashraf, M.; Iqbal, N.; Khaliq, A.; Ahmed, R. Physiological and biochemical aspects of pre-sowing heat stress on cotton seed. *Seed Sci. Technol.* 2004, 32, 765–774.
43. Chevrefils, G.; Caron, E.; Wright, H.; Sakamoto, G.; Payment, P.; Barbeau, B.; Cairns, B. UV dose required to achieve incremental log inactivation of bacteria, protozoa and viruses. *IUVA News* 2006, 8, 38–45.
44. Bokhari, N.A.; Siddiqui, I.; Parveen, K.; Siddique, I.; Rizwana, H.; Soliman, D.A. Management of anthracnose of banana by UV radiation. *J. Anim. Plant Sci.* 2013, 23, 1211–1214.
45. González-Aguilar, G.A.; Wang, C.Y.; Buta, J.G.; Krizek, D.T. Use of UV- C radiation to prevent decay and maintain postharvest quality of ripe ‘Tommy Atkins’ mangoes. *Int. J. Food Sci. Technol.* 2001, 36, 767–773.
46. Wilson, C.L.; Heidinger, G.F. Ultraviolet Light Treatments for Increasing Seed Yields. U.S. Patent WO2010085513 A1, 29 July 2010.

47. Darras, A.I.; Demopoulos, V.; Tiniakou, C. UV-C radiation induces defense responses and improves vase-life of cut gerbera flowers. *Postharvest Biol. Technol.* 2012, 64, 168–174.
48. Shetta, N.D.; Areaf, I.M. Impact of ultraviolet-C radiation on seed germination and chlorophyll concentration of some woody trees grown in Saudi Arabia. *J. Agric. Env. Sci* 2009, 8, 1–20.
49. Middleton Middleton, E.M.; Teramura, A.H. The role of flavonol glycosides and carotenoids in protecting soybean from ultraviolet-B damage. *Plant Physiol.* 1993, 103, 741–752.
50. Peykarestan, B.; Seify, M. UV irradiation effects on seed germination and growth, protein content, peroxidase and protease activity in redbean. *Int. Res. J. Appl. Basic. Sci.* 2012, 3, 92–102.
51. Hieng, B.; Ugrinovic, K.; Sustar-vozlic, J.; Kidric, M. Different classes of proteases are involved in the response to drought of *Phaseolus vulgaris* L. cultivars differing in sensitivity. *J. Plant Physiol.* 2004, 161, 519–530.
52. Salama, H.M.H.; Watban, S.A.; Al-Fughom, A.T. Effect of ultraviolet radiation on chlorophyll, carotenoid, protein and proline contents of some annual desert plants. *Saudi J. Biol. Sci.* 2011, 18, 79–86.
53. Sharma, P.; Jha, A.B.; Dubey, R.S.; Pessarakli, M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *J. Bot.* 2012, 12, 1–26.
54. Rivera-Pastrana, D.M.; Gardea, A.A.; Yahia, E.M.; Martínez-Téllez, M.A.; González-Aguilar, G.A. Effect of UV-C irradiation and low temperature storage on bioactive compounds, antioxidant enzymes and radical scavenging activity of papa fruit. *J. Food Sci. Technol.* 2014, 51, 3821–3829.
55. Glazener, J.A. Accumulation of phenolic compounds in cells and formation of lignin-like polymers in cell walls of young tomato fruits after inoculation with *Botrytis cinerea*. *Physiol. Plant Pathol.* 1982, 20, 11–25.
56. Anand, T.A.; Chandrasekaran, A.; Kuttalam, S.; Raguchander, T.; Prakasam, V.; Samiyappan, R. Association of some plant defense enzyme activities with systemic resistance to early leaf blight and leaf spot induced in tomato plants by azoxystrobin and *Pseudomonas fluorescens*. *J. Plant Interact.* 2007, 2, 233–244.
57. Kanetis, L.; Holmes, G.J.; Ojiambo, P.S. Survival of *Pseudoperonospora cubensis* sporangia exposed to solar radiation. *Plant Pathol.* 2010, 59, 313–323.
58. Rotem, J.; Wooding, B.; Aylor, D.E. The role of solar radiation, especially ultraviolet, in the mortality of fungal spores. *Phytopathology* 1985, 75, 510–514.
59. Mizubuti, E.S.G.; Aylor, D.E.; Fry, W.E. Survival of *Phytophthora infestans* sporangia exposed to solar radiation. *Phytopathology* 2000, 90, 78–84.
60. Diffey, B.L. Solar ultraviolet radiation effects on biological systems. *Phys. Med. Biol.* 1991, 36, 299–328.

61. De Menezes, H.D.; Massola, N.S., Jr.; Flint, S.D.; Silva, G.J., Jr.; Bachmann, L.; Rangel, D.E.N.; Braga, G.U.L. Growth under visible light increases conidia and mucilage production and tolerance to UV-B radiation in the plant pathogenic fungus *Colletotrichum acutatum*. *Photochem. Photobiol.* 2015, 91, 397–402.
62. Begum, M.; Sariah, M.; Puteh, A.; Zainal, A. Detection of seed-borne fungi and site of infection by *Colletotrichum truncatum* in naturally-infected soybean seeds. *Int. J. Agric. Res.* 2007, 2, 812–819. Available online: <https://scialert.net/abstract/?doi=ijar.2007.812.819> (accessed on 15 January 2022).
63. Tepfer, D.; Leach, S. Survival and DNA damage in plant seeds exposed for 558 and 682 days outside the International Space Station. *Astrobiology* 2017, 17, 205–215.
64. Noble, R.E. Effects of UV-irradiance on seed germination. *Sci. Total Environ.* 2002, 299, 173–176.
65. Shaukat, S.; Farooq, M.; Siddiqui, M.; Zaidi, S. Effect of enhanced UV-B radiation on germination, seedling growth and biochemical responses of *Vigna mungo* (L.) Hepper. *Pak. J. Bot* 2013, 45, 779–785.
66. Araújo, S.S.; Paparella, S.; Dondi, D.; Bentivoglio, A.; Carbonera, D.; Balestrazzi, A. Physical methods for seed invigoration: Advantages and challenges in seed technology. *Front. Plant Sci.* 2016, 7, 646.
67. Sugimoto, K. Seed germination under UV-B irradiation. *Bull. Minamikyushu Univ.* 2013, 43, 1–9.
68. Hideg, É.; Hideg, E.; Rosenqvist, E.; Váradi, G.; Bornman, J.F.; Vincze, E. A comparison of UV-B induced stress responses in three barley cultivars. *Funct. Plant Biol.* 2006, 33, 77–90.
69. Foyer, C.H.; Noctor, G. Oxidant and antioxidant signaling in plants: A re-evaluation of the concept of oxidative stress in a physiological context. *Plant Cell Environ.* 2005, 28, 1056–1071.
70. Caldwell, M.M.; Caldwell, M.M.; Bornman, J.F.; Ballaré, C.L.; Flint, S.D.; Kulandaivelu, G. Terrestrial ecosystems, increased solar ultraviolet radiation, and interactions with other climate change factors. *Photochem. Photobiol. Sci.* 2007, 6, 252–266.
71. Ravindran, K.C.; Indrajith, A.; Pratheesh, P.V.; Sanjiviraja, K.; Balakrishnan, V. Effect of ultraviolet-B radiation on biochemical and antioxidant defense system in *Indigofera tinctoria* L. seedlings. *Int. J. Eng. Sci. Technol.* 2012, 2, 226–232.
72. Yáñez-Mendizábal, V.; Zeriouh, H.; Viñas, I.; Torres, R.; Usall, J.; De Vicente, A.; Pérez-García, A.; Teixido, N. Biological control of peach brown rot (*Monilinia* spp.) by *Bacillus subtilis* CPA-8 is based on production of fengycin-like lipopeptides. *Eur. J. Plant Pathol.* 2012, 132, 609–619.
73. Ongena, M.; Jacques, P. *Bacillus* lipopeptides: Versatile weapons for plant disease biocontrol. *Trends Microbiol.* 2008, 16, 115–125.

74. Romero, D.; de Vicente, A.; Rakotoaly, R.H.; Dufour, S.E.; Veening, J.W.; Arrebola, E.; Cazorla, F.M.; Kuipers, O.P.; Paquot, M.; Pérez-García, A. The iturin and fengycin families of lipopeptides are key factors in antagonism of *Bacillus subtilis* towards *Podosphaera fusca*. *Mol. Plant Microbe Interact.* 2007, 20, 430–440.
75. Ongena, M.; Henry, G.; Thonart, P. The roles of cyclic lipopeptides in the biocontrol activity of *Bacillus subtilis*. In *Recent Developments in Management of Plant Diseases, Plant Pathology in the 21st Century, Proceedings of the 9th International Congress of Plant Pathology, Torino, Italy, 24–29 August 2008*; Gisi, U., Chet, L., Gullino, M.L., Eds.; Springer: Dordrecht, The Netherlands, 2009.
76. Miles, T.; Nair, M.; Vandervoort, C.; Schilder, A.M.C. Characterization and biological activity of flavonoids from ripe fruit of an anthracnose resistant blueberry cultivar. *Physiol. Mol. Plant Pathol.* 2013, 83, 8–16.
77. Meazza, G.; Dayan, F.; Wedge, D.E. Activity of Quinones on *Colletotrichum* Species. *J. Agric. Food Chem.* 2003, 51, 3824–3828.
78. Deleu, M.; Paquot, M.; Nylander, T. Fengycin interaction with lipid monolayers at the air–aqueous interface—implications for the effect of fengycin on biological membranes. *J. Colloid Interface Sci.* 2005, 283, 358–365.
79. Vanittanakom, N.; Loeffler, W.; Koch, U.; Jung, G. Fengycin—A novel antifungal lipopeptide antibiotic produced by *Bacillus subtilis* F-29-3. *J. Antibiot.* 1986, 39, 888–901.
80. Yasmeen, S.; Sariah, M. Effect of Seed Bacterization on Plant Growth Response and Induction of Disease Resistance in Chilli. *Agric. Sci. China* 2009, 8, 963–971.
81. Manjula, K.; Podile, A.R. Production of fungal cell wall degrading enzymes by a biocontrol strain of *Bacillus subtilis* AF1. *Indian J. Exp. Biol.* 2005, 43, 892–896.
82. Singh, S.K.; Pathak, R.; Choudhary, V. Plant Growth-Promoting Rhizobacteria-Mediated Acquired Systemic Resistance in Plants against Pest and Diseases. In *Microbial-Mediated Induced Systemic Resistance in Plants*; Choudhary, D.K., Varma, A., Eds.; Springer: Singapore, 2016; pp. 125–134.
83. Park, J.W.; Balaraju, K.; Kim, J.W.; Lee, S.W.; Park, K. Systemic resistance and growth promotion of chili pepper induced by an antibiotic producing *Bacillus vallismortis* strain BS07. *Biol. Control* 2013, 65, 246–257.
84. Miljković, D.; Marinković, J.; Balešević-Tubić, S. The significance of *Bacillus* spp. in disease suppression and growth promotion of field and vegetable crops. *Microorganism* 2020, 8, 1037.
85. Chowdappa, P.; Kumar, S.M.; Lakshmi, M.J.; Mohan, S.P.; Upreti, K.K. Growth stimulation and induction of systemic resistance in tomato against early and late blight by *Bacillus subtilis* OTPB1 or *Trichoderma harzianum* OTPB3. *Biol. Control* 2013, 65, 109–117.

86. Jayapala, N.; Mallikarjunaiah, N.; Puttaswamy, H.; Gavirangappa, H.; Ramachandrappa, N.S. Rhizobacteria *Bacillus* spp. induce resistance against anthracnose disease in chili (*Capsicum annuum* L.) through activating host defense response. *Egypt. J. Biol. Pest Control* 2019, 29, 45.
87. Chen, F.; Wang, M.; Zheng, Y.; Luo, J.; Yang, X.; Wang, X. Quantitative changes of plant defense enzymes and phytohormone in biocontrol of cucumber Fusarium wilt by *Bacillus subtilis* B579. *World J. Microbiol. Biotechnol.* 2010, 26, 675–684.
88. Falconí, C.E.; Yáñez-Mendizábal, V.; Claudio, D. Native *Bacillus subtilis* strains efficiently control lupin anthracnose both under greenhouse and in field conditions. *Ecol. Environ. Conserv.* 2022; accepted.
89. Lindbeck, K.; Fanning, J.; Jason Brand, J. Diseases of Lupins. Updated on 22 June 2020. Available online: <https://extensionaus.com.au/FieldCropDiseasesVic/docs/identification-management-of-field-crop-diseases-in-victoria/lupins/summary/> (accessed on 7 February 2022).
90. Wang, X.Q.; Zhao, D.L.; Shen, L.L.; Jing, C.L.; Hang, C.S. Application and Mechanisms of *Bacillus subtilis* in Biological Control of Plant Disease. In *Role of Rhizospheric Microbes in Soil*; Springer: Singapore, 2018; pp. 225–250.
91. Hojin, R.; Hoon, P.; Dong-Sang, S.; Gun, H.J.; Kyungseok, P.; Byung, D.L. Biological control of *Colletotrichum panacicola* on *Panax ginseng* by *Bacillus subtilis* HK-CSM-1. *Ginseng Res.* 2014, 38, 215–219.
92. Falconi, C.E.; Visser, R.G.F.; van Heusden, A.W. Influence of Plant Stage on Resistance to Anthracnose in Andean Lupin (*Lupinus mutabilis* Sweet). *Crop Pasture Sci.* 2015, 66, 729–734.
93. Wei, F.; Hu, X.; Xu, X. Dispersal of *Bacillus subtilis* and its effect on strawberry phyllosphere microbiota under open field and protection conditions. *Sci. Rep.* 2016, 6, 22611.
94. Demoz, B.T.; Korsten, L. *Bacillus subtilis* attachment, colonization, and survival on avocado flowers and its mode of action on stemend rot pathogens. *BioControl* 2006, 37, 68–74.
95. Yadav, R.K.; Kakamanoli, K.; Vokou, D. Estimating bacterial population on the phyllosphere by serial dilution plating and leaf imprint method. *Biol. Ecoprint: Int. J. Ecol.* 2010, 17, 47–52.
96. Falconi, C.E. Control Biológico de Enfermedades de las Plantas en Ecuador. Capítulo 9. In *Control Biológico de Enfermedades en los Cultivos en Latinoamérica y el Caribe*; Belliot, W., Mondino, P., Comenarez, Y., Vero, S., Montealegre, J., Rivera, M., Eds.; Editorial Universidad de la República: Montevideo, Uruguay, 2014; 402p, pp. 214–242.
97. Falconí, C.E.; Peralvo, D.; Saavedra, L.; Yáñez-Mendizábal, V. Validación de biopreparados en base a bacterias epifitas para el control de la moniliasis en cacao fino de aroma en el cantón Valencia de la provincia de Los Ríos. In *Validación de Biopesticidas para el Control de la Moniliasis y Manejo Sustentable del Cacao Fino de Aroma en el Ecuador*; Falconí, C., Yáñez, V., Eds.; Artes Gráficas Señal: Quito, Ecuador, 2007; pp. 37–49.

98. Cruz-Martín, M.; Acosta-Suárez, M.; Mena, E.; Roque, B.; Pichardo, T.; Alvarado-Capó, Y. Antifungal activity of *Musa* phyllosphere *Bacillus pumilus* strain against *Mycosphaerella fijiensis*. *Trop. Plant Pathol.* 2017, 42, 121–125.
99. Thomas, G.J.; Sweetingham, M.W.; Adcock, K.G. Application of fungicides to reduce yield loss in anthracnose-infected lupins. *Crop Prot.* 2008, 27, 1071–1077.
100. Cambo, E.I. Evaluación de Cuatro Fungicidas Sobre la Antracnosis (*Colletotrichum acutatum*) en el Cultivo de Chocho (*Lupinus mutabilis*) variedad INIAP—450. Bachelor's Thesis, Universidad de las Fuerzas Armadas ESPE, Sangolquí, Ecuador, 2016; p. 54.
101. Jílková, B.; Víchová, J.; Pokorný, R.; Vejražka, K. Sensitivity of *Colletotrichum acutatum* Isolates to Selected Fungicides. *Acta Univ. Agric. Silvic. Mendel. Brun.* 2015, 63, 1111–1119.
102. Vega, L.E.; Rodríguez, D.G.; Murillo, A.R.; Mazón, N.G. Cuantificación del Daño y Validación de Una Estrategia de Control Químico de la Antracnosis (*Colletotrichum acutatum*) en el Cultivo del Chocho (*Lupinus mutabilis* Sweet) en Ecuador; 1er Congreso Internacional Ciencia y Tecnología Agropecuaria; INIAP—Estación Experimental Santa Catalina: Quito, Ecuador, 2018; pp. 80–82.
103. Shea, G.; Thomas, G.; Buirchell, B.; Salam, M.; McKirdy, S.; Sweetingham, M. Case study: Industry response to the lupin anthracnose incursion in Western Australia. In *Lupins for Health and Wealth, Proceedings of the 12th International Lupin Conference, Fremantle, Australia, 14–18 September 2008*; Palta, J.A., Berger, J.B., Eds.; International Lupin Association: Canterbury, New Zealand, 2008; pp. 425–431.
104. Silva-Junior, G.J.; Spósito, M.S.; Marin, D.R.; Amorim, L. Efficacy and timing of application of fungicides for control of citrus postbloom fruit drop. *Crop Prot.* 2014, 59, 51–56.
105. FRAC Code List ©*2021: Fungal Control Agents Sorted by Cross Resistance Pattern and Mode of Action (Including Coding for FRAC Groups on Product Labels). Available online: https://www.frac.info/docs/default-source/publications/frac-code-list/frac-code-list-2021--final.pdf?sfvrsn=f7ec499a_2 (accessed on 22 January 2022).
106. Hamaoka, K.; Aoki, Y.; Suzuki, S. Isolation and Characterization of Endophyte *Bacillus velezensis* KOF112 from Grapevine Shoot Xylem as Biological Control Agent for Fungal Diseases. *Plants* 2021, 10, 1815.
107. Saxena, A.; Raghuwanshi, R.; Gupta, V.K.; Singh, H.B. Chilli Anthracnose: The Epidemiology and Management. *Front. Microbiol.* 2016, 7, 1–18.
108. Förster, H.; Driever, G.F.; Thompson, D.C.; Adaskaveg, J.E. Postharvest decay management for stone fruit crops in California using the “reduced-risk” fungicides fludioxonil and fenhexamid. *Plant Dis.* 2007, 91, 209–215.
109. Falconi, C.E.; Visser, R.G.F.; van Heusden, A.W. Phenotypic, molecular, and pathological characterization of *Colletotrichum acutatum* associated with lupine and tamarillo in the

- Ecuadorian Andes. *Plant Dis.* 2013, 97, 819–827.
110. Wirtz, L.; Massola-Júnior, N.S.; Linhares de Castro, R.R.; Ruge-Wehling, B.; Schaffrath, U.; Loehrer, M. *Colletotrichum* spp. from Soybean Cause Disease on Lupin and Can Induce Plant Growth-Promoting Effects. *Microorganisms* 2021, 9, 1130.
111. Ruales, C. *Las Colecciones Botánicas de Joseph de Jussieu (1736–1747); Serie Monográfica “Plantas de Quito—La vegetación original de una ciudad siempre verde”*; Universidad San Francisco de Quito: Quito, Ecuador, 2013; Volume 1, p. 293.
112. Falconi, C.E.; van Heusden, A.W. Morphological, pathological and molecular characterization of lupin anthracnose and its relationship with tamarillo anthracnose in Ecuadorian Andes. *American Phytopathological Society Annual Meeting, Honolulu—Hawaii. Phytopathology* 2011, 101, S9–S23. Available online: <https://apsjournals.apsnet.org/doi/pdf/10.1094/PHYTO.2011.101.6.S1> (accessed on 12 January 2022).
113. Ficke, A.; Gadoury, D.M.; Seem, R.C. Ontogenic resistance and plant disease management: A case study of grape powdery mildew. *Phytopathology* 2002, 92, 671–675.
114. Beate Tschöpe, P.R. SIMONTO-Lupin: An ontogenetic simulation model for lupin species (*Lupinus angustifolius*, *L. luteus* and *L. albus*). *J. Für Kult.* 2011, 63, 1626–1632.
115. Sinde-González, I.; Falconí, C.E.; Luna-Granizo, P.; Godoy-Guanín, L.; Gil-Docampo, M.; Manguashca, J.; Nato, R. Spectral analysis of the phenological stages of *Lupinus mutabilis* through spectroradiometry and unmanned aerial vehicle imaging with different physical disinfection pretreatments of seeds. *Geocarto Int.* 2021, 1–18.
116. Thomas, G.J.; Sweetingham, M.W. Cultivar and environment influence the development of lupin anthracnose caused by *Colletotrichum lupini*. *Australas. Plant Pathol.* 2004, 33, 571–577.
117. Yáñez-Mendizábal, V.; Usall, J.; Viñas, I.; Casals, C.; Marín, S.; Solsona, C.; Teixidó, N. Potential of a new strain of *Bacillus subtilis* CPA-8 to control the major postharvest diseases of fruit. *Biocontrol. Sci. Technol.* 2011, 21, 409–426.
118. Yáñez-Mendizábal, V.; Viñas, Y.; Usall, J.; Torres, R.; Solsana, C.; Teixidó, N. Production of the postharvest biocontrol agent *Bacillus subtilis* CPA-8 using low cost commercial products and by-products. *Biol. Control* 2012, 60, 280–289.
119. Yáñez-Mendizábal, V.; Viñas, I.; Usall, J.; Teixidó, N. Endospore production allows using spray-drying. As a possible formulation system of the biocontrol agent *Bacillus subtilis* CPA-8. *Biotechnol. Lett.* 2012, 34, 729–735.
120. Yáñez-Mendizábal, V.; Viñas, I.; Usall, J.; Torres, R.; Solsona, C.; Abadias, M.; Teixidó, N. Formulation development of the biocontrol agent *Bacillus subtilis* strain CPA-8 by spray-drying. *J. Appl. Microbiol.* 2012, 112, 954–965.

Retrieved from <https://encyclopedia.pub/entry/history/show/51773>