DPPH· Stable Free Radical

Subjects: Chemistry, Organic Contributor: Petre Ionita

DPPH· is 2,2-diphenyl-1-(2,4,6-trinitrophenyl)hydrazyl stable free radical. It was discovered in 1922.

Keywords: hydrazyl ; free radical ; EPR ; ESR ; spin-trapping ; diradical

1. Introduction

The chemistry and physics of stable or persistent free radicals are well represented by several books ^{[1][2][3]}, two of which place a greater emphasis on their organic reactions, the first one being published in 1968 ^[4] and the second one in 2010 ^[5]. However, as mentioned in the literature ^[6], in the most recent there is a notable absence of hydrazyl (and also aminyl) free radicals. Many of free radicals derived from the most encountered stable hydrazyl (DPPH, Figure 1).



Figure 1. Chemical structure of DPPH, a stable hydrazyl free radical.

1.1. Free Radicals

A free radical is a chemical entity that contains an unpaired electron (free electron) that possess a quantum-mechanical property called spin. Such an entity typically has a high reactivity due to its open-shell structure. However, nowadays there are a lot of such known compounds that are stable under usual laboratory conditions (room temperature and presence of air) ^{[3][4][5]}. Their high stability is a sum of several structural characteristics, the most important being steric hindrance and conjugation. The push–pull effect that contributes to the greater stability of several classes of such free radicals was postulated in the 1960s ^[2].

1.2. Hydrazyls

A hydrazyl free radical contains the chemical moiety denoted in Figure 1 Left, where the dots represents the unpaired electron. The most known hydrazyl free radical is 2,2-diphenyl-1-(2,4,6-trinitrophenyl) hydrazyl, usually encountered as DPPH⁻ (the 2,4,6-trinitrophenyl fragment is also frequently named picryl).

The history of hydrazyl free radicals starts about 100 years ago, when Goldschmidt firstly observed that oxidation of triphenylhydrazine led to an intense blue color that fades rapidly ^[8]. Between many congeners of triphenylhydrazine, he found that the oxidation of 2,2-diphenyl-1-picrylhydrazine, a yellow compound, also gives an intense violet solution, but in this particular case the color is stable. In this way, Goldschmidt isolated in high yields the first stable hydrazyl free radical ^[9]. It was found that this free radical does not dimerize or react with oxygen and is stable either in solution or in solid state. However, it is still a reactive compound, active in many types of reactions, as will later be shown.

1.3. DPPH[·] Free Radical

The stability of such hydrazyl free radicals may be easily evidenced in the case of DPPH⁻. As mentioned before, steric hindrance and conjugation play a dominant role. The push–pull effect (as picryl moiety plays the role of the electron-acceptor part of the molecule and the diphenylamino moiety plays the role of the electron-donor part of the molecule) is

clearly evidenced by the resonance structures (Figure 2) and is in accordance with Linnet theory ^[10]. Another important aspect that is worth remembering is the dipole moment of DPPH⁻ (4.88 D), higher than DPPH-H (3.59 D), emphasizing the importance of the polar resonance structures [^[11]. The N-N bond also has a higher bond order, as was directly demonstrated by the X-ray structure ^[12]. Due to its extreme stability, DPPH⁻ is nowadays used as a standard in Electron Spin Resonance (ESR) spectroscopy ^{[3][13]}.



Figure 2. Some resonance structures of DPPH' free radical.

The extreme stability of the DPPH free radical is easily demonstrated by the comparison with its congeners, 2,2-diphenyl-1-(2,4-dinitrophenyl) hydrazyl **1** and 2,2-diphenyl-1-(2,6-dinitrophenyl)-hydrazyl **2** free radicals (Figure 3). The most important nitro groups are situated in the ortho-position with regard to the nitrogen atom; thus, between the two 2,4-dinitro and 2,6-dinitro isomers **1** and **2** (Figure 3), there is a huge difference in terms of their stability ^{[14][15]}. Accordingly, isomer **1** (2,4-dinitro) can exist only in solution (with a half-life of about 90 h) and not in solid state, while isomer **2** (2,6-dinitro) is stable both in solution and in solid state ^[16]. As a consequence, most of the free radicals of such type that contain any other group in position 6 (para-) are stable ^[4].



Figure 3. Stability of two hydrazyl isomers.

The hydrogen atom bound to the hydrazine group cannot only to be removed by oxidation, leading to the hydrazyl free radical, but also by a base, because it has an acidic character. Most of the hydrazyl free radicals have a violet color, while the parent hydrazines are yellow; the corresponding anion is usually red-brown, but it is possible also to be green, depending on the substituents. All these acid-base or redox processes are reversible, as shown in Figure 4 ^[17].





2. Application of Hydrazyls

2.1. Acid-Base and Redox Processes

It was shown before that mainly the p-substituents on the phenyl ring has the most influence on the oxidation capacity of the DPPH⁻ congeners, as well as on their acidity (for parent hydrazines). Concerted electron–proton transfer may occur $\frac{[18]}{18}$. For example, each supplementary p-nitro group increases the oxidation potential of the free radicals by 0.1–0.2 V,

while, for parent hydrazines, each group lowers the pK_a value by about 1 unit. This means that hydrazyl free radicals become stronger oxidants, whereas hydrazines become stronger acids. Table 1 shows these values.

Compound	λradical	λanion	λhydrazine	рКа	Eox	BDE
DPPH	518	424	322	8.54	0.30	75.3
1	512	383	337	11.3	0.13	74.5
9	500	505	352	7.35	0.49	
10	500	485	352	6.49	0.60	-
12	524	413	353	8.20	0.34	82
13	506	441	363	10.7	0.25	76.4
23	500	365	346	11.1	0.24	76.3
25	524	655	324	12.5	0.87	76.81
26	517	495	330	11.6	1.17	76.96
27	517	492	332	10.9	1.23	70.67
28	486	620	392	7.82	0.173	75.45
29	507	645	404	8.07	0.107	75.08
30	510	654	482	8.33	0.130	75.30

Table 1. λ_{max} (nm), pK_a, E_{ox} (V) and bond dissociation energy (BDE) values (kcal/mol) for selected compounds $\frac{[17][19][20]}{[21][22]}$

All these redox or acid-base processes can be followed by color change ^[23]. Table 1 also compiles the wavelength values where hydrazyl free radicals, their parent hydrazines or the corresponding anions have the maximum absorption. One important aspect that can be easily evaluated, knowing the pK_a and E_{ox} values, is the bond dissociation energy- BDE for the N-H bond. This can be evaluated following the Equation (1) ^[24].

As a remark on Table 1, it can be concluded that all the BDE values are closer to 75 kcal/mol and this is due to the fact that values of pK_a and E_{ox} compensate each other (DPPH has the literature BDE value ~75–80 kcal/mol ^[25]; a revised and accurate value was reported as 79 kcal/mol ^[26]). These BDE values can be regarded as an important tool in elucidation of the H-abstraction mechanism of DPPH congeners. As mentioned by Bordweel ^[24], most substituents on DPPH congeners will play a dual role in affecting the stability of such radicals, inducing both stabilization by the delocalizing property and at the same time destabilization by the electron-withdrawing ability. Recently, an electrografting method employing diazonium chemistry was used for the isolation of the aryl radical/DPPH coupling product ^[27].

2.2. Generators of Short-Lived Radicals

The oxidant capacity of hydrazyl free radicals can be used in the generation of short-lived radicals. It was stated before that DPPH can abstract one electron from an anion X⁻ yielding the radical X⁻. In a similar way, DPPH can abstract one hydrogen atom from other compounds, yielding again unstable free radicals (for example, from phenols). Such processes were used in the generation of several radicals of different types, O-, N-, S, C-, or P-centered ^[28]. These short-lived radicals are best evidenced by the ESR spin-trapping technique that uses a diamagnetic unsaturated compound that reacts with the short-lived free radical forming a persistent nitroxide, with a half-life of minutes-hours. These compounds are called spin-traps, and most of them are nitrones or nitroso derivatives ^[29].

Because the spin-trapping process requires both the presence of a short-lived free radical and a spin-trap, and also because it is a necessary a system that generates those short-lived free radicals, a step-forwarding idea was to design hybrid molecules that contain a hydrazyl and a nitrone moiety covalently linked. Such compounds (**28–30**) are shown in Figure 5 $\frac{[22]}{2}$.



Figure 5. Hybrid hydrazyl-nitrones.

These compounds have the great advantage of finding direct applications as sensors or probes in ESR spectroscopy, as simultaneous generators and traps for short-lived radicals. The so-called DPPH method for the total antioxidant capacity measurement is one of the most used [30][31].

3. Conclusions and Outlook

Besides all these general and particular aspects, hydrazyl free radicals are still a particular domain in chemistry that contains unexplored fields. Recently, DPPH⁻ found applications in very different areas, such as catalysis ^{[32][33]}; it may also may find similar applications in organic battery technologies ^{[34][35]}. Another possibility of employing hydrazyl radicals and their congeners is related to the structure of betaines **15** and **20** to exist as diradicals (similar to Thiele, Tschitschibabin or Muller hydrocarbons) ^{[36][37][38]}. Although most recent literature data is about the use of DPPH free radical as scavenger in antioxidant measurements ^{[30][31][39]}, there is a lot of room to develop new compounds and processes involving such stable open-shell structures, as their multifunctionality ^{[22][40]} provides real working opportunities. Along with the well-known class of nitroxide free radicals ^[41], hydrazyl radicals stabilized by the captodative effect ^[42] will fulfill and complete novel and important possibilities. Future work will bring out more interesting and unexpected results.

References

- 1. Weil, J.A.; Bolton, J.R. Electron Paramagnetic Resonance Spectroscopy: Elementary Theory and Applications; Wiley-Interscience: Hoboken, NJ, USA, 2007.
- 2. Brustolon, M.; Giamello, E. Electron Paramagnetic Resonance: A Practitioner's Toolkit; John Wiley & Sons: Hoboken, NJ, USA, 2009.
- 3. Chechik, V.; Carter, E.; Murphy, D.M. Electron Paramagnetic Resonance; Oxford Chemistry Primers; Oxford University Press: Oxford, UK, 2016.
- 4. Forrester, A.R.; Hay, J.M.; Thomson, R.H. Organic Chemistry of Stable Free Radicals; Academic Press: London, UK, 1968.
- 5. Hicks, R.G. Stable Radicals: Fundamentals and Applied Aspects of Odd-Electron Compounds; John Wiley & Sons, Ltd.: Chichester, UK, 2010.
- Balaban, A.T. Stable Hydrazyls and Push–Pull (Capto-Dative) Aminyl Free Radicals; PATAI's Chemistry of Functional Groups, John Wiley & Sons: Hoboken, NJ, USA, 2017.
- Balaban, A.T.; Frangopol, P.T.; Frangopol, M.; Negoita, N. Stability and equilibria of free radicals. III: Preparation of stable, sterically shielded, diarylnitrogen radicals with donor and acceptor aryl groups in the same molecule. Tetrahedron 1967, 23, 4661–4676.
- 8. Goldschmidt, S. Uber Zweiwertiger Stickstoff. Das Triphenyl hydrazyl. Ber. Dtsch. Chem. Ges. 1920, 53, 44-62.
- Goldschmidt, S.; Renn, K. Zweiwertiger Stickstoff: Über das α,α-Diphenyl-β-trinitrophenylhydrazyl. Ber. Dtsch. Chem. Ges. 1922, 55, 628–643.
- 10. Linnett, J.W. A modification of the Lewis-Langmuir octet rule. J. Am. Chem. Soc. 1961, 83, 2643–2653.
- 11. Turkevich, J.; Osper, P.F.; Smyth, C.P. The dipole moment of a free radical. J. Am. Chem. Soc. 1942, 64, 1179–1180.
- 12. Williams, D.E. Crystal structure of 2,2-diphenyl-1-picrylhydrazyl free radical. J. Am. Chem. Soc. 1967, 89, 4280–4287.
- Chen, O.; Zhuang, J.; Guzzetta, F.; Lynch, J.; Angerhofer, A.; Cao, Y.C. Synthesis of water-soluble 2,2'-diphenyl-1picrylhydrazyl nanoparticles: A new standard for electron paramagnetic resonance spectroscopy. J. Am. Chem. Soc. 2009, 131, 12542–12543.

- 14. Balaban, A.T.; Frangopol, P.T.; Marculescu, M.; Bally, J. Factors affecting stability and equilibria of free radicals: Steric factors in hydrazyls. Tetrahedron 1961, 13, 258–267.
- 15. Tanaseichuk, B.S.; Tomilin, O.B.; Pryanichnikova, M.K.; Boyarkina, O.V.; Burtasov, A.A. Relative stability of 1-(4-R-2,6dinitrophenyl)-2,2-diphenylhydrazides. Russ. J. Org. Chem. 2017, 53, 679–685.
- 16. Balaban, A.T.; Marculescu, M.; Pascaru, I.; Rotaru, M.; Valeriu, A.; Weiner, M. Electron-spin resonance (E.S.R.) hyperfine structure in hydrazyl free radicals. Z. Phys. Chem. 1962, 219, 285–294.
- Patrascu, B.; Lete, C.; Popescu, C.; Matache, M.; Paun, A.; Madalan, A.M.; Ionita, P. Synthesis and spectral comparison of electronic and molecular properties of some hydrazines and hydrazyl free radicals. Arkivoc 2020, vi, 1– 10.
- 18. Baciocchi, E.; Calcagni, A.; Lanzalunga, O. A kinetic study of the reaction of N,N-dimethylanilines with 2,2-diphenyl-1picrylhydrazyl radical: A concerted proton–electron transfer? J. Org. Chem. 2008, 73, 4110–4115.
- 19. Ionita, P.; Caproiu, M.T.; Balaban, A.T. New sulfonyl derivatives of 2,2-diphenyl-1-picrylhydrazyl and their supramolecular complexes with crown ethers or kryptands. Rev. Roum. Chim. 2000, 45, 935–941.
- 20. Ionita, P.; Constantinescu, T.; Luca, C.; Caldararu, H.; Caproiu, M.T.; Balaban, A.T. The reaction between the DPPH free radical and potassium cyanide in the presence of crown ether 18-C-6. New J. Chem. 1997, 21, 511–515.
- 21. Ionita, P.; Gilbert, B.C.; Whitwood, A.C. Synthesis and characterisation of some novel hetero-diradicals containing linked hydrazyl and nitroxide moieties. Perkin Trans. II 2001, 1453–1462.
- 22. Ionita, P. Hydrazyl-nitrones, novel hybrid molecules in free radical research. Free Radic. Res. 2006, 40, 59–65.
- 23. Gubanov, V.A.; Pereliaeva, L.A.; Chirkov, A.K.; Yastchenko, G.N.; Matevosian, R.O. Electronic absorption spectra for free hydrazyl radicals, their ionic forms and initial hydrazines. Int. J. Quantum Chem. 1971, 5, 513–524.
- Bordwell, F.G.; Zhang, X.M. From equilibrium acidities to radical stabilization energies. Acc. Chem. Res. 1993, 26, 510– 517.
- Bordwell, F.G.; Cheng, J.P.; Harrelson, J.A. Homolytic bond dissociation energies in solution from equilibrium acidity and electrochemical data. J. Am. Chem. Soc. 1988, 110, 1229–1231.
- 26. Foti, M.C.; Daquino, C. Kinetic and thermodynamic parameters for the equilibrium reactions of phenols with the dpph radical. Chem. Commun. 2006, 3252–3254.
- 27. Menanteau, T.; Levillain, E.; Breton, T. Electrografting via diazonium chemistry: From multilayer tomonolayer using radical scavenger. Chem. Mater. 2013, 25, 2905–2909.
- 28. Ionita, P.; Gilbert, B.C.; Whitwood, A.C. Generation of oxygen-, sulfur, nitrogen- and phosphorus-centred short-lived radicals via one-electron oxidation with stable hydrazyl radicals. Lett. Org. Chem. 2004, 1, 70–74.
- 29. Buettner, G.R. Spin trapping: ESR parameters of spin adducts. Free Radic. Biol. Med. 1987, 3, 259–303.
- Tabart, J.; Kevers, C.; Pincemail, J.; Defraigne, J.; Dommesa, J. Comparative antioxidant capacities of phenolic compounds measured by various tests. Food Chem. 2009, 113, 1226–1233.
- 31. Yeo, J.; Shahidi, F. Critical re-evaluation of DPPH assay: Presence of pigments affects the results. J. Agric. Food Chem. 2019, 67, 7526–7529.
- 32. Canillas, M.; Chinarro, E.; Freitas, M.; Pêgo, A.P.; Moreno, B. Titanium dioxide catalytic activity contributes to the process of free radical scavenging. J. Catal. 2020, 381, 186–192.
- 33. Zhu, Y.; Xu, J.; Lu, M. Oxidation of primary and secondary alcohols to the corresponding carbonyl compounds with molecular oxygen using 1,1-diphenyl-2-picrylhydrazyl and WO3/Al2O3 as catalysts. Catal. Comm. 2014, 48, 78–84.
- Wang, S.; Li, F.; Easley, A.D.; Lutkenhaus, J.L. Real-time insight into the doping mechanism of redox-active organic radical polymers. Nat. Mater. 2019, 18, 69–75.
- 35. Bai, W.L.; Zhang, Z.; Chen, X.; Wei, X.; Zhang, Q.; Xu, Z.X.; Liu, Y.S.; Chang, B.; Wang, K.X.; Chen, J.S. Boosting the electrochemical performance of Li–O2 batteries with DPPH redox mediator and graphene-luteolin-protected lithium anode. Energy Storage Mater. 2020, 31, 373–381.
- 36. Thiele, J.; Balhorn, H. Ueber einen chinoïden Kohlenwasserstoff. Ber. Dtsch. Chem. Ges. 1904, 37, 1463–1470.
- 37. Tschitschibabin, A.E. Über einige phenylierte Derivate des p,p-Ditolyls. Ber. Dtsch. Chem. Ges. 1907, 40, 1810–1819.
- 38. Muller, E.; Pfanz, H. Über biradikaloide Terphenylderivate. Ber. Dtsch. Chem. Ges. 1941, 74, 1051–1074.
- 39. Pyrzynska, K.; Pekal, A. Application of free radical diphenylpicrylhydrazyl (DPPH) to estimate the antioxidant capacity of food samples. Analyt. Methods 2013, 5, 4288–4295.

- 40. Tudose, M.; Angelescu, D.; Ionita, G.; Caproiu, M.T.; Ionita, P. New hydrazyl derivatives with multiple properties. Let. Org. Chem. 2010, 7, 182–185.
- 41. Likhtenshtein, G. Nitroxides—Brief History, Fundamentals, and Recent Developments; Springer International Publishing: Berlin/Heidelberg, Germany, 2020.
- 42. Peterson, J.P.; Winter, A.H. Solvent effects on the stability and delocalization of aryl dicyanomethyl radicals: The captodative effect revisited. J. Am. Chem. Soc. 2019, 141, 12901–12906.

Retrieved from https://encyclopedia.pub/entry/history/show/17803