## **Biogenic amines in Fish**

Subjects: Microbiology | Agriculture, Dairy & Animal Science Contributor: Pierina Visciano , Maria Schirone , Antonello Paparella

The occurrence of biogenic amines in fish is directly associated with microorganisms with decarboxylase activity. These compounds are generally detoxified by oxidases in the intestinal tract of humans, but some conditions, such as alcohol consumption, enzyme deficiency, or monoamino-oxidase antidepressant use, can make their intake by food dangerous. Due to its toxicity, histamine is a unique biogenic amine with regulatory limits for fishery products. This review focuses on biogenic amines in fish, with a detailed picture of the number of alert notifications or intoxication events reported in the last years. The favoring conditions for their formation, as well as the main preventive and control measures to ensure public health, are also reviewed.

biogenic amines

histamine food poisoning

outbreaks

seafood safety

## **1. Introduction**

Biogenic amines (BAs) are nitrogenous compounds resulting from the free amino acid decarboxylation or the amination of carbonyl-containing organic compounds through the metabolism of different microorganisms. Thus, their accumulation in food can be considered a good indicator of spoilage <sup>[1]</sup>. BAs are distinguished based on chemical structure into heterocyclic, aliphatic, or aromatic compounds (Table 1), or based on the number of amine groups into monoamines (tyramine and phenylethylamine), diamines (histamine, putrescine, and cadaverine), or polyamines (spermidine and spermine) <sup>[2]</sup>.

**Table 1.** Classification of biogenic amines based on chemical structure.

Precursors	Biogenic Amines			
	Aliphatic	Aromatic	Heterocyclic	
Arginine	Agmatine			
Lysine	Cadaverine			

Ornithine	Putrescine		
Phenylalanine		Phenylethylamine	
Tyrosine		Tyramine	
Histidine			Histamine
Tryptophan			Tryptamine

Fish products constitute an important part of the human diet because they are an excellent source of nutrients, including proteins, vitamins, salt minerals, and polyunsaturated fatty acids <sup>[3]</sup>. Nevertheless, they are very perishable due to postmortem modifications followed by the formation of spoilage compounds, such as organic acids, aldehydes and ketones, alcohols, sulfides, and BAs <sup>[4]</sup>. The inappropriate storage of fish and/or temperature abuse can lead to BA formation due to microbial enzymatic activities.

Gram-positive and -negative bacteria associated with fish spoilage can produce BAs. They are generally located on skin, gills, or in the gastrointestinal tract <sup>[5][6][7][8][9][10][11]</sup> (Table 2), and can spread to muscle tissue during butchering or gutting through rupture or spillage of gastric contents <sup>[12]</sup>. The most frequent species belong to *Enterobacteriaceae* and include mesophilic and psychrotolerant bacteria, such as *Morganella*, *Enterobacter*, *Hafnia*, *Proteus*, and *Photobacterium* <sup>[13]</sup>. Also, *Pseudomonas* spp. and lactic acid bacteria belonging to *Lactobacillus* and *Enterococcus* genera can cause BA formation <sup>[14]</sup>.

**Table 2.** Microorganisms producing biogenic amines in seafood.

Biogenic Amine	Microorganisms	References
Histamine	Morganella morganii, Morganella psychrotolerans, Hafnia alvei, Photobacterium phosphoreum	Hungerforf, 2010 <sup>[5]</sup>
	Photobacterium psychrotolerans, Klebsiella	Biji et al., 2016 <sup>[2]</sup>
	pneumoniae, Clostridium spp., Pseudomonas fluorescens, Pseudomonas putida, Pseudomonas	Doeun et al., 2017 <sup>[6]</sup>
	cepaciae, Aeromonas spp., Aeromonas hydrophila, Acinetobacter lowffi, Plesiomonas shigelloides,	Barbieri et al., 2019 [7]
	Proteus vulgaris, Proteus mirabilis, Serratia fonticola,	Xu et al., 2020 <sup>[8]</sup>

	Serratia liquefaciens, Enterobacter cloacae, Enterobacter aerogenes, Klebsiella oxytoca, Citrobacter freundii, Raoultella planticola, Staphylococcus xylosus, Staphylococcus epidermidis, Bacillus spp., Vibrio alginolyticus, Vibrio spp., Escherichia spp.,	
Tiramine	Lactic acid bacteria (including <i>lactobacilli</i> , <i>lactococci</i> , <i>enterococci</i> and <i>carnobacteria</i> )	Marcobal et al., 2012 <sup>[9]</sup>
Putrescine	Enterobacter spp., Hafnia alvei, Pantoea agglomerans, Serratia liquefaciens, Photobacterium phosphoreum, Aeromonas spp., Lactobacillus curvatus, Lactobacillus sakei, Carnobacterium divergens	Wunderlichová et al., 2014 [ <u>10]</u>
Cadaverine	Pseudomonads, Enterobacteriaceae	Paleologos et al., 2004 <sup>[11]</sup> Kuley et al., 2017 <sup>[4]</sup>

The amounts of BAs ingested by food are regulated in the human organism by a detoxification system formed mainly by mono and diamino oxidases. High intake of BAs or the presence of factors such as the use of alcohol or medication that reduce the effectiveness of such detoxifying enzymes, rather than a genetic deficiency, can lead to intoxication with different symptoms depending on the type of BA. Histamine is the most toxic, as it can act as neurotransmitter and vasodilator, causing headache, hypotension, heart palpitations, asthma attacks, and cutaneous (edema and flushing of the face, neck, and upper arms) or gastrointestinal (difficulties in swallowing, vomiting, and diarrhea) effects <sup>[Z]</sup>. Bronchospasm, respiratory distress, and vasodilatory shock are also described <sup>[15]</sup>. A detailed summary of histamine intoxication is reported in Table 3. Instead, tyramine may increase the cardiac frequency or cause respiratory disorders, but also nausea and vomiting, while phenylethylamine can be a migraine inductor. Further symptoms associated with tyramine and phenylethylamine are hypertension and cerebral hemorrhage. Even if putrescine and cadaverine are not toxic, they can potentiate the adverse effects of the above cited BAs, as they favor their adsorption or interfere with the detoxification system <sup>[16][17]</sup>.

In the literature, several reviews are reported on BAs in fish. Some authors <sup>[2][6][18]</sup> described in detail the content of histidine in some fish species, the decarboxylation reactions, some indexes of quality, and toxicological effects of many BAs, not only histamine, alongside the presence of these compounds in various foods. Instead, the focus of this review is the statement of outbreaks reported in the Member States of European Union (EU) and the

notifications of the Rapid Alert System of Food and Feed (RASFF). In addition, factors affecting BA formation are also discussed.

Apparatus	Symptoms
Integumentary	Face, neck ,and upper arm flushing, itchy rash, hives, localized swelling, redness, urticaria, pruritus
Cardiovascular	Hypotension with distributive shock, cardiac arrhythmias, myocardial disfunction, acute pulmonary edema, oral numbness, tingling
Gastrointestinal	Abdominal pain, stomach cramps, nausea, vomiting, diarrhea
Neurological	Throbbing headache, migraines, loss of sight, dizziness, faintness, anxiety, tremor
Respiratory	Asthma attacks, respiratory distress, rhinitis, bronchoconstriction, dyspnea
Other	Metallic or peppery taste, oral numbness, difficulties in swallowing and thirst, feeling of warmth around the mouth

## 2. Biogenic Amines in Seafood: Focus on Scombroid Poisoning Outbreaks

Histamine, tyramine, putrescine, and cadaverine are the most common BAs found in fish and derive from decarboxylation of corresponding free amino acids by microorganisms <sup>[18]</sup>. Their accumulation depends on the presence of the precursor amino acids, the growth or activity of decarboxylating bacteria, and a favorable environment. The main factors that influence the microbiota and its enzymatic activity are temperature, pH, water activity, oxygen availability, concentration of NaCl, some additives, and competition among microorganisms. The combination of these factors can be responsible of the variability of BA content within the same batch, and also within individual fish <sup>[12][19]</sup>.

In the flesh of fish, BAs can be considered an indicator of good handling and storage procedures, and some authors reported different chemical indices for fish quality as combinations of amines <sup>[20]</sup>. However, the levels of

different BAs are associated with the predominant muscle type in fish. Fish with dark muscles have more histidine content compared to those with white muscles, and therefore they accumulate more histamine when kept under elevated temperatures. On the contrary, fish belonging to the second category (white muscles) can show high cadaverine and putrescine concentrations due to poor handling, but also temperature abuse. For cephalopods, agmatine is used as a guality indicator, while in the case of crustaceans, shrimp, and lobsters, putrescine and cadaverine are considered <sup>[21]</sup>. Individual BA or a combination of various amines are considered as a quality index of fish freshness. Some authors reported that a histamine content lower than a value of 10 mg/kg denotes fish of good guality, while concentrations between 30 and 50 mg/kg represent important and definite deterioration, respectively <sup>[22]</sup>. Instead, the consumption of meals with histamine concentrations of 8–40 mg can cause only slight intoxication, while values of 40-100 mg or higher than 100 mg are associated with intermediate and severe intoxication, respectively <sup>[2]</sup>. The Food and Drug Administration (FDA) established a defect action level of 50 mg/kg for histamine (according to the revised Guidance in year 2020) <sup>[23]</sup>, whereas the Commission Regulation (EC) No 2073/2005, which is currently in force, fixed maximum levels of 200 and 400 mg/kg for raw fish and fishery products subjected to enzyme maturation treatment in brine, respectively. These limits apply to the fish families Scombridae, Scomberosocidae, Engraulidae, Clupeidae, Coriphaenidae, and Pomatomidae, with high concentrations of free histidine in the muscle tissue. Some outbreak reports <sup>[24]</sup> and scientific studies confirmed that fish with low levels of the precursor amino acid showed no toxic histamine content in their flesh, whereas the opposite situation was described by Visciano et al. <sup>[25]</sup> in fish experimentally subjected to temperature abuse and belonging to the above cited families with regulatory limits.

Scombroid poisoning occurs worldwide and the largest numbers of such events are described in the United States, the United Kingdom, Australia, and Japan. Up to 40% of foodborne outbreaks reported in Europe and the United States can be ascribed to histamine intoxication <sup>[26]</sup>. In Table 4, the histamine poisoning and human cases in the EU from 2010 to 2018 are shown. Even if a positive trend was observed during the years 2010–2017, a sudden decrease was described in 2018, probably due to the lack of case reports from some Member States <sup>[27][28][29][30]</sup> [31][32][33][34][35].

Outbreaks (N)	Cases (N)

 Table 4. Number of histamine poisoning outbreaks reported in the European Union during the years 2010–2018 [27]

 [28][29][30][31][32][33][34][35]

Year					
	Strong-Evidence	Weak-Evidence	Total	Human Cases	Hospitalized
2018	24	56	80	488	115

2017	56	61	117	572	51
2016 *	28	78	106	489	74
2015	23	57	80	437	43
2014	35	_ **	35	164	15
2013	42	-	42	231	30
2012	34	-	34	241	14
2011	58	-	58	259	31
2010	33	-	33	185	12

Legend: \* Data reported as other causative agents that include chemical agents, histamine, lectin, marine biotoxins, mushroom toxins, and scombrotoxin; \*\* data of weak-evidence outbreaks not reported.

A Rapid Alert System for Food and Feed RASFF was organized among the EU Member States for the notification of risks to human health was established by the Regulation (EC) No 178/2002, with histamine representing one of the most common causes of such notifications. The list of notifying countries and the countries of origin of notifications for histamine presence in fish products from 2015 to 2020 are shown in Figures 1 and 2, respectively. Italy is the major notifying Member State, with a total of 74 alert or information for attention notifications, followed by France, with 33. Instead, the main origin of fish and fish products contaminated by histamine is linked to Spain (56 cases). Most notifications derived from the analysis of products showed high concentrations of histamine (Table 4), whereas some cases (n = 22) were associated with foodborne outbreaks generally occurring in the notifying country. The highest percentage of both alert (30.2%) and information for attention (34.4%) notifications reported a histamine content (Table 5) that slightly exceeded the maximum limits set by the Commission Regulation (EC) No 2073/2005. Instead, the highest values, i.e., >1000 or even >2000 mg/kg, referred to scombroid poisoning caused by chilled or canned fish, particularly regarding tuna species. In Figures 3 and 4, the fish species and types of fish products (i.e., raw or processed) involved in the RASFF reports from 2015 to August 2020 are shown. Tuna is the most representative species because it presents high concentrations of histidine in muscle tissue for its high speed and long duration swimming as predator fish <sup>[36]</sup>. Moreover, harvesting practices, such as longlining and gillnetting,

can contribute to histamine formation due to the long period in which the fish remains in the sea before it is brought onboard the vessel. Such conditions are particularly dangerous for tuna species, which can generate heat into their body that exceeds the environmental temperature, thereby favoring the growth of histamine-forming microorganisms <sup>[23]</sup>.



**Figure 1.** List of notifying countries and number of notifications for histamine presence in "fish and fish products" by the Rapid Alert System for Food and Feed (RASFF) from 1 January 2015 to 31 August 2020.



**Figure 2.** List of countries of origin with notifications of histamine presence in "fish and fish products" by the Rapid Alert System for Food and Feed (RASFF) from 1 January 2015 to 31 August 2020.



**Figure 3.** Distribution of histamine in fish species according to the Rapid Alert System for Food and Feed (RASFF) from 1 January 2015 to 31 August 2020.



**Figure 4.** Presence of histamine in different fish products according to the Rapid Alert System for Food and Feed (RASFF) from 1 January 2015 to 31 August 2020.

**Table 5.** Histamine concentrations reported in "fish and fish products" by the Rapid Alert System for Food and Feed(RASFF) from 1 January 2015 to 31 August 2020.

Notification	Histamine Concentrations (mg/kg)					
	≤200	>200 and ≤500	>500 and ≤1000	>1000 and ≤2000	>2000	Total
IA*	31 (20.6%)	52 (34.4%)	23 (15.2%)	29 (19.2%)	16 (10.6%)	151 (100%)
Alert	15 (12.9%)	35 (30.2%)	17 (14.7%)	26 (22.4%)	23 (19.8%)	116 (100%)
BR **	8 (22.2%)	18 (50%)	9 (25%)	-	1 (2.8%)	36 (100%)

Legend: \* information for attention; \*\* border rejection.

## References

- 1. Zhong, X.; Huo, D.; Fa, H.; Luo, X.; Wang, Y.; Zhao, Y.; Hou, C. Rapid and ultrasensitive detection of biogenic amines with colorimetric sensor array. Actuators B Chem 2018, 274, 464–471.
- 2. Biji, K.B.; Ravishankar, C.N.; Venkateswarlu, R.; Mohan, C.O.; Srinivasa Gopal, T.K. Biogenic amines in seafood: A review. Food Sci. Technol. 2016, 53, 2210–2218.
- Visciano, P. Chemicals and Safety of Chemical Contaminants in Seafood. In Food Safety Chemistry. Toxicant Occurrence, Analysis and Mitigation; Liangli, L.Y., Shuo, W.; Bao-Guo, S., Eds.; CRC Press; Taylor & Francis Group: Boca Raton, FL, USA, 2015; pp. 215–236.
- Kuley, E.; Durmus, M.; Balikci, E.; Ucar, Y.; Regenstein, J.M.; Özoğul, F. Fish spoilage bacterial growth and their biogenic amine accumulation: Inhibitory effect of olive by-products. J. Food Prop. 2017, 20, 1029–1043.
- 5. Hungerford, J.M. Scombroid poisoning: A review. Toxicon 2010, 56, 231–243.
- Doeun, D.; Davaatseren, M.; Chung, M.S. Biogenic amines in foods. Food Sci. Biotechnol. 2017, 26, 1463–1474.
- 7. Barbieri, F.; Montanari, C.; Gardini, F.; Tabanelli, G. Biogenic amine production by Lactic Acid Bacteria: A review. Foods 2019, 8, 1–27.
- 8. Xu, Y.; Zang, J.; Regenstein, J.M.; Xia, W. Technological roles of microorganisms in fish fermentation: A review. Rev. Food Sci. Nutr. 2020, doi:10.1080/10408398.2020.1750342.
- 9. Marcobal, A.; De Las Rivas, B.; Landete, J.M.; Tabera, L., Muñoz, R. Tyramine and phenylethylamine biosynthesis by food bacteria. Rev. Food Sci. Nutr. 2012, 52, 448–467.
- Wunderlichová, L.; Buňková, L.; Koutný, M.; Jančová, P.; Buňka, F. Formation, degradation, and detoxification of putrescine by foodborne bacteria: A review. Rev. Food Sci. Food Saf. 2014, 13, 1012–1030.
- Paleologos, E.K.; Savvaidis, I.N.; Kontominas, M.G. Biogenic amines formation and its relation to microbiological and sensory attributes in ice-stored whole, gutted and filleted Mediterranean Sea bass (Dicentrarchus labrax). Food Microbiol. 2004, 21, 549–557.
- FAO/WHO (Food and Agriculture Organization of the United Nations/World Health Organization).
   Public Health Risks of Histamine and Other Biogenic Amines from Fish and Fishery Products.
   Meeting Report; FAO Headquarters: Rome, Italy, 2012.

- Comas-Basté, O.; Latorre-Moratalla, M.L.; Sánchez-Pérez, S.; Veciana-Nogués, M.T.; Vidal-Carou, M.C. Histamine and other biogenic amines in food. From Scombroid Poisoning to histamine intolerance. In Biogenic Amines; Proestos, C., Ed.; IntechOpen: London, UK, 2019.
- 14. Fusek, M.; Michálek, J.; Buñková, L.; Buñka, F. Modelling biogenic amines in fish meat in Central Europe using censored distributions. Chemosphere 2020, 251, 126390.
- 15. Stratta, P.; Badino, G. Scombroid poisoning. CMAJ 2012, 184, 674.
- Palomino-Vasco, M.; Acedo-Valenzuela, M.I.; Rodríguez-Cáceres, M.I.; Mora-Diez, N. Automated chromatographic method with fluorescent detection to determine biogenic amines and amino acids. Application to craft beer brewing process. Chromatogr. A 2019, 1601, 155–163.
- Silva, I.P.; Dias, L.G.; da Silva, M.O.; Machado, C.S.; Paula, V.M.B.P.; Evangelista-Barreto, N.S.; de Carvalho, C.A.L.; Estevinho, L.M. Detection of biogenic amines in mead of social bee. LWT Food Sci. Technol. 2020, 121, 108969.
- Prester, L. Biogenic amines in fish, fish products and shellfish: A review. Food Addit. Contam. 2011, 28, 1547–1560.
- 19. Visciano, P.; Schirone, M.; Tofalo, R.; Suzzi, G. Biogenic amines in raw and processed seafood. Microbiol. 2012, 3, 1–10.
- 20. Mattsson, L.; Xu, J.; Preininger, C.; Bui, B.T.S.; Haupt, K. Competitive fluorescent pseudoimmunoassay exploiting molecularly imprinted polymers for the detection of biogenic amines in fish matrix. Talanta 2018, 181, 190–196.
- 21. Sedaghati, M.; Mooraki, N. Biogenic amines in sea products. Surv. Fish. Sci. 2019, 6, 1–8.
- 22. Muscarella, M.; Lo Magro, S.; Campaniello, M.; Armentano, A.; Stacchini, P. Survey of histamine levels in fresh fish and fish products collected in Puglia (Italy) by ELISA and HPLC with fluorometric detection. Food Control 2013, 31, 211–217.
- FDA (Food and Drug Administration). Fish and Fishery Products Hazards and Controls Guidance, 4th ed.; FDA: Silver Spring, MD, USA, 2020; Chapter 7—Scombrotoxin (Histamine) Formation; pp. 113–152.
- 24. Colombo, F.M.; Cattaneo, P.; Confalonieri, E.; Bernardi, C. Histamine food poisonings: A systematic review and meta-analysis. Rev. Food Sci. 2018, 58, 1131–1151.
- 25. Visciano, P.; Campana, G.; Annunziata, L.; Vergara, A.; Ianieri, A. Effect of storage temperature on histamine formation in Sardina pilchardus and Engraulis encrasicolus after catch. Food Biochem. 2007, 31, 577–588.
- 26. Guergué-Díaz de Cerio, O.; Barrutia-Borque, A.; Gardeazabal-García, J. Scombroid poisoning: A practical approach. Actas Dermodifiliogr. 2016, 107, 567–571.

- 27. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union One Health 2018 Zoonoses Report. EFSA J. 2019, 17, 5926.
- 28. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2017. EFSA J. 2018, 16, 5500.
- 29. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2016. EFSA J. 2017, 15, 55077.
- 30. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2015. EFSA J. 2016, 14, 4634.
- 31. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2014. EFSA J. 2015, 13, 4329.
- 32. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2013. EFSA J. 2015, 13, 3991.
- 33. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2012. EFSA J. 2014, 12, 3547.
- 34. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2011. EFSA J. 2013, 11, 3129.
- 35. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2010. EFSA J. 2012, 10, 2597.
- 36. Mercogliano, R.; Santonicola, S. Scombroid fish poisoning: Factors influencing the production of histamine in tuna supply chain. A review. LWT Food Sci. Technol. 2014, 114, 108374.

Retrieved from https://www.encyclopedia.pub/entry/history/show/10886