

# Bagging on Fruit Quality

Subjects: Horticulture

Contributor: Muhammad Moaaz Ali

Fruit bagging consists essentially of enclosing a young fruit in a food bag by capping the bag with a ribbon or a clamp on the fruit stalk. Isolating the fruit from the external environment protects it during development from mechanical or biotic damage, especially in regions where fruits are prone to attacks by fungi, bacteria, insects and even birds.

fruit skin color

light-induced coloration

fruit appearance

bag material

anthocyanin content

texture


## 1. Introduction

Fruits are a source of numerous compounds essential for the human body and are included in a well-balanced healthy diet. Although fruits and vegetables are low in calories, the nutritive value of fruits has gained interest nowadays, being the source of health-promoting vitamins, fibers, minerals [1][2], phytochemicals and bioactive compounds, which help to prevent cancers and cardiovascular risks [3][4][5]. Sufficient intake of fruits and vegetables replaces harmful saturated fats and sugars from the body and enhances healthy nutrients and dietary fiber [6]. The Food and Agriculture Organization of the United Nations (FAO) and the World Health Organization (WHO) recommend consuming at least 400 g of fresh fruits and vegetables each day [7]. However, physical and biochemical attributes of fruits are greatly influenced by environmental factors [8][9].

The bagging technique, which was first utilized in Japan in the 20th century for pears and grapes, is now widely applied in Asian countries (Japan, China, Korea), Australia and the USA, protecting fruits from the surrounding environment (mainly from light and pathogens, then stresses related to temperature, water/humidity, and air movement) with a sort of shield—a physical barrier around the fruit [9]. In fact, bagging consists essentially of enclosing a young fruit in a food bag by capping the bag with a ribbon or a clamp on the fruit stalk. Isolating the fruit from the external environment protects it during development from mechanical or biotic damage, especially in regions where fruits are prone to attacks by fungi, bacteria, insects and even birds [9]. The purpose is to obtain fruits without external imperfections, and with desired shape and color depending on the regional or national consumer preferences for the specific fruit. The expected color changes in comparison to non-bagged fruits can therefore correspond either to a reduction or an increase in color or even a greater homogeneity of the color itself. This is particularly important in markets, e.g., Japan, where aesthetic factors represent an important competitive factor.

## 2. The Role of Bagging on Fruit Quality

Bagging technique is used specifically to enhance fruit appearance and quality, especially in Asia. There are different types of bags/bagging material (Figure 1). Initially, newspaper bags were used to wrap fruits to prevent damage from pests and diseases in Korea, but around 1985, artificially manufactured bags were introduced.



**Figure 1.** Different types of bagging materials used to improve the quality of fruits: (a) transparent paraffin bag; (b) nylon bags; (c) blue colored plastic bags; (d) two-layered bag (yellow paper outside and plastic inside); (e) cellophane bags; (f) organza bags; (g) paper bags; (h) two-layered paper bag (brown outside and grey inside).

Though the bag production cost is high and the practice labor intensive, bagging with new materials has shown excellent results. A bag around a fruit controls sunlight, temperature, humidity, evaporation and mechanical damage. Bagging may also regulate harvesting time [10], and it can control pest attacks, especially fruit flies, minimizing residues of pesticides [11][12][13], which is particularly important during the rainy-season [14]. Thus, bagging is an excellent method to yield fruits with a very low input or residues of pesticide. In addition, bagging is able to promote the production of high-value organic fruits, as demonstrated for organic peaches in the southeastern United States by Allran et al. [15], who showed that fruit quality (size, Brix degree, acidity) was similar between bagged and control fruits, and by Campbell et al. [16], who reported that bagging protects against various pests and diseases but has minimal effects on organic peach quality. Similar findings were obtained by Araújo Neto et al. [17] after a bagging treatment of organic guava fruits. In addition, for organic fruits, bags can be doubled [18], or, in conventional farming, impregnated with insecticide [19] or sprayed with insecticides/fungicide before bagging [20].

Bagging can determine numerous changes in the physiology of the fruit and in the preservation of its characteristics, and particular attention has been paid to tropical fruits, for which there are numerous applications (Table 1), often found also for other types of fruits.

**Table 1.** The effects of bagging on color, quality and physiological disorders of some tropical fruits.

Crop/Cultivar	Bagging Start	Bagging Material	Effect	Ref.
Mango "Langra" and "Khirshapat"	30 d before harvest	black polybag, transparent polybag, brown paper	higher total soluble sugars and better physical quality of fruit	[21]
Mango "Nam Dok Mai #4"	48 d after full bloom	two-layered paper (brown outside and black inside)	improvement in fruit weight and skin appearance	[22]
Mango "Harumanis"	56 d before harvest	brown and black paper	improvement in skin color	[23]
Mango "Nam Dok Mai #4"	45 d after full bloom	low-density polyethylene	improvement in fruit weight and skin glossiness	[24]
Mango "Apple"	40–50 d before harvest	standard Kraft paper	reduction in lenticel discoloration	[25]
Mango "Khirsapat"	42 d before harvest	brown paper	reduced significantly post-harvest losses	[26]
Carambola "Malaysia"	10–31 d after full bloom	plastic, newspaper, and non-woven cloth	increase in fruit size, fruit weight and soluble solid content	[27]
Guava "Allahabad Safeda"	30 days after pollination	nylon nets, non-woven polypropylene, butter paper and brown paper	advanced fruit maturity, improved fruit weight, texture, visual appeal, quality and functional attributes	[14]
Guava "Tai-Kuo"	for 146 and 175 d during fruit development	waxed paper, nylon, Taiwan bag, telephone book paper	protection against pests and mechanical damage	[28]
Litchi "Feizixiao"	15 and 30 d after full bloom	cellophane or fabric	better fruit coloration/appearance	[29]
Litchi "Rose Scented"	14 d before harvest	perforated transparent polyethylene	reduction in fruit drop. increase in fruit size, higher soluble solids content	[30]
Loquat "Baiyu" and "Ninghaibai"	after fruit thinning (early April)	white paper (50% light transmittance) and two-layered paper (out grey, inside black—0% light transmittance)	improvement in appearance decrease in fruit weight	[31]
Loquat "Qingzhong"	after fruit thinning	paper	promotion in appearance, increased sucrose, glucose and soluble solids content, decreased	[32]

Crop/Cultivar	Bagging Start	Bagging Material	Effect	Ref.
			fructose, sorbitol and titratable acidity content	
Longan "Chuliang"	34 d after full bloom	perforated plastic, white or black adhesive-bonded	increased fruit size and fruit retention rate, reduced fruit cracking incidence	[33]
Persimmon "Shinsyu"	35–50 d before harvest	paper	no black stain	[34]
Persimmon "Fuyu"	1–4 months before harvest	white paper (40% shade)	reduced fruit blemishing (increase of blemishing with early removal)	[35]
Yuzu ( <i>Citrus junos</i> )	early September	recycled Japanese phone book paper, grey colored paper and black polyester	significant reduction in fruit spot injury	[36]
Date Palm "Zaghoul"	at pollination time	transparent and blue polyethylene	reduction in tip cracked fruit	[37]
Date Palm "Succary" and "Khalas"	28 d after pollination	black, white blue, yellow plastic	acceleration fruit ripening	[38]
Date Palm "Helali"	30 d after pollination	black and blue polyethylene, paper	[41] increased rate of fruit ripening	[39]

proven to be successful against chilling injury, preventing browning of the banana pulp [19][42].

Bagging can increase fruit sugars and organic acid contents, two significant determinants of fruit organoleptic quality [43], although the response to bagging varies according to the fruits considered. Indeed, Zhou et al. [44] reported a decrease in sugar content after bagging of Chinese white olives (*Canarium album* (Lour.) Räusch.), as it was found for apple [45] and also date [38]. Conversely, Sarker et al. [21] and Islam et al. [46] reported an increase in sugar contents in bagged mango fruits, while Bently and Viveros [47] registered an improvement of fruit sweetness in Granny Smith apple. Huang et al. [48] stated that bagging has a non-significant effect on soluble sugars but decreases organic acids in pear fruits. Kim et al. [10] reported that peach fruits bagged with yellow paper (Figure 1d) showed an increase in total titratable acids due to low light, and white-colored bags determined an increase of soluble solid contents, chlorophyll and anthocyanins.

Xu et al. [31] investigated the effects of different light transmitting paper bags on fruits of two different cultivars of loquat ("Baiyu" and "Ninghaibai"); bagging materials included one layer white paper bags with ~50% light transmittance ( $T_1$ ), and paper bags with a black inner layer and a grey outer layer with ~0% light transmittance ( $T_2$ ). Fruit weight decreased, but fruit appearance improved with bagging, whereas total sugar content was higher in fruits subjected to  $T_1$  treatment than  $T_2$  and control. Both bagging materials reduced phenolics and flavonoids, with the lowest contents in  $T_2$  fruits [31]. Sharma et al. [49] reported that bag color also influences total fruit sugars in Red Delicious apples; Asrey et al. [50] indicated that red cellulosic bags applied 60 days after flowering are successful in producing high-quality pomegranate fruits (characterized by high consumer acceptability) in terms of

total anthocyanin and ascorbic acid content, although with slightly lower calcium and total phenol; instead, Pantone® 1205C bags determined a delay in pomegranate fruit development and ripening, which were outweighed by a reduced incidence of peel sunburn and higher antioxidant activity [51]. Yang et al. [33] observed that in longan fruits, sugar content was not significantly affected by bag type but resulted in an increase of fruit size and reduced cracking.

In apple, bagging determined a better absorption of calcium by the fruits with a lower incidence of bitter pit in the cultivars "Red Fuji", "Fuji Suprema", "Imperial Gala" and "Gamhong" [52][53][54][55].

Bagging technique leads to the production of more attractive fruits due to fewer blemishes and visible marks [9], particularly in apple [47][49][56][57], pear [12][58][59][60][61], peach fruits [10][62], pomegranate [41], mango [21][22][23][24][25], carambola [27], guava [14][28], litchi [29][30], loquat [31][32], persimmon [34][35] and yuzu [36]. In addition, post-harvest losses are significantly reduced for mango [26].

However, some studies have also reported a negative impact of fruit bagging, for example reduced concentration of essential elements in mango [63]; smaller fruit size for loquat, pear, pomegranate and apple [9]; lower content in sugars and organic acids in apple [45]; ascorbate decline in pear [64]; and a reduced level of total carotenoids in peach [65].

### 3. Light and Fruit Flavonoids


Light is required for the photosynthetic process that provides the chemical energy needed for plant growth and productivity. Moreover, plant metabolism, gene expression and plant processes (e.g., movement of stomatal guard cells, abscission, mineral absorption, phototropism) are regulated or conditioned by light [66][67][68][69].

Concerning fruits, several researchers have proposed that solar radiation can induce changes in the flavonoid levels in terms of both quality and quantity [70][71][72]. Others have observed that light can elicit the expression of genes such as phenylalanine ammonia-lyase (*PAL*), chalcone synthase (*CHS*) or flavanone 3-hydroxylase (*F3H*), which are involved in the biosynthesis of flavonoids [73][74][75]. *F3H* catalyzes the stereospecific 3b-hydroxylation of (2S)-flavanones to the dihydroflavonols and is required for the biosynthesis of flavonols and anthocyanins [73][74], representing antioxidant compounds able to protect leaves from high light intensity and other stressful conditions [75].

In *Arabidopsis*, the *BANYLUS* (*BAN*) gene encodes anthocyanin reductase, which converts anthocyanidins to their corresponding 2,3-cis-flavan-3-ols on the pathway of condensed tannins; in fact, a mutation in the *BAN* gene leads to the accumulation of anthocyanins and a loss of condensed tannins in *Arabidopsis* seeds [76]. A correlation between the expression of the flavonoid pathway genes and the anthocyanin accumulation was demonstrated in bilberry ripening fruits [77]; in addition, the upper bilberry leaves exposed to direct sunlight showed an increase in the expression of flavonoid pathway genes and a higher concentration of anthocyanins, catechins and flavonols in

comparison with lower shaded leaves [78]. These data support a protective role of flavonoids against excess solar radiation, and that high light conditions increase the accumulation of anthocyanins [73][79].

Interestingly, Zhao et al. [74] irradiated with UVA or UVB peach fruits, following 60–70 days of bagging, and proposed that UV light regulates the biosynthesis of anthocyanins, altering expression of several light receptors and in turn up-regulating several genes of the biosynthetic pathway; the working hypothesis was that photoreceptors interact with signal transduction elements of photomorphogenesis (constitutive photomorphogenic 1 (COP1), constitutive photomorphogenic 10 (COP10), phytochrome-interacting basic helix-loop-helix transcription factor (PIF), suppressor of phytochrome A (PHYA) (SPA), squamosa promoter-binding protein-like (SPL), which impact light-reaction effectors downstream (elongated hypocotyl 5 (HY5), elongated hypocotyl homologue 5 (HYH)) and the MYB–bHLH–WD40 (MBW) complex (myeloblastosis (MYB)/basic helix-loop-helix (bHLH)/WD40 domain-containing protein (WD40)) to regulate the transcription of the genes involved in the anthocyanin biosynthesis in response to light, as summarized in [Figure 2](#) [73][80]. Especially, the “Granny Smith” apple underwent red pigmentation after bag removal, whereas both unbagged and bagged until harvest fruits did not acquire any tone of red; moreover, the expression of *PHYE*, *phototropin2 (PHOT2)* and of the *UVB photoreceptors UV resistance locus8 (UVR8)*, *DE-ETIOLATED (DET)*, two *phytochrome kinase substrates (PKS1 and PKS3)* and *COP1* tightly correlated with anthocyanin levels in apple skin [81].



**Figure 2.** A model showing fruit peel coloration induced by light. UVR8—UV resistance locus 8; CRY—cryptochrome; PHOT—phototropin; PHY—phytochrome; SPL—SQUAMOSA promoter-binding protein-like; PIF—phytochrome-interacting basic helix-loop-helix transcription factors; COP—constitutive photomorphogenic; SPA—suppressor of PHYA; PKS—phytochrome kinase substrate; NAC—NAM (no apical meristem)/ATAF (Arabidopsis transcription activation factor)/CUC (cup-shaped cotyledon) transcription factor; DET—DE-ETIOLATED; HY5—elongated hypocotyl 5; HYH—HY5 homolog; WD40—WD40 domain-containing protein; bHLH—basic helix-loop-helix; MYB—myeloblastosis (modified after Chaves-Silva et al. [73], Zhao et al. [74], Ma et al. [81] and Zhou et al. [82]).

Concerning the transcriptional regulation of anthocyanin biosynthesis, the most studied fruits are apple, strawberry and grape [73]. Particularly, in red-fleshed apple, two fruit variants have been identified; type I shows pigmentation in plant vegetative organs, and fruits exhibit a more intense color at early stages of development, reducing the color at ripening, whereas in type II apple pigmentation occurs only in fruit tissues (peel and flesh), which acquire color during maturation [83]. This means that light may regulate the biosynthesis of anthocyanins at different development stages in the two apple types.

## 4. Bagging and the Color of Fruits

Since color is the main attractor for the purchase of fruit, bagging has been mainly used to obtain a specific color of the fruit skin and as a simple method to study the gene expression of the anthocyanin pathway in fruits [84]. Although some experiments have also been conducted on tropical fruits (as reported in the previous section), great attention has been paid to some pome fruit, stone fruit or vines.

In apple, besides the pigmented type I and type II varieties, other important commercial cultivars are typically acyanic, such as "Granny Smith" and "Golden Delicious", but fruits turn to pink/red after bag removal [85]. The red coloration ten days after bag removal is more intense for "Granny Smith" than for "Golden Delicious"; this was associated with a different level of *MdMYB1* gene expression, which seems to be the consequence of hypomethylation of the *MdMYB1* promoter in "Granny Smith" [85]. Further investigation analyzing differential expressed genes between unbagged, bagged and bag removed (before harvest) confirmed the importance of *MdMYB1* and other genes as *PHYE*, *PHOT2*, *UVR8*, *DET*, *PKS1*, *PKS3* and *COP1* for perception and transduction of the light signal after a dark period inside the bag [81]. From a practical point of view, the conclusion is the opportunity to realize the bagging of apples with materials that allow the passage of a substantial part of light radiation to maintain unaltered the color of red apples [86] and to avoid the blush of the skin in acyan apples [47][81]. Alternatively, bags must be removed a few weeks before harvest to avoid the red color reduction in cyan apples [57] ([Table 2](#)).

**Table 2.** The influence of bagging on physiological disorders, color and quality of apple fruits.

Apple Cultivar	Bagging Start	Bagging Material	Effect	Ref.
"Granny Smith"	40 d after full bloom (removed at 160 d after full bloom)	two-layer paper (outer brown, inner red)	increase in anthocyanin content after bag removal, increased expression of genes involved in light signal perception and transduction	[81]
"Qinguan" (deep-red cultivar), "Cripps Pink" (pale-red cultivar), and "Golden Delicious" (non-red cultivar)	45 d after full bloom	double layer paper (outer yellow, inner red paper coated with wax)	reduced anthocyanin accumulation in red cultivars, reduced sugar and organic acid contents	[45]
"Granny Smith"	114–118 d before harvest	brown paper	improvement of sweetness, sunburn reduction, 30 to 40% additional yield	[47]
"Delicious"	30 d before harvest	light yellow fabric	improvement in fruit color, firmness, and reduction in postharvest disorders	[49]

Apple Cultivar	Bagging Start	Bagging Material	Effect	Ref.
“Red Fuji”	40 d after full bloom	paper	better absorption of calcium in fruit	[52]
“Gamhong”	28–35 d after full bloom	Ca-coated paper	reduction in bitter pit	[53]
“Fuji Suprema”	40 d after full bloom	transparent micro-holed plastic and non-textured fabric	lower incidence of bitter pit, higher incidence of russetting, improvement in Ca content	[54]
“Imperial Gala”	40 d after full bloom	transparent micro-perforated plastic or non-textured fabric bags	reduction in bitter pit incidence	[55]
“Golden Delicious”	113 d before harvesting	two double layer paper: (a) outside grey–inside yellow; (b) outside newspaper–inside yellow	improved fruit skin, slightly decrease in size and weight	[56]
“Kurenainoyume”	39–54 days after full bloom (removed 29–48 d before harvesting)	light impermeable double-layered paper	incidence of cork spot in non-bagged fruits, no decrease in flesh firmness during storage	[57]
“Golden Delicious” and “Granny Smith”	40 d after full bloom (removed at 120 d or 160 d after full bloom)	two-layer paper (outer brown, inner red)	red/pink pigmentation after bag removal, more intense in “Granny Smith”	[85]
“Idared”	40 d after full bloom	1–3 layers of black hail net	small increase in mechanical properties Increase in russet susceptibility	[86]
“Fuji Raku Raku”	60–75 d after full bloom	double layer paper (outer grey, inner red)	lower internal browning with more rotting, lower phenolic content	[87]

*P. pyrifolia*) fruits are subjected to light within 10 days from bag removal [48]. Interestingly, the pigmentation patterns of *P. pyrifolia* (cultivar “Mantianhong”) differs from *P. communis* (cultivar “Cascade”) [88]; the first one develops a red color after bagging removal or postharvest UV/VIS irradiation. At the same time “Cascade” did not respond to light or UV exposure [88]. Additionally, the same authors indicated *PyMYB10* as the key regulator of anthocyanin biosynthesis in response to light [88]. Kim et al. [89] confirmed that in *P. communis* (cultivar “Kalle”), the anthocyanin contents in unbagged fruits remain higher than in bagged fruit. Qian et al. [90] employed bagging to study the light control of anthocyanin biosynthesis in pear fruit, demonstrating that *miR156* was expressed in peels, increased after removing the bags, targeted four *SPL* genes and, additionally, *PpSPL10* and *PpSPL13* interact with *PpMYB10*. More recently Zhu et al. [91] have investigated the light-response patterns of 27 pear cultivars after

bagging confirming that *MYB10*, *bHLH33* and *WD40* genes regulate the anthocyanin biosynthesis and that the expression of *HY5*, *PHYA*, *COP1*, *DET* and *PIF3* genes are also part of the color regulatory mechanisms in response to light.

**Table 3.** The influence of bagging on physiological disorders, color and quality of pear fruits.

Pear Cultivar/Species	Bagging Start	Bagging Material	Effect	Ref.
“Meirensu” and “Yunhongli No. 1” ( <i>P. pyrifolia</i> )	20 d after full bloom (removed 1–3 weeks before harvest)	single- or two-layer paper with different levels of light reduction	improvement of anthocyanins accumulation removing bags 2–3 weeks before harvest	[48]
“Housui” ( <i>P. pyrifolia</i> )	34 d and/or 83 d after full bloom	several colored paper combinations or transparent paraffin	improved fruit appearance (uniform, shine and smooth skin color with small lenticels)	[58]
“Carmen” ( <i>P. communis</i> )	66 d before harvest (removed 13 d before harvesting)	paper bags: (1) white; (2) yellow; (3) black; (4) outside grey–inside yellow; (5) outside newspaper–inside yellow	red over-color formation removing bags before harvest, fruits were slightly smaller, improved quality of the skin	[59]
“Conference” ( <i>P. communis</i> )	30 d after full bloom	micro-perforated polyethylene	reduction in skin blemish and russet	[60]
“Cuiguan” ( <i>P. pyrifolia</i> )	20 d (changing the bag at day 45) or 35 d after full bloom	paper	fruits bagged earlier were brighter, with less russet, fewer dots and stone cells	[61]
“Cuiguan” ( <i>P. pyrifolia</i> )	20 d after full bloom	double-layer paper (yellow outside, red inside)	ascorbate decline	[64]
“Mantianhong” ( <i>P. pyrifolia</i> ) and “Cascade” ( <i>P. communis</i> )	20 d after full bloom (removed 10 d before harvest)	double layers of yellow–black paper	red skin coloration in response to light/UV irradiation	[88]
“Kalle” ( <i>P. communis</i> )	20 d after full bloom	white, yellow and double layered black paper	reduced skin color intensity, best performance with white bags	[89]
“Meirensu” ( <i>P. pyrifolia</i> )	40 d after full bloom (removed 10 d before harvest)	double-layered yellow–black paper	anthocyanin accumulation and expression of	[90]

Pear Cultivar/Species	Bagging Start	Bagging Material	Effect	Ref.
			miR156 and its target <i>PpSPL</i> genes,	
27 different cultivars ( <i>P. pyrifolia</i> , <i>P. communis</i> , <i>P. bretschneideri</i> , <i>P. ussuriensis</i> )	40 d after full bloom, harvest 10 d before commercial maturity, then treatment with artificial light	double-layered paper (outer layer yellow outside and black inside, inner layer red)	increasing levels of anthocyanin under artificial light conditions.	[91]
“Chili” ( <i>P. bretschneideri</i> )	77 d after full bloom	polyethylene and non-woven fabric	prevention of scald with non-woven fabric, higher scald with polyethylene	[92]

~~PPSPL1~~. As with apples and pears, peach fruit bagging gives different results depending on the cultivar and the bag material [94]. The naturally deeply colored “Hujingmilu” peach and the light colored “Yulu” developed a deeper color when bagged with white non-woven polypropylene instead of yellow paper because the first type of envelope does not reduce the incoming of UV and blue light. The same authors proposed white non-woven polypropylene as a perfect replacement of yellow paper for peach bagging [94].

Later, Zhao et al. [74], still employing bagging on “Hujingmilu” and “Yulu” peach cultivars, demonstrated that both UVA and UVB induce pigmentation in “Hujingmilu”, while only UVB has an effect on “Yulu” fruits. In addition, Zhao et al. [74] supported the role of the light receptor as COP10 and HYH, and of the transcription factors belonging to gene families *MYB*, *bHLH*, *bZIP* and *NAC*, as summarized above.

The intensity of the color tends to decrease in bagged fruit but, as for apples and pears, unbagging peach fruits ten days before harvest restores a blush comparable to the control [15]. Zhou et al. [95] indicated that shortening the bagging period increases the anthocyanin level in peach peel but reduces peel brightness and chlorophyll content. Additionally, the effects of bagging on carotenoid content were studied in yellow-fleshed peach [65], for which the use of yellow–black double-layered bags reduced the carotenoid level significantly (Table 4).

**Table 4.** The influence of bagging on color and quality of peach and grape fruits.

Crop/Cultivar	Bagging Start	Bagging Material	Effect	Ref.
Peach “Hujingmilu” and “Yulu”	42 days after full bloom	yellow paper	UV-light induction of anthocyanin biosynthesis	[74]
Peach “Janghowon Hwangdo”	after final thinning (early June)	coated white paper, coated yellow paper, white paper, yellow paper and newspaper	improvement in the appearance and in the accumulation of anthocyanins	[10]

Crop/Cultivar	Bagging Start	Bagging Material	Effect	Ref.
Peach "Hakuho"	before pit hardening, and 15 days before harvest	orange paper or orange triple and single parchment paper, 15%, 50%, 80% transmittance	decrease of the color intensity proportionally to the light reduction. Increase in aroma volatile content.	[62]
Peach "3D-8" and "C18"	50 d after full bloom, harvest at 70, 80 and 90 d after full bloom	double-layer paper (yellow outside and black inside)	reduced content in total carotenoids, low quality	[65]
Peach "Hujingmilu" and "Yulu"	96–100 days after full bloom, harvest at commercial maturity or 106–139 days after full bloom	yellow paper, and black, white, blue and grey nonwoven polypropylene bags	non-woven polypropylene bags determined the highest anthocyanin content in peel.	[94]
Peach "Hujingmilu"	50 days after flowering, bags removed at 90 or 105 days	paper single-layer, yellow	a short bagging period improves and stabilizes peel anthocyanin content reducing peel brightness and chlorophyll	[95]
Grape "Cabernet Sauvignon"	3 weeks after full bloom (with different timing) to harvest	two-layer paper (yellow outside, black coated with wax inside), with a bent straw	limited effects on berry quality positive correlation of phenolics to different light regimes	[96]
Grape "Shenhua" and "Shenfeng"	45 days after full bloom	white (light 25%) or shading light bags (light 0%)	incomplete color development, lower content of soluble sugar	[97]
Grape "Italia", "Autumn Royal", and "Regal Seedless"	berries at pea size (bagged at least 90 days)	paper	increased yield for the three cultivars and increased berry hardness for "Autumn Royal", and "Regal Seedless"	[98]
Grape "Muscat Hamburg"	after fruit set	non-woven UV stabilized polypropylene of different colors	improved yield (both in summer and winter season)	[99]
Grape "Kyoho" ( <i>V. vinifera</i> × <i>V. labrusca</i> )	5 weeks after full bloom	white, green, blue and red paper	promotion of accumulation of esters, inhibition of synthesis of aldehydes, alcohols, terpenes, ketones and acids	[100]
Grape "Cabernet Sauvignon" and "Carignan"	from fruit set	fruit bags with a black double-layer inside	promotion of melatonin biosynthesis in berry skins, delayed fruit coloring and ripening	[101]

The bagging treatments have low effects on grape berries because they inhibit anthocyanin accumulation in the skin and do not modify phenolic acid biosynthesis. A significantly elevated flavan-3-ol and flavonol syntheses were observed in re-exposed berries after early stages of bagging [96]. Moreover, bagging allowed for the detection of changes in the expression of *CRY2*, *HY5/HYHs* and *MYBA1* that matched with the biosynthesis of flavonoids in

response to light [96]. A reduction of the color development and lower sugar contents in bagged grape berries was confirmed by Zha et al. [97] in "Shenhua" and "Shenfeng" cultivars, while fruit color and sugar content were rapidly restored by re-exposing the fruits to the light. Quite recently, Pisciotta et al. [98] reported that a bagging treatment is effective both in red and white table grapes; in fact, bagging led to a yield increase for the white varieties "Italia" and "Regal Seedless" and also for the red cultivar "Autumn Royal", whose bunches, although of a slightly lighter skin color, showed increased color uniformity, reduced color variation and improved berry hardness. Furthermore, the bagging with white color non-woven polypropylene of "Muscat Hamburg", which is a black berried grape, suitable both for wine-making and as table grape, determined a higher yield in terms of bunch weight, berry weight and wine yield [99].

Other results of grape bagging were a different production of volatile compounds and melatonin production. Ji et al. [100], investigating the influence of colored paper bags on the production of volatile compounds in "Kyoho" grape berries, indicated that the fruit bagging represents an effective technique to improve the grape aroma. Recently, Guo et al. [101] confirming that grape bagging delayed fruit coloring, sugar content, weight and ripening of the berries, and discovered that bagged berries of both "Cabernet Sauvignon" and "Carignan" cultivars synthesized more melatonin than did unbagged berries, suggesting a new interesting treatment in viticulture ([Table 4](#)).

Additionally, the bagging was recently employed to investigate the red blushed skin formation in apricot and kiwifruit (*Actinidia arguta*). Two blushed and two non-blushed apricot cultivars were compared; blush was not detected on the skin of bagged fruits, while transgenic experiments demonstrated the regulator role of PaMYB10 in apricot anthocyanin biosynthesis [102]. Bagging treatment on kiwifruit demonstrated that also in this fruit, light is necessary for normal skin coloration and that bagging suppression of anthocyanin biosynthesis occurs through inhibition of AaMYB1 expression [103].

Finally, bagging screenings were employed to obtain non-photosensitive eggplants still able to produce an apparently average level of anthocyanins in the peel after bagging (with double-layer paper bags) treatment. These data allowed He et al. [104] to identify 22 transcription factors and 4 transduction elements as putative key regulators of the anthocyanin synthesis in the dark confirming bagging as a tool to study the fruit response to light.

## References

1. Slavin, J.L.; Lloyd, B. Health Benefits of Fruits and Vegetables. *Adv. Nutr.* 2012, 3, 506–516.
2. Sommer, A.; Vyas, K.S. A global clinical view on vitamin A and carotenoids. *Am. J. Clin. Nutr.* 2012, 96, 1204S–1206S.
3. Wang, X.; Ouyang, Y.; Liu, J.; Zhu, M.; Zhao, G.; Bao, W.; Hu, F.B. Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: Systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ* 2014, 349, g4490.

4. Aune, D.; Giovannucci, E.; Boffetta, P.; Fadnes, L.T.; Keum, N.; Norat, T.; Greenwood, D.C.; Riboli, E.; Vatten, L.J.; Tonstad, S. Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality—A systematic review and dose-response meta-analysis of prospective studies. *Int. J. Epidemiol.* 2017, **46**, 1029–1056.
5. Kaur, C.; Kapoor, H.C. Antioxidants in fruits and vegetables—the millennium’s health. *Int. J. Food Sci. Technol.* 2008, **36**, 703–725.
6. Pem, D.; Jeewon, R. Fruit and vegetable intake: Benefits and progress of nutrition education interventions-narrative review article. *Iran. J. Public Health* 2015, **44**, 1309–1321.
7. Food and Agriculture Organization of the United Nations; World Health Organization. Sustainable Healthy Diets—Guiding Principles. Available online: (accessed on 20 December 2020).
8. Nunes, M.C.N. Impact of environmental conditions on fruit and vegetable quality. *Stewart Postharvest Rev.* 2008, **4**, 1–14.
9. Sharma, R.R.; Reddy, S.V.R.; Jhalegar, M.J. Pre-harvest fruit bagging: A useful approach for plant protection and improved post-harvest fruit quality—A review. *J. Hortic. Sci. Biotechnol.* 2014, **89**, 101–113.
10. Kim, Y.H.; Kim, H.H.; Youn, C.K.; Kweon, S.J.; Jung, H.J.; Lee, C.H. Effects of bagging material on fruit coloration and quality of “janghowon hwangdo” peach. *Acta Hortic.* 2008, **772**, 81–86.
11. Frank, D. Evaluation of Fruit Bagging as a Pest Management Option for Direct Pests of Apple. *Insects* 2018, **9**, 178.
12. Jing, L.; Youhong, C.; Zhimei, Y.; Xiaogang, L. Effects of bagging on the quality of pear fruit and pesticide residues. *Acta Hortic.* 2008, **772**, 315–318.
13. Xu, G.; Nie, J.; Wu, Y.; Yan, Z.; Ye, M. The effects of fruit bagging on residue behavior and dietary risk for four pesticides in apple. *Sci. Rep.* 2018, **8**, 14348.
14. Sharma, R.R.; Nagaraja, A.; Goswami, A.K.; Thakre, M.; Kumar, R.; Varghese, E. Influence of on-the-tree fruit bagging on biotic stresses and postharvest quality of rainy-season crop of ‘Allahabad Safeda’ guava (*Psidium guajava* L.). *Crop Prot.* 2020, **135**, 105216.
15. Allran, J.; Schnabel, G.; Melgar, J.C. Peach bagging in the Southeastern US. *J. Am. Pomol. Soc.* 2019, **73**, 38–46.
16. Campbell, D.; Sarkhosh, A.; Brecht, J.K.; Gillett-Kaufman, J.L.; Liburd, O.; Melgar, J.; Treadwell, D. Bagging Organic Peaches Reduces Physical Injuries and Storage Decay with Minimal Effects on Fruit Quality. *HortScience* 2021, **56**, 52–58.
17. Araújo Neto, S.E.; Rocha, C.; de Farias, J.F.; Minosso, S.C.C.; Ferreira, R.L.F. Quality of guava fruits bagged with different materials in an organic system. *Comun. Sci.* 2020, **11**, e3206.

18. Shimizu, K.; Kawana, T.; Ohtani, T.; Mihira, T.; Furukawa, S. Meta-analysis of the damage suppression effect of a double-layer fruit protection bag on fruit bug injuries of loquat fruits and evaluation by explicit solution of odds ratio. *Appl. Entomol. Zool.* 2019, 54, 247–254.

19. Lima, J.D.; Engelking, E.W.; Rozane, D.E.; Gomes, E.N.; da Silva, S.H.M.G.; Kluge, R.A. Effect of bunch protection material and bagging time on the yield of 'Nanica' banana and chilling control. *Aust. J. Crop Sci.* 2020, 14, 574–580.

20. Issarakraisila, M. Effect of types of bagging materials on growth, quality and disease-insect damages in pummelo fruit in tropical humid conditions. *Acta Hortic.* 2018, 1208, 319–324.

21. Sarker, D.; Rahman, M.; Barman, J. Efficacy of different bagging materials for the control of mango fruit fly. *Bangladesh J. Agric. Res.* 2009, 34, 165–168.

22. Watanawan, A.; Watanawan, C.; Jarunate, J. Bagging "Nam Dok Mai #4" mango during development affects color and fruit quality. *Acta Hortic.* 2008, 787, 325–328.

23. Ding, P.; Syakirah, M.N. Influence of fruit bagging on postharvest quality of "Harumanis" mango (*Mangifera indica* L.). *Acta Hortic.* 2010, 877, 169–174.

24. Chonhencob, V.; Kamhangwong, D.; Kruenate, J.; Khongrat, K.; Tangchantra, N.; Wichai, U.; Singh, S.P. Preharvest bagging with wavelength-selective materials enhances development and quality of mango (*Mangifera indica* L.) cv. Nam Dok Mai #4. *J. Sci. Food Agric.* 2011, 91, 664–671.

25. Mathooko, F.M.; Kahangi, E.M.; Runkuab, J.M.; Onyangob, C.A.; Owinob, W.O. Pre-harvest mango (*Mangifera indica* L. 'Apple') fruit bagging controls lenticel discolouration and improves postharvest quality. *Acta Hortic.* 2011, 906, 55–62.

26. Rahman, M.A.; Alam, S.M.K.; Reza, M.H.; Uddin, M.S.; Amin, M.N.; Nasrin, T.A.A. Impact of pre-harvest fruit bagging and improved postharvest practices in reducing losses and managing quality of mango in the value chain system. *Int. J. Postharvest Technol. Innov.* 2019, 6, 117–136.

27. Xu, C.X.; Chen, H.B.; Huang, R.Y.; He, Y.J. Effects of bagging on fruit growth and quality of carambola. *Acta Hortic.* 2008, 773, 195–200.

28. Morera-Montoya, R.; Blanco-Metzler, H.; Luis-Loria, C. Evaluation of different bagging materials for the control of the fruit fly *Anastrepha* sp. (Diptera:Tephritidae) and fruit pathogens in Taiwanese guava fruits (*Psidium guajava* L.). *Acta Hortic.* 2010, 849, 283–292.

29. Hu, G.B.; Chen, D.C.; Li, P.; Ouyang, R.; Gao, F.F.; Wang, H.C.; Dong, J. Effects of bagging on fruit coloration and phenylalanine ammonia lyase and polyphenol oxidase in "Feizixiao" Litchi. *Acta Hortic.* 2001, 558, 273–278.

30. Pal, M.; Lal, R.L.; Nautiyal, P.; Joshi, P. Effect of chemicals and physical means on harvesting span, yield and quality of litchi (*Litchi chinensis* Sonn.) cv. Rose Scented. *J. Appl. Hortic.* 2016,

18, 71–75.

31. Xu, H.X.; Chen, J.W.; Xie, M. Effect of different light transmittance paper bags on fruit quality and antioxidant capacity in loquat. *J. Sci. Food Agric.* 2010, 90, 1783–1788.

32. Ni, Z.; Zhang, Z.; Gao, Z.; Gu, L.; Huang, L. Effects of bagging on sugar metabolism and the activity of sugar metabolism related enzymes during fruit development of Qingzhong loquat. *Afr. J. Biotechnol.* 2011, 10, 4212–4216.

33. Yang, W.H.; Zhu, X.C.; Bu, J.H.; Hu, G.B.; Wang, H.C.; Huang, X.M. Effects of bagging on fruit development and quality in cross-winter off-season longan. *Sci. Hortic.* 2009, 120, 194–200.

34. Fumuro, M.; Gamo, H. Effects of bagging on the occurrence of black stain on the skin of 'Shinsyu' persimmon (*Diospyros kaki* L.) grown under film. *J. Jpn. Soc. Hortic. Sci.* 2001, 70, 261–263.

35. Katagiri, T.; Satoh, Y.; Fukuda, T.; Kataoka, I. Improving marketability of 'Fuyu' persimmon fruit by bagging culture. *Acta Hortic.* 2003, 601, 213–217.

36. Kwan, K.D.; Soo, C.D.; Sik, K.E.; Hyuan, H.K.; Ho, L.K.; Sik, K.K.; Cheol, L.K. Fruit quality of Yuzu (*Citrus junos* L.) as influenced by bagging time and materials of bagging treatment. *J. Korean Soc. Hortic. Sci.* 2000, 41, 190–193.

37. Kassem, H.A.; Omar, A.K.H.; Ahmed, M.A. Response of 'Zaghloul' date palm productivity, ripening and quality to different polyethylene bagging treatments. *Am. Eurasian J. Agric. Environ. Sci.* 2011, 11, 616–621.

38. Al-Obeed, R.S.; Harhash, M.M. Effect of bunch bagging color on "sucary" and "khalas" date palm cultivars: Fruit chemical characteristics. *Acta Hortic.* 2010, 882, 1213–1217.

39. Awad, M.A. Increasing the rate of ripening of date palm fruit (*Phoenix dactylifera* L.) cv. Helali by preharvest and postharvest treatments. *Postharvest Biol. Technol.* 2007, 43, 121–127.

40. Hao, Y.; Zhao, Q.; Liu, Q.; Li, W. Effects of the micro-environment inside fruit bags on the structure of fruit peel in 'Fuji' apple. *Acta Ecol. Sin.* 2011, 31, 2831–2836.

41. Hamedi Sarkomi, F.; Moradinezhad, F.; Khayat, M. Pre-harvest bagging influences sunburn, cracking and quality of pomegranate fruits. *J. Hortic. Postharvest Res.* 2019, 2, 131–142.

42. Chaiwong, S.; Saengrayap, R.; Prahsarn, C. Effects of different materials for banana bunch covers during winter in Thailand. *Acta Hortic.* 2019, 1245, 21–28.

43. Zhang, B.B.; Guo, J.Y.; Ma, R.J.; Cai, Z.X.; Yan, J.; Zhang, C.H. Relationship between the bagging microenvironment and fruit quality in 'guibao' peach [*Prunus persica* (L.) Batsch]. *J. Hortic. Sci. Biotech.* 2015, 90, 303–310.

44. Zhou, J.; Zhong, G.; Lin, Z.; Xu, H. The effects of bagging on fresh fruit quality of *Canarium album*. *J. Food Agric. Environ.* 2012, 10, 505–508.

45. Jing, C.; Feng, D.; Zhao, Z.; Wu, X.; Chen, X. Effect of environmental factors on skin pigmentation and taste in three apple cultivars. *Acta Physiol. Plant.* 2020, 42, 69.

46. Islam, M.T.; Shamsuzzoha, M.; Rahman, M.S.; Bari, M.A.; Akter, M.M.; Khatun, A.; Huque, R.; Uddin, M.S. Effect of bagging time on fruit quality and shelf life of mango (*Mangifera indica* L.) Cv. Langra in Bangladesh. *Int. J. Agric. Environ. Biores.* 2019, 4, 279–289.

47. Bentley, W.J.; Viveros, M. Brown-bagging Granny Smith apples on trees stops codling moth damage. *Calif. Agric.* 1992, 46, 30–32.

48. Huang, C.; Yu, B.; Teng, Y.; Su, J.; Shu, Q.; Cheng, Z.; Zeng, L. Effects of fruit bagging on coloring and related physiology, and qualities of red Chinese sand pears during fruit maturation. *Sci. Hortic.* 2009, 121, 149–158.

49. Sharma, R.R.; Pal, R.K.; Asrey, R.; Sagar, V.R.; Dhiman, M.R.; Rana, M.R. Pre-harvest fruit bagging influences fruit color and quality of apple cv. Delicious. *Agric. Sci.* 2013, 4, 443–448.

50. Asrey, R.; Kumar, K.; Sharma, R.R.; Meena, N.K. Fruit bagging and bag color affects physico-chemical, nutraceutical quality and consumer acceptability of pomegranate (*Punica granatum* L.) arils. *J. Food Sci. Technol.* 2020, 57, 1469–1476.

51. Griñán, I.; Morales, D.; Galindo, A.; Torrecillas, A.; Pérez-López, D.; Moriana, A.; Collado-González, J.; Carbonell-Barrachina, Á.A.; Hernández, F. Effect of preharvest fruit bagging on fruit quality characteristics and incidence of fruit physiopathies in fully irrigated and water stressed pomegranate trees. *J. Sci. Food Agric.* 2019, 99, 1425–1433.

52. Dong, Z.F.; Wang, Y.Z.; Wang, L.; Liu, C.L.; Dong, X.Y.; Liu, G.S.; Yuan, Y.B. Effects of different bag treatments on the absorption of calcium in 'Red Fuji' apple fruit. *Acta Hortic. Sin.* 2007, 34, 835–840.

53. Kim, D.H.; Byun, J.K.; Choi, C.; Choi, D.G.; Kang, I.K. The effect of calcium chloride, prohexadione-Ca, and Ca-coated paper bagging on reduction of bitter pit in 'Gamhong' apple. *Korean J. Hortic. Sci. Technol.* 2008, 26, 367–371.

54. Teixeira, R.; Boff, M.I.C.; Amarante, C.V.T.; Steffens, C.A.; Boff, P. Effects of fruit bagging on pests and diseases control and on quality and maturity of 'Fuji Suprema' apples. *Bragantia* 2011, 70, 688–695.

55. Teixeira, R.; Amarante, C.V.T.; Boff, M.I.C.; Ribeiro, L.G. Control of insect pests and diseases, maturity and quality of 'Imperial Gala' apples submitted to bagging. *Braz. Mag. Fruit Cult.* 2011, 33, 394–401.

56. Rajametov, S.; Nurbekov, A. Effects of different types of paper bags on 'Golden Delicious' apple fruits. *Acta Hortic.* 2020, 1275, 415–418.

57. Matsumoto, K.; Fujita, T.; Sato, S. Effects of 1-MCP and Pre-harvest Fruit Bagging Treatments on Cold Storability of the Red-fleshed Apple 'Kurenainoyume'. *Hortic. J.* 2018, 87, 443–451.

58. Faoro, I.D.; Mondardo, M. Enpacamento de frutos de pereira cv. Housui. *Rev. Bras. Frutic.* 2004, 26, 86–88.

59. Rajametov, S.; Nurbekov, A. Effects of different types of paper bags on 'Carmen' pear fruits. *Acta Hortic.* 2020, 1275, 337–340.

60. Amarante, C.; Banks, N.H.; Max, S. Pre-harvest bagging improves packout and fruit quality of pears (*Pyrus communis*). *N. Z. J. Crop Hortic. Sci.* 2002, 30, 93–98.

61. Lin, J.; Wang, J.H.; Li, X.J.; Chang, Y.H. Effects of bagging twice and room temperature storage on quality of 'Cuiguan' pear fruit. *Acta Hortic.* 2012, 934, 837–840.

62. Jia, H.; Araki, A.; Okamoto, G. Influence of fruit bagging on aroma volatiles and skin coloration of 'Hakuho' peach (*Prunus persica* Batsch). *Postharvest Biol. Technol.* 2005, 35, 61–68.

63. Das, K.; Roy, M.; Roy, D.; Dutta, P. Influence of different sources of potassium on fruit quality and shelf-life of mango cv Himsagar. *Environ. Ecol.* 2017, 35, 1318–1322.

64. Zhang, H.P.; Qin, A.; Qi, K.J.; Tao, S.T.; Wu, J.Y.; Huang, X.S.; Zhang, S.L. The effect of bagging on ascorbate in *Pyrus* fruit. *N. Z. J. Crop Hortic. Sci.* 2019, 47, 19–31.

65. Zhu, M.; Fang, W.; Chen, C.; Wang, L.; Cao, K. Effects of Shading by Bagging on Carotenoid Accumulation in Peach Fruit Flesh. *J. Plant Growth Regul.* 2020.

66. De Wit, M.; Galvão, V.C.; Fankhauser, C. Light-Mediated Hormonal Regulation of Plant Growth and Development. *Annu. Rev. Plant Biol.* 2016, 67, 513–537.

67. Sakuraba, Y.; Yanagisawa, S. Light signalling-induced regulation of nutrient acquisition and utilisation in plants. *Semin. Cell Dev. Biol.* 2018, 83, 123–132.

68. Sanchez, S.E.; Rognone, M.L.; Kay, S.A. Light Perception: A Matter of Time. *Mol. Plant* 2020, 13, 363–385.

69. Teixeira, R.T. Distinct Responses to Light in Plants. *Plants* 2020, 9, 894.

70. Bakhshi, D.; Arakawa, O. Induction of phenolic compounds biosynthesis with light irradiation in the flesh of red and yellow apples. *J. Appl. Hortic.* 2006, 8, 101–104.

71. Awad, M.A.; Wagenmakers, P.S.; de Jager, A. Effects of light on flavonoid and chlorogenic acid levels in the skin of 'Jonagold' apples. *Sci. Hortic.* 2001, 88, 289–298.

72. Kolb, C.A.; Käser, M.A.; Kopecký, J.; Zott, G.; Riederer, M.; Pfündel, E.E. Effects of natural intensities of visible and ultraviolet radiation on epidermal ultraviolet screening and photosynthesis in grape leaves. *Plant Physiol.* 2001, 127, 863–875.

73. Chaves-Silva, S.; Santos, A.L.D.; Chalfun-Júnior, A.; Zhao, J.; Peres, L.E.P.; Benedito, V.A. Understanding the genetic regulation of anthocyanin biosynthesis in plants—Tools for breeding purple varieties of fruits and vegetables. *Phytochemistry* 2018, 153, 11–27.

74. Zhao, Y.; Dong, W.; Wang, K.; Zhang, B.; Allan, A.C.; Lin-Wang, K.; Chen, K.; Xu, C. Differential Sensitivity of Fruit Pigmentation to Ultraviolet Light between Two Peach Cultivars. *Front. Plant Sci.* 2017, 8, 1–15.

75. Gould, K.S. Nature’s Swiss army knife: The diverse protective roles of anthocyanins in leaves. *J. Biomed. Biotechnol.* 2004, 5, 314–320.

76. Xie, D.; Sharma, S.; Paiva, N.; Ferreira, D.; Dixon, R. Role of Anthocyanidin Reductase, Encoded by BANYULS in Plant Flavonoid Biosynthesis. *Science* 2003, 299, 396–399.

77. Jaakola, L.; Maatta, K.; Pirtila, A.M.; Torronen, R.; Karenlampi, S.; Hohtola, A. Expression of Genes Involved in Anthocyanin Biosynthesis in Relation to Anthocyanin, Proanthocyanidin, and Flavonol Levels during Bilberry Fruit Development. *Plant Physiol.* 2002, 130, 729–739.

78. Hohtola, A.; Jaakola, L.; Maatta-Riihinen, K.; Karenlampi, S. Activation of flavonoid biosynthesis by solar radiation in bilberry (*Vaccinium myrtillus* L.) leaves. *Planta* 2004, 218, 721–728.

79. Tattini, M.; Gravano, E.; Pinelli, P.; Mulinacci, N.; Romani, A. Flavonoids accumulate in leaves and glandular trichomes of *Phillyrea latifolia* exposed to excess solar radiation. *New Phytol.* 2000, 148, 69–77.

80. Ravaglia, D.; Espley, R.V.; Henry-Kirk, R.A.; Andreotti, C.; Ziosi, V.; Hellens, R.P.; Costa, G.; Allan, A.C. Transcriptional regulation of flavonoid biosynthesis in nectarine (*Prunus persica*) by a set of R2R3 MYB transcription factors. *BMC Plant Biol.* 2013, 13, 68.

81. Ma, C.; Liang, B.; Chang, B.; Yan, J.; Liu, L.; Wang, Y.; Yang, Y.; Zhao, Z. Transcriptome profiling of anthocyanin biosynthesis in the peel of ‘Granny Smith’ apples (*Malus domestica*) after bag removal. *BMC Genom.* 2019, 20, 353.

82. Zhou, H.; Lin-Wang, K.; Wang, H.; Gu, C.; Dare, A.P.; Espley, R.V.; He, H.; Allan, A.C.; Han, Y. Molecular genetics of blood-fleshed peach reveals activation of anthocyanin biosynthesis by NAC transcription factors. *Plant J.* 2015, 82, 105–121.

83. Volz, R.K.; Oraguzie, N.C.; Whitworth, C.J.; How, N.; Chagné, D.; Carlisle, C.M.; Gardiner, S.E.; Rikkerink, E.H.A.; Lawrence, T. Breeding for Red Flesh Colour in Apple: Progress and Challenges. *Acta Hortic.* 2009, 814, 337–342.

84. Wang, L.; Zhang, X.; Liu, Y.; Shi, X.; Wang, Y.; Zhang, C.; Zhao, Z. The effect of fruit bagging on the color, phenolic compounds and expression of the anthocyanin biosynthetic and regulatory genes on the ‘Granny Smith’ apples. *Eur. Food Res. Technol.* 2013, 237, 875–885.

85. Ma, C.; Jing, C.; Chang, B.; Yan, J.; Liang, B.; Liu, L.; Yang, Y.; Zhao, Z. The effect of promoter methylation on *MdMYB1* expression determines the level of anthocyanin accumulation in skins of two non-red apple cultivars. *BMC Plant Biol.* 2018, 18, 108.

86. Khanal, B.P.; Mekonnen, S.B.; Knoche, M. Shading affects fracture force and fracture strain of apple fruit skins. *Sci. Hortic.* 2020, 274, 109651.

87. Yuri, J.A.; Neira, A.; Fuentes, M.; Razmilic, I.; Lepe, V.; González, M.F. Bagging cv. Fuji, Raku Raku Apple Fruit Affects Their Phenolic Profile and Antioxidant Capacity. *Erwerbs Obstbau* 2020, 62, 221–229.

88. Qian, M.; Zhang, D.; Yue, X.; Wang, S.; Li, X.; Teng, Y. Analysis of different pigmentation patterns in 'Mantianhong' (*Pyrus pyrifolia* Nakai) and 'Cascade' (*Pyrus communis* L.) under bagging treatment and postharvest UV-B/visible irradiation conditions. *Sci. Hortic.* 2013, 151, 75–82.

89. Kim, Y.-K.; Kang, S.-S.; Choi, J.-J.; Park, K.-S.; Won, K.-H.; Lee, H.-C.; Han, T.-H. The effect of several paper bags on fruit skin coloration of red skin European pear 'Kalle'. *Korean J. Hortic. Sci. Technol.* 2014, 32, 10–17.

90. Qian, M.; Ni, J.; Niu, Q.; Bai, S.; Bao, L.; Li, J.; Sun, Y.; Zhang, D.; Teng, Y. Response of miR156-SPL module during the red peel coloration of bagging-treated Chinese sand pear (*Pyrus pyrifolia* Nakai). *Front. Physiol.* 2017, 8, 550.

91. Zhu, Y.-F.; Su, J.; Yao, G.-F.; Liu, H.-N.; Gu, C.; Qin, M.-F.; Bai, B.; Cai, S.-S.; Wang, G.-M.; Wang, R.-Z.; et al. Different light-response patterns of coloration and related gene expression in red pears (*Pyrus* L.). *Sci. Hortic.* 2018, 229, 240–251.

92. Li, Q.; Cheng, C.; Zhang, X.; Wang, C.; Yang, S. Preharvest bagging and postharvest calcium treatment affects superficial scald incidence and calcium nutrition during storage of 'Chili' pear (*Pyrus bretschneideri*) fruit. *Postharvest Biol. Technol.* 2020, 163, 111149.

93. Liu, B.; Wang, L.; Wang, S.; Li, W.; Liu, D.; Guo, X.; Qu, B. Transcriptomic analysis of bagging-treated 'Pingguo' pear shows that MYB4-like1, MYB4-like2, MYB1R1 and WDR involved in anthocyanin biosynthesis are up-regulated in fruit peels in response to light. *Sci. Hortic.* 2019, 244, 428–434.

94. Liu, T.; Song, S.; Yuan, Y.; Wu, D.; Chen, M.; Sun, Q.; Zhang, B.; Xu, C.; Chen, K. Improved peach peel color development by fruit bagging. Enhanced expression of anthocyanin biosynthetic and regulatory genes using white non-woven polypropylene as replacement for yellow paper. *Sci. Hortic.* 2015, 184, 142–148.

95. Zhou, H.; Yu, Z.; Ye, Z. Effect of bagging duration on peach fruit peel color and key protein changes based on iTRAQ quantitation. *Sci. Hortic.* 2019, 246, 217–226.

96. Sun, R.-Z.; Cheng, G.; Li, Q.; Zhu, Y.-R.; Zhang, X.; Wang, Y.; He, Y.-N.; Li, S.-Y.; He, L.; Chen, W.; et al. Comparative physiological, metabolomic, and transcriptomic analyses reveal

developmental stage-dependent effects of cluster bagging on phenolic metabolism in Cabernet Sauvignon grape berries. *BMC Plant Biol.* 2019, **19**, 583.

97. Zha, Q.; Xi, X.J.; He, Y.; Jiang, A.L. Bagging Affecting Sugar and Anthocyanin Metabolism in the Ripening Period of Grape Berries. *Not. Bot. Hortic. Agrobot.* 2019, **47**, 1194–1205.

98. Pisciotta, A.; Planeta, D.; Giacosa, S.; Paissoni, M.A.; Di Lorenzo, R.; Rolle, L. Quality of Grapes Grown Inside Paper Bags in Mediterranean Area. *Agronomy* 2020, **10**, 792.

99. Kiran, A.S.; Kavitha, C.; Soorianathasundaram, K.; Sritharan, N. Impact of Fruit Bagging with Different Coloured Non-woven Polypropylene Bags on Yield Attributes in Grapes. *Asian J. Dairy Food Res.* 2020, **39**, 359–362.

100. Ji, X.-H.; Wang, B.-L.; Wang, X.-D.; Shi, X.-B.; Liu, P.-P.; Liu, F.-Z.; Wang, H.-B. Effects of different color paper bags on aroma development of Kyoho grape berries. *J. Integr. Agric.* 2019, **18**, 70–82.

101. Guo, S.-H.; Xu, T.-F.; Shi, T.-C.; Jin, X.-Q.; Feng, M.-X.; Zhao, X.-H.; Zhang, Z.-W.; Meng, J.-F. Cluster bagging promotes melatonin biosynthesis in the berry skins of *Vitis vinifera* cv. Cabernet Sauvignon and Carignan during development and ripening. *Food Chem.* 2020, **305**, 125502.

102. Xi, W.; Feng, J.; Liu, Y.; Zhang, S.; Zhao, G. The R2R3-MYB transcription factor PaMYB10 is involved in anthocyanin biosynthesis in apricots and determines red blushed skin. *BMC Plant Biol.* 2019, **19**, 287.

103. Li, Y.; Qi, X.; Cui, W.; Lin, M.; Qiao, C.; Zhong, Y.; Fang, J.; Hu, C. Restraint of Bagging on Fruit Skin Coloration in on-Tree Kiwifruit (*Actinidia arguta*). *J. Plant Growth Regul.* 2020.

104. He, Y.; Chen, H.; Zhou, L.; Liu, Y.; Chen, H. Comparative transcription analysis of photosensitive and non-photosensitive eggplants to identify genes involved in dark regulated anthocyanin synthesis. *BMC Genom.* 2019, **20**, 678.

Retrieved from <https://encyclopedia.pub/entry/history/show/19074>