

Phenotyping Based on Sweet Cherries

Subjects: **Horticulture**

Contributor: Viola Devasirvatham , Daniel K. Y. Tan

Australia produces high-quality sweet cherries and generates revenue from local and export markets. Due to increased demand in the markets, the area of sweet cherry production has increased in Australia. Sweet cherry breeding and production have challenges such as self-incompatibility genotypes and phenotyping of agronomic, physiological, and quality traits. Phenotyping is identified as an interaction between cultivars and plant responses to different environmental conditions which determine the plant yield and fruit quality. Efficient phenotyping techniques are important to develop new sweet cherry cultivars in Australia with high yield potential, high fruit quality, temperature tolerance (cold and heat), and drought tolerance.

environmental factors

fruiting spur

fruit development

1. Phenotyping Based on Agronomic and Physiological Traits in Sweet Cherries

Phenotyping is identified as an interaction between cultivars and plant responses to different environmental conditions which determine the plant yield and fruit quality. Efficient phenotyping techniques are important to develop new sweet cherry cultivars in Australia with high yield potential, high fruit quality, temperature tolerance (cold and heat), and drought tolerance.

1.1. Agronomic Traits

The important agronomic traits of sweet cherry are blooming and fruit development and have a direct influence on maturity ^[1]. Fruits from early and late-maturity cultivars attract a higher price on the market ^[2]. For both maturity groups, the periods of bud breaking, blooming, and maturity are important traits in sweet cherry production ^[3]. The positive correlation between blooming and maturity period has been observed in peaches and apricots ^{[4][5]}. Breeders need knowledge of the blooming and fruit development for various sweet cherry germplasm and the impact of cold and high temperatures during that period.

Sweet cherry has three stages of fruit development (double sigmoid growth pattern). In Stage 1, cell division and enlargement of the pericarp occurs. In Stage 2, embryo development occurs, which causes pit hardening. Usually, Stage 2 is shorter in early cultivars than late cultivars and has a high chance for abortion ^[6]. Cold temperature during Stage 2 in early cultivars was the main reason for embryo abortion ^[7]. Day temperature ≤ 16 °C had an adverse effect on embryo development in sweet cherry ^[8]. The pit hardening stage was also susceptible to low temperatures in stone fruits ^[9]. Particularly, sensitivity of temperature in sweet cherry endures the transition period

from Stage 1 to Stage 2 and temperature also had an impact on fruit quality [8]. Fruit development was studied from peak blooming to fruit maturity using cumulative growing degree days (CGDD) [10]. The base temperature for the peak blooming period was 4 °C. For example, the most popular mid-season cultivar, Bing, needs ≥ 800 chilling hours and 683 positive chilling units during 30% bud breaking period in Orange (NSW), Australia. This indicates that the Bing cultivar has a high chilling requirement in Stage 1 [11]. Optimum (24 °C) and critical temperatures (>35 °C) for cherry fruit development were also identified [10]. Duration of Stage 2 was increased due to cold temperatures (4 °C to 15 °C) for late-maturing cultivars [10]. The critical temperature of >35 °C during fruit development for sweet cherry is similar to seed development in grain crops such as wheat [12], chickpea [13], and peanuts [14]. Stage 3 of cherry fruit development consists of continuation of fruit growth and ripening of the exocarp (fruit flesh). It is a critical period to achieve high yield and fruit quality. The optimum temperature during Stage 1 and Stage 2 fruit development has a major influence on Stage 3 fruit development [8]. Hence, temperature is a key factor influencing flowering and fruit development.

Rootstocks regulate the subsequent tree growth, management of the orchard, and crop yield. It is also important that the selection of rootstock has to match the environmental condition of the orchard [15]. In Australia, Mazzard rootstocks are compatible with most sweet cherry cultivars and the Mahaleb rootstock is incompatible with Australian sweet cherry cultivars [15]. The influence of rootstock on the sweet cherry cultivar affects the timing of flowering, the intensity of flowering, fruit quality, and yield [16]. The influence of rootstock on the flower bud sensitivity to frost (−5 °C to −7 °C) was studied in Belgrade. Rootstock affected the flower bud blooming time, flowers per bud, timing of harvest, fruit weight, and yield [16]. Rootstocks also influence the number of flowers and percentage of fruit set. However, certain cultivars had impaired pollen viability and pollen germination under frost [17]. For example, the 'Kordia' cultivar had small pollen grains and poor viability. The cultivar 'Regina' had more frost resistance than 'Kordia' [16]. These cultivars have a great impact on the selection of rootstock in the breeding programme. Therefore, the rootstocks influence the performance of the sweet cherry cultivars and have an effect on tree growth, crop yield, flowering, and pollination.

1.2. Physiological Traits

Australian sweet cherry tree growth stages and physiological process are outlined in **Figure 1**. Each stage is influenced by tree density, genetic material, temperature, and water availability. The key physiological drivers for fruit growth in sweet cherry are photosynthesis and carbon–nitrogen mobilisation of tissues. Different tree planting systems are being used commercially in Australia. Simplified tree structures encourage light penetration and photosynthesis. Therefore, leaf traits are important physiological traits in sweet cherry breeding. The leaf length and width had a positive correlation with fruit weight in sweet cherry [18]. Consequently, the role of photosynthesis and leaf size (length and width) are important in determining fruit size [19][20]. Higher photosynthetic productivity and leaf size could be achieved with higher light interception in the sweet cherry tree [21].




Figure 1. Australian cherry stage of development and physiological process.

On the other hand, sweet cherry trees are deciduous and leaves develop after blooming and this happens throughout fruit development. Net photosynthesis productivity was maintained from fruit development (Stage 2) to fruit maturity (Stage 3) [22]. Thus, maintaining a good tree canopy, including the leaf population, is important to achieve maximum photosynthesis. Different training systems using trellis have been adapted in the sweet cherry orchard to maintain the tree canopy and fruiting units. The knowledge of leaf population including size and fruiting units is vital to all training systems. Generally, leaves supply sugars to the fruits, which encourages them to produce fruiting units and influence fruit quality [23]. The fruiting units of a 4-year-old tree are enhanced by pruning. The pruning encourages the leaf to fruit ratio (crop load) that increases fruit size and quality [24][25]. Strategic pruning increases the source of photo-assimilates (carbohydrates from leaves of spurs from 2nd- and 3rd-year branch and current-year leaves) and nitrogen demand by sink (buds, flowers, fruits) [26][27]. In addition to annual pruning, hand thinning of buds, flowers, and young fruits improves crop load and progresses the photo-assimilate supply among developing fruits [28]. Pruning along with foliar N application on the mature leaves in Stage 3 of fruit development increased the fruit quality [27].

Photosynthesis and its impact on yield and quality indices such as fruit weight (g), fruit sugar content (sweetness), and fruit firmness (firm to very hard) are directly influenced by the rootstock and scion cultivars. Hence, the rootstocks (dwarf and semi-dwarf) influence tree density in the orchard. The sweet cherry cultivars from different maturity groups such as Burlet (early), Summit (mid), and Van (late) growing on five rootstocks (*Prunus avium* L.—vigorous; CAB11E—semi-vigorous; Maxma14—semi-dwarfing; Gisela5—dwarfing; Edabriz—very dwarfing) determined the physiological traits of stem water potential, stomatal conductance and photosynthesis, which directly influenced the fruit weight, firmness and sweetness [29]. Furthermore, the use of inter-stem (grafting a third genetic material) between rootstock and cultivar (scion) with planting system to understand photosynthesis and crop yield was attempted. Trees on inter-stem had higher yields due to the periphery leaves of the tree canopy, which produced higher dry-matter content per unit of leaf surface area and net photosynthetic productivity, which led to bigger fruits and better yield [21]. Therefore, physiological traits of photosynthesis, stem water potential, and stomatal conductance are controlled by rootstock, scion cultivar, and rootstock x scion cultivar interaction, which directly affect fruit quality.

1.3. Impact of Environmental Factors in Agronomic, Physiological and Reproductive Traits

Generally, each stage in fruit development is affected by temperature. Flower bud differentiation was slow when average temperatures reached above 18 °C, which indirectly caused shorter fruit development periods [30]. Temperature has a direct influence on the period of fruit development on Stage 1. Low temperature (<12 °C) combined with rain negatively influenced the pollination and activity of bees and fruit set [31][32]. Generally, honey bees are active pollinators at or above 19 °C and completely cease their activity at <13 °C. Wind speeds at >8 km/h and rain also stop honey bee activity. In Tasmania, cherry growers use bumblebees as a pollinator to achieve effective pollination because bumblebees are active under cold and wet conditions [15]. Five bee species that are active at low temperatures (≤12 °C) combined with rainy days were identified in Turkey [31]. Hence, there are potential alternative pollinators which can be active in adverse climatic conditions to achieve economic returns from the cherry orchard. Simultaneously, high temperatures (>30 °C) during flowering time, particularly the differentiation of pistil primordia (stigma, style, and ovary) created double fruits [32][33]. High temperatures (29 to 35 °C) during the harvest period caused an increase in the number of double fruits [34]. Furthermore, high temperatures accelerate pollen hydration and ovule senescence in sweet cherry [35]. Therefore, extreme temperatures (too cold—>0 °C and too hot—≥35 °C) have a negative impact on reproductive organs in sweet cherry and reduce crop yield by shortening the period of fruit development.

Another factor which influences agronomic and physiological traits in sweet cherry production is water supply. Most Australian cherry production areas have limited water supply and need careful water management in the cherry orchards [15]. Generally, the harvest period for sweet cherries in Australia relies on a warm summer with less rain. In the fruiting period, the tree continually develops leaves and flower buds for the following season (**Figure 1**). It is common for fruit growers to apply water during the harvest period to match summer temperatures. The amount of water supply at this stage depends on crop age, planting density, and climatic conditions of the orchard. For example, the average daily water uptake of the 4 years old trees (early cultivar 'Rita') in a high-density orchard (spacing 4 m × 2 m) during summer (May to July) in Hungary was 23–24 L per tree [36]. Water stress (where tree demand exceeds the supply) at harvest can cause interruption in photosynthesis, leading to changes in biochemical processes in leaves such as leaf sugar content [37]. Regulated deficit irrigation (RDI) 65% of crop evapotranspiration during harvest was monitored by stem water potential rather than fruit water potential. RDI (65% of crop evapotranspiration) during harvest did not affect the current year's fruit yield and quality [38]. Irrigation with 85% of crop evapotranspiration during preharvest and 55% of crop evapotranspiration at postharvest did not affect flowering, fruit set, and fruit growth [34]. However, irrigation during preharvest and postharvest periods which did not meet plant water requirements decreased vegetative growth and leaf area: fruit ratio, which subsequently affects fruit size. Consequently, regulated deficit irrigation which meets plant water requirements in critical periods of flowering and fruit development saved up to 40% irrigation water without decreasing fruit yield [39]. Sweet cherry phytochemicals (total sugars, total organic acids, and anthocyanins) in fruits were higher in irrigated trees (rain water plus 100% evapotranspiration) than in non-irrigated trees (only rain water) [40]. Therefore, adequate chilling hours during blooming, optimum temperatures during fruit development, water supply based on tree demand during flowering, and fruit development encourage good sweet cherry fruit quality and crop yield.

2. Phenotyping Based on Fruit Quality Traits in Sweet Cherries

From the commercial point of view, the key sweet cherry fruit quality traits are fruit size, fruit colour, firmness, and sweetness [24][41][42]. The Australian cherry production guide explains the quality descriptions for fruit size, fruit colour, sweetness, and firmness tester. The fruit size varies from 22 mm to 32 mm (in diameter) (Figure 2). The price of cherries varies with size, along with its firmness, fruit colour, and sweetness. Fruit firmness is an important trait in cherry quality, which is usually tested by fruit firmness tester (or) penetrometer and expressed as g/m². Fruit sweetness is determined by soluble solid concentration using a digital refractometer (% brix). The fruit firmness and sweetness by commercial standards from the Australian cherry production guide is explained in Table 1.



Figure 2. Australian commercial cherry size (24, 26, 28 mm in diameter).

Table 1. Sweet cherry traits of firmness and sweetness in commercial production [15][43].

Firmness	g/m ²	Sweetness	Brix %
Hard	>350	Super sweet	≥21
Very firm	300–350	Very sweet	19–20

Firmness	g/m ²	Sweetness	Brix %
Firm	240–300	Sweet	17–18
Less firm	<240	Moderate sweet	16

Generally, plant growth regulators have been reported to improve fruit quality traits of sweetness, firmness, and flavour in sweet cherries. Many studies showed that the growth of pericarp can be altered using plant growth regulators (e.g., gibberellic acid (GA)), consequently changing carbon partitioning into fruit and fruit quality [44][45]. The rate and time of GA application in cherry is profound. The application of GA during high rainfall produced increased fruit crack because of reduced transpiration for two days from application [46]. Commonly, the application of GA at 15–30 mg/L at the beginning of Stage 3 fruit development in early maturing sweet cherry cultivars improves fruit firmness, sweetness, and fruit colour [47][48]. For late-maturing sweet cherry cultivars, the application of GA at the end of Stage 2 at a low rate (between 10 and 25 ppm) improves fruit firmness [49] to obtain a premium price at the end of the sweet cherry season in local markets as well as for export.

Fruit cracking has been identified as a major consequence of excess amount of water (25 mL above field capacity) from rain or irrigation [46][50]. Larger-sized fruits are also susceptible to fruit cracking [46]. Reasons for sweet cherry fruit cracking include high water content within the plant and rain or precipitation on the fruit. Water content from precipitation on the fruit influences the osmotic potential in the fruit. Water moves through the xylem from the pedicel or stem. Water uptake results in three types of fruit cracking in sweet cherry [51][52]. Fruit cracking happens around the stem end, causing deep cracks inside the fruit and apical cracking at the end of the fruit.

In sweet cherry, calcium content in fruit is reported as a major reason for fruit cracking due to rain [53]. Generally, calcium in the cherry mesocarp or fruit flesh (0.41 to 0.64 mg per fruit in the 'sweetheart' cultivar) increases in the fruit development stage 2 to stage 3. However, the dry mass ratio (2.45 to 1.25 mg Ca per g of dry mass in 'sweetheart' cultivar) decreases from Stage 2 to Stage 3 [54]. This has been reported in many fruits, including apples [55] and kiwifruit [56]. The calcium dry mass ratios were two to threefold higher at the stem end than at the apical (blossom) end [54]. This indicates that the distribution of calcium is a gradient from the stem to the apical end. Furthermore, xylem fluxes from the cherry stem play a key role in importing water to the developing fruits [54]. A similar trend was observed in tomatoes [57]. Additionally, the cracking in the skin or exocarp happened due to tangential stress (stress in horizontal direction) experienced by the fruit skin in cherries [52]. Ultimately, due to stress, the weakest point of skin breaks [58]. In sweet cherry, cracking mostly happened when trees received rain, hail, and wind during the period of three weeks before commercial harvest [52]. Recently, an effort was made to avoid fruit cracking using high tunnels in sweet cherry. However, high tunnels encouraged high temperature (>27 °C) and humidity (96%) inside the tunnel and affected fruit firmness [59]. In Australia, some regions (e.g., Adelaide Hills Orchard) birds can do considerable damage to fruit. Those regions are encouraged to use net covers to protect cherries from birds [15]. In the future, more cover protection studies combined with microclimate management across different production areas and genotypes will add support to sweet cherry production. Hence, quality traits are affected by rain, hail, wind, temperature, and birds [53][59].

References

1. Hardner, C.M.; Hayes, B.J.; Kumar, S.; Vanderzande, S.; Cai, L.; Piaskowski, J.; Quero-Garcia, J.; Campoy, J.A.; Barreneche, T.; Giovannini, D.; et al. Prediction of genetic values of sweet cherry fruit maturity among environments using a 6K SNP array. *Hortic. Res.* 2019, 6, 6.
2. Yue, C.R.; Gallardo, K.; Luby, J.J.; Rihn, A.L.; McFerson, J.R.; McCracken, V.; Oraguzie, N.; Weebadde, C.; Sebolt, A.; Iezzoni, A. An evaluation of U.S. tart and sweet cherry producer's trait prioritization: Evidence from audience surveys. *HortScience* 2014, 49, 931–937.
3. Quero-Garcia, J.; Fodor, A.; Reignier, A.; Capdeville, G.; Joly, J.; Tauzin, Y.; Fouilhaux, L.; Direlwanger, E. QTL detection of important agronomic traits for sweet cherry production. *Acta Hortic.* 2014, 1020, 57–64.
4. Etienne, C.; Rothan, C.; Moing, A.; Plomion, C. Candidate genes and QTLs for sugar and organic acid content in peach. *Theor. Appl. Genet.* 2002, 105, 145–159.
5. Salazar, J.A.; Ruiz, D.; Campoy, J.A.; Tartarini, S.; Dondini, L.; Martinez-Gomez, P. Inheritance of reproductive phenology traits and related QTL identification in apricot. *Tree Genet. Genomes* 2016, 12, 71.
6. Tukey, H.B. Growth of the peach embryo in relation to growth of fruit and season of ripening. *Proc. Amer. Soc. Hort. Sci.* 1933, 30, 208–218.
7. Tukey, H.B. Embryo abortion in early-ripening varieties of *Prunus avium*. *Bot. Gaz.* 1933, 94, 433–468.
8. Braak, J.P. The effect of flowering date and temperature on embryo development in sweet cherry *Prunus avium* L. *Neth. J. Agric. Sci.* 1978, 26, 13–30.
9. Marti, A.F.; Athanson, B.; Koepke, T.; Font-Forcada, C.; Dhingra, A.; Oraguzie, N. Genetic diversity and relatedness of sweet cherry (*Prunus avium* L.) cultivars based on single nucleotide polymorphic markers. *Front. Plant Sci.* 2012, 3, 116.
10. Azarenko, A.N.; Chozinski, A.; Brewer, L.J. Fruit growth curve analysis of seven sweet cherry cultivars. *Acta Hortic.* 2008, 795, 561–565.
11. Brunt, C. *Cherry Cultivar Selection: Chill Hours and Climate Change*; Horticulture Australia Ltd.: North Sydney, Australia, 2012.
12. Prasad, P.V.V.; Djanaguiraman, M. Response of floret fertility and individual grain weight of wheat to high temperature stress: Sensitive stages and thresholds for temperature and duration. *Funct. Plant Biol.* 2014, 41, 1261–1269.
13. Devasirvatham, V.; Gaur, P.M.; Mallikarjuna, N.; Raju, T.N.; Trethowan, R.M.; Tan, D.K.Y. Effect of high temperature on the reproductive development of chickpea genotypes under controlled

environments. *Funct. Plant Biol.* 2012, 39, 1009–1018.

14. Prasad, P.V.V.; Craufurd, P.Q.; Summerfield, R.J. Sensitivity of peanut to timing of heat stress during reproductive development. *Crop Sci.* 1999, 39, 1352–1359.

15. James, P. Australian Cherry Production guide. Cherry Growers of Australia Inc. 2011. Available online: www.cherrygrowers.org.au (accessed on 28 January 2022).

16. Dordevic, B.; Durovic, D.B.; Zec, G.N.; Boskov, D.D. The influence of rootstocks on the sensitivity of flower buds to frost and the main properties of the Carmen sweetcherry cultivar. *J. Agri. Sci.* 2021, 66, 351–358.

17. Dziedzic, E.; Bieniasz, M.; Kowalczyk, B. Morphological and physiological features of sweet cherry floral organ affecting the potential fruit crop in relation to the rootstock. *Sci. Hortic.* 2019, 251, 127–135.

18. Baji, M.E.; Hafida, H.; En-Nahli, S.; Company, R.S.; Kodad, O. Morphological and Pomological Characteristics of Sweet Cherry (*Prunus avium* L.) Grown In-situ under South Mediterranean Climate in Morocco. *Int. J. Fruit Sci.* 2021, 21, 52–65.

19. Rakonjac, V.; Mratinic, E.; Jovkovic, R.; Aksic, F.M. Analysis of morphological variability in wild cherry (*Prunus avium* L.) genetic resources from Central Serbia. *J. Agric. Sci. Tech.* 2014, 16, 151–162.

20. Ganopoulos, I.; Moysiadis, T.A.; Xanthopoulou, M.; Ganopoulou, E.; Avramidou, F.A.; Aravanopoulos, E.; Tani, P.; Madesis, A.; Tsafaris, A.; Kazantzis, K. Diversity of morphophysiological traits in worldwide sweet cherry cultivars of Gene Bank collection using multivariate analysis. *Sci. Hortic.* 2015, 197, 381–391.

21. Bondarenko, P. Physiological basis of sweet cherry productivity depending on rootstocks, interstems and plant density. *Open Agric.* 2019, 6, 267–274.

22. Roper, T.R.; Keller, J.D.; Loescher, W.H.; Rom, C.R. Photosynthesis and carbohydrate partitioning in sweet cherry: Fruiting effects. *Physi. Plan.* 2006, 71, 42–47.

23. Demirsoy, H.; Demirsoy, L. A study on the relationships between some fruit characteristics in cherries. *Fruits* 2004, 59, 219–223.

24. Kappel, F.; Fisher-Fleming, B.; Hogue, E. Fruit characteristics and sensory attributes of an ideal sweet cherry. *HortScience* 1996, 31, 443–446.

25. Ayala, M.; Lang, G.A. ¹³C-Photoassimilate partitioning in sweet cherry (*Prunus avium*) during early spring. *Sci. Agr. Res.* 2015, 42, 191–203.

26. Ayala, M.; Lang, G.A. Examining the influence of different leaf populations in sweet cherry fruit quality. *Acta Hortic.* 2004, 636, 481–488.

27. Ayala, M.; Mora, L.; Torreblanca, J. Effect of pre-bloom pruning on ^{13}C and ^{15}N distribution during early spring in sweet cherry. *HortScience* 2018, 53, 805–809.

28. Macit, I.; Lang, G.A.; Demirsoy, H. Bud management affects fruit wood, growth, and precocity of cherry trees. *Turk. J. Agr. For.* 2017, 41, 42–49.

29. Goncalves, B.; Moutinho-Pereira, J.; Santos, A.; Silva, A.P.; Bacelar, E.; Correia, C.; Rosa, E. Scion-rootstock interaction affects the physiology and fruit quality of sweet cherry. *Tree Physiol.* 2006, 26, 93–104.

30. Li, B.; Xie, Z.; Zhang, A.; Xu, W.; Zhang, C.; Liu, Q.; Liu, C.; Wang, S. Tree growth characteristics and flower bud differentiation of sweet cherry (*Prunus avium*) under different climate conditions in China. *Hortic. Sci.* 2010, 27, 6–13.

31. Guler, Y.; Dikmen, F. Potential bee pollinators of sweet cherry in inclement weather conditions. *J. Ento. Res. Soc.* 2013, 15, 9–19. Available online: <https://www.researchgate.net/publication/286268023> (accessed on 20 May 2022).

32. Predieri, S.; Dris, R.; Sekse, L.; Rapparini, F. Influence of environmental factors and orchard management on yield and quality of sweet cherry. *Food Agri. Envi.* 2003, 1, 263–266. Available online: <https://www.researchgate.net/publication/266226841> (accessed on 20 May 2022).

33. Beppu, K.; Ikeda, T.; Kataoka, I. Effect of high temperature exposure time during flower bud formation on the occurrence of double pistils in ‘Satohnishiki’ sweet cherry. *Sci. Hortic.* 2001, 87, 77–84.

34. Blanco, V.; Blayo-rose, P.J.; Torres-Sánchez, R.; Domingo, R. Influence of regulated deficit irrigation and environmental conditions on reproductive response of sweet cherry trees. *Plants* 2020, 9, 94.

35. Zang, L.; Ferguson, L.; Whiting, M.D. Temperature effect on pistil viability and fruit set in sweet cherry. *Sci. Hortic.* 2018, 241, 8–17.

36. Juhasz, A.; Sepsi, P.; Nagy, Z.; Tokei, L.; Hrotko, K. Water consumption of sweet cherry trees estimated by sap flow measurement. *Sci. Hortic.* 2013, 164, 41–49.

37. Ranney, T.G.; Bassuk, N.L.; Whitlow, T.H. Osmotic adjustment and solute constituents in leaves and roots of water-stressed cherry (*Prunus*) Trees. *J. Am. Soc. Hortic. Sci.* 1991, 116, 684–688.

38. Blanco, V.; Martínez-Hernández, G.B.; Artés-Hernández, F.; Blaya-Ros, P.J.; Torres-Sánchez, R.; Domingo, R. Water relations and quality changes throughout fruit development and shelf life of sweet cherry grown under regulated deficit irrigation. *Agric. Water Manag.* 2019, 217, 243–254.

39. Ruiz-Sánchez, M.C.; Domingo, R.; Castel, J.R. Review. Deficit irrigation in fruit trees and vines in Spain. *Span. J. Agric. Res.* 2010, 8, 5.

40. Vosnjak, M.; Mrzlic, D.; Hudina, M.; Usenik, V. The effect of water supply on sweet cherry phytochemicals in buds, leaf and fruit. *Plants* **2021**, *10*, 1131.

41. Romano, G.S.; Cittadini, E.D.; Pugh, B.; Schouten, R. Sweet cherry quality in the horticultural production chain. *Stewart Post Harv. Rev.* **2006**, *6*, 2.

42. Kazantzis, K.; Chatzicharissis, I.; Papachatzis, A.; Sotiropoulos, T.; Kalorizou, H.; Koutinas, N. Evaluation of sweet cherry cultivars introduced in Greece. *J. Univ. Craiova Ser. Hortic.* **2011**, *16*, 293–296. Available online: <http://pomologyinstitute.gr> (accessed on 28 March 2022).

43. Cherry growers Australia Inc. 2021, Varieties and Management, Cherry Quality. Available online: <http://cherrygrowers.org.au> (accessed on 21 May 2022).

44. Clayton, M.; Biasi, W.V.; Agar, I.T.; Southwick, S.M.; Mitcham, E.I. Sensory quality of Bing sweet cherries following preharvest treatment with hydrogen cyanamide, calcium ammonium nitrate or gibberellic acid. *HortScience* **2006**, *41*, 745–748.

45. Zang, C.; Whitting, M. Plant growth regulators improve sweet cherry fruit quality without reducing endocarp growth. *Sci. Hortic.* **2011**, *150*, 73–79.

46. Kaiser, C.; Long, L.; Brewer, L. Understanding and Preventing Sweet Cherry Cracking. Oregon State University EM 9227. 2019. Available online: <https://www.researchgate.net/publication/331358391> (accessed on 15 February 2022).

47. Looney, N.E.; Lidster, P.D. Some growth regulator effects on fruit quality, mesocarp composition, and susceptibility to postharvest surface marking of sweet cherries. *J. Amer. Soc. Hortic. Sci.* **1980**, *105*, 130–134.

48. Kappel, F.; MacDonald, I.K. Gibberellic acid increases fruit firmness, fruit size, and delays maturity of ‘sweetheart’ sweet cherry. *Isr. J. Plant Sci.* **2002**, *56*, 219–222.

49. Einhorn, T.C.; Wang, Y.; Turner, J. Sweet cherry fruit firmness and postharvest quality of late maturing cultivars are improved with low rate and single application of Gibberellic Acid. *HortScience* **2013**, *48*, 1010–1017.

50. Christensen, J.V. Rain-induced cracking of sweet cherries: Its causes and prevention. In *Cherries: Crop Physiology, Production and Uses*; Webster, A.D., Looney, N.E., Eds.; CAB International: Wallingford, UK, 1996; pp. 297–330. ISBN 9780851989365.

51. Christensen, J.V. Cracking in cherries. III. Determination of cracking susceptibility. *Acta Agric. Scand.* **1972**, *22*, 128–136.

52. Measham, P.F.; Bound, S.A.; Gracie, A.J.; Wilson, S.J. Incidence and type of cracking in sweet cherry (*Prunus avium* L.) are affected by genotype and season. *Crop Pasture Sci.* **2009**, *60*, 1002–1008.

53. Winkler, A.; Knoche, M. Calcium and the physiology of sweet cherries: A review. *Sci. Hortic.* 2019, 245, 107–115.

54. Winkler, A.; Fiedler, B.; Knoche, M. Calcium physiology of sweet cherry fruits. *Trees* 2020, 34, 1157–1167.

55. Amarante, C.V.T.; Miqueloto, A.; Stefens, C.A.; Dos Santos, A.; Argenta, L.C. Changes in xylem functionality during apple fruit development: Implications on calcium concentration and incidence of bitter pit. *Acta Hortic.* 2013, 1012, 135–140.

56. Xiloyannis, C.; Celano, G.; Montanaro, G.; Dichio, B.; Sebastiani, L.; Minnoci, A. Water relations, calcium and potassium concentration in fruits and leaves during annual growth in mature kiwifruit plants. *Acta Hortic.* 2001, 564, 129–134.

57. Windt, C.W.; Gerkema, E.; Van As, H. Most water in the tomato truss is imported through the Xylem, not the Phloem: A nuclear magnetic resonance flow imaging study. *Plant Physiol.* 2009, 151, 830–842.

58. Winkler, A.; Peschel, S.; Kohrs, K.; Knoche, M. Rain cracking in sweet cherry is not due to excess water but localized skin phenomena. *J. Am. Soc. Hortic. Sci.* 2016, 141, 653–660.

59. Blanco, V.; Zoffoli, J.P.; Ayala, M. Influence of high tunnel microclimate on fruit quality and calcium concentration in Santina sweet cherries in a Mediterranean climate. *Agronomy* 2021, 11, 1186.

Retrieved from <https://encyclopedia.pub/entry/history/show/63384>