#### **Small RNA GcvB**

Subjects: Biology | Environmental Sciences Contributor: Conghui You

Small non-translated regulatory RNAs control plenty of bacterial vital activities. The small RNA GcvB has been extensively studied, indicating the multifaceted roles of GcvB beyond amino acid metabolism.

Escherichia coli small RNA GcvB oxidative stress OxyR

#### 1. Introduction

The *Escherichia coli* chromosome encodes more than 80 small non-translated regulatory RNAs <sup>[1]</sup> that control plenty of bacterial vital activities, including biofilm formation <sup>[2]</sup>, transcription termination <sup>[3]</sup>, cell signaling <sup>[4]</sup> and cellular responses to growth conditions of various factors of oxidative stress, osmolarities, temperature or iron levels <sup>[5][6][7][8][9][10][11][12]</sup>.

The small RNA GcvB has been extensively studied since its discovery two decades ago <sup>[5]</sup>. GcvB is approximately 200 nt in length and is highly conserved among Enterobacteriaceae [13][14]. GcvB expression is tightly related to alvcine metabolism <sup>[15][16][17][18][19]</sup> and GcvA activates the transcription of GcvB <sup>[20]</sup> when glycine is abundant. Intriguingly, GcvB expression shows a strong growth condition-dependent feature. GcvB is abundant when cells are grown in rich medium during the exponential phase but is deficient when cells reach the stationary phase or are grown in minimal medium <sup>[5][13][15][21]</sup>. GcvB is depicted as a regulator of 1–2% of all the mRNAs in Salmonella Typhimurium and E. coli by using genome-wide experimental approaches including microarray  $\frac{16}{14}$  and RIL-seg  $\frac{14}{14}$ , or by applying network <sup>[22]</sup> or in silico prediction <sup>[16][17]</sup>. Most of the known targets negatively regulated by GcvB are amino acid biosynthesis proteins and transporters of amino acids or peptides, as exampled by the two asparagine synthetases of AsnA and AsnB<sup>[21]</sup>, the periplasmic transporter components of DppA and OppA<sup>[20]</sup>, the serine/threonine transporter of SstT<sup>[23]</sup>, and a certain amino acids transporter of CycA<sup>[15]</sup>, suggesting the primary physiology role of GcvB could be to limit the uptake and biosynthesis of energy-expensive amino acid under nutrient-rich conditions [13][20]. In parallel, GcvB also negatively regulates the expression of several transcription factors including Lrp <sup>[24]</sup>, a global regulator responding to the intracellular leucine level <sup>[25]</sup>; PhoP <sup>[26]</sup> in the PhoOP two-component system, which controls the expression of genes with functions in magnesium transport, acid resistance and lipopolysaccharide modification <sup>[27][28][29]</sup>; and CsqD <sup>[30]</sup>, the master regulator of curli synthesis <sup>[31]</sup> Moreover, in a direct or an indirect mode, GcvB is positively involved in the acid stress response [32] and stressinduced DNA mutagenic break repair <sup>[33]</sup>. All this evidence indicates the relatively multifaceted roles of GcvB that are beyond amino acid metabolism.

In most cases, GcvB functions as a repressor. It blocks translational initiation by classically interfering with the 30S ribosome subunit binding to a ribosome-binding site or inducing an active mRNA decay via the recruitment of ribonuclease E after pairing to the target mRNA <sup>[21]</sup>. GcvB also works as an activator. GcvB maintains the stability of the *rbn* mRNA with the help of Hfq and protects it from RNase E cleavage <sup>[34]</sup>. GcvB up-regulates sigma factors RpoS <sup>[32]</sup> and RpoE <sup>[33]</sup>, although the precise nature of the interactions between GcvB and these sigma factors is to be determined.

# 2. The Expression of Flu Was Highly Induced in the gcvB Deletion Strain

To characterize the putative target genes controlled by the small RNA GcvB (GenBank acc. n: CP011496.1, region 4,184,713–4,184,917), we compared the transcriptomes of the wild-type *E. coli* K-12 strain NCM3722 and its *gcvB* deletion derivative CY325 grown in glucose minimal medium by high resolution RNA-seq <sup>[35]</sup>. We first found that both the *gcvB* deletion and wild-type strains showed similar doubling times in the glucose minimal medium, indicative of a relatively limited role of GcvB regulation in this condition (Supplementary Figure S1). In total, five genes were retained as significant with a two-fold cutoff (with an FDR smaller than 0.01) (Figure 1A, Supplementary Table S3) in the transcriptome study. Compared to their expression in NCM3722, the genes of *flu* (GenBank acc. n: CP011496.1, region 3,331,422–3,334,541) and *yeeR* (GenBank acc. n: CP011496.1, region 3,334,662–3,336,223) exhibited highly increased expression in CY325, respectively, showing a fold change of 63.6 and 4.6, whereas the remaining three genes showed an approximately two-fold decreased expression in CY325. Since *flu* and *yeeR* showed the biggest fold of expression changes, we further tested their expression in the two strains using the RT-qPCR assay. Consistent with the RNA-seq data, the mRNA levels of *flu* and *yeeR* in CY325 were significantly higher than that in the wild type when cells were grown in glucose minimal medium (Figure 1B,C). Similar results were also observed when cells were grown in LB rich medium (Figure 1D,E). Thus, *flu* and *yeeR* could be negatively regulated by GcvB in a direct or an indirect mode.



**Figure 1.** Genes responding to GcvB. (**A**) Volcano map comparing the expression of genes in the *gcvB* knockout strain CY325 versus that in the wild-type strain NCM3722. The horizontal dashed line points to *q*-value (FDR) = 0.01 on *y*-axis and the vertical dashed lines point to 2-fold cutoff of the expression on *x*-axis. Red dots indicate up-expressed genes; green dots indicate down-expressed genes; grey dots indicate genes showed no significant changes. Note that the data points of the remaining genomic genes carrying a *q*-value (FDR) = 1 with a corresponding  $-\log_{10}$ FDR = 0 were not shown. (**B**,**E**) Gene expression of *flu* and *yeeR* detected using RT-qPCR in the *gcvB* knockout strain CY325 and in the wild-type strain NCM3722 grown in glucose minimal medium (**B**,**C**) or in LB rich medium (**D**,**E**). The mRNA level of each gene in NCM3722 was normalized to 1 and that in CY325 was determined relative to this value. The relative expression was shown as the average ± S.D. of three independent experiments. \*\*\* *p* < 0.001 using Student's *t*-test.

In the genome of NCM3722, the gene of *flu* localized right upstream of the gene of *yeeR* in the same DNA strand. The gene of *yeeR* is a pseudogene and the gene of *flu* encodes antigen 43 (Ag43). Ag43 is an abundant outer membrane protein that belongs to the auto-transporter family and is a major determinant of auto-aggregation in *E. coli* [36][37]. Thus, we studied the auto-aggregation of both the *gcvB* knockout and wild-type strains. As reported, bacterial motility and chemotaxis were required for the Ag43-dependent aggregation of *E. coli* [37]. However, the

two strains applied in the transcriptome study were non-motile <sup>[38]</sup> because the flagellar filament structural protein <sup>[39]</sup> encoded by *fliC* (GenBank acc. n: CP011496.1, region 3,261,993–3,263,489) carries a lone mis-sense mutation (N87K) <sup>[40]</sup>. We complemented the mutated *fliC* of NCM3722 to the wild-type *fliC* to obtain the motile strain CY713, and the gene of *gcvB* was deleted in CY713 to obtain the *gcvB* knockout strain CY1027. Consequently, the auto-aggregation ability of CY713 and CY1027 was then compared using microscopy. Indeed, we observed that the *gcvB* knockout strain CY1027 aggregated more efficiently than the *gcvB* wild-type strain CY713 (**Figure 2**), supporting the highly induced *flu* expression in the *gcvB* knockout strain.



**Figure 2.** Aggregation assays of three *E. coli* strains. The aggregation of CY713 (WT), CY1027 (*gcvB*::cm) and CY1038 (*oxyR*::cm) was detected at 37 °C into a 24-well polystyrene plate using microscopy as described in methods. Bacterial aggregation was monitored at various time points (0–3 h). The formation of the white cluster indicates aggregation of cells. One representative result of two independent observations is shown.

### 3. OxyR Showed Decreased Expression in the gcvB Deletion Strain

As reported, the transcription of *flu* was repressed by OxyR <sup>[41]</sup>. Accordingly, we identified the highly induced expression of *flu* in the *oxyR* (GenBank acc. n: CP011496.1, region 736,030–736,947) deletion strain (<u>Supplementary Figure S2A</u>). In parallel, the *oxyR* deletion strain was observed to exhibit the high auto-aggregation efficiency (**Figure 2**) indicative of the highly expressed *flu*. Moreover, we monitored the expression of *yeeR* in the *oxyR* deleted background and found that *yeeR* showed a similar induced expression as *flu* (<u>Supplementary Figure S2B</u>). Given their congruent responses and their concatenated locations in the genome, it was very likely that *flu* and *yeeR* formed an operon and both were under the control of OxyR.

Thus, the protein levels of OxyR in the *gcvB* knockout and wild-type strains grown in the LB rich medium or the glucose minimal medium were determined using Western blot. We identified that the protein amount of OxyR in CY455 ( $\Delta gcvB$ ) was approximately half that in the *gcvB* wild-type strain CY454 in both conditions (**Figure 3**). As a result, the decreased expression of the OxyR repressor in the *gcvB* knockout strain could induce the high expression of *flu*.



**Figure 3.** The protein levels of OxyR in the *gcvB* knockout and wild-type strains. The protein levels of OxyR in the *gcvB* wild-type strain CY454 and in the *gcvB* knockout strain CY455 cultured in LB rich medium (**A**) or in glucose minimal medium (**B**) were tested using Western blot. RpoB was applied as a loading control. The 0.025  $OD_{600}$  (**A**) or 0.05  $OD_{600}$  (**B**) cells were loaded. One representative result of six repeats in two independent Western blots is shown on the left. The relative protein level of three repeats in one Western blot is shown as the average ± S.D. on the right. The protein level of OxyR in CY455 was normalized to 1 and that in CY454 was determined relative to this value. \*\* *p* < 0.01, \* *p* < 0.05 using Student's *t*-test.

# 4. The Small RNA GcvB Enhanced the Expression of OxyR at the Translational Level

We next explored how GcvB stimulated the expression of OxyR. The mRNA level of *oxyR* did not show significant changes in the two transcriptomes of the *gcvB* wild-type and knockout strains (<u>Supplementary Figure S3A</u>) and this

finding was further demonstrated using the RT-qPCR assay (<u>Supplementary Figure S3B</u>). Moreover, we made an *oxyR* promoter with *lacZ* transcriptional fusion (<u>Supplementary Figure S3C</u>) in both the *gcvB* wild-type and knockout strains and observed that the  $\beta$ -galactosidase activity showed no significant changes in the two backgrounds (<u>Supplementary Figure S3D</u>). As a result, it was most likely that the regulation of GcvB on OxyR existed at the post-transcriptional level. To substantiate this hypothesis, we constructed the *oxyR* promoter with *lacZ* translational fusions in both the *gcvB* wild-type and knockout strains. We made two fusion constructions, with P1 and P2, respectively, carrying 99 and 45 nt after the translational start codon of *oxyR* (**Figure 4**A). Supporting the Western blot result (**Figure 3**), both translational fusions showed significantly decreased  $\beta$ galactosidase activity in the *gcvB* knockout strain when being compared to that in the *gcvB* wild-type strain (**Figure 4**B,C), indicating GcvB activated the expression of OxyR at the translational level.



**Figure 4.** β-galactosidase activities of the *oxyR* promoter with *lacZ* translational fusions in the *gcvB* knockout and wild-type strains. (**A**) Composition diagrams of the promoters of *oxyR* with *lacZ* translational fusions. P<sub>1</sub>*oxyR* covers the DNA region from –266 bp to +99 bp relative to *oxyR* translational start point and P<sub>2</sub>*oxyR* covers the DNA region from –266 bp to +45 bp relative to *oxyR* translational start point. Yellow arrow indicates the entire coding region of *lacZ*. TSS indicates the transcriptional start site of *oxyR*. (**B**) β-galactosidase activities of P<sub>1</sub>*oxyR*-*lacZ* fusion in the *gcvB* wild-type (CY1037) and knockout strains (CY1041). (**C**) β-galactosidase activities of P<sub>2</sub>*oxyR*-*lacZ* fusion in the *gcvB* wild-type (CY1059) and knockout strains (CY1060). The LacZ activity was shown as the average ± S.D. of three independent experiments. \*\* *p* < 0.01, \* *p* < 0.05 using Student's *t*-test.

#### **5. The gcvB Deletion Strain Was Sensitive to Oxidative Stress**

Since OxyR is the transcriptional regulator that induces the expression of antioxidant genes in response to oxidative stress <sup>[42][43]</sup>, the reduced expression of OxyR when *gcvB* was deleted suggested that the *gcvB* deletion strain could be sensitive to oxidative stress. Therefore, we tested the response of the *gcvB* wild-type and knockout strains to oxidative stress. Indeed, when additional  $H_2O_2$  was supplied to the growth medium, the *gcvB* mutant grew more slowly and showed less resistance to  $H_2O_2$  than the *gcvB* wild-type strain (**Figure 5**A). Moreover, we also tested the response of the *oxyR* deletion strain to  $H_2O_2$  treatment. As expected, the *oxyR* deletion strain was sensitive to the oxidative stress. In the same line, both the *gcvB* and the *oxyR* mutants showed significantly lower survival rates than the wild type in the survival assay (**Figure 5**B).



**Figure 5.** The sensitivity of the *gcvB* deletion strain to oxidative stress. (**A**) The growth of strains with  $H_2O_2$  supplied to the medium. The growth curve of each strain was recorded after 3 mM  $H_2O_2$  was supplied into the growth medium. One representative result of three independent experiments is shown. NCM3722 (WT); CY325 ( $\Delta gcvB$ ); CY1057 ( $\Delta oxyR$ ). (**B**) Survival rate of the three strains after a 10-minute challenge of 20 mM  $H_2O_2$ . The

survival rate is shown as the average  $\pm$  S.D. of the six replicates in two independent experiments. \*\*\* p < 0.001 using Student's *t*-test.

The endogenous ROS were generated as byproducts of aerobic respiration <sup>[44]</sup>. Given the decreased expression of OxyR in the *gcvB* mutant, the *gcvB* mutant could accumulate more endogenous ROS. To verify this point, using fluorescence microscopy, we monitored the intracellular amount of ROS that remained in the *gcvB* wild-type and knockout strains by employing two kinds of ROS indicators, DHE (dihydroethidium) and H<sub>2</sub>DCFDA (2',7'-dichlorodihydro-fluorescein diacetate), both of which are membrane-permeant dyes that can be oxidized to show red and green fluorescence, respectively, and have been widely used as fluorescence probes for the detection of superoxide and peroxide due to their specificity to these radicals <sup>[45]</sup>. The phase contrast and fluorescence microphotographs of the two strains stained with DHE and H<sub>2</sub>DCFDA were exhibited (**Figure 6**A and **Figure 7**A). Compared to the *gcvB* wild-type cells, larger numbers of *gcvB* knockout cells showed stronger fluorescence using both indicators indicative of higher levels of ROS remaining in the *gcvB* knockout cells. The mean fluorescence intensity of CY325 cells, respectively, stained with DHE or H<sub>2</sub>DCFDA was approximately two- to six-fold stronger than that of NCM3722 cells (**Figure 6**B and **Figure 7**B). These observations suggested that, compared to the *gcvB* knockout cells, the *gcvB* knockout cells could lose some of their ability to eliminate endogenous ROS where the expression of OxyR was reduced.



**Figure 6.** Detection of superoxide. (**A**) Detection of superoxide by DHE staining. The phase contrast shows the cells under microscope and the red fluorescence represents the detection of superoxide. Two to three independent observations were performed and one image containing representative cells is shown. (**B**) Quantification of fluorescence intensity. Fluorescence intensity was expressed as the average  $\pm$  S.E.M., calculated from 393 to 416 individual cells. NCM3722 (WT); CY325 ( $\Delta gcvB$ ). \*\*\* *p* < 0.001 using Student's *t*-test.



**Figure 7.** Detection of peroxide. (**A**) Detection of peroxide by  $H_2DCFDA$  staining. The phase contrast shows the cells under microscope and the green fluorescence represents the detection of peroxide. Two to three independent observations were performed and one image containing representative cells is shown. (**B**) Quantification of fluorescence intensity. Fluorescence intensity was expressed as the average ± S.E.M., calculated from 468 to 741 individual cells. NCM3722 (WT); CY325 ( $\Delta gcvB$ ). \*\*\* *p* < 0.001 using Student's *t*-test.

#### References

- 1. Moon, K.; Gottesman, S. A PhoQ/P-regulated small RNA regulates sensitivity of Escherichia coli to antimicrobial peptides. Mol. Microbiol. 2009, 74, 1314–1330.
- Parker, A.; Cureoglu, S.; De Lay, N.; Majdalani, N.; Gottesman, S. Alternative pathways for Escherichia coli biofilm formation revealed by sRNA overproduction. Mol. Microbiol. 2017, 105, 309–325.

- Sedlyarova, N.; Shamovsky, I.; Bharati, B.K.; Epshtein, V.; Chen, J.; Gottesman, S.; Schroeder, R.; Nudler, E. sRNA-Mediated Control of Transcription Termination in E. coli. Cell 2016, 167, 111– 121.e13.
- Khan, M.A.; Durica-Mitic, S.; Gopel, Y.; Heermann, R.; Gorke, B. Small RNA-binding protein RapZ mediates cell envelope precursor sensing and signaling in Escherichia coli. EMBO J. 2020, 39, e103848.
- Argaman, L.; Hershberg, R.; Vogel, J.; Bejerano, G.; Wagner, E.G.H.; Margalit, H.; Altuvia, S. Novel small RNA-encoding genes in the intergenic regions of Escherichia coli. Curr. Biol. 2001, 11, 941–950.
- 6. Gottesman, S. The small RNA regulators of Escherichia coli: Roles and mechanisms. Annu. Rev. Microbiol. 2004, 58, 303–328.
- 7. Hershberg, R.; Altuvia, S.; Margalit, H. A survey of small RNA-encoding genes in Escherichia coli. Nucleic Acids Res. 2003, 31, 1813–1820.
- 8. Majdalani, N.; Vanderpool, C.K.; Gottesman, S. Bacterial small RNA regulators. Crit. Rev. Biochem. Mol. Biol. 2005, 40, 93–113.
- 9. Storz, G.; Altuvia, S.; Wassarman, K.M. An abundance of RNA regulators. Annu. Rev. Biochem. 2005, 74, 199–217.
- 10. Vogel, J.; Papenfort, K. Small non-coding RNAs and the bacterial outer membrane. Curr. Opin. Microbiol. 2006, 9, 605–611.
- 11. Wagner, E.G.H.; Flärdh, K. Antisense RNAs everywhere? TRENDS Genet. 2002, 18, 223–226.
- 12. Wassarman, K.M. Small RNAs in bacteria: Diverse regulators of gene expression in response to environmental changes. Cell 2002, 109, 141–144.
- Sharma, C.M.; Darfeuille, F.; Plantinga, T.H.; Vogel, J. A small RNA regulates multiple ABC transporter mRNAs by targeting C/A-rich elements inside and upstream of ribosome-binding sites. Genes Dev. 2007, 21, 2804–2817.
- Melamed, S.; Peer, A.; Faigenbaum-Romm, R.; Gatt, Y.E.; Reiss, N.; Bar, A.; Altuvia, Y.; Argaman, L.; Margalit, H. Global Mapping of Small RNA-Target Interactions in Bacteria. Mol. Cell 2016, 63, 884–897.
- 15. Pulvermacher, S.C.; Stauffer, L.T.; Stauffer, G.V. Role of the sRNA GcvB in regulation of cycA in Escherichia coli. Microbiology 2009, 155, 106–114.
- Sharma, C.M.; Papenfort, K.; Pernitzsch, S.R.; Mollenkopf, H.J.; Hinton, J.C.; Vogel, J. Pervasive post-transcriptional control of genes involved in amino acid metabolism by the Hfq-dependent GcvB small RNA. Mol. Microbiol. 2011, 81, 1144–1165.

- Wright, P.R.; Georg, J.; Mann, M.; Sorescu, D.A.; Richter, A.S.; Lott, S.; Kleinkauf, R.; Hess, W.R.; Backofen, R. CopraRNA and IntaRNA: Predicting small RNA targets, networks and interaction domains. Nucleic Acids Res. 2014, 42, W119–W123.
- 18. Ernst, D.C.; Downs, D.M. 2-Aminoacrylate Stress Induces a Context-Dependent Glycine Requirement in ridA Strains of Salmonella enterica. J. Bacteriol. 2016, 198, 536–543.
- 19. Rossi, C.C.; Bosse, J.T.; Li, Y.; Witney, A.A.; Gould, K.A.; Langford, P.R.; Bazzolli, D.M. A computational strategy for the search of regulatory small RNAs in Actinobacillus pleuropneumoniae. RNA 2016, 22, 1373–1385.
- Urbanowski, M.L.; Stauffer, L.T.; Stauffer, G.V. The gcvB gene encodes a small untranslated RNA involved in expression of the dipeptide and oligopeptide transport systems in Escherichia coli. Mol. Microbiol. 2000, 37, 856–868.
- 21. Lalaouna, D.; Eyraud, A.; Devinck, A.; Prevost, K.; Masse, E. GcvB small RNA uses two distinct seed regions to regulate an extensive targetome. Mol. Microbiol. 2019, 111, 473–486.
- Modi, S.R.; Camacho, D.M.; Kohanski, M.A.; Walker, G.C.; Collins, J.J. Functional characterization of bacterial sRNAs using a network biology approach. Proc. Natl. Acad. Sci. USA 2011, 108, 15522–15527.
- 23. Pulvermacher, S.C.; Stauffer, L.T.; Stauffer, G.V. The small RNA GcvB regulates sstT mRNA expression in Escherichia coli. J. Bacteriol. 2009, 191, 238–248.
- 24. Lee, H.J.; Gottesman, S. sRNA roles in regulating transcriptional regulators: Lrp and SoxS regulation by sRNAs. Nucleic Acids Res. 2016, 44, 6907–6923.
- 25. Tani, T.H.; Khodursky, A.; Blumenthal, R.M.; Brown, P.O.; Matthews, R.G. Adaptation to famine: A family of stationary-phase genes revealed by microarray analysis. Proc. Natl. Acad. Sci. USA 2002, 99, 13471–13476.
- 26. Coornaert, A.; Chiaruttini, C.; Springer, M.; Guillier, M. Post-transcriptional control of the Escherichia coli PhoQ-PhoP two-component system by multiple sRNAs involves a novel pairing region of GcvB. PLoS Genet. 2013, 9, e1003156.
- 27. Kato, A.; Tanabe, H.; Utsumi, R. Molecular characterization of the PhoP-PhoQ two-component system in Escherichia coli K-12: Identification of extracellular Mg2+-responsive promoters. J. Bacteriol. 1999, 181, 5516–5520.
- Minagawa, S.; Ogasawara, H.; Kato, A.; Yamamoto, K.; Eguchi, Y.; Oshima, T.; Mori, H.; Ishihama, A.; Utsumi, R. Identification and molecular characterization of the Mg2+ stimulon of Escherichia coli. J. Bacteriol. 2003, 185, 3696–3702.
- 29. Zwir, I.; Shin, D.; Kato, A.; Nishino, K.; Latifi, T.; Solomon, F.; Hare, J.M.; Huang, H.; Groisman, E.A. Dissecting the PhoP regulatory network of Escherichia coli and Salmonella enterica. Proc.

Natl. Acad. Sci. USA 2005, 102, 2862–2867.

- Jorgensen, M.G.; Nielsen, J.S.; Boysen, A.; Franch, T.; Moller-Jensen, J.; Valentin-Hansen, P. Small regulatory RNAs control the multi-cellular adhesive lifestyle of Escherichia coli. Mol. Microbiol. 2012, 84, 36–50.
- 31. Römling, U.; Bian, Z.; Hammar, M.; Sierralta, W.D.; Normark, S. Curli fibers are highly conserved between Salmonella typhimurium and Escherichia coli with respect to operon structure and regulation. J. Bacteriol. 1998, 180, 722–731.
- 32. Jin, Y.; Watt, R.M.; Danchin, A.; Huang, J.D. Small noncoding RNA GcvB is a novel regulator of acid resistance in Escherichia coli. BMC Genom. 2009, 10, 165.
- Barreto, B.; Rogers, E.; Xia, J.; Frisch, R.L.; Richters, M.; Fitzgerald, D.M.; Rosenberg, S.M. The Small RNA GcvB Promotes Mutagenic Break Repair by Opposing the Membrane Stress Response. J. Bacteriol. 2016, 198, 3296–3308.
- 34. Chen, H.; Previero, A.; Deutscher, M.P. A novel mechanism of ribonuclease regulation: GcvB and Hfq stabilize the mRNA that encodes RNase BN/Z during exponential phase. J. Biol. Chem. 2019, 294, 19997–20008.
- 35. Roberts, A.; Pimentel, H.; Trapnell, C.; Pachter, L. Identification of novel transcripts in annotated genomes using RNA-Seq. Bioinformatics 2011, 27, 2325–2329.
- 36. Van der Woude, M.W.; Henderson, I.R. Regulation and function of Ag43 (flu). Annu. Rev. Microbiol. 2008, 62, 153–169.
- 37. Laganenka, L.; Colin, R.; Sourjik, V. Chemotaxis towards autoinducer 2 mediates autoaggregation in Escherichia coli. Nat. Commun. 2016, 7, 12984.
- 38. Cremer, J.; Honda, T.; Tang, Y.; Wong-Ng, J.; Vergassola, M.; Hwa, T. Chemotaxis as a navigation strategy to boost range expansion. Nature 2019, 575, 658–663.
- 39. Berg, H.C. The rotary motor of bacterial flagella. Annu. Rev. Biochem. 2003, 72, 19–54.
- 40. Lyons, E.; Freeling, M.; Kustu, S.; Inwood, W. Using genomic sequencing for classical genetics in E. coli K12. PLoS ONE 2011, 6, e16717.
- 41. Henderson, I.R.; Owen, P. The major phase-variable outer membrane protein of Escherichia coli structurally resembles the immunoglobulin A1 protease class of exported protein and is regulated by a novel mechanism involving Dam and OxyR. J. Bacteriol. 1999, 181, 2132–2141.
- Anand, A.; Chen, K.; Catoiu, E.; Sastry, A.V.; Olson, C.A.; Sandberg, T.E.; Seif, Y.; Xu, S.; Szubin, R.; Yang, L. OxyR is a convergent target for mutations acquired during adaptation to oxidative stress-prone metabolic states. Mol. Biol. Evol. 2020, 37, 660–667.

- 43. Storz, G.; Tartaglia, L.A.; Ames, B.N. Transcriptional regulator of oxidative stress-inducible genes: Direct activation by oxidation. Science 1990, 248, 189–194.
- 44. Rodríguez-Rojas, A.; Kim, J.J.; Johnston, P.R.; Makarova, O.; Eravci, M.; Weise, C.; Hengge, R.; Rolff, J. Non-lethal exposure to H2O2 boosts bacterial survival and evolvability against oxidative stress. PLoS Genet. 2020, 16, e1008649.
- 45. Marcén, M.; Ruiz, V.; Serrano, M.J.; Condón, S.; Mañas, P. Oxidative stress in E. coli cells upon exposure to heat treatments. Int. J. Food Microbiol. 2017, 241, 198–205.

Retrieved from https://encyclopedia.pub/entry/history/show/39249