

Heme oxygenase-1

Subjects: Genetics

Submitted by:  Stefan Ryter

Definition

Heme oxygenases (E.C. 1:14:99:33) are vital metabolic enzymes that catalyze the rate-limiting step in the degradation of heme, with the generation of carbon monoxide, biliverdin, and iron. The inducible form, heme oxygenase-1 (HO-1), is a stress protein, whose expression is responsive to a broad spectrum of adverse chemical and physical stimuli. HO-1 is known to provide cytoprotection and can exert anti-inflammatory and immunomodulatory effects in tissues, via heme removal. HO-1 is a potential therapeutic target in inflammatory diseases. The end-products of HO-1 activity, including carbon monoxide, may contribute to HO-1 mediated protection. Carbon monoxide delivery by inhalation at low concentration, as well as through application of carbon monoxide releasing molecules (CORMs), has been explored for therapeutic potential. Recently completed clinical trials have evaluated the safety and feasibility of inhaled CO as a therapy for acute and chronic lung disease,

1. Introduction

Heme oxygenase-1, a vital heme degradative enzyme and inducible stress protein Heme oxygenase (HO; E.C. 1:14:99:3) is a metabolic enzyme system first described in the late 1960s which provides the rate-limiting step in the oxidative degradation of heme^{[1][2]}. Heme catalysis by HO requires molecular oxygen (O₂) and NADPH: cytochrome p-450 reductase as the electron source^{[1][2]} (Figure 1). The HO reaction generates several products, each with biological sequelae, including carbon monoxide (CO) (which evolves from the α -methene bridge carbon of heme), the bile pigment biliverdin-IX α (BV), and free ferrous iron, Fe(II). BV is subsequently converted to its lipid soluble metabolite bilirubin-IX α (BR) by cytosolic NADPH: biliverdin reductase (BVR)^[3].

Heme degradative activity is catalyzed by two major isoforms of HO, a constitutive form (heme oxygenase 2, HO-2) and an inducible form (heme oxygenase 1, HO-1), which are products of distinct genes^{[4][5]}. The enzymatic HO substrate, heme, provides prosthetic groups for hemoglobin, the principal O₂ carrier in the blood, and serves as a cofactor for mitochondrial cytochromes and other metabolic enzymes^{[6][7]}. Heme, in free form, may also represent a hazardous pro-oxidant compound due to the presence of the central iron, a potential catalyst of harmful free radical generating processes^[8]. Indeed, heme accumulation has been implicated as a pathological mediator in various inflammatory conditions such as sepsis^{[9][10][11]}, malaria^[12], sickle cell disease^[13], and atherosclerosis^[14]; with its clearance by HO-1 implicated as a protective and anti-oxidative strategy in these and other diseases.

HO-1 is not only recognized as an essential metabolic enzyme, but also as a key intermediate in the stress response and in cellular adaptation to injury^[15]. The upregulation of HO activity, specifically via the transcriptional regulation of HO-1, has been widely associated with cytoprotection and the protective dampening of inflammation in many cellular and preclinical models of injury^{[16][17][18]}.

Under conditions where HO-1 is found to be associated with cytoprotection, a plausible hypothesis is that the individual HO reaction products contribute, individually or collectively, to the implicit cytoprotective properties of HO-1^[19]. This hypothesis does not exclude activity-independent functions of the protein, such as potential intramolecular interactions resulting in the regulation of other proteins^[20]. The iron release from the HO reaction drives the synthesis of ferritin, an iron sequestering macromolecule that may contribute to cytoprotection in vascular endothelium and other cell types^[21]. Both BV, and its metabolite BR, are known to exert natural antioxidant properties in serum and bile^{[22][23][24]}. CO evolution may contribute to the modulation of cellular function resulting in cellular adaptation, and this notion has provided a rationale for the application of pharmacological or gaseous CO to harness its therapeutic potential^{[25][26]}. In contrast, pro-pathogenic and detrimental roles of excess HO-1 activation have also been proposed^{[27][28]}. For example, the cytoprotective properties of HO-1 may confer a survival advantage to tumor cells, and thus contribute to cancer progression^[29]. In neurodegenerative diseases, the excessive release of pro-oxidant iron by HO activity may contribute to neural injury and disease pathogenesis^[30].




Figure 1. Heme oxygenase activity and end-product generation. Heme oxygenase (HO) is a metabolic enzyme that catalyzes the rate-limiting step in the oxidative catabolism of heme to generate the bile pigment biliverdin-IX α (BV) [1][2]. The HO reaction requires molecular oxygen (O₂) and reducing equivalents from NADPH cytochrome p-450 reductase. The HO reaction releases the α -methene bridge carbon of heme as carbon monoxide (CO). The heme iron is released as ferrous iron (Fe II). BV is further metabolized to bilirubin-IX α (BR) by NAD(P)H biliverdin reductase [1][2][3]. Both BV and BR are potent antioxidants [22][23][24]. The metabolic fate of iron released by HO activity remains incompletely understood. Iron may act as a catalyst for the Haber-Weiss cycle, leading to the production of the highly reactive and deleterious hydroxyl radical, and may promote other free radical generating reactions such as the decomposition of organic peroxides [27]. Seminal studies suggested that HO-derived iron can trigger the synthesis of ferritin via modulation of iron binding protein activity [21]. Ferritin can in turn act as a potential cytoprotectant in the endothelium. Endogenously produced HO-1 derived CO has a myriad of possible biological effects, including vasoregulation, as well as cytoprotective effects that include regulation of inflammation, apoptosis, and cell proliferation, and modulation of immune responses [17][18][25][26].

The crucial role of HO-1 in systemic homeostasis, has been uncovered by rare cases of HO-1 deficiency, which have been associated with endothelial cell injury, anemia, abnormal tissue iron accumulation, inflammation, and nephritis [31][32]. HO-1 knockout mice (*Hmox1*^{-/-}) show evidence of hepatic and renal iron deposition, serum anemia, and increased vulnerability to oxidative stress [33]. HO-1 is now recognized as a ubiquitous stress protein whose transcriptional regulation can be induced in cells and tissues by numerous environmental toxins and exogenous effectors [16][34]. These include both the enzyme substrate heme and its downstream catabolic product CO, physical stresses (i.e., heat shock and ultraviolet-A radiation), as well as chemical agents such as heavy metals, thiol (-SH) reactive substances, nitric oxide (NO), natural or synthetic electrophilic antioxidant compounds (e.g., curcumin, resveratrol), high and low oxygen tension (i.e., hyperoxia, hypoxia), pro-inflammatory and pro-oxidant states, pro-inflammatory cytokines, and hormones [18]. The transcriptional regulation of HO-1, while influenced by multiple nuclear factors is primarily governed by nuclear factor erythroid 2-related factor 2 (Nrf2), a master regulator of the cellular antioxidant response, which may act in combination with cell- and inducer-specific accessory transcription factors [35][36].

2. Therapeutic Modulation of HO-1 in Human Disease

The therapeutic modulation of HO-1 for amelioration of human diseases has been proposed by many investigative groups. Metalloporphyrins (i.e., tin and zinc protoporphyrin) which as competitive inhibitors of HO activity, have been extensively studied as an alternative to phototherapy for BR accumulation in neonatal jaundice [145]. More selective HO-1 inhibitors have been described [37][38]. Novel genetic knockdown methods using the CRISPR/Cas9 system have proven effective at inhibiting HO-1 expression in vitro, albeit with noted effects on cell growth, proliferation, and sensitivity to oxidative stress [39].

The prospects of gene therapy approach to drive HO-1 expression are limited by current technology advances in this domain. Since HO-1 generates several reaction products with multiple effector functions, the effects of HO-1 gene

delivery will likely incur pleiotropic and off-target effects. These efforts may benefit from highly targeted cell- or tissue-specific approaches to gene delivery, as recently demonstrated in adipocytes^[40]. Natural antioxidants can be used to target systemic HO-1 expression, and the mechanisms of action of many such natural plant derived substances including polyphenolic antioxidants, has been related in part to enhanced HO-1 expression^{[41][42][43]}. Use of these compounds orally, will likely be non-selective. Synthetic inducers of HO-1 such as cobalt-protoporphyrin may also have pleiotropic effects and unsuitable toxicity profiles for human use/consumption. Dimethylfumarate, a drug developed for multiple sclerosis, activates Nrf2-dependent antioxidant responses, and may have further potential for clinical use^[44]. Among the natural HO-1 enzymatic byproducts, iCO remains closer to therapeutic development, as discussed below.

3. Inhaled Carbon Monoxide (iCO): Summary of Current Clinical Progress

Based on a solid foundation of preclinical studies that were performed over the last two decades in rodents and higher animals (including swine and NHPs), ongoing investigations continue to seek the development of inhaled CO (iCO) as a therapeutic agent for critical care medicine applications and other human diseases. Despite initial promise of preclinical studies, therapeutic development has yet to deliver an immediate breakthrough, due to funding, regulatory hurdles, and public misconceptions despite the demonstration of safety^{[45][46]}. Nevertheless, completed clinical trials to date have demonstrated that therapeutic application of iCO in humans is well-tolerated at low dose, and feasible^{[47][48]}. Early trials of patients with chronic lung disease demonstrated that iCO was safe and did not cause adverse events^[48]. A recent multicenter phase IIa, double-blinded, sham-controlled, clinical trial was completed, in which idiopathic pulmonary fibrosis patients were randomized to receive iCO or medical air for 12 weeks^[49]. The study reported no differences in physiological measures, incidence of acute exacerbations, hospitalization, death, patient-reported outcomes, or secondary endpoints. Furthermore, no differences were reported in the distribution of adverse events. This study concluded that iCO can be safely administered to IPF patients^[49].

A phase I dose escalation trial was recently performed to assess feasibility and safety of low-dose iCO administration in patients with sepsis-induced ARDS. Study subjects ($n = 12$) were randomized to iCO or placebo air 2:1 in two cohorts. In this trial, patients were administered iCO (100–200 ppm) or placebo air for 90 min for five days^[50]. No participants exceeded a CO-Hb level of 10%, and no adverse events were reported. iCO-treated participants (100–200 ppm) displayed significant increases in COHb (up to 5%) compared with placebo-treated subjects. These studies reported that the Coburn–Forster–Kane (CFK) equation was highly accurate at predicting CO-Hb levels in patients exposed to 200 ppm CO. The study concluded that precise administration of low-dose iCO is safe and feasible in patients with sepsis-induced ARDS. Furthermore, circulating mtDNA levels were reduced in the iCO treatment groups. The study provided a precedent for ongoing Phase II studies (NCT03799874) and planned future efficacy trials^[50].

References

1. Tenhunen, R.; Marver, H.S.; Schmid, R. The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc. Natl. Acad. Sci. USA* 1968, 61, 748–755.
2. Tenhunen, R.; Marver, H.S.; Schmid, R. Microsomal heme oxygenase. Characterization of the enzyme. *J. Biol. Chem.* 1969, 244, 6388–6394.
3. Tenhunen, R.; Ross, M.E.; Marver, H.S.; Schmid, R. Reduced nicotinamide-adenine dinucleotide phosphate dependent biliverdin reductase: Partial purification and characterization. *Biochemistry* 1970, 9, 298–303.
4. Maines, M.D.; Trakshel, G.M.; Kutty, R.K. Characterization of two constitutive forms of rat liver microsomal heme oxygenase. Only one molecular species of the enzyme is inducible. *J. Biol. Chem.* 1986, 261, 411–419.
5. Cruse, I.; Maines, M.D. Evidence suggesting that the two forms of heme oxygenase are products of different genes. *J. Biol. Chem.* 1988, 263, 3348–3353.
6. Immenschuh, S.; Vijayan, V.; Janciauskiene, S.; Gueler, F. Heme as a target for therapeutic interventions. *Front. Pharmacol.* 2017, 8, 146.
7. Donegan, R.K.; Moore, C.M.; Hanna, D.A.; Reddi, A.R. Handling heme: The mechanisms underlying the movement of heme within and between cells. *Free Radic. Biol. Med.* 2019, 133, 88–100.
8. Kumar, A.; Ganini, D.; Deterding, L.J.; Ehrenshaft, M.; Chatterjee, S.; Mason, R.P. Immuno-spin trapping of heme-induced protein radicals: Implications for heme oxygenase-1 induction and heme degradation. *Free Radic. Biol. Med.* 2013, 61, 265–272.
9. Larsen, R.; Gozzelino, R.; Jeney, V.; Tokaji, L.; Bozza, F.A.; Japiassú, A.M.; Bonaparte, D.; Cavalcante, M.M.; Chora, A.; Ferreira, A.; Marguti, I.; Cardoso, S.; Sepúlveda, N.; Smith, A.; Soares, M.P. A central role for free heme in the pathogenesis of severe sepsis. *Sci. Transl. Med.* 2010, 2, 51ra71.

10. James, J.; Srivastava, A.; Valuparampil Varghese, M.; Eccles, C.A.; Zemskova, M.; Rafikova, O.; Rafikov R. Heme induces rapid endothelial barrier dysfunction via the MKK3/p38MAPK axis. *Blood J. Am. Soc. Hematol.* 2020, 136, 749–754.
11. Englert, F.A.; Seidel, R.A.; Galler, K.; Gouveia, Z.; Soares, M.P.; Neugebauer, U.; Clemens, M.G.; Sponholz, C.; Heinemann, S.H.; Pohnert, G.; Bauer, M.; Weis, S. Labile heme impairs hepatic microcirculation and promotes hepatic injury. *Arch. Biochem. Biophys.* 2019, 672, 108075.
12. Pereira, M.L.M.; Marinho, C.R.F.; Epiphany, S. Could heme oxygenase-1 be a new target for therapeutic intervention in malaria-associated acute lung injury/acute respiratory distress syndrome? *Front. Cell Infect. Microbiol.* 2018, 8, 161.
13. Gomperts, E.; Belcher, J.D.; Otterbein, L.E.; Coates T.D.; Wood, J.; Skolnick, B.E.; Levy, H.; Vercellotti, G.M. The role of carbon monoxide and heme oxygenase in the prevention of sickle cell disease vaso-occlusive crises. *Am. J. Hematol.* 2017, 92, 569–582.
14. Morita, T. Heme oxygenase and atherosclerosis. *Arterioscler. Thromb. Vasc. Biol.* 2005, 25, 1786–1795.
15. Abraham, N.G.; Kappas, A. Pharmacological and clinical aspects of heme oxygenase *Pharmacol. Rev.* 2008, 60, 79–127.
16. Keyse, S.M.; Applegate, L.A.; Tromvoukis, Y.; Tyrrell, R.M. Oxidant stress leads to transcriptional activation of the human heme oxygenase gene in cultured skin fibroblasts. *Mol. Cell. Biol.* 1990, 10, 4967–4969.
17. Ryter, S.W.; Choi, A.M. Targeting heme oxygenase-1 and carbon monoxide for therapeutic modulation of inflammation. *Transl. Res.* 2016, 167, 7–34.
18. Ryter, S.W.; Alam, J.; Choi, A.M. Heme oxygenase-1/carbon monoxide: From basic science to therapeutic applications. *Physiol. Rev.* 2006, 86, 583–650.
19. Otterbein, L.E.; Soares, M.P.; Yamashita, K.; Bach, F.H. Heme oxygenase-1: Unleashing the protective properties of heme. *Trends Immunol.* 2003, 24, 449–455.
20. Dennerly, P.A. Signaling function of heme oxygenase proteins. *Antioxid. Redox Signal.* 2014, 20, 1743–1753.
21. Vile, G.F.; Tyrrell, R.M. Oxidative stress resulting from ultraviolet A irradiation of human skin fibroblasts leads to a heme oxygenase-dependent increase in ferritin. *J. Biol. Chem.* 1993, 268, 14678–14681.
22. Stocker, R.; Yamamoto, Y.; McDonagh, A.F.; Glazer, A.N.; Ames, B.N. Bilirubin is an antioxidant of possible physiological importance. *Science* 1987, 235, 1043–1046.
23. Neuzil, J.; Stocker, R. Bilirubin attenuates radical-mediated damage to serum albumin. *FEBS Lett.* 1993, 331, 281–284.
24. Stocker, R.; Ames, B.N. Potential role of conjugated bilirubin and copper in the metabolism of lipid peroxides in bile. *Proc. Natl. Acad. Sci. USA* 1987, 84, 8130–8134.
25. Motterlini, R.; Otterbein, L.E. The therapeutic potential of carbon monoxide. *Nat. Rev. Drug Discov.* 2010, 9, 728–743.
26. Ryter, S.W.; Ma, K.C.; Choi, A.M.K. Carbon monoxide in lung cell physiology and disease. *Am. J. Physiol. Cell Physiol.* 2018, 314, C211–C227.
27. Ryter, S.W.; Tyrrell, R.M. The heme synthesis and degradation pathways: Role in oxidant sensitivity. Heme oxygenase has both pro- and antioxidant properties. *Free Radic. Biol. Med.* 2000, 28, 289–309.
28. Waza, A.A.; Hamid, Z.; Ali, S.; Bhat, S.A.; Bhat, M.A. A review on heme oxygenase-1 induction: Is it a necessary evil. *Inflamm. Res.* 2018, 67, 579–588.
29. Chau, L.Y. Heme oxygenase-1: Emerging target of cancer therapy. *J. Biomed. Sci.* 2015, 22, 22.
30. Schipper, H.M.; Song, W.; Tavitian, A.; Cressatti, M. The sinister face of heme oxygenase-1 in brain aging and disease. *Prog. Neurobiol.* 2019, 172, 40–70.
31. Yachie, A.; Niida, Y.; Wada, T.; Igarashi, N.; Kaneda, H.; Toma, T.; Ohta, K.; Kasahara, Y.; Koizumi, S. Oxidative stress causes enhanced endothelial cell injury in human heme oxygenase-1 deficiency. *J. Clin. Invest.* 1999, 103, 129–135.
32. Radhakrishnan, N.; Yadav, S.P.; Sachdeva, A.; Pruthi, P.K.; Sawhney, S.; Piplani, T.; Wada, T.; Yachie, A. Human heme oxygenase-1 deficiency presenting with hemolysis, nephritis, and asplenia. *J. Pediatr. Hematol. Oncol.* 2011, 33, 74–78.
33. Poss, K.D.; Tonegawa, S. Heme oxygenase 1 is required for mammalian iron reutilization. *Proc. Natl. Acad. Sci. USA* 1997, 94, 10919–10924.
34. Keyse, S.M.; Tyrrell, R.M. Heme oxygenase is the major 32-kDa stress protein induced in human skin fibroblasts by UVA radiation, hydrogen peroxide, and sodium arsenite. *Proc. Natl. Acad. Sci. USA* 1989, 86, 99–103.
35. Alam, J.; Stewart, D.; Touchard, C.; Boinapally, S.; Choi, A.M.; Cook, J.L. Nrf2, a Cap'n'Collar transcription factor, regulates induction of the heme oxygenase-1 gene. *J. Biol. Chem.* 1999, 274, 26071–26078.
36. Alam, J.; Igarashi, K.; Immenschuh, S.; Shibahara, S.; Tyrrell, R.M. Regulation of heme oxygenase-1 gene transcription: Recent advances and highlights from the International Conference (Uppsala, 2003) on Heme Oxygenase. *Antioxid. Redox Signal.* 2004, 6, 924–933.
37. Kinobe, R.T.; Dercho, R.A.; Nakatsu, K. Inhibitors of the heme oxygenase—Carbon monoxide system: On the doorstep of the clinic? *Can. J. Physiol. Pharmacol.* 2008, 86, 577–599.
38. Kinobe, R.T.; Vlahakis, J.Z.; Vreman, H.J.; Stevenson, D.K.; Brien, J.F.; Szarek, W.A.; Nakatsu, K. Selectivity of imidazole-dioxolane compounds for in vitro inhibition of microsomal haem oxygenase isoforms. *Br. J. Pharmacol.* 2006, 147, 307–315.
39. Mucha, O.; Podkalicka, P.; Czarnek, M.; Biela, A.; Mieczkowski, M.; Kachamakova-Trojanowska, N.; Stepniewski, J.; Jozkowicz, A.; Dulak, J.; Loboda, A. Pharmacological versus genetic inhibition of heme oxygenase-1—the comparison of metalloporphyrins, shRNA and CRISPR/Cas9 system. *Acta Biochim. Pol.* 2018, 65, 277–286.
40. Singh, S.P.; Greenberg, M.; Glick, Y.; Singh, S.P.; Bellner, L.; Favero, G.; Rezzani, R.; Rodella, L.F.; Agostinucci, K.; Shapiro, J.I.; Abraham, N.G. Adipocyte specific HO-1 gene therapy is effective in antioxidant treatment of insulin resistance and vascular function in

an obese mice model. *Antioxidants* 2020, 9, 40.

41. Son, Y.; Lee, J.H.; Chung, H.T.; Pae, H.O. Therapeutic roles of heme oxygenase-1 in metabolic diseases: Curcumin and resveratrol analogues as possible inducers of heme oxygenase-1. *Oxid. Med. Cell. Longev.* 2013, 2013, 639541.
42. Hammad, A.S.A.; Ahmed, A.F.; Heeba, G.H.; Taye, A. Heme oxygenase-1 contributes to the protective effect of resveratrol against endothelial dysfunction in STZ-induced diabetes in rats. *Life Sci.* 2019, 239, 117065.
43. Funes, S.C.; Rios, M.; Fernández-Fierro, A.; Covián, C.; Bueno, S.M.; Riedel, C.A.; Mackern-Oberti, J.P.; Kalergis, A.M. Naturally derived heme-oxygenase 1 inducers and their therapeutic application to immune-mediated diseases. *Front. Immunol.* 2020, 11, 1467.
44. Belcher, J.D.; Chen, C.; Nguyen, J.; Zhang, P.; Abdulla, F.; Nguyen, P.; Killeen, T.; Xu P.; O'Sullivan, G.; Nath, K.A.; Vercellotti, G.M. Control of oxidative stress and inflammation in sickle cell disease with the Nrf2 activator dimethyl fumarate. *Antioxid. Redox Signal.* 2017, 26, 748–762.
45. Goebel, U.; Wollborn, J. Carbon monoxide in intensive care medicine-time to start the therapeutic application?! *Intensive Care Med. Exp.* 2020, 8, 2.
46. Hopper, C.P.; Meinel, L.; Steiger, C.; Otterbein, L.E. Where is the clinical breakthrough of heme oxygenase-1 / carbon monoxide therapeutics? *Curr. Pharm. Des.* 2018, 24, 2264–2282.
47. Mayr, F.B.; Spiel, A.; Leitner, J.; Marsik, C.; Germann, P.; Ullrich, R.; Wagner, O.; Jilma, B. Effects of carbon monoxide inhalation during experimental endotoxemia in humans. *Am. J. Respir. Crit. Care Med.* 2006, 171, 354–360.
48. Bathorn, E.; Slebos, D. J.; Postma, D. S.; Koeter, G. H.; van Oosterhout, A. J.; van der Toorn, M.; Boezen, H. M.; Kerstjens, H.A. Anti-inflammatory effects of inhaled carbon monoxide in patients with COPD: A pilot study. *Eur. Respir. J.* 2007, 30, 1131–1137.
49. Rosas, I.O.; Rosas, I. O., Goldberg, H. J., Collard, H. R., El-Chemaly, S., Flaherty, K., Hunninghake, G. M., Lasky, J. A., Lederer, D. J., Machado, R., Martinez, F. J., Maurer, R., Teller, D., Noth, I., Peters, E., Raghu, G., Garcia, J., Choi, A.M. A Phase II clinical trial of low-dose inhaled carbon monoxide in idiopathic pulmonary fibrosis. *Chest* 2018, 153, 94–104.
50. Fredenburgh, L. E.; Perrella, M. A.; Barragan-Bradford, D.; Hess, D. R.; Peters, E.; Welty-Wolf, K. E.; Kraft, B. D.; Harris, R. S.; Maurer, R.; Nakahira, K.; Oromendia, C.; Davies, J. D.; Higuera, A.; Schiffer, K. T.; Englert, J. A.; Dieffenbach, P. B.; Berlin, D. A.; Lagambina, S.; Bouthot, M.; Sullivan, A. I.; Nuccio, P.F.; Kone, M.T.; Malik, M.J.; Porras, M.A.P.; Finkelsztejn, E.; Winkler, T.; Hurwitz, S.; Serhan, C.N.; Piantadosi, C.A.; Baron, R.M.; Thompson, B.T.; Choi, A.M. A phase I trial of low-dose inhaled carbon monoxide in sepsis-induced ARDS. *JCI Insight.* 2018, 3, e124039.

Keywords

Carbon monoxide;Heme oxygenase-1;Inflammation;Heme;Ischemia/reperfusion;lung disease