

# Recovery of Bioactive from Food Waste

Subjects: [Chemistry](#), [Analytical](#)

Contributor: Irene Dini

Biowastes, especially food wastes, contain bioactive compounds that are suitable for producing functional foods, supplements, and nutricosmetics. Vegetables and fruits have primary metabolites (e.g., amino acids, lipids, dietary fibers, cellulose, hemicellulose, lignin, and fatty acids), and secondary metabolites (e.g., flavonoids, phenols, alkaloids, glucosinolates, carotenoids, and terpenes). The extraction of bioactive compounds from biowastes depends on the source, functionality, chemical properties, and end-use. Various temperatures, pH values, electromagnetic waves, and extraction techniques are used (e.g., supercritical fluid, subcritical water, ultrasonic wave, microwave, and pulsed electric field).

[food waste](#)

[recycling](#)

[Food chemistry](#)

[nutraceutical](#)

[nutraceutical](#)

## 1. Primary Metabolites Recovered from Agri-Food

### 1.1. Proteins

Proteins are macronutrients with a central role in human nutrition. They are formed by amino acid units. Proteins from food waste sources are used as value-added ingredients and/or products, including human foods and animal feed. The high-value products are used as foaming, thickeners, and gel stabilizers [\[1\]](#), and the low-value products are used as fish and animal feed [\[2\]](#). Food waste protein sources can be classified into animal and plant sources. Plant byproducts used as protein sources include oat, rice, wheat bran protein [\[3\]](#)[\[4\]](#)[\[5\]](#), and defatted meals from the oil industry. Wheat bran contains between 13% to 18% of proteins [\[3\]](#), the defatted meals obtained from the oil industry (e.g., canola, sunflower, palm, rapeseed, and peanuts) have between 15% to 50%, and soybean curd residue contain 27% protein [\[6\]](#). Sugar beet and mushroom flakes are used as a feed ingredient source since they contain 40% essential amino acids [\[7\]](#). Finally, food waste proteins obtained by animals (e.g., meat, fishmeal, bone meal, yogurt, and cheese) are considered good-quality protein sources that are of high biological value [\[8\]](#). Some extraction methods were used to isolate protein, including enzyme-assisted, cavitation-assisted, ultrasound-assisted, hydrodynamic cavitation, microwave-assisted, supercritical, liquid biphasic flotation, and hybrid extractions [\[9\]](#). In enzyme-assisted extraction, the protein recovery depends on the enzyme ratio, substrate characteristics, extraction time, and pH [\[10\]](#). Protein isolates were generally obtained by defatted pressed legume cakes and animal sources via precipitation at the isoelectric point [\[11\]](#). Hydrolysate from protein isolates is also used [\[12\]](#)[\[13\]](#)[\[14\]](#) since it produces higher solubility products and smaller peptides [\[14\]](#)[\[15\]](#). Cavitation-assisted extraction is used in large-scale protein extraction. Low frequency (20 to 100 kHz), temperature, sonication power, and treatment time affect the protein yield [\[16\]](#). Ultrasound-assisted extraction is coupled with enzyme-assisted or microwave-assisted extraction technologies to improve protein extraction efficiency [\[12\]](#). Microwave-assisted

extraction of proteins can depend on nonuniform temperature distribution and closed- or open-type vessel systems [17][18]. It enhances the proteins' functional properties (e.g., water absorption, emulsifying, foam activity, and foam stability indexes) [12]. Supercritical extraction of proteins depends on temperature [19] and solvent concentration [20]. Chemical dehydration and/or evaporation are required to remove moisture. These procedures can affect protein purity [12]. Liquid biphasic flotation has high separation efficiency and determines the minimal protein loss [21][22]. Cell receptors, drug residues in food, and wastewater treatments were extracted using this technology [23].

## Possible Uses of the Recovered Proteins

The food waste proteins can be utilized in feed supplements to enhance the food products' functional properties [24]. Milk protein and whey protein are used to enrich ice cream [25], improve the mixture's viscosity, and decelerate the melting time [26]. The animal proteins can be used as a foaming agent with recycled PET aggregates to produce cementitious concrete composites [27]. Whey protein can be employed to produce plastic films for food packaging materials [28].

## 1.2. Pectins

Pectins are polysaccharides that are formed by d-galacturonic acid, d-galactose, or l-arabinose units, and are found in the cell walls of plant tissue [29]. The degree of pectin esterification affects the pectins' functional properties as a thickening and gelling agent. Conventional (e.g., extraction with the mineral acids) and innovative techniques (e.g., ultrasound- or enzyme-assisted microwave- extraction) were used to extract them from biowaste. Traditionally, pectin is extracted via continuous stirring with water that is acidified (e.g., in nitric, 0.05–2M sulfuric, phosphoric, hydrochloric, or acetic acid) for 1 h under controlled temperature (80 and 100 °C) [30]. The maximum pectin yield is obtained using hydrochloric acid at pH 2.0 [31]. Innovative extraction methods help in the extraction of pectins, disrupting the cell membrane's structure by electromagnetic or sound waves and facilitating the contact between solvent and bioactive molecules. Among the most innovative approaches, ultrasound-assisted technology improves (+20%) the pectins' molecular weight and extraction yield compared to the traditional method under the same temperature, pH, and time conditions [32]. The microwave-assisted extraction of pectins is affected by the weight of the biomaterial, the power of the wave, the time of extraction, and the pH. For example, the optimum processing conditions to extract pectins from lime bagasse are a sample weight of 6 g, a wave power of 400 W, a time of extraction of 500 s, and a pH of 1 [33]. Finally, enzymes can enhance the extraction process by hydrolyzing the plant cell wall matrix (enzyme-assisted extraction). The enzymes used to extract pectins are protease, cellulase, alcalase, hemicellulase, xylase,  $\alpha$ -amylase, polygalacturonase,  $\beta$ -glucosidase, endopolygalacturonase, neurase, and pectinesterase [34].

## Possible Uses of the Recovered Pectins

The food industry employs pectins as emulsifiers, stabilizers, thickeners, and gelling agents.

The pharmaceutical industry uses them as drug-controlled release matrices and prebiotic, hypoglycemic, hypocholesterolemic, and metal-binding agents [35].

Finally, the functionalization of pectins with nanomaterials and phenolics can produce active packaging films with antimicrobial properties [36].

### 1.3. Omega-3 from Fish Waste

Omega-3 fatty acids (e.g., eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)) have the first double bond on carbon 3, counting from the terminal carbon. Fish are a good source of omega-3. They accumulate them from plankton, algae, and prey fish [37]. The omega-3 fatty acids regulate cell membranes' architecture and permeability, produce energy and eicosanoids, and modulate the human body's pulmonary, cardiovascular, immune, reproductive, and endocrine systems [37]. Their potential health benefits include the prevention of cancer, cardiovascular disease (CVD), Alzheimer's disease, depression, rheumatoid arthritis, attention deficit hyperactivity disorder (ADHD), dry eyes, and macular degeneration [38]. Numerous apparatuses and techniques were proposed to extract omega-3 fatty acids from fishes. Traditional extraction techniques use organic solvents (e.g., hexane, methanol, petroleum ether, and chloroform), which cannot be employed on an industry scale [39][40]. Soxhlet extractor, ultrasounds, or microwave-assisted extractions decrease the time and use of solvents [41]. On an industrial scale, fish oil extraction is achieved through a wet-reduction or wet-rendering process [42]. Supercritical fluid extraction (SFC) [43] solves the problem of n-hexane use for extraction in traditional extraction methods, uses low temperature to reduce the oxidation of polyunsaturated fatty acids, decreases residual solvent contaminants (polychlorinated biphenyls and heavy metals), does not modify the biomass, and allows other bioactive molecules to recover. The ethanol used as a co-solvent is much more food-compatible than hexane [44]. The disadvantage of this method is the extract's smell due to volatile compounds. Encapsulation decreases this effect and improves the extracts' palatability [45].

#### Possible Uses of the Recovered Omega-3

Omega-3 fatty acids are employed in supplements and fortified food and feed [46]. Moreover, they can be used to ameliorate cutaneous abnormalities and maintain skin homeostasis in cosmetics [47].

## 2. Secondary Metabolites Recovered from Agri-Food

### 2.1. Phenolic Compounds from Biowaste

Phenolic compounds (PC) are secondary metabolites characterized by an aromatic ring with more hydroxyl substituents. Plants produce them in response to environmental stimuli. They vary greatly by type and level based on target species and biotic and abiotic factors. Biocontrol agents, such as *Trichoderma* strains, can interfere with phenolic production in plants [48][49][50][51][52]. Phenolic compounds have antioxidant and antimicrobial properties [53] and beneficial potential for human health, such as antitumor, anti-obesity, anti-inflammatory, and ultraviolet (UV) radiation protective activities. Isolated and dosed phenolic compounds [54] were used to prepare food supplements [55][56], nutricosmetic formulations [57][58][59], and nutraceutical food [60][61]. The process used to recover PC from biowaste consists of a pre-treatment (thermal treatment of the sample followed by electro-osmotic dewatering,

foam mat, pulverization, or micro-filtration) [62], extraction (obtained by maceration, or assisted by microwave, ultrasound, supercritical fluid, pressurized liquid, pulsed electric field, high voltage electrical discharges, or multi-technique) [63], concentration (obtained by distillation, or steam distillation), and purification (obtained by microfiltrations, ultrafiltration, resins, or chromatography) [63].

## Anthocyanins from Agri-Food Waste

Anthocyanins are water-soluble antioxidants pigments that belong to the flavonoid family. They have a C6-C3-C6 skeleton [64]. The flavylium cation gives the red color to the anthocyanins [65]. Temperature, enzymes, pH, light, metallic ions, oxygen, sulfites, and interaction with flavonoids, phenolics, ascorbic acid, and sugars can affect anthocyanins' integrity according to their chemical structure and food concentration. Anthocyanins subjected to high temperatures and prolonged heating oxidize. Moreover, the deglycosylation process, nucleophilic attack of water, cleavage, and polymerization determine the structure breakdown. These structural changes decrease anthocyanin content in the final product [64]. Sulfite adds colorless adducts interrupting the conjugated  $\pi$ -electron system [65]. Anthocyanin association reactions (e.g., self-association between anthocyanins via hydrophobic interactions, co-pigmentation between anthocyanins and other phenols through van der Waals interactions between the planar polarizable nuclei or binding the sugar and the flavylium nucleus covalently, and metal complexing between anthocyanins and with metals, via their *o*-hydroxy groups) defend the flavylium chromophore from the water's nucleophilic bond, enhancing their stability and color [66]. The anthocyanins can manage and/or prevent chronic degenerative diseases, including cancers, cardiovascular diseases, type 2 diabetes mellitus, dyslipidemias, and neurodegenerative diseases [67]. Several biochemical parameters involved in the inflammatory responses (e.g., interleukins (ILs), tumor necrosis factor-alpha (TNF- $\alpha$ ), nuclear factor-kappa B (NF- $\kappa$ B), and cyclooxygenase 2), which are able to improve malonaldehyde and reactive oxygen species, and to reduce the activity/expression of antioxidant enzymes (e.g., catalase and superoxide dismutase), have been related to the prevention or development of these diseases [68][69][70]. The anthocyanins improve the sensitivity, secretion, and lipid profile of insulin by inhibiting the limiting enzyme of cholesterol synthesis (3-hydroxy-3-methylglutaryl-coenzyme A) or adipose triglyceride lipase engaged in triglyceride breakdown in diabetics and prediabetic subjects [71][72]. Moreover, anthocyanins regulate the expression of adipokines, thereby enabling the avoidance of insulin resistance and the progression of type 2 diabetes mellitus [73]. Anthocyanins can reduce triglycerides, cholesterol, LDL-cholesterol, and inflammatory biomarkers regulating the expression/activity of pro-inflammatory cytokines (e.g., IL-6, TNF- $\alpha$ , and IL-1A), pro-inflammatory enzymes (e.g., COX-2), and NF- $\kappa$ B signaling pathways, decreasing the production of pro-inflammatory molecules (e.g., C-reactive protein), and improving SOD and proliferator-activated receptor- $\gamma$  expression [74][75]. Finally, anthocyanin supplementation can enhance cardiovascular function [76][77], regulate gut microbiota composition [78][79], improve exercise recovery effectiveness [80], decrease ulcerative colitis symptoms [81], reduce ocular fatigue [82], and promote healthy facial skin conditions [80]. Some innovative processes are proposed for the extraction of anthocyanin from biowastes, such as the extractions assisted by a pulsed electric field, microwave, and ultrasound technologies. The enhancement of the extraction of anthocyanin, helped by pulsed electric field technology, is related to permanent (irreversible) or temporary (reversible) pores in the cell membranes, which facilitate the anthocyanins release into the medium [64]. The target compounds' extraction does not constantly improve with the increasing of the electric field strength,

specific energy input, treatment time, temperature, and pulse number [83][84]. For example, studies on blueberry extraction showed an improvement (+75%) in the extraction of anthocyanin when the specific energy input was increased [85]. Instead, studies showed that improving electric field intensity and specific energy input does not increase the anthocyanins content extracted from the sweet cherry byproduct [86]. The application of high intensity can lead to anthocyanin degradation. instead, the application of low/moderate-intensity enhanced anthocyanin recovery without anthocyanin's degradation/modification. The pulsed electric field technology allows the selective extraction of the single anthocyanin classes [87]. The pulsed electric field technology improves the monoglucoside anthocyanin extraction compared to acylated glucoside anthocyanins from grape pomace [88] and the extraction of cyanidin, delphinidin, and petunidin glycosides from blueberry byproducts [89]. The anthocyanin recovery from grape pomace enhances when the microwave and irradiation time improve [90], and longer irradiation times enhance anthocyanin recovery from wine lees [90], sour cherry pomace [91], and saffron floral bio-residues [92]. Nevertheless, the excessive intensification of MW extraction process parameters can decrease the extraction of anthocyanin from biowaste due to their degradation (anthocyanins are thermolabile compounds) [93]. Thus, it is recommended to use extraction temperatures below 60 °C to minimize the anthocyanin's losses. The extraction time also impacts MW extraction. The excess time causes the degradation of anthocyanin due to higher exposure to microwave powers and high temperatures [94]. The high microwave power determines internal overheating, leading to carbonization and isomerization, and/or degradation of molecules [95]. According to some authors, anthocyanins' thermal degradation determines the loss of sugar moieties, the formation of a carbinol pseudo base, and chalcone by hydrolysis of the remaining sugar moiety and cut between C2 and C3 [84]. According to others, the degradation of anthocyanin is due to decomposition reactions of water molecules and the production of reactive oxygen species [96]. Studies on anthocyanin recovery from eggplant peel and fig peel showed that the recovery of anthocyanin decreased when the microwave powers and irradiation times improved [97]. Studies on grape pomace [87], blackcurrant bagasse [98], blueberry peel [99], black rice bran [100], and corn husk [101] showed that the extraction of anthocyanin decreased when the irradiation times were long. Finally, anthocyanins' structures impact their recovery from bio matrices. For example, anthocyanin analogs that are unsubstituted at C3 of the C-ring are more stable to MW treatment than other anthocyanins [102], as well as acylated anthocyanins than non-acylated ones [103]. Another technique used to improve the extraction of anthocyanin is ultrasound. Acoustic cavitation can determine the thermal and chemical degradations of the anthocyanins since the acoustic cavitation phenomenon can determine thermal stress and free radical formation [104]. Long processing times can cause severe degradations in anthocyanins [105]. For example, the anthocyanin extracted from black chokeberry wastes degrades when an ultrasound water bath (30.8 kHz) for 60 min at 70 °C, with a nominal ultrasound power of 100 W, and 50% ethanol in water are used [106]. Using enzymes (pectinase compound and pectinase) combined with ultrasound can improve the extraction technique's performance [105].

## Possible Uses of the Recovered Phenolics

Phenolics are used as functional food additives. Their antimicrobial and antioxidant activities enhance the shelf-life of foodstuffs [107]. Anthocyanins can be employed as a coloring additive (EFSA code E163).

Phenolic compounds can be employed as supplement ingredients [108], pharmaceutical [107], and cosmeceutical agents [59].

Caffeic and gallic acids can be used in chitosan-based biofilms to inhibit the growth of *Bacillus subtilis* and *Staphylococcus aureus* and enhance the film's oxygen and vapor permeability [109].

Tannins could develop protein-based biofilms since they can interact with proteins through non-covalent bonds and hydrogen bonding [110].

Finally, the phenolics might be helpful to the textile industry as natural dyes with antimicrobial properties [110].

## 2.2. Carotenoids from Agri-Food Waste

Carotenoids are fat-soluble pigments with excellent antioxidant activity (e.g., singlet oxygen-quenching capacity and free radical activity) [111][112]. The two main classes of carotenoids are xanthophylls (yellow color) that contain oxygen and carotenes (orange color) that consist of linear hydrocarbons, which can cyclize at both ends of the molecule [59]. Carotene supplementation is related to the inhibition of atherosclerosis-related multiple sclerosis [113], cardiovascular diseases [114], macular degeneration [115], and degenerative diseases [116]. Carotenes are most bioavailable in their natural trans-form [117][118]. Light, metals, heat, and pro-oxidants can isomerize the trans-form into cis-form [119]. The extraction of carotenoids is achieved using organic solvents (e.g., hexane, methanol, acetone, and ethanol or solvent combinations). The supercritical fluid extraction with CO<sub>2</sub> improves the extraction's efficiency, as is also the case for the extraction of carotenoids from carrot peels (+86.1% at 349 bar) [120]. In addition, the microwave [121] and ultrasound increase the carotenoids' recovery [122]. Finally, an extraction method of lycopene and pectin from tomato pomace [123] and pink guava decanter [124], which relied on a simple water-induced hydrocolloidal complexation, was tested. The pH, temperature, solid loading, and stirring affected the complexation of carotenoids and pectin [125].

### Possible Use of the Recovered Carotenoids

Carotenoids can be used as food and feed additives. Supplementation of the animal diet with carotenoids improves the nutritional quality of animal products [126].

Moreover, they can be employed as a food colorant to improve food desirability and acceptability [127].

## 2.3. Essential Oil from Agri-Food Waste

Essential oils are lipophilic substances, including terpenic hydrocarbons (e.g., monoterpenes and sesquiterpenes) and oxygenated derivatives (e.g., aldehydes, phenols, esters, and alcohols) [128]. The traditional methods used to extract the essential oils from plant matrices are liquid–solvent extraction and steam- and hydro-distillation. High temperatures (around 100 °C) can lead to the decomposition of essential oils, and the use of pentane and hexane can determine toxic organic residues. Supercritical fluid extraction (using carbon dioxide) is the most widely applied

innovative technological process. Carbon dioxide is suitable for the extraction of lipophilic compounds since it has a polarity similar to pentane as well as other desirable characteristics such as being non-flammable, non-toxic, available in high purity, and easily removed from the extract, as well as having low temperature (32 °C) and critical pressure (74 bar). Favorable extraction conditions are related to the solubility of essential oil compounds in supercritical CO<sub>2</sub> [113], the plant's pretreatment (used to facilitate the extraction by improving solvent contact and breaking cells), and online fractionation (used to achieve the separation of the essential oil from cuticular waxes) [129]. Essential oils from *Lamiaceae* (e.g., oregano, thyme, rosemary, sage, basil, and marjoram) and citrus (e.g., lemon, orange, etc.) family plants [129] were extracted using this technology. The fractionization, which can be achieved in the mode of multi-step fractionation (successive steps in which CO<sub>2</sub> density improves) or online fractionation (using a cascade decompression system consisting of two or three separators in series), can improve the selectivity of supercritical fluid extraction. This strategy can be employed when molecules with different solubilities in supercritical CO<sub>2</sub> are extracted from the same matrix [129].

## Possible Uses of the Recovered Essential Oils

Essential oils can be used as flavors, fragrances [130], and antimicrobial agents [131] in food and cosmetic products.

Moreover, they can be incorporated into food packaging to improve the UV barrier property and surface hydrophobicity to protect foods against oxidation and microbial injuries [132].

Finally, they can be used as green pesticides in agriculture [133].

## 2.4. Organosulfur Compounds from Agri-Food Waste

Organosulfur compounds are thiols with sulfur in their structure. They include glucosinolates (isothiocyanates), allyl sulfides, indoles, and sulforaphane [134]. Glucosinolates are thioglucosides that are produced in cruciferous vegetables of the *Brassica* family (e.g., broccoli, radish, cabbage, and cauliflower). They are responsible for the plant's defense against insects and pathogens. Moreover, they have some health-beneficial properties, including antioxidant (e.g., scavenge free radicals) [135], anti-inflammatory (e.g., activating detoxification enzymes, suppression of interferon regulatory factor 3, and macrophage migration inhibitory factor) [136][137], cardioprotective (e.g., they reduce low-density lipoproteins), neuroprotective, and anti-carcinogenic attributes (detoxifying carcinogens and toxicants) [138][139]. The conventional technologies used to extract them employ boiling water or aqueous organic solvent extraction [140] in a single extraction process or repetitive cycles [141]. Among the non-conventional extraction technologies, ultrasound techniques (20 kHz and 400 W) were used to improve the extraction's yield of the sinigrin from Indian mustard (+70.67% recovery than conventional techniques), microwave techniques were used to decrease the extraction time of sulforaphane from cabbages (30 min for conventional extraction to 1.5–3 min for microwave-assisted technique) [142][143], and supercritical technology was used to extract allyl isothiocyanates from wasabi (SC-CO<sub>2</sub>, 35 °C, and 25 Mpa) [144].

Allyl sulfides, produced by alliaceous vegetables (e.g., shallots, chives, leeks, and scallions), originate from S-alk(en)yl-L-cysteine sulfoxides and γ-glutamyl-S-alk(en)yl-L-cysteines. Among these, diallyl sulfides are responsible

for the pungent aroma [145][146]. Organosulfur compounds have some health-beneficial properties, including anticancer (e.g., they promote apoptosis, xenobiotic-metabolizing enzyme production, and carcinogen detoxification, in addition to the production of the enzymes that are responsible for DNA repair, are engaged in cell cycle arrest, and decrease the metabolism of nitrosamines and hydrocarbons), antioxidant (e.g., they scavenge free radicals), and antimicrobial properties [147][148]. They are thermally unstable and are lost during sterilization, pasteurization, drying, and cooking [149]. High-temperature processing decreases their bioavailability [150][151], while high-pressure processing reduces their anticancer, antimicrobial, and antioxidative properties, decreases their enzyme alliinase activity [152] and improves their enzyme alliinase activity [153][154]. Moreover, freeze-drying and infrared-drying technologies [155] and microwave-assisted and pressurized liquid extractions have negative thermal effects. Therefore, only supercritical fluid extraction is suitable for efficient extraction among the new technologies used to replace conventional ones [156].

## Possible Uses of the Recovered Organosulfur Compounds

Organosulfur compounds can be used as supplements, food additives (because of their aromatic taste and smell) [157], and biopesticides [158].

## References

1. Van Dyk, J.S.; Gama, R.; Morrison, D.; Swart, S.; Pletschke, B.I.; Morrison, D.; Swart, S.; Pletschke, B.I. Food processing waste: Problems, current management and prospects for utilisation of the lignocellulose component through enzyme synergistic degradation. *Renew. Sustain. Energy Rev.* 2013, 26, 521–531.
2. Wong, M.H.; Mo, W.Y.; Choi, W.M.; Cheng, Z.; Man, Y.B. Recycle food wastes into high quality fish feeds for safe and quality fish production. *Environ. Pollut.* 2016, 219, 631–638.
3. Apprich, S.; Tirpanalan, Ö.; Hell, J.; Reisinger, M.; Böhmdorfer, S.; Siebenhandl-Ehn, S.; Novalin, S.; Kneifel, W. Wheat bran-based biorefinery 2: Valorization of products. *LWT Food Sci. Technol.* 2014, 56, 222–231.
4. Guan, X.; Li, L.; Liu, J.; Li, S. Effects of Ultrasonic-microwave-assisted technology on hordein extraction from barley and optimization of process parameters using response surface methodology. *J. Food Qual.* 2018, 2018, 9280241.
5. Tang, S.; Hettiarachchy, N.S.; Horax, R.; Eswaranandam, S. Physicochemical properties and functionality of rice bran protein hydrolyzate prepared from heat-stabilized defatted rice bran with the aid of enzymes. *J. Food Sci.* 2003, 68, 152–157.
6. Li, S.; Zhu, D.; Li, K.; Yang, Y.; Lei, Z.; Zhang, Z. Soybean curd residue: Composition, utilization, and related limiting factors. *ISRN Ind. Eng.* 2013, 2013, 1–8.

7. Prandi, B.; Faccini, A.; Lambertini, F.; Bencivenni, M.; Jorba, M.; Van Droogenbroek, B.; Bruggeman, G.; Schöber, J.; Petrusan, J.; Elst, K.; et al. Food wastes from agrifood industry as possible sources of proteins: A detailed molecular view on the composition of the nitrogen fraction, amino acid profile and racemisation degree of 39 food waste streams. *Food Chem.* 2019, 286, 567–575.

8. Chadd, S.A.; Davies, W.P.; Koivisto, J.M. Practical production of protein for food animals. In Proceedings of the Protein Sources for the Animal Feed Industry, Proceedings of the FAO Expert Consultation and Workshop of Conference, Bangkok, Thailand, 29 April–3 May 2002; FAO: Rome, Italy; pp. 77–121.

9. Kamal, H.; Le, C.F.; Salter, A.M.; Ali, A. Extraction of protein from food waste: An overview of current status and opportunities. *CRFSFS* 2021, 20, 2455–2475.

10. Demuez, M.; Mahdy, A.; Tomás-Pejó, E.; González-Fernández, C.; Ballesteros, M. Enzymatic cell disruption of microalgae biomass in biorefinery processes. *Biotechnol. Bioeng.* 2015, 112, 1955–1966.

11. Garba, U.; Kaur, S.; Kaur, S. Protein isolates: Production, functional properties and application. *Int. J. Curr. Res. Rev.* 2014, 6, 35–45.

12. Kamal, H.; Jafar, S.; Mudgil, P.; Murali, C.; Amin, A.; Maqsood, S. Inhibitory properties of camel whey protein hydrolysates toward liver cancer cells, dipeptidyl peptidase-IV, and inflammation. *J. Dairy Sci.* 2018, 101, 8711–8720.

13. Mudgil, P.; Jobe, B.; Kamal, H.; Alameri, M.; Al Ahbabi, N.; Maqsood, S.; Jobe, B.; Kamal, H.; Alameri, M.; Al Ahbabi, N.; et al. Dipeptidyl peptidase-IV,  $\alpha$ -amylase, and angiotensin I converting enzyme inhibitory properties of novel camel skin gelatin hydrolysates. *LWT* 2019, 101, 251–258.

14. Mudgil, P.; Omar, L.S.; Omar, L.S.; Kamal, H.; Kilari, B.P.; Maqsood, S. Multi-functional bioactive properties of intact and enzymatically hydrolysed quinoa and amaranth proteins. *LWT* 2019, 110, 207–213.

15. Jafar, S.; Kamal, H.; Mudgil, P.; Hassan, H.M.; Maqsood, S.; Kamal, H.; Mudgil, P.; Hassan, H.M.; Maqsood, S. Camel whey protein hydrolysates displayed enhanced cholesteryl esterase and lipase inhibitory, anti-hypertensive and anti-haemolytic properties. *LWT* 2018, 98, 212–218.

16. Ly, H.L.; Tran, T.M.C.; Tran, T.T.T.; Ton, N.M.N.; Le, V.V.M. Application of ultrasound to protein extraction from defatted rice bran. *Int. Food Res. J.* 2018, 25, 695–701.

17. Sarker, S.D.; Latif, Z.; Gray, A.I. Natural products isolation: An overview. In *Natural Products Isolation*; Humana Press: Totowa, NJ, USA, 2006; pp. 1–25.

18. Martins, C.P.C.; Cavalcanti, R.N.; Couto, S.M.; Moraes, J.; Esmerino, E.A.; Silva, M.C.; Raices, R.S.L.; Gut, J.A.W.; Ramaswamy, H.S.; Tadini, C.C.; et al. Microwave processing: Current

background and effects on the physicochemical and microbiological aspects of dairy products. *CRFSFS* 2019, 18, 67–83.

19. Hata, S.; Wiboonsirikul, J.; Maeda, A.; Kimura, Y.; Adachi, S.; Adachi, S. Extraction of defatted rice bran by subcritical water treatment. *Biochem. Eng. J.* 2008, 40, 44–53.

20. Chiou, T.Y.; Neoh, T.L.; Kobayashi, T.; Adachi, S.; Kobayashi, T.; Adachi, S. Extraction of defatted rice bran with subcritical aqueous acetone. *Biosci. Biotechnol. Biochem.* 2012, 76, 1535–1539.

21. Sankaran, R.; Show, P.L.; Cheng, Y.S.; Tao, Y.; Ao, X.; Nguyen, T.D.P.; Show, P.L.; Cheng, Y.S.; Tao, Y.; Ao, X.; et al. Integration process for protein extraction from microalgae using liquid biphasic electric flotation (LBEF) system. *Mol. Biotechnol.* 2018, 60, 749–761.

22. Chia, S.R.; Mak, K.Y.; Khaw, Y.J.; Suhaidi, N.; Chew, K.W.; Show, P.L.; Chew, K.W.; Show, P.L. An efficient and rapid method to extract and purify protein—Liquid triphasic flotation system. *Biores. Technol.* 2019, 294, 122158.

23. Iqbal, M.; Tao, Y.; Xie, S.; Zhu, Y.; Chen, D.; Wang, X.; Huang, L.; Peng, D.; Sattar, A.; Shabbir, M.A.B.; et al. Aqueous two-phase system (ATPS): An overview and advances in its applications. *Biol. Proced. Online* 2016, 18, 1–18.

24. Fausto-Castro, L.; Rivas-García, P.; Gómez-Nafte, J.A.; Rico-Martínez, R.; Rico-Ramírez, V.; Gomez-Gonzalez, R.; Cuarón-Ibargüengoytia, J.A.; Botello-Álvarez, J.E. Selection of food waste with low moisture and high protein content from Mexican restaurants as a supplement to swine feed. *J. Clean. Prod.* 2020, 256, 120137.

25. Daw, E.; Hartel, R.W. Fat destabilization and melt-down of ice creams with increased protein content. *Intern. Dairy J.* 2015, 43, 33–41.

26. Tekin, E.; Sahin, S.; Sumnu, G. Physicochemical, rheological, and sensory properties of low-fat ice cream designed by double emulsions. *Eur. J. Lipid Sci. Technol.* 2017, 119, 1600505.

27. Remadnia, A.; Dheilly, R.; Laidoudi, B.; Quéneudec, M. Use of animal proteins as foaming agent in cementitious concrete composites manufactured with recycled PET aggregates. *Constr. Build. Mater.* 2009, 23, 3118–3123.

28. Bugnicourt, E.; Schmid, M.; Nerney, O.M.; Wildner, J.; Smykala, L.; Lazzeri, A.; Cinelli, P. Processing and validation of whey-protein-coated films and laminates at semi-industrial scale as novel recyclable food packaging materials with excellent barrier properties. *Adv. Mater. Sci. Eng.* 2013, 2013, 496207:1–496207:10.

29. Mohnen, D. Pectin structure and biosynthesis. *Curr. Opin. Plant Biol.* 2008, 11, 266–277.

30. Georgiev, Y.; Ognyanov, M.; Yanakieva, I.; Kussovski, V.; Kratchanova, M. Isolation, characterization and modification of citrus pectins. *J. Biosc. Biotechnol.* 2012, 1, 223–233.

31. Kulkarni, S.G.; Vijayanand, P. Effect of extraction conditions on the quality characteristics of pectin from passion fruit peel (*Passiflora edulis* f *flavicarpa* L.). *LWT Food Sci. Technol.* 2010, 43, 1026–1031.

32. Minjares-Fuentes, R.; Femenia, A.; Garau, M.C.; Meza-Velázquez, J.A.; Simal, S.; Rosselló, C. Ultrasound-assisted extraction of pectins from grape pomace using citric acid: A response surface methodology approach. *Carbohyd. Polym.* 2014, 106, 179–189.

33. Thirugnanasambandham, K.; Sivakumar, V. Application of D-optimal design to extract the pectin from lime bagasse using microwave green irradiation. *Inter. J. Biol. Macromol.* 2015, 72, 1351–1357.

34. Jeong, H.-S.; Kim, H.-Y.; Ahn, S.H.; Oh, S.C.; Yang, I.; Choi, I.-G. Optimization of enzymatic hydrolysis conditions for extraction of pectin from rapeseed cake (*Brassica napus* L.) using commercial enzymes. *Food Chem.* 2014, 157, 332–338.

35. Lara-Espinoza, C.; Carvajal-Millán, E.; Balandrán-Quintana, R.; López-Franco, Y.; Rascón-Chu, A. Pectin and Pectin-Based Composite Materials: Beyond Food Texture. *Molecules* 2018, 23, 942.

36. Kumar, M.; Tomar, M.; Saurabh, V.; Mahajan, T.; Punia, S.; Contreras, M.d.M.; Rudra, S.G.; Kaur, C.; Kennedy, J.F. Emerging trends in pectin extraction and its anti-microbial functionalization using natural bioactives for application in food packaging. *Trends Food Sci. Technol.* 2020, 105, 223–237.

37. Zhang, H.; Zhao, H.; Zhang, Y.; Shen, Y.; Jin, J.; Jin, Q.; Wang, X. Characterization of Positional Distribution of Fatty Acids and Triacylglycerol Molecular Compositions of Marine Fish Oils Rich in Omega-3 Polyunsaturated Fatty Acids. *BioMed Res. Int.* 2018, 2018, 3529682.

38. Alfio, V.G.; Manzo, C.; Micillo, R. From Fish Waste to Value: An Overview of the Sustainable Recovery of Omega-3 for Food Supplements. *Molecules* 2021, 26, 1002.

39. Fiori, L.; Solana, M.; Tosi, P.; Manfrini, M.; Strim, C.; Guella, G. Lipid profiles of oil from trout (*Oncorhynchus mykiss*) heads, spines and viscera: Trout byproducts as a possible source of omega-3 lipids? *Food Chem.* 2012, 134, 1088–1095.

40. Adeoti, I.A.; Hawboldt, K. A review of lipid extraction from fish processing byproduct for use as a biofuel. *Biomass Bioenergy* 2014, 63, 330–340.

41. Baiano, A. Recovery of biomolecules from food wastes—A review. *Molecules* 2014, 19, 14821.

42. Food and Agriculture Organization. The Production of Fish Meal and Oil. The Process. Available online: <http://www.fao.org/3/x6899e/x6899e00.htm7bcd94a78e/> (accessed on 30 November 2020).

43. Al Khawli, F.; Pateiro, M.; Domínguez, R.; Lorenzo, J.M.; Gullón, P.; Kousoulaki, K.; Ferrer, E.; Berrada, H.; Barba, F.J. Innovative green technologies of intensification for valorization of seafood and their byproducts. *Mar. Drugs* 2019, 17, 689.

44. Caruso, G.; Floris, R.; Serangeli, C.; Di Paola, L. Fishery Wastes as a Yet Undiscovered Treasure from the Sea: Biomolecules Sources, Extraction Methods and Valorization. *Mar. Drugs* 2020, 18, 622.

45. Ruiz, J.C.R.; Vazquez, E.D.L.L.O.; Campos, M.R.S. Encapsulation of vegetable oils as source of omega-3 fatty acids for enriched functional foods. *Crit. Rev. Food Sci. Nutr.* 2017, 57, 1423–1434.

46. Saini, R.K.; Keum, Y. Omega-3 and omega-6 polyunsaturated fatty acids: Dietary sources, metabolism, and significance—A review. *Life Sci.* 2018, 203, 255–267.

47. Huang, T.-H.; Wang, P.-W.; Yang, S.-C.; Chou, W.-L.; Fang, J.-Y. Cosmetic and Therapeutic Applications of Fish Oil's Fatty Acids on the Skin. *Mar. Drugs* 2018, 16, 256.

48. Dini, I.; Marra, R.; Cavallo, P.; Pironti, A.; Sepe, I.; Troisi, J.; Scala, G.; Lombari, P.; Vinale, F. Trichoderma Strains and Metabolites Selectively Increase the Production of Volatile Organic Compounds (VOCs) in Olive Trees. *Metabolites* 2021, 11, 213.

49. Dini, I.; Pascale, M.; Staropoli, A.; Marra, R.; Vinale, F. Effect of Selected Trichoderma Strains and Metabolites on Olive Drupes. *Appl. Sci.* 2021, 11, 8710.

50. Dini, I.; Graziani, G.; Fedele, F.L.; Sicari, A.; Vinale, F.; Castaldo, L.; Ritieni, A. Effects of Trichoderma Biostimulation on the Phenolic Profile of Extra-Virgin Olive Oil and Olive Oil By-Products. *Antioxidants* 2020, 9, 284.

51. Dini, I.; Graziani, G.; Gaspari, A.; Fedele, F.L.; Sicari, A.; Vinale, F.; Cavallo, P.; Lorito, M.; Ritieni, A. New Strategies in the Cultivation of Olive Trees and Repercussions on the Nutritional Value of the Extra Virgin Olive Oil. *Molecules* 2020, 25, 2345.

52. Dini, I.; Graziani, G.; Fedele, F.L.; Sicari, A.; Vinale, F.; Castaldo, L.; Ritieni, A. An environmentally friendly practice used in olive cultivation capable of increasing commercial interest in waste products from oil processing. *Antioxidants* 2020, 9, 466.

53. Dini, I.; Laneri, S. Nutricosmetics: A brief overview. *Phytother. Res.* 2019, 33, 3054–3063.

54. Dini, I.; Seccia, S.; Senatore, A.; Coppola, D.; Morelli, E. Development and Validation of an Analytical Method for Total Polyphenols Quantification in Extra Virgin Olive Oils. *Food Anal. Methods* 2019, 13, 457–464.

55. Dini, I. Spices and herbs as therapeutic foods. In *Food Quality: Balancing Health and Disease*; Holban, A.M., Grumezescu, A.M., Eds.; Academic Press/Elsevier: London, UK, 2018; pp. 433–469.

56. Dini, I.; Laneri, S. Spices, Condiments, Extra Virgin Olive Oil and Aromas as Not Only Flavorings, but Precious Allies for Our Wellbeing. *Antioxidants* 2021, 10, 868.

57. Laneri, S.; Di Lorenzo, R.M.; Bernardi, A.; Sacchi, A.; Dini, I. *Aloe barbadensis*: A plant of nutricosmetic interest. *Nat. Prod. Comun.* 2020, 15, 1–6.

58. Dini, I.; Falanga, D.; Di Lorenzo, R.; Tito, A.; Carotenuto, G.; Zappelli, C.; Grumetto, L.; Sacchi, A.; Laneri, S.; Apone, F. An Extract from *Ficus carica* Cell Cultures Works as an Anti-Stress Ingredient for the Skin. *Antioxidants* 2021, 10, 515.

59. Dini, I.; Laneri, S. The New Challenge of Green Cosmetics: Natural Food Ingredients for Cosmetic Formulations. *Molecules* 2021, 26, 3921.

60. Cavallo, P.; Dini, I.; Sepe, I.; Galasso, G.; Fedele, F.L.; Sicari, A.; Bolletti Censi, S.; Gaspari, A.; Ritieni, A.; Lorito, M.; et al. An Innovative Olive Pâté with Nutraceutical Properties. *Antioxidants* 2020, 9, 581.

61. Dini, I.; Izzo, L.; Graziani, G.; Ritieni, A. The Nutraceutical Properties of “Pizza Napoletana Marinara TSG” a Traditional Food Rich in Bioaccessible Antioxidants. *Antioxidants* 2021, 10, 495.

62. Galanakis, C.M. Recovery of high added-value components from food wastes: Conventional, emerging technologies and commercialized applications. *Trends Food Sci. Technol.* 2012, 26, 68–87.

63. Jimenez-Lopez, C.; Fraga-Corral, M.; Carpena, M.; Garcia-Oliveira, P.; Echave, J.; Pereira, A.; Lourenço-Lopes, C.; Prieto, M.; Simal-Gandara, J. Agriculture wastes valorisation as a source of antioxidant phenolic compounds within a circular and sustainable bio-economy. *Food Funct.* 2020, 11, 4853–4877.

64. Echegaray, N.; Munekata, P.E.S.; Gullón, P.; Dzuvor, C.K.O.; Gullón, B.; Kubi, F.; Lorenzo, J.M. Recent advances in food products fortification with anthocyanins. *Crit. Rev. Food Sci. Nutr.* 2020, 16, 1–15.

65. Amogne, N.Y.; Ayele, D.W.; Tsigie, Y.A. Recent advances in anthocyanin dyes extracted from plants for dye sensitized solar cell. *Mater. Renew. Sustain. Energy* 2020, 9, 23.

66. Rodriguez-Amaya, D.B. Update on natural food pigments—A mini-review on carotenoids, anthocyanins, and betalains. *Food Res. Int.* 2019, 124, 200–205.

67. Weber, F.; Larsen, L.R. Influence of fruit juice processing on anthocyanin stability. *Food Res. Int.* 2017, 100, 354–365.

68. Kšonžeková, P.; Mariychuk, R.; Eliašová, A.; Mudroňová, D.; Csank, T.; Király, J.; Marcinčáková, D.; Pistl, J.; Tkáčiková, L. In vitro study of biological activities of anthocyanin-rich berry extracts on porcine intestinal epithelial cells. *J. Sci. Food Agric.* 2016, 96, 1093–1100.

69. Zhang, H.; Xu, Z.; Zhao, H.; Wang, X.; Pang, J.; Li, Q.; Yang, Y.; Ling, W. Anthocyanin supplementation improves anti-oxidative and anti-inflammatory capacity in a dose–response manner in subjects with dyslipidemia. *Redox Biol.* 2020, 32, 101474.

70. Hurst, R.D.; Lyall, K.A.; Wells, R.W.; Sawyer, G.M.; Lomiwes, D.; Ngametua, N.; Hurst, S.M. Daily Consumption of an Anthocyanin-Rich Extract Made From New Zealand Blackcurrants for 5 Weeks Supports Exercise Recovery Through the Management of Oxidative Stress and Inflammation: A Randomized Placebo Controlled Pilot Study. *Front. Nutr.* 2020, 7, 16.

71. Roth, S.; Spalinger, M.R.; Gottier, C.; Biedermann, L.; Zeitz, J.; Lang, S.; Weber, A.; Rogler, G.; Scharl, M. Bilberry-Derived Anthocyanins Modulate Cytokine Expression in the Intestine of Patients with Ulcerative Colitis. *PLoS ONE* 2016, 11, e0154817.

72. Yang, L.; Ling, W.; Yang, Y.; Chen, Y.; Tian, Z.; Du, Z.; Chen, J.; Xie, Y.; Liu, Z.; Yang, L. Role of Purified Anthocyanins in Improving Cardiometabolic Risk Factors in Chinese Men and Women with Prediabetes or Early Untreated Diabetes—A Randomized Controlled Trial. *Nutrients* 2017, 9, 1104.

73. Yang, L.; Qiu, Y.; Ling, W.; Liu, Z.; Yang, L.; Wang, C.; Peng, X.; Wang, L.; Chen, J. Anthocyanins regulate serum adipsin and visfatin in patients with prediabetes or newly diagnosed diabetes: A randomized controlled trial. *Eur. J. Nutr.* 2021, 60, 1935–1944.

74. Aboonabi, A.; Meyer, R.R.; Gaiz, A.; Singh, I. Anthocyanins in berries exhibited anti-atherogenicity and antiplatelet activities in a metabolic syndrome population. *Nutr. Res.* 2020, 76, 82–93.

75. Aboonabi, A.; Aboonabi, A. Anthocyanins reduce inflammation and improve glucose and lipid metabolism associated with inhibiting nuclear factor-kappaB activation and increasing PPAR-γ gene expression in metabolic syndrome subjects. *Free Radic. Biol. Med.* 2020, 150, 30–39.

76. Thompson, K.; Pederick, W.; Singh, I.; Santhakumar, A.B. Anthocyanin supplementation in alleviating thrombogenesis in overweight and obese population: A randomized, double-blind, placebo-controlled study. *J. Funct. Foods* 2017, 32, 131–138.

77. Thompson, K.; Hosking, H.; Pederick, W.; Singh, I.; Santhakumar, A.B. The effect of anthocyanin supplementation in modulating platelet function in sedentary population: A randomised, double-blind, placebo-controlled, cross-over trial. *Br. J. Nutr.* 2017, 118, 368–374.

78. Groh, I.A.M.; Riva, A.; Braun, D.; Sutherland, H.G.; Williams, O.; Bakuradze, T.; Pahlke, G.; Richling, E.; Haupt, L.M.; Griffiths, L.R.; et al. Long-Term Consumption of Anthocyanin-Rich Fruit Juice: Impact on Gut Microbiota and Antioxidant Markers in Lymphocytes of Healthy Males. *Antioxidants* 2020, 10, 27.

79. Istan, G.; Wood, E.; Le Sayec, M.; Rawlings, C.; Yoon, J.; Dandavate, V.; Cera, D.; Rampelli, S.; Costabile, A.; Fromentin, E.; et al. Effects of aronia berry (poly)phenols on vascular function and

gut microbiota: A double-blind randomized controlled trial in adult men. *Am. J. Clin. Nutr.* 2019, 110, 316–329.

80. Shimizu, N.; Yamada, W.; Miyasaka, K.; Shimoda, H. Ameliorating Effects of Delphinol®, Anthocyanin Standardized Maqui Berry Extract, on Skin Brightness and Redness in Japanese Females: A Randomized Double-Blind Placebo-Controlled Pilot Study. *J. Cosmet. Dermatol. Sci. Appl.* 2020, 10, 149–162.

81. Xi, J.; Li, Z.; Fan, Y. Recent advances in continuous extraction of bioactive ingredients from food-processing wastes by pulsed electric fields. *Crit. Rev. Food Sci. Nutr.* 2020, 1–13.

82. Kosehira, M.; Machida, N.; Kitaichi, N. A 12-Week-Long Intake of Bilberry Extract (*Vaccinium myrtillus* L.) Improved Objective Findings of Ciliary Muscle Contraction of the Eye: A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Comparison Trial. *Nutrients* 2020, 12, 600.

83. Martínez, J.M.; Delso, C.; Álvarez, I.; Raso, J. Pulsed electric field-assisted extraction of valuable compounds from microorganisms. *Compr. Rev. Food Sci. Food Saf.* 2020, 19, 530–552.

84. Leong, S.Y.; Treadwell, M.; Liu, T.; Hochberg, M.; Sack, M.; Mueller, G.; Sigler, J.; Silcock, P.; Oey, I. Influence of Pulsed Electric Fields processing at high-intensity electric field strength on the relationship between anthocyanins composition and colour intensity of Merlot (*Vitis vinifera* L.) musts during cold maceration. *Innov. Food Sci. Emerg. Technol.* 2020, 59, 102243.

85. Corrales, M.; Toepfl, S.; Butz, P.; Knorr, D.; Tauscher, B. Extraction of anthocyanins from grape byproducts assisted by ultrasonics, high hydrostatic pressure or pulsed electric fields: A comparison. *Innov. Food Sci. Emerg. Technol.* 2008, 9, 85–91.

86. Pataro, G.; Carullo, D.; Bobinaite, R.; Donsì, G.; Ferrari, G. Improving the extraction yield of juice and bioactive compounds from sweet cherries and their byproducts by pulsed electric fields. *Chem. Eng. Trans.* 2017, 57, 1717–1722.

87. Arruda, H.S.; Pereira, G.A.; Pastore, G.M. Optimization of Extraction Parameters of Total Phenolics from *Annona crassiflora* Mart. (*Araticum*) Fruits Using Response Surface Methodology. *Food Anal. Methods* 2017, 10, 100–110.

88. Da Rocha, C.B.; Noreña, C.P.Z. Microwave-Assisted Extraction and Ultrasound-Assisted Extraction of Bioactive Compounds from Grape Pomace. *Int. J. Food Eng.* 2020, 16, 20190191.

89. Bobinaitė, R.; Pataro, G.; Raudonis, R.; Vškelis, P.; Bobinas, C.; Šatkauskas, S.; Ferrari, G. Improving the extraction of juice and anthocyanin compounds from blueberry fruits and their byproducts by pulsed electric fields. In Proceedings of the 1st World Congress on Electroporation and Pulsed Electric Fields in Biology, Medicine and Food & Environmental Technologies (WC2015), Portorož, Slovenia, 6–10 September 2015; Jarm, T., Kramar, P., Eds.; Springer: Singapore, 2016; Volume 53, pp. 363–366.

90. Romero-Díez, R.; Matos, M.; Rodrigues, L.; Bronze, M.R.; Rodríguez-Rojo, S.; Cocero, M.J.; Matias, A.A. Microwave and ultrasound pretreatments to enhance anthocyanins extraction from different wine lees. *Food Chem.* 2019, 272, 258–266.

91. Okur, İ.; Baltacıoğlu, C.; Ağçam, E.; Baltacıoğlu, H.; Alpas, H. Evaluation of the Effect of Different Extraction Techniques on Sour Cherry Pomace Phenolic Content and Antioxidant Activity and Determination of Phenolic Compounds by FTIR and HPLC. *Waste Biomass Valoriz.* 2019, 10, 3545–3555.

92. Da Porto, C.; Natolino, A. Extraction kinetic modelling of total polyphenols and total anthocyanins from saffron floral bio-residues: Comparison of extraction methods. *Food Chem.* 2018, 258, 137–143.

93. Puértolas, E.; Barba, F.J. Electrotechnologies applied to valorization of byproducts from food industry: Main findings, energy and economic cost of their industrialization. *Food Bioprod. Process.* 2016, 100, 172–184.

94. Zhao, M.; Luo, Y.; Li, Y.; Liu, X.; Wua, J.; Liao, X.; Chen, F. The identification of degradation products and degradation pathway of malvidin-3-glucoside and malvidin-3,5-diglucoside under microwave treatment. *Food Chem.* 2013, 141, 3260–3267.

95. Ciriminna, R.; Carnaroglio, D.; Delisi, R.; Arvati, S.; Tamburino, A.; Pagliaro, M. Industrial feasibility of natural products extraction with microwave technology. *Chem. Sel. Rev.* 2016, 1, 549–555.

96. Doulabi, M.; Golmakani, M.; Ansari, S. Evaluation and optimization of microwave-assisted extraction of bioactive compounds from eggplant peel byproduct. *J. Food Process. Preserv.* 2020, 44, e14853.

97. Varadharajan, V.; Shanmugam, S.; Ramaswamy, A. Model generation and process optimization of microwave-assisted aqueous extraction of anthocyanins from grape juice waste. *J. Food Process Eng.* 2017, 40, e12486.

98. Li, X.; Chen, F.; Li, S.; Jia, J.; Gu, H.; Yang, L. An efficient homogenate-microwave-assisted extraction of flavonols and anthocyanins from blackcurrant marc: Optimization using combination of Plackett-Burman design and Box-Behnken design. *Ind. Crops Prod.* 2016, 94, 834–847.

99. Grillo, G.; Gunjević, V.; Radošević, K.; Redovniković, I.R.; Cravotto, G. Deep Eutectic Solvents and Nonconventional Technologies for Blueberry-Peel Extraction: Kinetics, Anthocyanin Stability, and Antiproliferative Activity. *Antioxidants* 2020, 9, 1069.

100. Halee, A.; Supavititpatana, P.; Ruttarattanamongkol, K.; Jittrepotch, N.; Rojsuntornkitti, K.; Kongbangkerd, T. Optimisation of the microwave-assisted extraction of natural antioxidants from defatted black rice bran of *Oryza sativa* L. cv. Homnin. *J. Microbiol. Biotechnol. Food Sci.* 2020, 9, 1134–1140.

101. Fernandez-Aulis, F.; Hernandez-Vazquez, L.; Aguilar-Osorio, G.; Arrieta-Baez, D.; Navarro-Ocana, A. Extraction and Identification of Anthocyanins in Corn Cob and Corn Husk from Cacahuacintle Maize. *J. Food Sci.* 2019, 84, 954–962.

102. Herrman, D.A.; Brantsen, J.F.; Ravisankar, S.; Lee, K.M.; Awika, J.M. Stability of 3-deoxyanthocyanin pigment structure relative to anthocyanins from grains under microwave assisted extraction. *Food Chem.* 2020, 333, 127494.

103. Tiwari, B.K.; Patras, A.; Brunton, N.; Cullen, P.J.; O'Donnell, C.P. Effect of ultrasound processing on anthocyanins and color of red grape juice. *Ultrason. Sonochem.* 2010, 17, 598–604.

104. Xue, H.; Tan, J.; Li, Q.; Tang, J.; Cai, X. Ultrasound-Assisted Enzymatic Extraction of Anthocyanins from Raspberry Wine Residues: Process Optimization, Isolation, Purification, and Bioactivity Determination. *Food Anal. Methods* 2021, 10, 279.

105. Galván D'Alessandro, L.; Dimitrov, K.; Vauchel, P.; Nikov, I. Kinetics of ultrasound assisted extraction of anthocyanins from Aronia melanocarpa (black chokeberry) wastes. *Chem. Eng. Res. Des.* 2014, 92, 1818–1826.

106. Mark, R.; Lyu, X.; Lee, J.J.L.; Parra-Saldívar, R.; Chen, W.N. Sustainable production of natural phenolics for functional food applications. *J. Funct. Foods* 2019, 57, 233–254.

107. Wang, Y.; Du, H.; Xie, M.; Ma, G.; Yang, W.; Hu, Q.; Pei, F. Characterization of the physical properties and biological activity of chitosan films grafted with gallic acid and caffeic acid: A comparison study. *Food Packag. Shelf Life* 2019, 22, 100401.

108. Albuquerque, B.R.; Héleno, S.A.; Oliveira, M.B.P.; Barros, L.; Ferreira, I.C. Phenolic compounds: Current industrial applications, limitations and future challenges. *Food Funct.* 2021, 12, 14–29.

109. Cano, A.; Andres, M.; Chiralt, A.; González-Martínez, C. Use of tannins to enhance the functional properties of protein based films. *Food Hydrocoll.* 2020, 100, 105443.

110. Jia, Y.; Liu, B.; Cheng, D.; Li, J.; Huang, F.; Lu, Y. Dyeing characteristics and functionability of tussah silk fabric with oak bark extract. *Text. Res. J.* 2017, 87, 1806–1817.

111. Merhan, O. Biochemistry and antioxidant properties of carotenoids. *Carotenoids* 2017, 5, 51.

112. Nguyen, V.T.; Scarlett, C.J. Mass proportion, bioactive compounds and antioxidant capacity of carrot peel as affected by various solvents. *Technologies* 2016, 4, 36.

113. Cerna, J.; Athari, N.; Robbs, C.; Walk, A.; Edwards, C.; Adamson, B.; Khan, N. Macular Carotenoids, Retinal Morphometry, and Cognitive Function in Multiple Sclerosis (OR05–07–19). *Curr. Dev. Nutr.* 2019, 3 (Suppl. S1), nzz029-OR05.

114. Wang, Y.; Chung, S.J.; McCullough, M.L.; Song, W.O.; Fernandez, M.L.; Koo, S.I.; Chun, O.K. Dietary carotenoids are associated with cardiovascular disease risk biomarkers mediated by serum carotenoid concentrations. *J. Nutr.* 2014, 144, 1067–1074.

115. Wu, J.; Cho, E.; Willett, W.C.; Sastry, S.M.; Schaumberg, D.A. Intakes of Lutein, Zeaxanthin, and Other Carotenoids and Age-Related Macular Degeneration during 2 Decades of Prospective Follow-up. *JAMA Ophthalmol.* 2015, 133, 1415–1424.

116. Hwang, S.; Lim, J.W.; Kim, H. Inhibitory effect of lycopene on amyloid- $\beta$ -induced apoptosis in neuronal cells. *Nutrients* 2017, 9, 883.

117. Gul, K.; Tak, A.; Singh, A.K.; Singh, P.; Yousuf, B.; Wani, A.A. Chemistry, encapsulation, and health benefits of  $\beta$ -carotene-A review. *Cogent Food Agric.* 2015, 1, 1018696.

118. Khoo, H.E.; Prasad, K.N.; Kong, K.W.; Jiang, Y.; Ismail, A. Carotenoids and their isomers: Color pigments in fruits and vegetables. *Molecules* 2011, 16, 1710.

119. Pénicaud, C.; Achir, N.; Dhuique-Mayer, C.; Dornier, M.; Bohuon, P. Degradation of  $\beta$ -carotene during fruit and vegetable processing or storage: Reaction mechanisms and kinetic aspects: A review. *Fruits* 2011, 66, 417–440.

120. de Andrade Lima, M.; Charalampopoulos, D.; Chatzifragkou, A. Optimisation and modelling of supercritical CO<sub>2</sub> extraction process of carotenoids from carrot peels. *J. Supercrit. Fluids* 2018, 133, 94–102.

121. Hiranvarachat, B.; Devahastin, S. Enhancement of microwave-assisted extraction via intermittent radiation: Extraction of carotenoids from carrot peels. *J. Food Eng.* 2014, 126, 17–26.

122. Saini, R.K.; Keum, Y.-S. Carotenoid extraction methods: A review of recent developments. *Food Chem.* 2018, 240, 90–103.

123. Nagarajan, J.; Pui Kay, H.; Krishnamurthy, N.P.; Ramakrishnan, N.R.; Aldawoud, T.M.S.; Galanakis, C.M.; Ooi, C.W. Extraction of carotenoids from tomato pomace via water-induced hydrocolloidal complexation. *Biomolecules* 2020, 10, 1019.

124. Nagarajan, J.; Krishnamurthy, N.P.; Nagasundara Ramanan, R.; Raghunandan, M.E.; Galanakis, C.M.; Ooi, C.W. A facile water-induced complexation of lycopene and pectin from pink guava byproduct: Extraction, characterization and kinetic studies. *Food Chem.* 2019, 296, 47–55.

125. Nagarajan, J.; Kay, H.P.; Arumugam, P.; Krishnamurthy, N.P.; Ramakrishnan, N.R.; Aldawoud, T.M.; Galanakis, C.M.; Wei, O.C. Valorisation of carrot peel waste by water-induced hydrocolloidal complexation for extraction of carotenoid and pectin. *Chemosphere* 2021, 272, 129919.

126. Moreno, J.A.; Díaz-Gómez, J.; Nogareda, C.; Angulo, E.; Sandmann, G.; Portero-Otin, M.; Serrano, J.C.; Twyman, R.M.; Capell, T.; Zhu, C.; et al. The distribution of carotenoids in hens fed on biofortified maize is influenced by feed composition, absorption, resource allocation and storage. *Sci. Rep.* 2016, 6, 35346.

127. Cataldo, V.F.; López, J.; Cárcamo, M.; Agosin, E. Chemical vs. biotechnological synthesis of C 13-apocarotenoids: Current methods, applications and perspectives. *Appl. Microb. Biotechnol.*

2016, 100, 5703–5718.

128. Arce, A.; Marchiaro, A.; Martínez-Ageitos, J.M.; Soto, A. Citrus essential oil deterpenation by liquid-liquid extraction. *Can. J. Chem. Eng.* 2005, 83, 366–370.

129. Fornari, T.; Vicente, G.; Vázquez, E.; García-Risco, M.R.; Reglero, G. Isolation of essential oil from different plants and herbs by supercritical fluid extraction. *J. Chromatogr. A* 2012, 1250, 34–48.

130. Diniz do Nascimento, L.; Moraes, A.A.B.d.; Costa, K.S.d.; Pereira Galúcio, J.M.; Taube, P.S.; Costa, C.M.L.; Neves Cruz, J.; de Aguiar Andrade, E.H.; Faria, L.J.G.d. Bioactive Natural Compounds and Antioxidant Activity of Essential Oils from Spice Plants: New Findings and Potential Applications. *Biomolecules* 2020, 10, 988.

131. Khorshidian, N.; Yousefi, M.; Khanniri, E.; Mortazavian, A.M. Potential application of essential oils as antimicrobial preservatives in cheese. *Innov. Food Sci. Emerg. Technol.* 2018, 45, 62–72.

132. Sharma, S.; Barkauskaite, S.; Jaiswal, A.K.; Jaiswal, S. Essential oils as additives in active food packaging. *Food Chem.* 2020, 343, 128403.

133. Abdelatti, Z.A.S.; Hartbauer, M. Plant oil mixtures as a novel botanical pesticide to control gregarious locusts. *J. Pest Sci.* 2020, 93, 341–353.

134. Ruhee, R.T.; Roberts, L.A.; Ma, S.; Suzuki, K. Organosulfur Compounds: A Review of Their Anti-inflammatory Effects in Human Health. *Front Nutr.* 2020, 7, 64.

135. Kim, S.; Kim, D.B.; Jin, W.; Park, J.; Yoon, W.; Lee, Y.; Kim, S.; Lee, S.; Kim, S.; Lee, O.H.; et al. Comparative studies of bioactive organosulphur compounds and antioxidant activities in garlic (*Allium sativum* L.), elephant garlic (*Allium ampeloprasum* L.) and onion (*Allium cepa* L.). *Nat Prod Res.* 2018, 32, 1193–1197.

136. Putnik, P.; Gabric, D.; Roohinejad, S.; Barba, F.J.; Granato, D.; Mallikarjunan, K.; Lorenzo, J.M.; Bursać Kovačević, D. An overview of organosulfur compounds from *Allium* spp.: From processing and preservation to evaluation of their bioavailability, antimicrobial, and anti-inflammatory properties. *Food Chem.* 2019, 276, 680–691.

137. Nair, A.; Chattopadhyay, D.; Saha, B. Plant-derived immunomodulators. In *New Look to Phytomedicine*; Elsevier: Amsterdam, The Netherlands, 2019; pp. 435–499.

138. Bischoff, K.L. Glucosinolates. In *Nutraceuticals*; Elsevier: Amsterdam, The Netherlands, 2016; pp. 551–554.

139. Kris-Etherton, P.M.; Hecker, K.D.; Bonanome, A.; Coval, S.M.; Binkoski, A.E.; Hilpert, K.F.; Griell, A.E.; Etherton, T.D. Bioactive compounds in foods: Their role in the prevention of cardiovascular disease and cancer. *Am. J. Med.* 2002, 113 (Suppl. 9B), 71s–88s.

140. Wang, T.; Liang, H.; Yuan, Q. Optimization of ultrasonic-stimulated solvent extraction of sinigrin from Indian mustard seed (*Brassica juncea* L.) using response surface methodology. *Phytochem. Anal.* 2011, 22, 205–213.

141. Omirou, M.; Papastylianou, I.; Iori, R.; Papastefanou, C.; Papadopoulou, K.K.; Ehaliotis, C.; Karpouzas, D.G. Microwave-assisted extraction of glucosinolates from *Eruca sativa* seeds and soil: Comparison with existing methods. *Phytochem. Anal.* 2009, 20, 214–220.

142. Vilkhu, K.; Mawson, R.; Simons, L.; Bates, D. Applications and opportunities for ultrasound assisted extraction in the food industry: A review. *Innov. Food Sci. Emerg. Technol.* 2008, 9, 161–169.

143. Ares, A.M.; Bernal, J.; Martin, M.T.; Bernal, J.L.; Nozal, M.J. Optimized formation, extraction, and determination of sulforaphane in broccoli by liquid chromatography with diode array detection. *Food Anal. Methods* 2014, 7, 730–740.

144. Li, L.; Lee, W.; Lee, W.J.; Auh, J.H.; Kim, S.S.; Yoon, J. Extraction of allyl isothiocyanate from wasabi (*Wasabia japonica* Matsum) using supercritical carbon dioxide. *Food Sci. Biotechnol.* 2010, 19, 405–410.

145. Zhou, Y.; Yang, G.; Tian, H.; Hu, Y.; Wu, S.; Geng, Y.; Lin, K.; Wu, W. Sulforaphane metabolites cause apoptosis via microtubule disruption in cancer. *Endocr. Relat. Cancer* 2018, 25, 255–268.

146. Lawson, L.D.; Wang, Z.J. Allicin and allicin-derived garlic compounds increase breath acetone through allyl methyl sulfide: Use in measuring allicin bioavailability. *J. Agric. Food Chem.* 2005, 53, 1974–1983.

147. Wang, H.C.; Pao, J.; Lin, S.Y.; Sheen, L.Y. Molecular mechanisms of garlic-derived allyl sulfides in the inhibition of skin cancer progression. *Ann. N. Y. Acad. Sci.* 2012, 1271, 44–52.

148. Nuutila, A.M.; Puupponen-Pimiä, R.; Aarni, M.; Oksman-Caldentey, K.-M. Comparison of antioxidant activities of onion and garlic extracts by inhibition of lipid peroxidation and radical scavenging activity. *Food Chem.* 2003, 81, 485–493.

149. Putnik, P.; Barba, F.J.; Lucini, L.; Rocchetti, G.; Montesano, D. Conventional, non-conventional extraction techniques and new strategies for the recovery of bioactive compounds from plant material for human nutrition. *Food Res Int.* 2019, 123, 516–517.

150. Tomatis, I. Allium sativum Bulb Absolutes and Therapeutic or Cosmetic Uses. U.S. Patent 7,192,613, 20 March 2007.

151. Murtaza, A.; Iqbal, A.; Marszałek, K.; Iqbal, A.M.; Waseem Ali, S.; Xu, X.; Pan, S.; Hu, W. Enzymatic, Phyto-, and Physicochemical Evaluation of Apple Juice under High-Pressure Carbon Dioxide and Thermal Processing. *Foods* 2020, 9, 243.

152. Kim, K.W.; Kim, Y.-T.; Kim, M.; Noh, B.-S.; Choi, W.-S. Effect of high hydrostatic pressure (HHP) treatment on flavor, physicochemical properties and biological functionalities of garlic. *LWT Food Sci. Technol.* 2014, 55, 347–354.

153. Gonzalez, M.E.; Anthon, G.E.; Barrett, D.M. Onion cells after high pressure and thermal processing: Comparison of membrane integrity changes using different analytical methods and impact on tissue texture. *J. Food Sci.* 2010, 75, E426–E432.

154. Putnik, P.; Lorenzo, J.M.; Barba, F.J.; Roohinejad, S.; Režek Jambrak, A.; Granato, D.; Montesano, D.; Bursać Kovačević, D. Novel food processing and extraction technologies of high-added value compounds from plant materials. *Foods* 2018, 7, 106.

155. Zhou, L.; Guo, X.; Bi, J.; Yi, J.; Chen, Q.; Wu, X.; Zhou, M. Drying of garlic slices (*Allium sativum* L.) and its effect on thiosulfinates, total phenolic compounds and antioxidant activity during infrared drying. *J. Food Process. Preserv.* 2017, 41, e12734.

156. Poojary, M.M.; Putnik, P.; Kovačević, D.B.; Barba, F.J.; Lorenzo, J.M.; Dias, D.A.; Shpigelman, A. Stability and extraction of bioactive sulfur compounds from *Allium* genus processed by traditional and innovative technologies. *J. Food Compos. Anal.* 2017, 61, 28–39.

157. Zhao, M.M.; Xiong, S.H.; Zhao, G.D.; Yu, H. Organosulfur compounds in food. In *Handbook of Dietary Phytochemicals*; Xiao, J., Sarker, S.D., Asakawa, Y., Eds.; Springer: Singapore, 2020; pp. 1–21.

158. Durán-Lara, E.F.; Valderrama, A.; Marican, A. Natural Organic Compounds for Application in Organic Farming. *Agriculture* 2020, 10, 41.

Retrieved from <https://encyclopedia.pub/entry/history/show/37706>