

# Organoids and Aging

Subjects: Cell Biology | Geriatrics & Gerontology

Contributor: Marta Gonzalez-Freire

The biology of aging is focused on the identification of novel pathways that regulate the underlying processes of aging to develop interventions aimed at delaying the onset and progression of chronic diseases to extend lifespan. However, the research on the aging field has been conducted mainly in animal models, yeast, *Caenorhabditis elegans*, and cell cultures. Thus, it is unclear to what extent this knowledge is transferable to humans since they might not reflect the complexity of aging in people. An organoid culture is an in vitro 3D cell-culture technology that reproduces the physiological and cellular composition of the tissues and/or organs. This technology is being used in the cancer field to predict the response of a patient-derived tumor to a certain drug or treatment serving as patient stratification and drug-guidance approaches. Modeling aging with patient-derived organoids has a tremendous potential as a preclinical model tool to discover new biomarkers of aging, to predict adverse outcomes during aging, and to design personalized approaches for the prevention and treatment of aging-related diseases and geriatric syndromes. This could represent a novel approach to study chronological and/or biological aging, paving the way to personalized interventions targeting the biology of aging.

Keywords: organoids ; aging ; precision medicine

---

## 1. Introduction: Organoids, Spheroids, and Matrix-Embedded 3D Cultures

Patient-derived organoids (PDOs) are self-organized 3D tissue cultures that are derived from stem cells. Isolated patients' stem cells differentiate to form an organ-like tissue comprising multiple cell types. Organoids have self-renewal and self-organization capabilities and retain the characteristics of the physiological structure and function of their source [1][2]. Recent culturing advances aim to create the right environment for the stem cells so they can follow their own genetic instructions to self-organize, forming organoid structures that resemble miniature organs composed of many cell types. This approach provides tractable in vitro models of human physiology and pathology, thereby enabling interventional studies that are difficult or impossible to conduct in human subjects [1][3]. Attempts to model the biology of human organs—including the differentiation of human stem cells in 2D, in either the presence or absence of a 3D matrix; bio-printing of human cells; and the culture of cells in a microfluidic device ("organ-on-a-chip")—were made prior to the emergence of organoids and have shown some potential for drug screening or human disease research.

During the late twentieth and early twenty-first centuries, the use of classical cell lines and animal model systems in biomedical research has helped to improve our understanding of cellular signaling pathways, to identify potential drug targets and to guide the design of candidate drugs for pathologies including cancer and infectious diseases, among others [3][4]. Recent studies have identified biological processes that are specific to the human body, such as brain development, metabolism, and the test of drug efficacy that cannot be modeled in animal or cell models. Nevertheless, extrapolating results from these model systems to humans has become a major bottleneck in the drug discovery process. Therefore, the emergence of human in vitro 3D cell cultures, such as organoids, spheroids, and matrix-embedded 3D cultures has received widespread attention due to the potential to overcome these limitations [4][5][6]. These 3D structures of cultured cells recapitulate important aspects of in vivo organ development and biological function. Such cultures can be crafted to replicate much of the complexity of an organ or to express selected aspects of its physiology like producing only certain types of cells [1][2][3].

Spheroids form by spontaneous aggregation of cells followed by the binding of cell surface integrins to the extracellular matrix (ECM). After initial cell–cell contact, cells upregulate E-Cadherin, which accumulates on the cell surface and then the spheroid becomes a compact structure through strong intercellular E-cadherin interactions [6]. Different spheroid models have been described based on their cellular sources. Multicellular tumor spheroids (MCTS) are often made from cancer cell lines but rarely from tumor tissues. MCTS show little histological resemblance to the original tumor, but they mimic metabolic and proliferation gradients of the in vivo tumor and model clinically relevant resistance to chemotherapy.

The advantages of MCTS are that they are clonal, simple to expand into large cultures, and suitable for high-throughput systems [5][6][7].

In contrast, organoids grow from stem cells, which can divide indefinitely and produce different types of cells as part of their progeny. Organoids allow genetic and pharmacological manipulation in a complex cellular context that reflects human biology and enable investigations of the early stages of organ development and disease onset. They complement (and may eventually replace) animal models in many areas of preclinical drug development. Moreover, they provide patient-specific “avatars” for drug development and precision therapies, including treatments for cancer, rare genetic diseases (such as cystic fibrosis), and complex multifactorial disorders (such as epilepsy). Finally, they promise to contribute to regenerative medicine, with the goal of producing functional biological structures that can be transplanted into patients [3][4][5][6][7][8][9].

## 2. Approaches to Generate a Patient-Derived Human Organoid: Surgical Resections, Liquid Biopsy, and iPSC-Derivation

The first organoids were developed from tissues of animal models. However, some biological aspects are unique to humans, and as a consequence, these models show limitations recapitulating human pathology. In that sense, PDOs emerged as a model to study cancer, infectious diseases, and inheritable genetic disorders [10][11][12][13]. The generation of organoids requires the use of stem cells, which can be either (a) pluripotent stem cells (PSCs)—embryonic stem cells (ESC) and induced pluripotent stem cells (iPSCs)—or (b) adult stem cells (ASCs) [7][8][9][13]. For instance, the source of PSC is restricted to iPSCs that are generated through the reprogramming of somatic cells, avoiding the ethical concerns of the use of ESC. iPSCs have the potential to generate all three germ layers. Differentiation into distinct cell and tissue types can be controlled in vitro by the sequential use of different factors that mimic in vivo organ development [7][11]. In contrast, ASCs can be obtained from tissues with regenerative ability, and they have a limited differentiation potential. In that case, the starting material for the generation of the organoids is normal or malignant human tissue that can be obtained from surgical resection or biopsy [10]. In fact, the generation of organoids from these sources allow the expansion and maintenance of this valuable material. The development of the organoids requires the use of specific growth factors depending on the tissue of origin, and they are mainly restricted to the growth of epithelial cells [14]. For that reason, the ASC-derived organoids are less complex than the iPSCs-derived, which might include mesenchymal and epithelial constituents [12]. On the other hand, tissue-derived organoids may recapitulate the genetic and epigenetic signature of the original organ [14], but iPSCs can lose this kind of information due to the dedifferentiation process required for the establishment of the cell line [15], thus hampering the use of iPSCs for preclinical models. In addition, liquid biopsies contain circulating tumor cells (CTCs), which although scarce in material could be good candidates to generate 3D structures, expanding the approaches that could be used to generate organoids [16]. In any case, the procedures to establish organoids rely on the self-renewal and differentiation of tissue-resident stem cells that expand in culture and self-organize into complex three-dimensional structures. Once established, organoids can be initiated from cryopreserved material, cultured using largely traditional cell culture techniques and equipment, and then expanded and cryopreserved for future use [17].

## 3. Organoids as a Model to Study Aging Signature across Tissues

Aging is the major risk factor for most chronic diseases. As a consequence of an increase in lifespan over the years, the elderly population is growing [18]. Frequently, this extension in longevity is not being accompanied with an increase in health-span [19]. Therefore, studying the underlying mechanisms of aging and developing interventions that target the aging process has become a priority field of research for most of the governments and private research agencies worldwide [19].

Organoids might provide a new valuable tool to model the changes that occur during aging across tissues and to study the development of age-associated diseases. Impaired processes and/or damage at the molecular and cellular levels accumulate as we age, leading to a decrease in the reserve capacity or resilience, ultimately developing the aging phenotypes, which have been divided into four domains: body composition, energetic imbalance between availability and demand, homeostatic dysregulation, and neurodegeneration [18][19][20]. Genomic instability, telomere attrition, epigenetic alterations, loss of proteostasis, deregulated nutrient sensing, mitochondrial dysfunction, cellular senescence, stem cell exhaustion, and altered intercellular communication have been identified as important hallmarks of aging in mammals [20]. Modeling these hallmarks using organoids seems to be possible (Table 1).

**Table 1.** Summary of studies using organoids to model aging.

Type of Organoid	Addressed Hallmark of Aging	Main Findings	Reference
Gut	Stem cell exhaustion; deregulated nutrient sensing	Lower O.F.E.; altered crypt formation	[21][22][23][24][25]
	Epigenetic changes; cellular senescence	Increased senescence markers; altered DNA methylation	[21][26][27]
	Stem cell exhaustion; deregulated nutrient sensing	CR increased O.F.E.; reduced mTOR signaling	[28][29]
	Stem cell exhaustion; deregulated nutrient sensing	NR supplementation increased O.F.E.	[30]
	Altered intercellular communication	Chronic inflammation led to NF-κB activation and cellular transformation	[31]
Liver	Genomic instability	Tissue-specific mutational profile; tumor development	[32][33]
	Genomic instability	Tissue-specific mutational profile; tumor development	[32][34]
Skin	Cellular senescence; altered intercellular communication	Increased senescence markers; decreased ECM synthesis	[35][36][37]
	Cellular senescence; altered intercellular communication	Adipose stem cells prevent skin senescence	[38]
	Altered intercellular communication	Altered TGF-β/Smad signaling	[39]
Tendon	Stem cell exhaustion; cellular senescence; altered intercellular communication	Lower O.F.E; decreased ECM synthesis; increased senescent markers	[40]
Lung	Stem cell exhaustion; cellular senescence; telomer attrition	Lower O.F.E; shortened telomeres; increased senescent markers	[41]
Breast	Genomic instability	Tumor development	[42]
Gastric	Genomic instability; epigenetic changes; altered intercellular communication	PDO characterization; altered Wnt signaling	[43]
Pancreatic	Genomic instability; altered intercellular communication	PDO characterization; altered Wnt signaling	[44]
Brain	Loss of proteostasis	Amyloid plaques and tau aggregates	[45][46]

O.F.E.: organoid formation efficiency; CR: calorie restriction; NR: nicotinamide riboside; ECM: extracellular matrix; PDO: patient-derived organoids.

For instance, aging is associated with a progressive loss of muscle mass and strength and a decline in neurophysiological functions, due to a gradual loss of motor neurons [47]. Recently, neuromuscular 3D organoids in vitro have been developed successfully to examine the roles of human autoantibodies in the pathogenesis of myasthenia gravis [48]. Therefore, with this model we could test whether changes in the neuromuscular junction precede or follow the decline of muscle mass and strength associated with aging. One of the most common approaches to study aging with organoids is deriving organoids from young and old donors. Reduced organoid formation efficiency has been described for aged mice and humans compared to their younger counterparts, which was associated with stem cell dysfunction and epigenetic changes resulting in a reduction in Wnt signaling [21][22][23][24][25]. Aged organoids also show increased levels of senescence markers such as p21 and p16, as well as decreased DNA methyltransferases [21]. Fasting and calorie restriction have been proposed as an anti-aging strategy and have been validated in organoids. Mihaylova et al. [25] showed that fasting for 24 h increased organoid formation and self-renewal potential of organoids derived from old mice. Similarly, improved organoid formation efficiency was observed in calorie-restricted mice, identifying the mTORC1 signaling in Paneth cells [28] and SIRT1 in intestinal stem cells [29][30] as key pathways to ameliorate stem cell function. On the other hand, long-term inflammation has also been modeled in intestinal organoids, and NF-κB signaling has been proposed as a driver of cellular transformation [31].

Skin equivalents (SE) consisting of a 3D culture of fibroblasts and keratinocytes have been developed to study skin aging, either by inducing senescence in vitro or by isolating cells from aged donors. SE have successfully modeled some features of skin aging such as decreased ECM synthesis, cellular loss, and thinner epidermal layer [35]. SE derived from aged donors show some of these histological features of aging [36][38], and p16 has been identified as a driver of these

changes, as demonstrated by the modulation of p16 levels in young- and old-derived SE [36]. Extended cultures of SE also recapitulate normal skin aging, including p16 induction [37]. Moreover, Metral et al. [38] showed that the addition of adipose-derived stem cells delayed the expression of senescence markers in SE. SE have also been used to analyze the role of the microenvironment in age-associated changes in the skin. Collagen fragmentation has been described as an important factor for skin aging. Aged fibroblasts detach from the ECM, resulting in impaired TGF- $\beta$  signaling and ultimately leading to a decrease in the synthesis of ECM components [39][49], further aggravating skin aging and fragility. Senescent melanocytes contribute to skin aging by inducing telomere damage and decreased proliferation of keratinocytes in a paracrine manner [50], while senescent fibroblasts induce changes in the secretome of SE, increasing IL-6, IL-1 $\alpha$ , and granulocyte macrophage colony-stimulating factor levels, which contribute to a decreased epidermal layer [51]. ECM age-associated changes have also been analyzed with tendon-derived organoids. Yan et al. [40] observed decreased organoid formation efficiency in organoids from aged donors, lower cell density, and decreased matrix deposition. These changes were associated with higher levels of the senescent markers p21 and p16 and stem cell dysfunction.

Some reports show that human intestinal organoids preserve their original DNA methylation pattern and their epigenetic age when cultured in vitro [26][52]. Notably, Lewis et al. [26] found that the epigenetic age of colon-derived organoids matched the actual age of the donor, while organoids derived from the small intestine showed a slight decrease in epigenetic age. Moreover, cultured mouse colon organoids seem to recapitulate age-associated epigenetic changes [27], opening the possibility of using organoids as a plausible model to study changes in DNA methylation patterns with aging. A lower organoid formation efficiency was also observed in organoids derived from aged alveolar epithelial type II cells. Furthermore, fibrotic cells and cells with shortened telomeres also decreased the organoid formation rate, which was associated with increased senescence and decreased stem cell function due to the activation of the Wnt signaling pathway [41]. Organoids have also been used to model age-associated diseases. For instance, 3D cultures have been developed to study osteoarthritis [53], macular degeneration [54], uterine leiomyoma [55], or Alzheimer's disease [56], which will be discussed in the next section. Before organoids can become important aging models, some issues must be considered such as the lower efficiency found when reprogramming cells to form organoids from aged donors compared to young ones or the increase in senescence rate when cultured for long time [57][58][59].

## 4. Application of Organoids in Age-Related Diseases: Cancer, Alzheimer's, and Parkinson's Disease

As we have stated along the review, organoids have emerged as an invaluable tool in biomedical research and have been extensively developed in the cancer field [60][61]. Other age-related diseases, such as Alzheimer's Disease (AD) and Parkinson's Disease (PD), could benefit from this technology.

### 4.1. Cancer

Cancer is considered an age-related disease because its incidence might be explained by the combination of (a) the accumulation of mutations in tissues throughout life and (b) the alterations of the tissue microenvironment that play a role as a selective pressure over them [62]. Moreover, the technical developments primarily achieved in organoid-based cancer research paves the way to the study of other age-related diseases by a patient-derived organoid approach [63]. In that sense, the expansion of clonal organoids from a single stem cell generates distinct mutational signatures, which allow the study of tumor genomic evolution [61]. Blokzijl and colleagues assessed the mutation accumulation in ASCs of the colon, the small intestine, and the liver, showing similar rates among them (36 mutations per year) but with tissue-specific mutational profiles and different cycling rates [32].

Once the candidate driver mutations are described, organoids have the potential to functionally validate them. Thus, wild-type organoids can be genetically engineered to model cancer initiation and progression recapitulating the oncogenic process. CRISPR/Cas9 technology was first used to introduce mutations in the most frequently mutated genes in colorectal cancer (*APC*, *P53*, *KRAS*, and *SMAD4*) in human small intestinal organoids in order to recapitulate the development of colorectal cancer [33]. Similarly, the knockout of tumor suppressor genes can recapitulate tumorigenesis in liver and breast human-derived organoids [34][42]. Interestingly, organoids can also, in part, reproduce the growth factor dependency of tumor cells within their microenvironment. With this approach, the genotype–phenotype correlation of the growth factor dependency was described in gastric cancer and pancreatic tumors allowing the description of tumor subtypes [43][44]. Finally, it is known that a link exists between cancer and infectious agents to which we can be exposed during our lifespan. Thus, the co-culture of organoids with pathogens renders the opportunity to mimic the host–pathogen interactions and pathogenic-induced oncogenesis. In this regard, human gastric organoids were microinjected with *Helicobacter pylori* to model the role of the microorganism in epithelial signaling and proliferation [64]. Recently, the role of human papillomavirus (HPV) in carcinogenesis was studied in ecto- and endocervical-derived organoids [65].

Besides, organoids can be co-cultured with microbiota [66], which is relevant due to its role in cancer and other aging-related processes [67].

## 4.2. Alzheimer's Disease and Parkinson's Disease

Even though Alzheimer's Disease (AD) is one of the leading causes of death worldwide, especially in late life, to date there are currently no available drug treatments to cure the disease. Similarly, Parkinson's Disease (PD) is the second most frequent neurodegenerative disorder after AD. Organoid systems based on human pluripotent stem cells (hPSCs) and neural stem/precursor cells (NSCs) have shown a promising potential to model neurodegenerative diseases, including AD and PD [45][68][69][70]. 3D brain organoid systems generated from hPSCs can recapitulate important features of AD pathophysiology, such as amyloid plaques and neurofibrillary tangle-like structures [46][68]. Similarly, NSCs utilized to derive human midbrain-specific organoids (hMO) show abundant neurons with dopaminergic identity, thus electrophysiologically functional neurons, as well as astroglia and oligodendrocyte differentiation [70]. Most of the reports using brain organoids in PD or AD have been focused on genetic risk factors, relying on CRISPR/Cas9 gene editing. In both cases, these organoids models have been used to test the efficacy of pharmacological agents in disease progression [45][69].

---

## References

1. Barbuzano. Available online: <https://hsci.harvard.edu/organoids> (accessed on 10 July 2021).
2. Yang, H.; Sun, L.; Liu, M.; Mao, Y. Patient-derived organoids: A promising model for personalized cancer treatment. *Gastroenterol. Rep. (Oxf.)* 2018, 6, 243–245.
3. Michels, B.E.; Mosa, M.H.; Streibl, B.I.; Zhan, T.; Menche, C.; Abou-El-Ardat, K.; Darvishi, T.; Czlonka, E.; Wagner, S.; Winter, J.; et al. Pooled In Vitro and In Vivo CRISPR-Cas9 Screening Identifies Tumor Suppressors in Human Colon Organoids. *Cell Stem Cell* 2020, 26, 782–792.e7.
4. Bock, C.; Boutros, M.; Camp, J.G.; Clarke, L.; Clevers, H.; Knoblich, J.A.; Liberali, P.; Regev, A.; Rios, A.C.; Stegle, O.; et al. The Organoid Cell Atlas. *Nat. Biotechnol.* 2020, 39, 13–17.
5. Kim, J.; Koo, B.K.; Knoblich, J.A. Human organoids: Model systems for human biology and medicine. *Nat. Rev. Mol. Cell Biol.* 2020, 21, 571–584.
6. Gunti, S.; Hoke, A.T.K.; Vu, K.P.; London, N.R., Jr. Organoid and Spheroid Tumor Models: Techniques and Applications. *Cancers* 2021, 13, 874.
7. Nunes, A.S.; Barros, A.S.; Costa, E.C.; Moreira, A.F.; Correia, I.J. 3D tumor spheroids as in vitro models to mimic in vivo human solid tumors resistance to therapeutic drugs. *Biotechnol. Bioeng.* 2018, 10, 26845.
8. Rossi, G.; Manfrin, A.; Lutolf, M.P. Progress and potential in organoid research. *Nat. Rev. Genet.* 2018, 19, 671–687.
9. Yu, J.; Vodyanik, M.A.; Smuga-Otto, K.; Antosiewicz-Bourget, J.; Frane, J.L.; Tian, S.; Nie, J.; Jonsdottir, G.A.; Ruotti, V.; Stewart, R.; et al. Induced pluripotent stem cell lines derived from human somatic cells. *Science* 2007, 318, 1917–1920.
10. Lancaster, M.A.; Huch, M. Disease modelling in human organoids. *Dis. Model. Mech.* 2019, 12, 39347.
11. Tindle, C.; Fuller, M.; Fonseca, A.; Taheri, S.; Ibeawuchi, S.R.; Beutler, N.; Katkar, G.D.; Claire, A.; Castillo, V.; Hernandez, M.; et al. Adult stem cell-derived complete lung organoid models emulate lung disease in COVID-19. *eLife* 2021, 10, e66417.
12. Azar, J.; Bahmad, H.F.; Daher, D.; Moubarak, M.M.; Hadadeh, O.; Monzer, A.; Al Bitar, S.; Jamal, M.; Al-Sayegh, M.; Abou-Kheir, W. The Use of Stem Cell-Derived Organoids in Disease Modeling: An Update. *Int. J. Mol. Sci.* 2021, 22, 7667.
13. Clevers, H. Modeling Development and Disease with Organoids. *Cell* 2016, 165, 1586–1597.
14. Fujii, M.; Sato, T. Somatic cell-derived organoids as prototypes of human epithelial tissues and diseases. *Nat. Mater.* 2021, 20, 156–169.
15. Liang, G.; Zhang, Y. Genetic and epigenetic variations in iPSCs: Potential causes and implications for application. *Cell. Stem Cell* 2013, 13, 149–159.
16. Yang, C.; Xia, B.R.; Jin, W.L.; Lou, G. Circulating tumor cells in precision oncology: Clinical applications in liquid biopsy and 3D organoid model. *Cancer Cell Int.* 2019, 19, 12935.
17. Clinton, J.; McWilliams-Koeppen, P. Initiation, Expansion, and Cryopreservation of Human Primary Tissue-Derived Normal and Diseased Organoids in Embedded Three-Dimensional Culture. *Curr. Protoc. Cell Biol.* 2019, 82, e66.

18. Ferrucci, L.; Gonzalez-Freire, M.; Fabbri, E.; Tanaka, T.; Moore, Z.; Salimi, S.; Sierra, F.; de Cabo, R. Measuring biological aging in humans: A quest. *Aging Cell*. 2020, 19, e13080.
19. Gonzalez-Freire, M.; Diaz-Ruiz, A.; Hauser, D.; Martinez-Romero, J.; Ferrucci, L.; Bernier, M.; de Cabo, R. The road ahead for health and lifespan interventions. *Ageing Res. Rev.* 2020, 59, 101037.
20. López-Otín, C.; Blasco, M.A.; Partridge, L.; Serrano, M.; Kroemer, G. The hallmarks of aging. *Cell* 2013, 153, 1194–1217.
21. Uchida, R.; Saito, Y.; Nogami, K.; Kajiyama, Y.; Suzuki, Y.; Kawase, Y.; Nakaoka, T.; Muramatsu, T.; Kimura, M.; Saito, H. Epigenetic silencing of Lgr5 induces senescence of intestinal epithelial organoids during the process of aging. *NPJ Aging Mech. Dis.* 2019, 5, 1–5.
22. He, D.; Wu, H.; Xiang, J.; Ruan, X.; Peng, P.; Ruan, Y.; Chen, Y.G.; Wang, Y.; Yu, Q.; Zhang, H.; et al. Gut stem cell aging is driven by mTORC1 via a p38 MAPK-p53 pathway. *Nat. Commun.* 2020, 11, 1–13.
23. Nalapareddy, K.; Nattamai, K.J.; Kumar, R.S.; Karns, R.; Wikenheiser-Brokamp, K.A.; Sampson, L.L.; Mahe, M.M.; Sundaram, N.; Yacyshyn, M.B.; Yacyshyn, B.; et al. Canonical Wnt Signaling Ameliorates Aging of Intestinal Stem Cells. *Cell Rep.* 2017, 18, 2608–2621.
24. Pentimikko, N.; Iqbal, S.; Mana, M.; Andersson, S.; Cognetta, A.B.; Suci, R.M.; Roper, J.; Luopajarvi, K.; Markelin, E.; Gopalakrishnan, S.; et al. Notum produced by Paneth cells attenuates regeneration of aged intestinal epithelium. *Nature* 2019, 571, 398–402.
25. Mihaylova, M.M.; Cheng, C.W.; Cao, A.Q.; Tripathi, S.; Mana, M.D.; Bauer-Rowe, K.E.; Abu-Remaileh, M.; Clavain, L.; Erdemir, A.; Lewis, C.A.; et al. Fasting Activates Fatty Acid Oxidation to Enhance Intestinal Stem Cell Function during Homeostasis and Aging. *Cell. Stem Cell* 2018, 22, 769–778.e4.
26. Lewis, S.K.; Nachun, D.; Martin, M.G.; Horvath, S.; Coppola, G.; Jones, D.L. DNA Methylation Analysis Validates Organoids as a Viable Model for Studying Human Intestinal Aging. *CMGH* 2020, 9, 527–541.
27. Tao, Y.; Kang, B.; Petkovich, D.A.; Bhandari, Y.R.; In, J.; Stein-O'Brien, G.; Kong, X.; Xie, W.; Zachos, N.; Maegawa, S.; et al. Aging-like Spontaneous Epigenetic Silencing Facilitates Wnt Activation, Stemness, and Braf V600E-Induced Tumorigenesis. *Cancer Cell* 2019, 35, 315–328.e6.
28. Yilmaz, Ö.H.; Katajisto, P.; Lamming, D.W.; Gültekin, Y.; Bauer-Rowe, K.E.; Sengupta, S.; Birsoy, K.; Dursun, A.; Onur Yilmaz, V.; Selig, M.; et al. mTORC1 in the Paneth cell niche couples intestinal stem-cell function to calorie intake. *Nature* 2012, 486, 490–495.
29. Igarashi, M.; Guarente, L. mTORC1 and SIRT1 Cooperate to Foster Expansion of Gut Adult Stem Cells during Calorie Restriction. *Cell* 2016, 166, 436–450.
30. Igarashi, M.; Miura, M.; Williams, E.; Jaksch, F.; Kadowaki, T.; Yamauchi, T.; Guarente, L. NAD<sup>+</sup> supplementation rejuvenates aged gut adult stem cells. *Aging Cell* 2019, 18, 1–10.
31. Hibiya, S.; Tsuchiya, K.; Hayashi, R.; Fukushima, K.; Horita, N.; Watanabe, S.; Shirasaki, T.; Nishimura, R.; Kimura, N.; Nishimura, T.; et al. Long-term Inflammation Transforms Intestinal Epithelial Cells of Colonic Organoids. *J. Crohns. Colitis* 2017, 11, 621–630.
32. Blokzijl, F.; de Ligt, J.; Jager, M.; Sasselli, V.; Roerink, S.; Sasaki, N.; Huch, M.; Boymans, S.; Kuijk, E.; Prins, P. Tissue-specific mutation accumulation in human adult stem cells during life. *Nature* 2016, 538, 260–264.
33. Drost, J.; van Jaarsveld, R.H.; Ponsioen, B.; Zimmerlin, C.; van Boxtel, R.; Buijs, A.; Sachs, N.; Overmeer, R.M.; Offerhaus, G.J.; Begthel, H. Sequential cancer mutations in cultured human intestinal stem cells. *Nature* 2015, 521, 43–47.
34. Artigiani, B.; van Voorthuisen, L.; Lindeboom, R.G.; Seinstra, D.; Heo, I.; Tapia, P.; López-Iglesias, C.; Postrach, D.; Dayton, T.; Oka, R.; et al. Probing the Tumor Suppressor Function of BAP1 in CRISPR-Engineered Human Liver Organoids. *Cell Stem Cell*. 2019, 24, 927–943.e6.
35. Diekmann, J.; Alili, L.; Scholz, O.; Giesen, M.; Holtkötter, O.; Brenneisen, P. A three-dimensional skin equivalent reflecting some aspects of in vivo aged skin. *Exp. Dermatol.* 2016, 25, 56–61.
36. Adamus, J.; Aho, S.; Meldrum, H.; Bosko, C.; Lee, J.M. P16INK4A influences the aging phenotype in the living skin equivalent. *J. Invest. Dermatol.* 2014, 134, 1131–1133.
37. Dos Santos, M.; Metral, E.; Boher, A.; Rousselle, P.; Thepot, A.; Damour, O. In vitro 3-D model based on extending time of culture for studying chronological epidermis aging. *Matrix Biol.* 2015, 47, 85–97.
38. Metral, E.; Dos Santos, M.; Thépot, A.; Rachidi, W.; Mojallal, A.; Auxenfans, C.; Damour, O. Adipose-derived Stem Cells Promote Skin Homeostasis and Prevent its Senescence in an In vitro Skin Model. *J. Stem Cell Res. Ther.* 2014, 4, 194.

39. Fisher, G.J.; Shao, Y.; He, T.; Qin, Z.; Perry, D.; Voorhees, J.J.; Quan, T. Reduction of fibroblast size/mechanical force down-regulates TGF- $\beta$  type II receptor: Implications for human skin aging. *Aging Cell* 2016, 15, 67–76.
40. Yan, Z.; Yin, H.; Brochhausen, C.; Pfeifer, C.G.; Alt, V.; Docheva, D. Aged Tendon Stem/Progenitor Cells Are Less Competent to Form 3D Tendon Organoids Due to Cell Autonomous and Matrix Production Deficits. *Front. Bioeng. Biotechnol.* 2020, 8, 406.
41. Lehmann, M.; Hu, Q.; Hu, Y.; Hafner, K.; Costa, R.; van den Berg, A.; Königshoff, M. Chronic WNT/ $\beta$ -catenin signaling induces cellular senescence in lung epithelial cells. *Cell. Signal.* 2020, 70, 109588.
42. Artegiani, B.; van Voorthuisen, L.; Lindeboom, R.; Seinstra, D.; Heo, I.; Tapia, P.; López-Iglesias, C.; Postrach, D.; Dayton, T.; Oka, R. Modeling Breast Cancer Using CRISPR-Cas9-Mediated Engineering of Human Breast Organoids. *J. Natl. Cancer Inst.* 2020, 112, 540–544.
43. Nanki, K.; Toshimitsu, K.; Takano, A.; Fujii, M.; Shimokawa, M.; Ohta, Y.; Matano, M.; Seino, T.; Nishikori, S.; Ishikawa, K. Divergent Routes toward Wnt and R-spondin Niche Independency during Human Gastric Carcinogenesis. *Cell* 2018, 174, 856–869.e17.
44. Seino, T.; Kawasaki, S.; Shimokawa, M.; Tamagawa, H.; Toshimitsu, K.; Fujii, M.; Ohta, Y.; Matano, M.; Nanki, K.; Kawasaki, K. Human Pancreatic Tumor Organoids Reveal Loss of Stem Cell Niche Factor Dependence during Disease Progression. *Cell. Stem Cell* 2018, 22, 454–467.e6.
45. Gonzalez, C.; Armijo, E.; Bravo-Alegria, J.; Becerra-Calixto, A.; Mays, C.E.; Soto, C. Modeling amyloid beta and tau pathology in human cerebral organoids. *Mol. Psychiatry.* 2018, 23, 2363–2374.
46. Choi, S.H.; Kim, Y.H.; Hebisch, M.; Sliwinski, C.; Lee, S.; D'Avanzo, C.; Chen, H.; Hooli, B.; Asselin, C.; Muffat, J. A three-dimensional human neural cell culture model of Alzheimer's disease. *Nature* 2014, 515, 274–278.
47. Gonzalez-Freire, M.; de Cabo, R.; Studenski, S.A.; Ferrucci, L. The Neuromuscular Junction: Aging at the Crossroad between Nerves and Muscle. *Front. Aging Neurosci.* 2014, 6, 208.
48. Faustino Martins, J.M.; Fischer, C.; Urzi, A.; Vidal, R.; Kunz, S.; Ruffault, P.L.; Kabuss, L.; Hube, I.; Gazzero, E.; Birchmeier, C.; et al. Self-Organizing 3D Human Trunk Neuromuscular Organoids published correction appears in *Cell Stem Cell*. *Cell. Stem Cell* 2020, 26, 172–186.e6.
49. Quan, T.; Wang, F.; Shao, Y.; Rittié, L.; Xia, W.; Orringer, J.S.; Voorhees, J.J.; Fisher, G.J. Enhancing structural support of the dermal microenvironment activates fibroblasts, endothelial cells, and keratinocytes in aged human skin in vivo. *J. Investig. Dermatol.* 2013, 133, 658–667.
50. Victorelli, S.; Lagnado, A.; Halim, J.; Moore, W.; Talbot, D.; Barrett, K.; Chapman, J.; Birch, J.; Ogrodnik, M.; Meves, A.; et al. Senescent human melanocytes drive skin ageing via paracrine telomere dysfunction. *EMBO J.* 2019, 38, 1–18.
51. Weinmüllner, R.; Zbiral, B.; Becirovic, A.; Stelzer, E.M.; Nagelreiter, F.; Schosserer, M.; Lämmermann, I.; Liendl, L.; Lang, M.; Terlecki-Zaniewicz, L.; et al. Organotypic human skin culture models constructed with senescent fibroblasts show hallmarks of skin aging. *NPJ Aging Mech. Dis.* 2020, 6, s41514.
52. Kraiczy, J.; Nayak, K.M.; Howell, K.J.; Ross, A.; Forbester, J.; Salvestrini, C.; Mustata, R.; Perkins, S.; Andersson-Rolf, A.; Leenen, E.; et al. DNA methylation defines regional identity of human intestinal epithelial organoids and undergoes dynamic changes during development. *Gut* 2019, 68, 49–61.
53. Lozito, T.P.; Alexander, P.G.; Lin, H.; Gottardi, R.; Cheng, A.W.; Tuan, R.S. Osteoarthritis is a disease of the osteochondral junction. *Osteoarthr. Cartil.* 2010, 4, 1–6.
54. Shokoohmand, A.; Jeon, J.E.; Theodoropoulos, C.; Baldwin, J.G.; Huttmacher, D.W.; Feigl, B. A Novel 3D Cultured Model for Studying Early Changes in Age-Related Macular Degeneration. *Macromol. Biosci.* 2017, 17, 2–9.
55. Xie, J.; Xu, X.; Yin, P.; Li, Y.; Guo, H.; Kujawa, S.; Chakravarti, D.; Bulun, S.; Kim, J.J.; Wei, J.J. Application of ex-vivo spheroid model system for the analysis of senescence and senolytic phenotypes in uterine leiomyoma. *Lab. Invest.* 2018, 98, 1575–1587.
56. Papaspyropoulos, A.; Tsolaki, M.; Foroglou, N.; Pantazaki, A.A. Modeling and Targeting Alzheimer's Disease With Organoids. *Front. Pharmacol.* 2020, 11, 1–8.
57. Mahmoudi, S.; Brunet, A. Aging and reprogramming: A two-way street. *Curr. Opin. Cell Biol.* 2012, 24, 744–756.
58. Lo Sardo, V.; Ferguson, W.; Erikson, G.A.; Topol, E.J.; Baldwin, K.K.; Torkamani, A. Influence of donor age on induced pluripotent stem cells. *Nat. Biotechnol.* 2017, 35, 69–74.
59. Zhang, W.; Qu, J.; Liu, G.H.; Belmonte, J.C.I. The ageing epigenome and its rejuvenation. *Nat. Rev. Mol. Cell Biol.* 2020, 21, 137–150.
60. Drost, J.; Clevers, H. Organoids in cancer research. *Nat. Rev. Cancer* 2018, 18, 407–418.

61. Lo, Y.H.; Karlsson, K.; Kuo, C.J. Applications of Organoids for Cancer Biology and Precision Medicine. *Nat. Cancer* 2020, 1, 761–773.
62. Laconi, E.; Marongiu, F.; DeGregori, J. Cancer as a disease of old age: Changing mutational and microenvironmental landscapes. *Br. J. Cancer* 2020, 122, 943–952.
63. Xu, H.; Lyu, X.; Yi, M.; Zhao, W.; Song, Y.; Wu, K. Organoid technology and applications in cancer research. *J. Hematol. Oncol.* 2018, 11, 13045.
64. McCracken, K.W.; Catá, E.M.; Crawford, C.M.; Sinagoga, K.L.; Schumacher, M.; Rockich, B.E.; Tsai, Y.H.; Mayhew, C.N.; Spence, J.R.; Zavros, Y.; et al. Modelling human development and disease in pluripotent stem-cell-derived gastric organoids. *Nature* 2014, 516, 400–404.
65. Löhmußaar, K.; Oka, R.; Espejo Valle-Inclán, J.; Smits, M.; Wardak, H.; Korving, J.; Begthel, H.; Proost, N.; van de Ven, M.; Kranenburg, O.W. Patient-derived organoids model cervical tissue dynamics and viral oncogenesis in cervical Cancer. *Cell Stem Cell* 2021, 28, 1380–1396.e6.
66. Puschhof, J.; Pleguezuelos-Manzano, C.; Martínez-Silgado, A.; Akkerman, N.; Saftien, A.; Boot, C.; de Waal, A.; Beumer, J.; Dutta, D.; Heo, I.; et al. Intestinal organoid cocultures with microbes published online ahead of print. *Aug. Nat. Protoc.* 2021, 10, 41596.
67. Boehme, M.; Guzzetta, K.E.; Bastiaanssen, T.F.; van de Wouw, M.; Moloney, G.M.; Gual-Grau, A.; Spichak, S.; Olavarría-Ramírez, L.; Fitzgerald, P.; Morillas, E.; et al. Microbiota from young mice counteracts selective age-associated behavioral deficits. *Nat. Aging* 2021, 1, 666–676.
68. Bi, F.C.; Yang, X.H.; Cheng, X.Y.; Deng, W.B.; Guo, X.L.; Yang, H.; Wang, Y.; Li, J.; Yao, Y. Optimization of cerebral organoids: A more qualified model for Alzheimer's disease research. *Transl. Neurodegener.* 2021, 10, 27.
69. García-Leon, J.A.; Cáceres-Palomo, L.; Sánchez-Mejías, E.; Mejías-Ortega, M.; Nuñez-Díaz, C.; Fernández-Valenzuela, J.J.; Sánchez-Varo, R.; Davila, J.C.; Vitorica, J.; Gutierrez, A. Human Pluripotent Stem Cell-Derived Neural Cells as a Relevant Platform for Drug Screening in Alzheimer's Disease. *Int. J. Mol. Sci.* 2020, 21, 6867.
70. Smits, L.M.; Schwamborn, J.C. Midbrain Organoids: A New Tool to Investigate Parkinson's Disease. *Front. Cell Dev. Biol.* 2020, 8, 359.

---

Retrieved from <https://encyclopedia.pub/entry/history/show/35924>