

Mycorrhizae and Water Quality

Subjects: Environmental Sciences

Contributors:  Jess Rubin,  Josef Gorres



Submitted by:  Jess Rubin

Definition

Mycorrhizae fungi are 400 million-year-old plant symbionts whose evolutionary success has been attributed to their ability to expand the rhizosphere of plants, enabling greater uptake of nutrients from surrounding soils in exchange for photosynthate provided by their host plants.

1. Introduction

Early research indicates that the improved phosphorus uptake by mycorrhizal plants can reduce P leaching. This may come about because the uptake of P in roots colonized by mycorrhizal fungi can be 3–5 times higher than in non-mycorrhizal roots ^[1]. Mycorrhizal inoculation in agricultural production reduces the amount of P fertility amendments required for plant growth. So called legacy phosphorus, i.e. phosphorus stored in the soil from previous applications of fertility amendments, becomes the source of plant nutrition. Avoiding further applications of phosphorus reduces the amount reaching water resources where phosphorus is a pollutant that causes eutrophication in freshwater lakes. The effectiveness of mycorrhizae in agricultural landscapes, however, is variable given the wide variety of farm management systems and practices which affect successful colonization of host plants. Nevertheless, Rillig et al. ^[2] advocates for the development of mycorrhizal technologies to enhance agroecosystems sustainably.

Mycorrhizal fungi are keystone mutualists in terrestrial ecosystems ^[3] whose ecological role in assisting recovery of severely disturbed ecosystems ^[4] is evident because they enhance P plant uptake in both crops and woody plants. Thus, they could play an important role in myco-phytoremediation of soil phosphorus which is often present in excessive amounts. This involves ecological engineering which harnesses nutrient exchange networks, within which mycorrhizae play an important role, crucial to ecosystem succession and resilience ^[5]. This strategy, though still relatively novel to the field of remediation, has tremendous potential to be applied in the burgeoning field of reconciliation ecology ^[6], which acknowledges that, while ecosystems cannot be completely restored to their original state, mitigation of degradation can return them to a new balance ^[7].

Of the seven groups of mycorrhizae, the two most common in agricultural and forested lands ^[8] are also the most likely to be employed in myco-phytoremediation: arbuscular mycorrhizae fungi (AMF) ^[9] and ectomycorrhizae (ECM). While AMF and ECM provide similar services to the plant (i.e., improved access to P) ^[1], their hyphae differ in architecture and in how they transfer P to the plant ^[10]. In the AMF, the transfer is accomplished intercellularly and via intracellular arbuscules from extra-radical hyphae that extend directly into the soil beyond plant rhizosphere depletion zones ^[11]. In ECM, the transfer occurs via intercellular Hartig net hyphal networks surrounding epidermal and cortex cells while outside of the mantle, extra radical mycelia form extensive nutrient-absorbing networks in the soil ^{[12][13]}. It is well established that AMF and ECM greatly enhance the uptake of immobile soil nutrients such as P and water by plant root hosts in exchange for carbohydrates supplied by the host plant ^{[14][15]} and improve soil properties. They also increase below- and above-ground biodiversity and provide pathogen resistance. This results in improved tree and shrub survival, better growth and establishment on moisture-, nutrient- and salt-stressed soils ^{[16][17][18][19]}. In addition, they facilitate plant succession ^{[20][21]}.

Additionally, when planting into AMF grasslands, tree and shrub species' growth and survival is improved by inoculation with ECM specific to the species planted ^[22]. ECM presence can support native trees to endure aggressive non-native species' presence ^[23] as well as play a critical role in the restoration of

degraded sites [24]. Mycorrhizae can mitigate P pollution at each stage of the three-pronged paradigm of water resource protection: source reduction via decreasing P amendment amounts needed, contamination event reduction by decreasing erosion through improved soil structure and vegetation establishment, and pollutant interception via redirecting P into plant roots out of soil water.

2. Mycorrhizae, Landscapes and Soils

Any design of a phosphorus mitigation strategy that involves mycorrhizae has to consider landscape position and soils which affect P availability and fate. In an ideal agricultural landscape, production fields are separated from water courses by a forested (or otherwise vegetated) riparian buffer [25], that attenuates the increased P in leachate when high fertilizer or manure P is applied [26]. Each landscape element in the catena has a different role to play in P mitigation. Drainage class and vegetation need to be considered as variables for establishment of mycorrhizal communities. The mycorrhizal communities likely differ between high organic matter riparian forest including both AMF and ECM and the agricultural field dominated by AMF [27]. Soil drainage class, tantamount to location in a toposequence, *per se* may not affect mycorrhizal plant infections. In a study on soybean fields stretching across three soil drainage classes (poorly, somewhat poorly, and moderately well drained), more AMF spores were found in the more poorly drained than the better drained soils. But, there was no discernible difference in colonization of plant roots [28]. In agricultural systems where flooding diminishes vegetation, crops following the flood are P deficient early in the season. The lack of hosts during flooding may result in lower colonization rates by AMF and thus less P uptake [29]. Lack of vegetation during flooding is not likely to occur in forested riparian forests [30], and agricultural fields can be managed to avoid fallow conditions by planting rotations and cover crops that host mycorrhizae [31].

However, drainage class may still enter into any myco-phytoremediation design because prolonged flooding in wetland riparian buffer, remobilizes P adsorbed to soil colloids. In particular, under anaerobic conditions ferric iron is reduced, releasing phosphate that would otherwise be strongly sorbed to ferric oxides [32]. It is not clear whether mycorrhizae can help with recovering P released in this way.

In terms of the water mitigation paradigm, agricultural fields would be targets of source reduction as they are the primary recipients of P. However, in an area where agriculture was practiced for decades, it is likely the soil has sufficient P to be a source itself [33]. Here myco-phytoremediation may reduce legacy P.

High SRP concentrations in agricultural fields are likely to reduce mycorrhizal infections [34]. Therefore, the use and amount of fertilizer P should be judicious [35][36]. Management of agricultural lands should consider the use of alternatives to inorganic P fertilizer to promote mycorrhizal growth and colonization [37].

Consequently, managing the field for mycorrhizae can reduce the amount of P fertilizer needed to achieve yield goals [31]. This includes reducing tilling and maintaining hosts by implementing crop rotation, and also choosing crops with root architecture efficient in accessing sufficient P and forming a symbiosis with AMF [38].

Oka [39] found that P application on soy beans could be reduced from 150 to 50 kg P ha⁻¹ without yield loss when it followed wheat, an AMF mycorrhizal crop (*Triticum sativum*); than when followed by radish (*Raphanus sativus*), a non-mycorrhizal crop. The benefits may be due to better establishment of mycorrhizae-plant associations under the low soil-P supply in the early season with increased uptake of P ensuing [40]. Application of excessive fertilizer at this time of the growing season may inhibit mycorrhizal infections [40] and should be avoided. Mycorrhizal cover crops may thus have several benefits to the plant. First, they provide hosts for mycorrhizae and a source of organic P, scavenged between cash crops. In addition, over time, the amount of sediment-bound phosphorus lost by erosion will diminish. Consequently, downslope P accumulations in riparian areas are minimized.

Although agriculture can be regarded as a myco-phytoremediation system for legacy P, agricultural

practices affect mycorrhizae. The type and timing of tillage has been identified as one such factor. The role of fungi in plant nutrition and soil conservation is compromised when the formation and survival of propagules (i.e., spores, hyphae, colonized roots) are disrupted through tillage. Spores serve as “long-term” propagules when host plants are not present, whereas hyphae are the main source of inoculum when plants are present in undisturbed soil. Deep plowing can ‘dilute’ propagules, reducing plant root inoculation, especially in autumn when hyphae are detached from the host plant. Conservation tillage can protect survivability and inoculation, thereby improving soil aggregation and P uptake [42].

The structure and texture of soils is also an important factor in whether AMF has significant impacts on leaching and erosion. In agriculture, it is important to look at the relationship between fertilization and runoff. AMF significantly reduced nutrient leaching after rainfall events in sandy grassland soils [43]. This finding has important implications for soils with poor P sorption capacity such as sandy soils and other highly permeable soils or heavily manured soils [44], where P can be lost during rainfall events.

Furthermore, mycorrhizae can intercept P in soil solution before it leaves the root zone with deep percolation. In contrast to the many studies that assess the effect of mycorrhizae on plant uptake of P, only few of them report how mycorrhizae affect P leaching. This is usually not regarded as a major pathway of P export from a field because of the high affinity of phosphate [45] to soil surfaces. However, Asghari et al. [46] explained that sandy-textured soils are likely to provide little internal surfaces for P adsorption the major mechanism of P retention in soils. In addition, soils that receive high P fertilizer may also leach phosphate [26]. Water quality in freshwater bodies is sensitive to even small amounts of P [41] and thus leaching may have a significant effect. Asghari et al. [46] found that AMF can reduce leachate P from soil columns packed with a loamy sand. In another laboratory experiment Köhl and van der Heijden [47] found that the effect varied with AMF species probably due to differences in root colonization: the more root colonization the greater the growth of the plant and presumably the less P was leached. This is because AMF symbiosis assists plants with P uptake [48] through reaching beyond P depletion zones to access greater soil P reserves [49]. Plant response to mycorrhizal formation also depends upon the extent of mycorrhizal development [14]. It is not clear whether the results of these controlled laboratory studies are directly transferable to processes that occur in the field where many other factors are in play; more research is needed here.




Figure 1. Influence of mycorrhizae on phosphorus cycling processes and pools. Red and green arrows are processes influenced by mycorrhizae. Broken lines show the net direction of reactions due to mycorrhizal effects.

Mycorrhizae are involved in most aspects of P cycling as can be seen in Figure 1. Data from the literature show the effect of mycorrhizae on plant uptake, leachate and soil concentration. For example, plant uptake can be enhanced by between 40 and several 100s of percent, leachate P is reduced by up to 60% and extractable soil P by 15% in a growing season (Table 1). However, variations in both plant and

mycorrhizae species greatly influence P removal from soil and thus its concentration in leachate. The effects of mycorrhizae on phosphorus mitigation should be considered when investigating strategies for water quality improvement from upland source areas.

Table 1. The effect of mycorrhizae on plant uptake, leaching and soil P from studies carried out under different experimental conditions and with different objectives. Underscored show the physical quantity measured.

Study Context	Study Conditions	Phosphorus Quantity Measured	% Change with Mycorrhiza #	Location	Ref. #
Crop uptake	Agro ecosystem <i>Triticum aestivum</i> , AMF	<u>Phosphorus use efficiency</u>	+85-102%	Uttar Pradesh, Haryana, India	[50]
		<u>Shoot P concentration</u> [mg/g]			
		Field			
		<i>S. pulchra</i> ,	+22%		
		<i>A. barbata</i>	+68%		
		Greenhouse			
Growth of native grasses	Field ecosystem and pots in greenhouse, <i>Stipa pulchra</i> , <i>Avena barbata</i> , fungicide/no fungicide ***	<u>Shoot P concentration</u>		San Diego CA, USA	[22]
		<i>S. pulchra</i>	+1.6%		
		<i>A. barbata</i>	-11.8%		

Root concentration

S. pulchra +24%

A. barbata -15%

Plant P concentrations (%)

No Mulch +28%

Living Mulch +135%

Plant P (mg P/plant)

No mulch +17%

Living mulch +709%

Plant P concentration [mg/g]

Plant uptake [mg P/pot] +1525%

P plant uptake [mg P/pot]

Shoots +168%

Mulch Experiment

Pots, greenhouse
Trifolium repens
Zea Mays
Fungicide/no fungicide ***

Morioka, Japan [51]

Crop uptake

Pots, AMF, *Allium fistulosum*

Haguromachi, Japan [52]

Effect of mycorrhizosphere bacteria on plant uptake

Pots, corn (*Zea Mays*), AMF

Denmark [53]

		Roots	+234%		
Effect of AMF on P leaching	Packed columns, greenhouse, <i>Trifolium subterraneum</i> AMF	<u>Leachate P [mg]</u>			
		without added P	-60%		
		with added P.	0%		
		<u>Plant P [mg]</u>			
		without added P	+251%		
		with added P	-23%		
		<u>Plant uptake (mg P/plant)</u>			
		Hybrid			
		P3979	+8.4%		
		LRS	+19.1%		
LNS	+19.8%				
Effect of mycorrhizae on crop uptake and extractable soil P	Pot, greenhouse, corn (<i>Zea Mays</i>), AMF	<u>Mehlich 3 extractable Soil P Concentration [mg/kg]</u>			
		Hybrids, no P fertilizer			
				South Australia	[46]
				Quebec Canada	[54]

Leaching mitigation

Pots, greenhouses, *Phalaris aquatic*, AMF

P3979	-5.1%
LRS	-14.4%
LNS	-10.5%

Mehlich 3 extractable Soil P Concentration [mg/kg].

Hybrids, P fertilizer applied	ns
--------------------------------------	----

<u>Shoot P content (mg)</u>	+150%
-----------------------------	-------

<u>Root P content (mg)</u>	+168%
----------------------------	-------

Leachate [kg P/ha] ###

Particulate P	-11.1%
---------------	--------

Dissolved Organic P	-14.4%
---------------------	--------

SRP (PO ₄) *	-81%
--------------------------	------

Runoff [kg P/ha]

Particulate P	-11.1%
---------------	--------

Southeastern Australia [55]

P losses from field

Microcosms *Orya sativa* L AMF

Jiangsu, China [56]

		Dissolved Organic P	-4.95%	
		SRP (PO ₄) *	-11%	
		<u>P in leachate [mg] ###</u>		
		Pasture		
Nutrient cycling in presence of mycorrhizae	Microcosms, Heath and Pasture communities, AMF	Added NH ₄	-14.2%	
		Added NO ₃	-38.5%	[57]
		Heath		
		Added NH ₄	-68.4%	Switzerland
		Added NO ₃	-63.4%	
		<u>Reduction in leaching</u>		
Leaching from grasslands	Mesocosms, grassland, AMF	Low nutrient availability	~ 60%	[43]
		High nutrient availability	ns	
		<u>Leachate P [ug] ###</u>		
Climate Change Resilience	Mesocosms, grassland communities, AMF	Moderate rain	-149%	The Netherlands [58]

High rain -58%

Crop Uptake

Pots, *Allium fistulosum* (Welsh Onion) AMF

Shoot concentration

+88%

Tozawa, Japan

[31]

Plant P [mg/plant]**

Year 1 Sample days

22 +26.5%

48 +46.5%

Crop uptake

Agroecosystem *Zea Mays* AMF

72 +18.7

Quebec, Canada

[59]

Year 2 Sample days

22 +19.4%

48 +14.2%

72 +41.8%

Leachate Loss SRP [mg]

Lolium multiflora

Nutrient Leaching

Laboratory mesocosms. *Lolium multiflorum*, *Trifolium pratense*, sterilized soils AMF

Claroideoglossomus claroideum +14.2%

Funnelformis mosseae -19.5%

Rhizoglossomus irregular +45.0%

Trifolium pratense

Claroideoglossomus claroideum ns

Funnelformis mosseae ns

Rhizoglossomus irregular ns

Unreactive P

Lolium multiflorum

Claroideoglossomus claroideum -10.8%

Funnelformis mosseae +3.9%

Rhizoglossomus irregular ns

Zürich, Switzerland

[60]

<i>Trifolium pratense</i>					
		<i>Claroideoglossum claroideum</i>	+29.9%		
		<i>Funnelformis mosseae</i>	+19.1%		
		<i>Rhizoglossum irregular</i>	+62.4%		
Vegetative buffers	Pot, <i>Salix</i> , <i>Populus</i> AMF	<u>P stem content</u>	+33%	Southern Quebec, Canada	[61]
Bioretention	Field mesocosms, <i>Carex stipata</i> , AMF/ECM commercial mix	<u>Leachate mass rate (mg/hour) ###</u>	-34%	Portland, Oregon, USA	[62]

ns = no significant difference; calculation of % change = (treatment - control)/control; ## also used leeks, but P uptake was 0, leaving the % change undefined; ### digitized from graphs using Image J (NIH, Bethesda, Maryland); ++ only the effect of AMF considered; * % difference represents an approximate estimate due to difficult digitization for PO₄. Authors state that the differences were significantly different; ** data analyzed for unfertilized plots, fungicide treatment used as control; *** treatments consisted of fungicide (no to low mycorrhizal colonization) and no fungicide (high mycorrhizal colonization).

References

- Sanders, F.E.; Tinker, P.B. Phosphate flow into mycorrhizal roots. *Pestic. Sci.* 1973, 4, 385-395.
- Rillig, M.C.; Sosa-Hernández, M.A.; Roy, J.; Aguilar-Trigueros, C.A.; Vályi, K.; Lehmann, A. Towards an Integrated Mycorrhizal Technology: Harnessing Mycorrhiza for Sustainable Intensification in Agriculture. *Front. Plant Sci.* 2016, 7.
- O'Neill, E.G.; O'Neill, R.V.; Norby, R.J. Hierarchy theory as a guide to mycorrhizal research on large-scale problems. *Environ. Pollut.* 1991, 73, 271-284.
- Dogan, I.; Ozyigit, I.I. Plant-Microbe Interactions in Phytoremediation. In *Soil Remediation in Plants, Prospects and Challenges*, 1st ed.; Hakeem, K.R., Sabir, M., Öztürk, M.A., Eds.; Academic Press: Cambridge, MA, USA, 2015.
- Zalewski, M. Ecohydrology—The scientific background to use ecosystem properties as management tools toward sustainability of water resources. *Ecol. Eng.* 2000, 16, 1-8.
- Dudgeon, D.; Arthington, A.H.; Gessner, M.O.; Kawabata, Z.-I.; Knowler, D.J.; Lévêque, C.; Naiman, R.J.; Prieur-Richard, A.-H.; Soto, D.; Stiassny, M.L.J.; et al. Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biol. Rev.* 2006, 81, 163-182.

7. Michener, W. Win-Win Ecology: How the Earth's Species Can Survive in the Midst of Human Enterprise. *Restor. Ecol.* 2004, 12, 306–307.
8. Ramakrishnan, K.G. Bhuvanewari Influence on Different Types of Mycorrhizal Fungi on Crop Productivity in Ecosystem. *Int. Lett. Nat. Sci.* 2015, 38, 9–15.
9. Khan, A.G. Mycorrhizoremediation—An enhanced form of phytoremediation. *J. Zhejiang Univ. Sci. B* 2006, 7, 503–514.
10. Bücking, H.; Liepold, E.; Ambilwade, P. The Role of the Mycorrhizal Symbiosis in Nutrient Uptake of Plants and the Regulatory Mechanisms Underlying These Transport Processes. *Plant Sci.* 2012.
11. Lin, C.; Wang, Y.; Liu, M.; Li, Q.; Xiao, W.; Song, X. Effects of nitrogen deposition and phosphorus addition on arbuscular mycorrhizal fungi of Chinese fir (*Cunninghamia lanceolata*). *Sci. Rep.* 2020, 10, 12260.
12. Smith, S.E.; Jakobsen, I.; Grønlund, M.; Smith, F.A. Roles of Arbuscular Mycorrhizas in Plant Phosphorus Nutrition: Interactions between Pathways of Phosphorus Uptake in Arbuscular Mycorrhizal Roots Have Important Implications for Understanding and Manipulating Plant Phosphorus Acquisition. *Plant Physiol.* 2011, 156, 1050–1057.
13. Hawkins, B.J.; Jones, M.D.; Kranabetter, J.M. Ectomycorrhizae and tree seedling nitrogen nutrition in forest restoration. *New For.* 2015, 46, 747–771.
14. Smith, S.E.; Read, D.J. *Mycorrhizal Symbiosis*; Academic Press: Cambridge, MA, USA, 2010.
15. Policelli, N.; Horton, T.R.; Hudon, A.T.; Patterson, T.; Bhatnagar, J.M. Back to roots: The role of ectomycorrhizal fungi in boreal and temperate forest restoration. *Front. For. Glob. Chang.* 2020, 3, 97.
16. Djighaly, P.I.; Ndiaye, S.; Diarra, A.M.; Dramé, F.A. Inoculation with arbuscular mycorrhizal fungi improves salt tolerance in *C. glauca* (Sieb). *J. Mater. Environ. Sci.* 2020, 11, 1616–1625.
17. Djighaly, P.I.; Ngom, D.; Diagne, N.; Fall, D.; Ngom, M.; Diouf, D.; Hocher, V.; Laplaze, L.; Champion, A.; Farrant, J.M.; et al. Effect of Casuarina Plantations Inoculated with Arbuscular Mycorrhizal Fungi and Frankia on the Diversity of Herbaceous Vegetation in Saline Environments in Senegal. *Diversity* 2020, 12, 293.
18. Begum, N.; Qin, C.; Ahanger, M.A.; Raza, S.; Khan, M.I.; Ashraf, M.; Ahmed, N.; Zhang, L. Role of Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress Tolerance. *Front. Plant Sci.* 2019, 10.
19. Diagne, N.; Ngom, M.; Djighaly, P.I.; Fall, D.; Hocher, V.; Svistoonoff, S. Roles of arbuscular mycorrhizal fungi on plant growth and performance: Importance in biotic and abiotic stressed regulation. *Diversity* 2020, 12, 370.
20. Asmelash, F.; Bekele, T.; Birhane, E. The Potential Role of Arbuscular Mycorrhizal Fungi in the Restoration of Degraded Lands. *Front. Microbiol.* 2016, 7.
21. Ortaş, I.; Rafique, M. The Mechanisms of Nutrient Uptake by Arbuscular Mycorrhizae. In *Mycorrhiza—Nutrient Uptake, Biocontrol, Ecorestoration*; Varma, A., Prasad, R., Tuteja, N., Eds.; Springer International Publishing: Cham, Switzerland, 2017; pp. 1–19.
22. Nelson, L.L.; Allen, E.B. Restoration of *Stipa pulchra* Grasslands: Effects of Mycorrhizae and Competition from *Avena barbata*. *Restor. Ecol.* 1993, 1, 40–50.
23. Policelli, N.; Horton, T.R.; García, R.A.; Naour, M.; Pauchard, A.; Nuñez, M.A. Native and non-native trees can find compatible mycorrhizal partners in each other's dominated areas. *Plant Soil* 2020, 454, 285–297.
24. Policelli, N.; Horton, T.R.; Hudon, A.T.; Patterson, T.; Bhatnagar, J.M. Back to roots: The role of ectomycorrhizal fungi in boreal and temperate forest restoration. *Front. For. Glob. Chang.* 2020, 3, 97.
25. Broadmeadow, S.; Nisbet, T.R. The effects of riparian forest management on the freshwater environment: A literature review of best management practice. *Hydrol. Earth Syst. Sci. Discuss.* 2004, 8, 286–305.
26. Heckrath, G.; Brookes, P.C.; Poulton, P.R.; Goulding, K.W.T. Phosphorus leaching from soils containing different phosphorus concentrations in the Broadbalk Experiment. *J. Environ. Qual.* 1995, 24, 904–910.
27. Holste, E.K.; Kobe, R.K.; Gehring, C.A. Plant species differ in early seedling growth and tissue nutrient responses to arbuscular and ectomycorrhizal fungi. *Mycorrhiza* 2017, 27, 211–223.
28. Khalil, S.; Loynachan, T.E. Soil drainage and distribution of VAM fungi in two toposequences. *Soil Biol. Biochem.* 1994, 26, 929–934.
29. Ellis, J.R. Post Flood Syndrome and Vesicular-Arbuscular Mycorrhizal Fungi. *J. Prod. Agric.* 1998, 11, 200–204.
30. Stevens, K.J.; Wellner, M.R.; Acevedo, M.F. Dark septate endophyte and arbuscular mycorrhizal status of vegetation colonizing a bottomland hardwood forest after a 100 year flood. *Aquat. Bot.* 2010, 92, 105–111.
31. Tawaraya, K.; Hirose, R.; Wagatsuma, T. Inoculation of arbuscular mycorrhizal fungi can substantially reduce phosphate fertilizer application to *Allium fistulosum* L. and achieve marketable yield under field condition. *Biol. Fertil. Soils* 2012, 48, 839–843.
32. Shenker, M.; Seitelbach, S.; Brand, S.; Haim, A.; Litaor, M.I. Redox reactions and phosphorus release in re-flooded soils of an altered wetland. *Eur. J. Soil Sci.* 2005, 56, 515–525.
33. Rubæk, G.H.; Kristensen, K.; Olesen, S.E.; Østergaard, H.S.; Heckrath, G. Phosphorus accumulation and spatial distribution in agricultural soils in Denmark. *Geoderma* 2013, 209–210, 241–250.
34. Fornara, D.A.; Flynn, D.; Caruso, T. Improving phosphorus sustainability in intensively managed grasslands: The potential role of arbuscular mycorrhizal fungi. *Sci. Total Environ.* 2020, 706, 135744.
35. Ngosong, C.; Jarosch, M.; Raupp, J.; Neumann, E.; Ruess, L. The impact of farming practice on soil microorganisms and arbuscular mycorrhizal fungi: Crop type versus long-term mineral and organic fertilization. *Appl. Soil Ecol.* 2010, 46,

36. Sheng, M.; Lalande, R.; Hamel, C.; Ziadi, N. Effect of long-term tillage and mineral phosphorus fertilization on arbuscular mycorrhizal fungi in a humid continental zone of Eastern Canada. *Plant Soil* 2013, 369, 599–613.
37. Thirkell, T.J.; Charters, M.D.; Elliott, A.J.; Sait, S.M.; Field, K.J. Are mycorrhizal fungi our sustainable saviours? Considerations for achieving food security. *J. Ecol.* 2017, 105, 921–929.
38. Liu, A.; Hamel, C.; Begna, S.H.; Ma, B.L.; Smith, D.L. Soil phosphorus depletion capacity of arbuscular mycorrhizae formed by maize hybrids. *Can. J. Soil Sci.* 2003, 83, 337–342.
39. Oka, N.; Karasawa, T.; Okazaki, K.; Takebe, M. Maintenance of soybean yield with reduced phosphorus application by previous cropping with mycorrhizal plants. *Soil Sci. Plant Nutr.* 2010, 56, 824–830.
40. Grant, C.; Bittman, S.; Montreal, M.; Plenchette, C.; Morel, C. Soil and fertilizer phosphorus: Effects on plant P supply and mycorrhizal development. *Can. J. Plant Sci.* 2005, 85, 3–14.
41. Djodjic, F. Phosphorus Leaching in Relation to Soil Type and Soil Phosphorus Content. *J. Environ. Qual.* 2004, 33, 7.
42. Kabir, Z. Tillage or no-tillage: Impact on mycorrhizae. *Can. J. Plant Sci.* 2005, 85, 23–29.
43. Heijden, M.G.A. van der Mycorrhizal fungi reduce nutrient loss from model grassland ecosystems. *Ecology* 2010, 91, 1163–1171.
44. Castán, E.; Satti, P.; González-Polo, M.; Iglesias, M.C.; Mazzarino, M.J. Managing the value of composts as organic amendments and fertilizers in sandy soils. *Agric. Ecosyst. Environ.* 2016, 224, 29–38.
45. Köhl, L.; Van Der Heijden, M.G. Arbuscular mycorrhizal fungal species differ in their effect on nutrient leaching. *Soil Biol. Biochem.* 2016, 94, 191–199.
46. Asghari, H.R.; Chittleborough, D.J.; Smith, F.A.; Smith, S.E. Influence of Arbuscular Mycorrhizal (AM) Symbiosis on Phosphorus Leaching through Soil Cores. *Plant Soil* 2005, 275, 181–193.
47. Köhl, L.; Van Der Heijden, M.G. Arbuscular mycorrhizal fungal species differ in their effect on nutrient leaching. *Soil Biol. Biochem.* 2016, 94, 191–199.
48. Landry, C.P.; Hamel, C.; Vanasse, A. Influence of arbuscular mycorrhizae on soil P dynamics, corn P-nutrition and growth in a ridge-tilled commercial field. *Can. J. Soil Sci.* 2008, 88, 283–294.
49. Bolan, N.S. A critical review on the role of mycorrhizal fungi in the uptake of phosphorus by plants. *Plant Soil* 1991, 134, 189–207.
50. Mäder, P.; Kaiser, F.; Adholeya, A.; Singh, R.; Uppal, H.S.; Sharma, A.K.; Srivastava, R.; Sahai, V.; Aragno, M.; Wiemken, A.; et al. Inoculation of root microorganisms for sustainable wheat-rice and wheat-black gram rotations in India. *Soil Biol. Biochem.* 2011, 43, 609–619.
51. Deguchi, S.; Uozumi, S.; Tuono, E.; Kaneko, M.; Tawraya, K. Arbuscular mycorrhizal colonization increases phosphorus uptake and growth of corn in a white clover living mulch system. *Soil Sci. Plant Nutr.* 2012, 58, 169–172.
52. Sato, T.; Ezawa, T.; Cheng, W.; Tawaraya, K. Release of acid phosphatase from extraradical hyphae of arbuscular mycorrhizal fungus *Rhizophagus clarus*. *Soil Sci. Plant Nutr.* 2015, 61, 269–274.
53. Battini, F.; Grønlund, M.; Agnolucci, M.; Giovannetti, M.; Jakobsen, I. Facilitation of phosphorus uptake in maize plants by mycorrhizosphere bacteria. *Sci. Rep.* 2017, 7, 4686.
54. Liu, A.; Hamel, C.; Begna, S.H.; Ma, B.L.; Smith, D.L. Soil phosphorus depletion capacity of arbuscular mycorrhizae formed by maize hybrids. *Can. J. Soil Sci.* 2003, 83, 337–342.
55. Asghari, H.R.; Cavagnaro, T.R. Arbuscular mycorrhizas enhance plant interception of leached nutrients. *Funct. Plant Biol.* 2011, 38, 219–226.
56. Zhang, S.; Guo, X.; Yun, W.; Xia, Y.; You, Z.; Rillig, M.C. Arbuscular mycorrhiza contributes to the control of phosphorus loss in paddy fields. *Plant Soil* 2020, 447, 623–636.
57. Bender, S.F.; Conen, F.; Van der Heijden, M.G.A. Mycorrhizal effects on nutrient cycling, nutrient leaching and N₂O production in experimental grassland. *Soil Biol. Biochem.* 2015, 80, 283–292.
58. Martinez-Garcia, L.B.; de Deyn, G.B.; Pugnaire, F.I.; Kothamasi, D.; van der Heijden, M.G.A. Symbiotic soil fungi enhance ecosystem resilience to climate change. *Glob. Chang. Biol.* 2017, 23, 5228–5236.
59. Broadmeadow, S.; Nisbet, T.R. The effects of riparian forest management on the freshwater environment: A literature review of best management practice. *Hydrol. Earth Syst. Sci. Discuss.* 2004, 8, 286–305.
60. Heckrath, G.; Brookes, P.C.; Poulton, P.R.; Goulding, K.W.T. Phosphorus leaching from soils containing different phosphorus concentrations in the Broadbalk Experiment. *J. Environ. Qual.* 1995, 24, 904–910.
61. Kieta, K.A.; Owens, P.N.; Lobb, D.A.; Vanrobaeys, J.A.; Flaten, D.N. Phosphorus dynamics in vegetated buffer strips in cold climates: A review. *Environ. Rev.* 2018, 26, 255–272.
62. Polomski, R.F.; Taylor, M.D.; Bielenberg, D.G.; Bridges, W.C.; Klaine, S.J.; Whitwell, T. Nitrogen and Phosphorus Remediation by Three Floating Aquatic Macrophytes in Greenhouse-Based Laboratory-Scale Subsurface Constructed Wetlands. *Water, Air, Soil Pollut.* 2009, 197, 223–232.

Keywords

mycorrhizae;phosphorus;water quality;mycoremedi

Retrieved from <https://encyclopedia.pub/7475>