

# Improving Biomass Quality in Miscanthus

Subjects: **Others**

Contributor: Oene Dolstra , Luisa Trindade , Kasper van der Cruijsen , Mohamad Al Hassan

Lignocellulosic crops are attractive bioresources for energy and chemicals production within a sustainable, carbon circular society. Miscanthus is one of the perennial grasses that exhibits great potential as a dedicated feedstock for conversion to biobased products in integrated biorefineries. The current biorefinery strategies are primarily focused on polysaccharide valorization and require severe pretreatments to overcome the lignin barrier. The need for such pretreatments represents an economic burden and impacts the overall sustainability of the biorefinery. Hence, increasing its efficiency has been a topic of great interest. Inversely, though pretreatment will remain an essential step, there is room to reduce its severity by optimizing the biomass composition rendering it more exploitable. Extensive studies have examined the miscanthus cell wall structures in great detail, and pinpointed those components that affect biomass digestibility under various pretreatments. Although lignin content has been identified as the most important factor limiting cell wall deconstruction, the effect of polysaccharides and interaction between the different constituents play an important role as well. The natural variation that is available within different miscanthus species and increased understanding of biosynthetic cell wall pathways have specified the potential to create novel accessions with improved digestibility through breeding or genetic modification.

miscanthus

lignocellulosic biomass

saccharification

cell wall

pretreatment

breeding

cellulose

hemicellulose

lignin

biomass quality

## 1. Introduction

Increased carbon dioxide levels are the foremost cause of anthropogenic climate change leading to global warming [\[1\]](#). The goal of keeping the global average temperature well below a 2 °C increase from the pre-industrial levels has been set in the Paris Agreement, in order to halt global warming. To reach this goal, it is important to develop methods or set measures to reduce CO<sub>2</sub> levels [\[2\]](#)[\[3\]](#). The majority of CO<sub>2</sub> emissions are associated with the production of energy and synthetic polymers from fossil resources as oil, coal and natural gas [\[4\]](#)[\[5\]](#). Therefore, there is an increasing demand for less polluting alternatives for the production of energy and chemicals. While there are several renewable sources for energy production, such as solar and wind power, only biomass can serve both purposes. Although skepticism exists about the capacity of biomass usage regarding mitigation of CO<sub>2</sub> emissions [\[6\]](#), thorough assessments have shown that biomass has the potential to offset greenhouse gas emissions when properly used, and could be an integral part of a wider strategy in order to meet global climate goals [\[7\]](#)[\[8\]](#). Transition towards a biobased economy requires change across the whole production chain, with particular importance for advancements regarding efficient biomass production, conversion into different products and utilization [\[9\]](#)[\[10\]](#).

Biomass itself is an attractive renewable energy source due to its potential to be carbon neutral and its global abundance. Carbon neutrality depends on the basic principle that the amount of CO<sub>2</sub> released upon combustion and or conversion is equal to the amount fixed by the crop during its lifetime [11]. As such, replacement of traditional fossil energy sources that heavily contribute to elevated levels of CO<sub>2</sub> with biomass energy can alleviate the associated effect on global warming [12]. Additionally, biomass provides an essentially unlimited source of natural and renewable building blocks for utilization as alternatives to oil-derived chemicals and materials [13].

Lignocellulosic biomass, retrieved from agricultural and forest side-streams or dedicated feedstocks is expected to become an essential resource for the production of energy, chemicals and materials in the near future. Dedicated biomass crops are needed next to agricultural and forest streams, because the contribution of the latter alone would be insufficient for meeting the energy demands [14][15][16][17]. Perennial C4 grasses have been considered as especially promising feedstocks due to their more efficient photosynthetic capacity relative to C3 plants. This is in most cases associated with higher biomass yield potential and increased nitrogen and water use efficiencies [18]. Moreover, their perennial nature also contributes to higher nutrient use efficiency in comparison to annual crops [19]. These features enable perennial grasses, like switchgrass and miscanthus, to achieve substantial yields even when cultivated on marginal and degraded lands [20][21][22]. Limiting cultivation to marginal lands avoids competition for arable land with food crops and will therefore not present a threat to food prices and security or induce land use change; both were points of concern and criticism accompanying the use of edible parts of food crops for the production of first generation biofuels [21][23][24]. Additionally, due to the perennial grasses' capacity to sequester CO<sub>2</sub> [25][26] and thereby tilt the carbon balance more favorably, it can be assumed that the detrimental effects of large scale use of forestry biomass on net CO<sub>2</sub> emissions [27][28] would not apply to these crops. Therefore, cultivation of perennial grasses can be seen as a sustainable alternative without any obvious negative societal impacts. From the available candidate biomass crops, miscanthus is seen as one of the most promising as it is able to utilize external resources even more efficiently than other C4 grasses [19][29].

## 2. Miscanthus for Industrial Use: Advantages, Challenges and Applications

Miscanthus is a genus of rhizomatous perennial grasses originating from Eastern Asia, which comprises around 12 different species [30][31]. The species have been adapted to a broad range of different climate conditions and hold substantial amounts of genetic diversity for key traits [32][33]. Interest in miscanthus has been, for a large part, due to excellent biomass yields that are provided on a yearly basis and could be achieved without the need of additional irrigation in Northern Europe [34]. Such yield potential is achieved due to its ability to maintain photosynthetic capacity at moderate temperatures [35][36]. Furthermore, low input requirements due to its high levels of water [29][37][38] and nutrient use efficiency [39][40] are also highly favorable characteristics of miscanthus species.

Initially, most research has focused on *M. x giganteus*, an interspecific sterile hybrid between *M. sinensis* and *M. sacchariflorus*. Cultivation of *M. x giganteus* is possible in areas where temperatures remain sufficiently high during winter, and high yields (18.7–36.8 t/ha) have been achieved [41]. Moreover, substantial yields (13–21 t/ha) were

reported when cultivated on marginal soils [42][43][44]. The potential for phytoremediation of heavy metal-contaminated soils [45][46] and its ability to act as a carbon sink during its cultivational lifespan [25][47][48] clearly add to why *M. x giganteus* is considered as one of the most promising biomass crops. However, costly rhizome propagation [49], vulnerability to potential pests and diseases due to the absence of genetic variability [50][51][52][53] and lack of cold tolerance leading to severe losses in the first winter after establishment [41][54][55][56] are notable drawbacks of this specific accession.

Despite being a highly promising energy crop, a major drawback surrounding *M. x giganteus* biomass is the resistance of its cell wall against deconstruction, making it recalcitrant towards targeted conversion and valorization through biorefinery. The recalcitrance of the cell wall is directly related to the composition, structure and architecture of the molecules it contains. *M. sinensis* and *M. sacchariflorus* genotypes with lower recalcitrance, performing up to 50% better, have been identified [57].

A large number of applications have been described for miscanthus biomass, with the suitability for a given application ultimately being determined by the cell wall composition. Especially the composition of the secondary cell wall, consisting of cellulose, hemicellulose and lignin, is of importance, as it accounts for >90% of the dry matter of the plant biomass.

Some applications aim to use the whole biomass fraction, such as energy generation through combustion or fast pyrolysis [58][59] or the production of biomaterials such as composite polymers, concrete or fiber boards [60]. Other applications only target a specific fraction of the cell wall for conversion into high value products. Polysaccharide-driven biorefineries are the most well-known example in this context, striving to hydrolyze cell wall polysaccharides into constituent monosaccharides to be fermented to ethanol or methane [61][62] or converted to platform chemicals such as furfural or 5-hydroxymethylfurfural [63]. Alternatively, cellulose could also be used for manufacturing nanocrystals [64]. Although the (hemi)cellulosic parts are still mainly targeted in most lignocellulosic refinery processes, and lignin is therefore generally considered an inconvenient barrier against the conversion of the biomass polysaccharides, lignin valorization is expected to become increasingly important, with it being the most abundant natural resource of aromatic building blocks [65][66].

Ideally, each biomass component could be efficiently separated and isolated for further processing [67][68]. However, this requires pretreatment of the lignocellulosic biomass, since it is recalcitrant to this fractionation and degradation. The required pretreatment stringency remains the first and foremost bottleneck for the design of a green and economically feasible production chain [69][70], requiring both high biomass digestibility and pretreatment efficiency. A useful measure to this end is enzymatic saccharification, since it allows evaluation of the amount of released monosaccharides.

Production of bioethanol and methane are among the best studied applications for lignocellulose feedstocks, as they were initially identified as the most promising value chains [71]. Their production makes use of different ways of enzymatic saccharification, either through the application of enzymatic cocktails containing endo- and exoglucanases and  $\beta$ -glucosidases of fungal origin or through exposure of biomass to hydrolytic bacteria [72][73][74][75].

After the saccharification step monosaccharides generated for bioethanol production are fermented, followed by distillation of the produced ethanol [70]. Alternatively, for methane production the monosaccharides are converted into organic acids and alcohols by acidogenic bacteria, which are subsequently converted into acetate, that serves as a substrate for methanogenic bacteria to produce methane and carbon dioxide [72][73][74]. Genetic studies in the field of biomass digestibility or pretreatment optimization for miscanthus or other lignocellulose grasses often use one of these approaches as a way to assess the performance differences among diverse accessions or pretreatment conditions.

### 3. Improving Biomass Quality in Miscanthus and Breeding Efforts

Interest in breeding of miscanthus is relatively recent, especially when compared to other crops [30]. It is a time-consuming and laborious process as, due to the perennial nature of the crop, agronomically relevant traits, such as plant yield and biomass quality, can only be evaluated in a representative manner after a growth period of at least 2–3 years [76]. Within the genus *Miscanthus*, a large variability for the different traits contributing to cell wall quality is present, enabling selection and breeding for reduced cell wall recalcitrance based on knowledge of cell wall composition. However, cell wall quality is not easily assessed or captured during the breeding process as it is determined by many different traits that are polygenic in nature.

Breeding starts with the availability, generation and search of genotypes with promising quality properties. In general, such genotypes do not have the best overall agronomic characteristics and need to be crossed with advanced breeding material to combine quality with other desirable characteristics, such as high-yielding potential. In practice, this implies a recurrent cyclic approach of crossings and selection to improve quality and agronomic performance and requires appropriate screening tools. The highly diverse germplasm available in *Miscanthus* are attractive sources for desirable cell-wall properties. For instance, natural *M. sinensis* populations from different geographical origins include six distinct genetic clusters and thereby the existence of potential heterotic groups that have so far remained unutilized [33]. Alternatively, deliberately created mutants obtained through targeted genetic modification or undirected mutagenesis could become an additional beneficial source of variation for cell wall genes. To discover useful quality characteristics from selected mutated plants, advanced breeding material and/or wild germplasm, they have to be clonally propagated, through plant splitting or tissue culture to establish replicated field trials for evaluation. The best performing genotypes could either be tested in multi-location trials for their potential as a clonally propagated cultivar, used as parental lines for production of hybrid seeds or serve as a source of beneficial genes for recurrent selection breeding [77][78].

The mating system of fertile species like *M. sinensis*, being gametophytic self-incompatibility (SI), influences the actual breeding in different ways. The system, most likely based on two multiallelic genes as is commonly found among grasses [79], prevents self-pollination but on the other hand it enables the use of heterosis. The breeding program at Wageningen University focusses on the latter and aims to breed for seed-based *M. sinensis* experimental hybrids through pair-wise crossings in isolation among selected genotypes. The SI system limits the full potential of hybrid breeding, but mating between either full sibs or half sibs can circumvent this limitation [80].

Emphasis of the Wageningen breeding program is on selection of candidate clones/individuals for making biparental crosses and on subsequent testing of full-sib families, in particular [81]. The ultimate goal is the creation of hybrid families suitable for commercial use. To remake the original families, the parental clones are maintained. Other breeding programs use similar approaches but instead aim mainly for interspecific seed-based hybrids [78]. Alternatively, creation of new clonally propagated “giganteus” varieties (*M. sinensis* x *M. sacchariflorus*) is also ongoing [82][83].

The use of molecular tools has been explored in miscanthus and resulted in the identification of genetic markers that could potentially speed up the breeding process dramatically. Mapping populations have successfully identified numerous QTLs contributing to important traits such as biomass yield and quality [84][85][86][87]. Additionally, genome-wide association studies in an experimental *M. sinensis* population showed the potential of genomic prediction and selection [88][89]. The use of sufficiently large populations to find SNPs corresponding to traits of interest requires phenotyping of a large number of plants. Analytical protocols for analysis of the main cell wall components and structural sugars have been commonly used for many years. Additionally, there are many more protocols available that make it possible to obtain detailed insight into the composition and structure of these components. While these methods provide information that could be critical for further advancing selection, they often require specialized equipment and are not considered as high throughput. Many of these traits have been successfully analyzed using infrared and near-infrared spectroscopy techniques, that are capable to predict the content of many cell wall structures and can be considered as high-throughput alternatives once appropriate calibration models have been created [90][91].

## References

1. Lacis, A.A.; Schmidt, G.A.; Rind, D.; Ruedy, R.A. Atmospheric CO<sub>2</sub>: Principal Control Knob Governing Earth's Temperature. *Science* 2010, 330, 356–359.
2. Huang, J.; Yu, H.; Dai, A.; Wei, Y.; Kang, L. Drylands face potential threat under 2 °C global warming target. *Nat. Clim. Chang.* 2017, 7, 417–422.
3. Rogelj, J.; Den Elzen, M.; Höhne, N.; Fransen, T.; Fekete, H.; Winkler, H.; Schaeffer, R.; Sha, F.; Riahi, K.; Meinshausen, M. Paris Agreement climate proposals need a boost to keep warming well below 2 °C. *Nature* 2016, 534, 631–639.
4. Heede, R. Tracing anthropogenic carbon dioxide and methane emissions to fossil fuel and cement producers, 1854–2010. *Clim. Chang.* 2014, 122, 229–241.
5. Liu, Z.; Feng, K.; Davis, S.J.; Guan, D.; Chen, B.; Hubacek, K.; Yan, J. Understanding the energy consumption and greenhouse gas emissions and the implication for achieving climate change mitigation targets. *Appl. Energy* 2016, 184, 741–747.
6. Anderson, K.; Peters, G. The trouble with negative emissions. *Science* 2016, 354, 182–183.

7. Souza, G.M.; Ballester, M.V.R.; de Brito Cruz, C.H.; Chum, H.; Dale, B.; Dale, V.H.; Fernandes, E.C.M.; Foust, T.; Karp, A.; Lynd, L.; et al. The role of bioenergy in a climate-changing world. *Environ. Dev.* 2017, 23, 57–64.
8. Fajard, M.; Mac Dowell, N. Can BECCS deliver sustainable and resource efficient negative emissions? *Energy Environ. Sci.* 2017, 10, 1389–1426.
9. Laibach, N.; Börner, J.; Bröring, S. Exploring the future of the bioeconomy: An expert-based scoping study examining key enabling technology fields with potential to foster the transition toward a bio-based economy. *Technol. Soc.* 2019, 58, 101118.
10. Vandermeulen, V.; der Steen, M.; Stevens, C.V.; Van Huylenbroeck, G. Industry expectations regarding the transition toward a biobased economy. *Biofuels Bioprod. Biorefining* 2012, 6, 453–464.
11. Zanchi, G.; Pena, N.; Bird, N. Is woody bioenergy carbon neutral? A comparative assessment of emissions from consumption of woody bioenergy and fossil fuel. *GCB Bioenergy* 2012, 4, 761–772.
12. Dhillon, R.S.; von Wuehlisch, G. Mitigation of global warming through renewable biomass. *Biomass Bioenergy* 2013, 48, 75–89.
13. Saba, N.; Jawaied, M.; Sultan, M.T.H.; Alothman, O.Y. Green Biocomposites for Structural Applications. In *Green Biocomposites: Design and Applications*; Springer: Berlin, Germany, 2017; ISBN 9783319493817.
14. McKendry, P. Energy production from biomass (part 1): Overview of biomass. *Bioresour. Technol.* 2002, 83, 37–46.
15. Lal, R. Crop residues as soil amendments and feedstock for bioethanol production. *Waste Manag.* 2008, 28, 747–758.
16. Scarlat, N.; Martinov, M.; Dallemand, J.F. Assessment of the availability of agricultural crop residues in the European Union: Potential and limitations for bioenergy use. *Waste Manag.* 2010, 30, 1889–1897.
17. Mitchell, R.B.; Schmer, M.R.; Anderson, W.F.; Jin, V.; Balkcom, K.S.; Kiniry, J.; Coffin, A.; White, P. Dedicated Energy Crops and Crop Residues for Bioenergy Feedstocks in the Central and Eastern USA. *Bioenergy Res.* 2016, 9, 384–398.
18. Van der Weijde, T.; Alvim Kamei, C.L.; Torres, A.F.; Vermerris, W.; Dolstra, O.; Visser, R.G.F.; Trindade, L.M. The potential of C4 grasses for cellulosic biofuel production. *Front. Plant Sci.* 2013, 4, 1–18.
19. Masters, M.D.; Black, C.K.; Kantola, I.B.; Woli, K.P.; Voigt, T.; David, M.B.; DeLucia, E.H. Soil nutrient removal by four potential bioenergy crops: *Zea mays*, *Panicum virgatum*, *Miscanthus* ×

giganteus, and prairie. *Agric. Ecosyst. Environ.* 2016, 216, 51–60.

20. Tilman, D.; Socolow, R.; Foley, J.A.; Hill, J.; Larson, E.; Lynd, L.; Pacala, S.; Reilly, J.; Searchinger, T.; Somerville, C.; et al. Beneficial biofuels-The food, energy, and environment trilemma. *Science* 2009, 325, 270–271.

21. Sims, R.E.H.; Mabee, W.; Saddler, J.N.; Taylor, M. An overview of second generation biofuel technologies. *Bioresour. Technol.* 2010, 101, 1570–1580.

22. Pancaldi, F.; Trindade, L.M. Marginal Lands to Grow Novel Bio-Based Crops: A Plant Breeding Perspective. *Front. Plant Sci.* 2020, 11, 227.

23. Field, C.B.; Campbell, J.E.; Lobell, D.B. Biomass energy: The scale of the potential resource. *Trends Ecol. Evol.* 2008, 23, 65–72.

24. Searchinger, T.; Heimlich, R.; Houghton, R.A.; Dong, F.; Elobeid, A.; Fabiosa, J.; Tokgoz, S.; Hayes, D.; Yu, T.-H. Use of U.S. Croplands for Biofuels Increases Greenhouse Gases Through Emissions from Land-Use Change. *Science* 2008, 319, 1238–1240.

25. Zeri, M.; Anderson-Teixeira, K.; Hickman, G.; Masters, M.; DeLucia, E.; Bernacchi, C.J. Carbon exchange by establishing biofuel crops in Central Illinois. *Agric. Ecosyst. Environ.* 2011, 144, 319–329.

26. Qin, Z.; Zhuang, Q.; Zhu, X. Carbon and nitrogen dynamics in bioenergy ecosystems: 2. Potential greenhouse gas emissions and global warming intensity in the conterminous United States. *GCB Bioenergy* 2015, 7, 25–39.

27. Fanous, J.; Moomaw, W.R. A Critical Look at Forest Bioenergy: Exposing a high carbon “climate solution”. *GDAE Clim. Policy Br.* 2018, 1–8.

28. Norton, M.; Baldi, A.; Buda, V.; Carli, B.; Cudlin, P.; Jones, M.B.; Korhola, A.; Michalski, R.; Novo, F.; Oszlányi, J.; et al. Serious mismatches continue between science and policy in forest bioenergy. *GCB Bioenergy* 2019, 1256–1263.

29. Zhuang, Q.; Qin, Z.; Chen, M. Biofuel, land and water: Maize, switchgrass or Miscanthus? *Environ. Res. Lett.* 2013, 8.

30. Clifton-Brown, J.; Chiang, Y.-C.; Hodkinson, T. Miscanthus: Genetic Resources and Breeding Potential to Enhance Bioenergy Production. In *Genetic Improvement of Bioenergy Crops*; Springer: Berlin, Germany, 2008; pp. 273–294. ISBN 978-0-387-70804-1.

31. Hodkinson, T.R.; Klaas, M.; Jones, M.B.; Prickett, R.; Barth, S. Miscanthus: A case study for the utilization of natural genetic variation. *Plant Genet. Resour. Characterisation Util.* 2015, 13, 219–237.

32. Hodkinson, T.R.; Chase, M.W.; Lledó, M.D.; Salamin, N.; Renvoize, S.A. Phylogenetics of Miscanthus, Saccharum and related genera (Saccharinae, Andropogoneae, Poaceae) based on

DNA sequences from ITS nuclear ribosomal DNA and plastid *trnL* intron and *trnL-F* intergenic spacers. *J. Plant Res.* 2002, **115**, 381–392.

33. Clark, L.V.; Brummer, J.E.; Główacka, K.; Hall, M.C.; Heo, K.; Peng, J.; Yamada, T.; Yoo, J.H.; Yu, C.Y.; Zhao, H.; et al. A footprint of past climate change on the diversity and population structure of *Miscanthus sinensis*. *Ann. Bot.* 2014, **114**, 97–107.

34. Lewandowski, I.; Clifton-Brown, J.C.; Scurlock, J.M.O.; Huisman, W. *Miscanthus*: European experience with a novel energy crop. *Biomass Bioenergy* 2000, **19**, 209–227.

35. Beale, C.V.; Long, S.P. Can perennial C4 grasses attain high efficiencies of radiant energy conversion in cool climates? *Plant. Cell Environ.* 1995, **18**, 641–650.

36. Naidu, S.L.; Moose, S.P.; AL-Shoaibi, A.K.; Raines, C.A.; Long, S.P. Cold Tolerance of C 4 photosynthesis in *Miscanthus × giganteus*: Adaptation in Amounts and Sequence of C 4 Photosynthetic Enzymes. *Plant Physiol.* 2003, **132**, 1688–1697.

37. Hamilton, S.K.; Hussain, M.Z.; Bhardwaj, A.K.; Basso, B.; Robertson, G.P. Comparative water use by maize, perennial crops, restored prairie, and poplar trees in the US Midwest. *Environ. Res. Lett.* 2015, **10**, 64015.

38. Kørup, K.; Lærke, P.E.; Baadsgaard, H.; Andersen, M.N.; Kristensen, K.; Münnich, C.; Didion, T.; Jensen, E.S.; Mårtensson, L.M.; Jørgensen, U. Biomass production and water use efficiency in perennial grasses during and after drought stress. *GCB Bioenergy* 2018, **10**, 12–27.

39. Oliveira, J.A.; West, C.P.; Afif, E.; Palencia, P. Comparison of miscanthus and switchgrass cultivars for biomass yield, soil nutrients, and nutrient removal in northwest Spain. *Agron. J.* 2017, **109**, 122–130.

40. Chung, J.-H.; Kim, D.-S. *Miscanthus* as a potential bioenergy crop in East Asia. *J. Crop Sci. Biotechnol.* 2012, **15**, 65–77.

41. Clifton-Brown, J.C.; Lewandowski, I.; Andersson, B.; Basch, G.; Christian, D.G.; Kjeldsen, J.B.; Jørgensen, U.; Mortensen, J.V.; Riche, A.B.; Schwarz, K.U.; et al. Performance of 15 *Miscanthus* genotypes at five sites in Europe. *Agron. J.* 2001, **93**, 1013–1019.

42. Jeżowski, S.; Mos, M.; Buckby, S.; Cerazy-Waliszewska, J.; Owczarzak, W.; Mocek, A.; Kaczmarek, Z.; McCalmont, J.P. Establishment, growth, and yield potential of the perennial grass *Miscanthus × Giganteus* on degraded coal mine soils. *Front. Plant Sci.* 2017, **8**, 1–8.

43. Yost, M.A.; Randall, B.K.; Kitchen, N.R.; Heaton, E.A.; Myers, R.L. Yield potential and nitrogen requirements of *miscanthus × giganteus* on eroded soil. *Agron. J.* 2017, **109**, 684–695.

44. Amaducci, S.; Facciotto, G.; Bergante, S.; Perego, A.; Serra, P.; Ferrarini, A.; Chimento, C. Biomass production and energy balance of herbaceous and woody crops on marginal soils in the Po Valley. *GCB Bioenergy* 2017, **9**, 31–45.

45. Barbosa, B.; Boléo, S.; Sidella, S.; Costa, J.; Duarte, M.P.; Mendes, B.; Cosentino, S.L.; Fernando, A.L. Phytoremediation of Heavy Metal-Contaminated Soils Using the Perennial Energy Crops *Miscanthus* spp. and *Arundo donax* L. *Bioenergy Res.* 2015, 8, 1500–1511.

46. Pidlisnyuk, V.; Stefanovska, T.; Lewis, E.E.; Erickson, L.E.; Davis, L.C. *Miscanthus* as a Productive Biofuel Crop for Phytoremediation. *CRC. Crit. Rev. Plant Sci.* 2014, 33, 1–19.

47. Nakajima, T.; Yamada, T.; Anzoua, K.G.; Kokubo, R.; Noborio, K. Carbon sequestration and yield performances of *Miscanthus × giganteus* and *Miscanthus sinensis*. *Carbon Manag.* 2018, 9, 415–423.

48. Robertson, A.D.; Whitaker, J.; Morrison, R.; Davies, C.A.; Smith, P.; McNamara, N.P. A *Miscanthus* plantation can be carbon neutral without increasing soil carbon stocks. *GCB Bioenergy* 2017, 9, 645–661.

49. Xue, S.; Kalinina, O.; Lewandowski, I. Present and future options for *Miscanthus* propagation and establishment. *Renew. Sustain. Energy Rev.* 2015, 49, 1233–1246.

50. Beccari, G.; Covarelli, L.; Balmas, V.; Tosi, L. First report of *Miscanthus giganteus* rhizome rot caused by *Fusarium avenaceum*, *Fusarium oxysporum* and *Mucor hiemalis*. *Australas. Plant Dis. Notes* 2010, 5, 28–29.

51. Falter, C.; Voigt, C.A. Comparative Cellular Analysis of Pathogenic Fungi with a Disease Incidence in *Brachypodium distachyon* and *Miscanthus × giganteus*. *Bioenergy Res.* 2014, 7, 958–973.

52. Scauflaire, J.; Gourgue, M.; Foucart, G.; Renard, F.; Vandepitte, F.; Munaut, F. *Fusarium miscanthi* and other *Fusarium* species as causal agents of *Miscanthus × giganteus* rhizome rot. *Eur. J. Plant Pathol.* 2013, 137, 1–3.

53. Mekete, T.; Reynolds, K.; Lopez-Nicora, H.D.; Gray, M.E.; Niblack, T.L. Plant-parasitic nematodes are potential pathogens of *Miscanthus × giganteus* and *Panicum virgatum* used for biofuels. *Plant Dis.* 2011, 95, 413–418.

54. Clifton-Brown, J.C.; Lewandowski, I. Overwintering problems of newly established *Miscanthus* plantations can be overcome by identifying genotypes with improved rhizome cold tolerance. *New Phytol.* 2000, 148, 287–294.

55. Kucharik, C.J.; VanLoocke, A.; Lengers, J.D.; Motew, M.M. *Miscanthus* Establishment and Overwintering in the Midwest USA: A Regional Modeling Study of Crop Residue Management on Critical Minimum Soil Temperatures. *PLoS ONE* 2013, 8.

56. De Melo Peixoto, M.; Friesen, P.C.; Sage, R.F. Winter cold-tolerance thresholds in field-grown *Miscanthus* hybrid rhizomes. *J. Exp. Bot.* 2015, 66, 4415–4425.

57. Van der Weijde, T.; Dolstra, O.; Visser, R.G.F.; Trindade, L.M. Stability of Cell Wall Composition and Saccharification Efficiency in Miscanthus across Diverse Environments. *Front. Plant Sci.* 2017, 7, 1–14.

58. Hodgson, E.M.; Fahmi, R.; Yates, N.; Barraclough, T.; Shield, I.; Allison, G.; Bridgwater, A.V.; Donnison, I.S. Miscanthus as a feedstock for fast-pyrolysis: Does agronomic treatment affect quality? *Bioresour. Technol.* 2010, 101, 6185–6191.

59. Lewandowski, I.; Kicherer, A. Combustion quality of biomass: Practical relevance and experiments to modify the biomass quality of *Miscanthus x giganteus*. *Eur. J. Agron.* 1997, 6, 163–177.

60. Moll, L.; Wever, C.; Völkerling, G.; Pude, R. Increase of *Miscanthus* cultivation with new roles in materials production—A review. *Agronomy* 2020, 10, 308.

61. Van der Weijde, T.; Kiesel, A.; Iqbal, Y.; Muylle, H.; Dolstra, O.; Visser, R.G.F.; Lewandowski, I.; Trindade, L.M. Evaluation of *Miscanthus sinensis* biomass quality as feedstock for conversion into different bioenergy products. *GCB Bioenergy* 2017, 9, 176–190.

62. Kiesel, A.; Wagner, M.; Lewandowski, I. Environmental performance of miscanthus, switchgrass and maize: Can C4 perennials increase the sustainability of biogas production? *Sustainability* 2017, 9, 5.

63. Świątek, K.; Gaag, S.; Klier, A.; Kruse, A.; Sauer, J.; Steinbach, D. Acid hydrolysis of lignocellulosic biomass: Sugars and furfurals formation. *Catalysts* 2020, 10, 437.

64. Cudjoe, E.; Hunsen, M.; Xue, Z.; Way, A.E.; Barrios, E.; Olson, R.A.; Hore, M.J.A.; Rowan, S.J. *Miscanthus Giganteus*: A commercially viable sustainable source of cellulose nanocrystals. *Carbohydr. Polym.* 2017, 155, 230–241.

65. Ragauskas, A.J.; Beckham, G.T.; Biddy, M.J.; Chandra, R.; Chen, F.; Davis, M.F.; Davison, B.H.; Dixon, R.A.; Gilna, P.; Keller, M.; et al. Lignin valorization: Improving lignin processing in the biorefinery. *Science* 2014, 344.

66. Rinaldi, R.; Jastrzebski, R.; Clough, M.T.; Ralph, J.; Kennema, M.; Bruijnincx, P.C.A.; Weckhuysen, B.M. Paving the Way for Lignin Valorisation: Recent Advances in Bioengineering, Biorefining and Catalysis. *Angew. Chem. Int. Ed.* 2016, 55, 8164–8215.

67. Wang, K.T.; Jing, C.; Wood, C.; Nagardeolekar, A.; Kohan, N.; Dongre, P.; Amidon, T.E.; Bujanovic, B.M. Toward complete utilization of miscanthus in a hot-water extraction-based biorefinery. *Energies* 2018, 11, 39.

68. Messaoudi, Y.; Smichi, N.; Bouachir, F.; Gargouri, M. Fractionation and Biotransformation of Lignocelluloses-Based Wastes for Bioethanol, Xylose and Vanillin Production. *Waste Biomass Valorization* 2019, 10, 357–367.

69. Tu, W.C.; Hallett, J.P. Recent advances in the pretreatment of lignocellulosic biomass. *Curr. Opin. Green Sustain. Chem.* 2019, 20, 11–17.

70. Limayem, A.; Ricke, S.C. Lignocellulosic biomass for bioethanol production: Current perspectives, potential issues and future prospects. *Prog. Energy Combust. Sci.* 2012, 38, 449–467.

71. Hassan, S.S.; Williams, G.A.; Jaiswal, A.K. Lignocellulosic Biorefineries in Europe: Current State and Prospects. *Trends Biotechnol.* 2019, 37, 231–234.

72. Adney, W.S.; Rivard, C.J.; Shiang, M.; Himmel, M.E. Anaerobic digestion of lignocellulosic biomass and wastes - Cellulases and related enzymes. *Appl. Biochem. Biotechnol.* 1991, 30, 165–183.

73. Chandra, R.; Takeuchi, H.; Hasegawa, T. Methane production from lignocellulosic agricultural crop wastes: A review in context to second generation of biofuel production. *Renew. Sustain. Energy Rev.* 2012, 16, 1462–1476.

74. Zheng, Y.; Zhao, J.; Xu, F.; Li, Y. Pretreatment of lignocellulosic biomass for enhanced biogas production. *Prog. Energy Combust. Sci.* 2014, 42, 35–53.

75. Saini, J.K.; Saini, R.; Tewari, L. Lignocellulosic agriculture wastes as biomass feedstocks for second-generation bioethanol production: Concepts and recent developments. *3 Biotech* 2015, 5, 337–353.

76. Lewandowski, I.; Clifton-Brown, J.; Trindade, L.M.; Van Der Linden, G.C.; Schwarz, K.U.; Müller-Sämann, K.; Anisimov, A.; Chen, C.L.; Dolstra, O.; Donnison, I.S.; et al. Progress on optimizing miscanthus biomass production for the european bioeconomy: Results of the EU FP7 project OPTIMISC. *Front. Plant Sci.* 2016, 7, 1–23.

77. Clifton-Brown, J.; Harfouche, A.; Casler, M.D.; Dylan Jones, H.; Macalpine, W.J.; Murphy-Bokern, D.; Smart, L.B.; Adler, A.; Ashman, C.; Awty-Carroll, D.; et al. Breeding progress and preparedness for mass-scale deployment of perennial lignocellulosic biomass crops switchgrass, miscanthus, willow and poplar. *GCB Bioenergy* 2019, 11, 118–151.

78. Clifton-Brown, J.; Schwarz, K.U.; Awty-Carroll, D.; Iurato, A.; Meyer, H.; Greef, J.; Gwyn, J.; Mos, M.; Ashman, C.; Hayes, C.; et al. Breeding strategies to improve Miscanthus as a sustainable source of biomass for bioenergy and biorenewable products. *Agronomy* 2019, 9, 673.

79. Jiang, J.; Guan, Y.; McCormick, S.; Juvik, J.; Lubberstedt, T.; Fei, S.Z. Gametophytic self-incompatibility is operative in *Miscanthus sinensis* (poaceae) and is affected by pistil age. *Crop Sci.* 2017, 57, 1948–1956.

80. Sacks, E.J.; Juvik, J.A.; Lin, Q.; Stewart, J.R.; Yamada, T. The Gene Pool of *Miscanthus* Species and Its Improvement. In *Genomics of the Saccharinae*; Paterson, A.H., Ed.; Springer: New York, NY, USA, 2013; pp. 73–101. ISBN 978-1-4419-5947-8.

81. Trindade, L.M.; Dolstra, O.; Van Loo, E.N.; Visser, R.G.F. Plant breeding and its role in a biobased economy. In *The Biobased Economy: Biofuels, Materials and Chemicals in the Post-oil Era*; Earthscan Ltd.: London, UK, 2010; pp. 67–82.

82. Dong, H.; Green, S.V.; Nishiwaki, A.; Yamada, T.; Stewart, J.R.; Deuter, M.; Sacks, E.J. Winter hardiness of Miscanthus (I): Overwintering ability and yield of new *Miscanthus ×giganteus* genotypes in Illinois and Arkansas. *GCB Bioenergy* 2019, **11**, 691–705.

83. Clark, L.V.; Dwiyanti, M.S.; Anzoua, K.G.; Brummer, J.E.; Ghimire, B.K.; Głowacka, K.; Hall, M.; Heo, K.; Jin, X.; Lipka, A.E.; et al. Biomass yield in a genetically diverse *Miscanthus sinensis* germplasm panel evaluated at five locations revealed individuals with exceptional potential. *GCB Bioenergy* 2019, **11**, 1125–1145.

84. Atienza, S.G.; Satovic, Z.; Petersen, K.K.; Dolstra, O.; Martin, A. Influencing combustion quality in *Miscanthus sinensis* Anderss.: Identification of QTLs for calcium, phosphorus and sulphur content. *Plant Breed.* 2003, **122**, 141–145.

85. Gifford, J.M.; Chae, W.B.; Swaminathan, K.; Moose, S.P.; Juvik, J.A. Mapping the genome of *Miscanthus sinensis* for QTL associated with biomass productivity. *GCB Bioenergy* 2015, **7**, 797–810.

86. Van der Weijde, T.; Kamei, C.L.A.; Severing, E.I.; Torres, A.F.; Gomez, L.D.; Dolstra, O.; Maliepaard, C.A.; McQueen-Mason, S.J.; Visser, R.G.F.; Trindade, L.M. Genetic complexity of miscanthus cell wall composition and biomass quality for biofuels. *BMC Genom.* 2017, **18**, 1–15.

87. Dong, H.; Liu, S.; Clark, L.V.; Sharma, S.; Gifford, J.M.; Juvik, J.A.; Lipka, A.E.; Sacks, E.J. Genetic mapping of biomass yield in three interconnected *Miscanthus* populations. *GCB Bioenergy* 2018, **10**, 165–185.

88. Slavov, G.T.; Nipper, R.; Robson, P.; Farrar, K.; Allison, G.G.; Bosch, M.; Clifton-Brown, J.C.; Donnison, I.S.; Jensen, E. Genome-wide association studies and prediction of 17 traits related to phenology, biomass and cell wall composition in the energy grass *Miscanthus sinensis*. *New Phytol.* 2014, **201**, 1227–1239.

89. Clark, L.V.; Dwiyanti, M.S.; Anzoua, K.G.; Brummer, J.E.; Ghimire, B.K.; Głowacka, K.; Hall, M.; Heo, K.; Jin, X.; Lipka, A.E.; et al. Genome-wide association and genomic prediction for biomass yield in a genetically diverse *Miscanthus sinensis* germplasm panel phenotyped at five locations in Asia and North America. *GCB Bioenergy* 2019, **988–1007**.

90. Xu, F.; Yu, J.; Tesso, T.; Dowell, F.; Wang, D. Qualitative and quantitative analysis of lignocellulosic biomass using infrared techniques: A mini-review. *Appl. Energy* 2013, **104**, 801–809.

91. Chadwick, D.T.; McDonnell, K.P.; Brennan, L.P.; Fagan, C.C.; Everard, C.D. Evaluation of infrared techniques for the assessment of biomass and biofuel quality parameters and conversion

technology processes: A review. *Renew. Sustain. Energy Rev.* 2014, 30, 672–681.

---

Retrieved from <https://encyclopedia.pub/entry/history/show/17824>