

Rhipicephalus Tick in Southeast Asia

Subjects: **Parasitology**

Contributor: LI PENG TAN

Rhipicephalus species are distributed globally with a notifiable presence in Southeast Asia (SEA) within animal and human populations. The *Rhipicephalus* species are highly adaptive and have established successful coexistence within human dwellings and are known to be active all year round, predominantly in tropical and subtropical climates existing in SEA.

Southeast Asia

Rhipicephalus tick

Tick and tick-borne diseases

Susceptibility Host Responses

Host Range

Economical impacts

1. Background

Southeast Asia (SEA) covers about 4.5 million km² of landmass, with a human population hovering around 670 million [1]. This region comprises 11 countries, and it is a vast Asian region situated east of the Indian subcontinent and South of China (Figure 1). All 11 countries fall within the tropical and subtropical climatic zones. The enormous variety of landscapes and climatic complexities have given rise to a considerable diversity of animals throughout the region, including ticks. With the consistent growth in the average annual gross domestic product (GDP), the concurrent expansion of SEA's livestock sector naturally occurred [2]. Several adverse effects have accompanied this spectacular change in—the “Livestock Revolution”—the phenomenal rise in demand for foods of animal origin in society [3]. Examples include the existing threats of outbreaks of zoonotic diseases that can compromise both animal and human health [4][5], cause economic losses due to diseases [6], and result in environmental pollutions from the usage of disease control drugs and pesticides [7][8]. Small-scale livestock farming (i.e., backyard and village farms) remain the predominant practice in most low-income countries in SEA [9]. This practice requires intensive contact between livestock and farmers, which creates ideal conditions for cross-transfer of pathogens associated with potential zoonosis, in addition to ticks [10].




Figure 1. Geographic depiction of Southeast Asia: SEA comprises countries within the Indo-Chinese peninsula of continental Asia, including Myanmar (Burma), Laos, Vietnam, Thailand, Cambodia, Malaysia, Singapore, Indonesia, Timor-Leste, Brunei and the Philippines (<https://aseanup.com/free-maps-asean-southeast-asia/>, accessed on 4 January 2021).

Ticks are second only to mosquitoes as vectors of disease of medical and veterinary importance. They transmit the widest variety of pathogens for any known arthropod vector, viz. viruses, bacteria, rickettsia, protozoa, or even certain helminths (microfilaria) [11][12]. *Rhipicephalus*. Being the genus most frequently associated with both human and domesticated animals, *Rhipicephalus* is thus the utmost studied genus.

2. Genus *Rhipicephalus* and Its Common Species in Southeast Asia

Ixodidae, also known as hard ticks, are exclusively parasitic arthropods. *Rhipicephalus* is one of the 12 extant genera of Ixodidae and comprises 84 described species [13][14]. *Rhipicephalus* falls under the subfamily of Rhipicephalinae (Metastriata) (Figure 2).




Figure 2. Phylogenetic tree based on maximum-likelihood analysis of the subfamilies of ticks from a 16S ribosomal RNA gene sequence alignment dataset. Branch support value on nodes indicates the bootstrap values of maximum-likelihood and Bayesian posterior probabilities. The highlighted names are *Rhipicephalus* spp. tick sequences from several countries.

Tick species under this genus are found globally even in regions they may not be necessarily 'indigenous' to. Animal trade across the SEA region and other parts of the world enhances the rapid distribution and establishment of tick species such as *Rhipicephalus*. *Rhipicephalus* species are associated with the infestation of livestock or domesticated animals, primarily cattle and dogs [15][16][17][18] imported into or exported out of the SEA region. They are mainly two- and three-host ticks (*Rhipicephalus*) or one host ticks for all the five species under *Boophilus*.

Morphology-based taxonomic classification of *R. microplus* and *R. sanguineus* s.l. has been challenging even for the most experienced taxonomists. The intra-species variations within the *R. microplus* species complex led to the description of multiple sub-species. However, many were later considered synonyms to *R. microplus* or *R. australis* [19]. In recent years, molecular-based phylogenetic analyses added a great deal of insight into the species diversity within the *R. microplus* species complex. Based on studies of mitochondrial cytochrome c oxidase subunit I (COI) gene marker, there are five different phylogenetic clades within the *R. microplus* species complex viz. *R. annulatus*, *R. australis* and three *R. microplus* sensu stricto (s.s.) clades [18][19][20]. These species are not possible to be differentiated based on morphology alone. *Rhipicephalus sanguineus* s.l. on the other hand was shown to have two major phylogenetic clades, the northern (tropical) and southern (temperate) lineages [21]. Besides, several other phylogenetic clades, or operational taxonomic units (OTUs), also exist, representing separate species and needs to be confirmed in further genetic characterization [21].

Rhipicephalus microplus has been reported to occur in Cambodia [22], Laos [22][23], Myanmar [19], Vietnam [24][25], Thailand [26][27], Malaysia [18], the Philippines [28][29] and Indonesia [30][22]. *Rhipicephalus microplus* is frequently found on livestock animals such as cattle [30], water buffaloes [29] and goats [18]. *Rhipicephalus microplus* is widely researched as it is a significant pest of cattle with substantial economic impact [31]. *Rhipicephalus sanguineus* s.l. refers to a group of closely related species associated with dogs worldwide [32]. In SEA, it has been recorded in Laos [23][33], Myanmar [34], Vietnam [35], Thailand [36], Malaysia [37][38], the Philippines [39] and Indonesia [40]. So far, the *R. sanguineus* s.l. identified in SEA fall within the tropical lineage [37]. Nevertheless, the genetic diversity of *R. microplus* and *R. sanguineus* s.l. ticks in SEA is still largely unexplored. Not to mention that there are other species of *Rhipicephalus* whose molecular work are comparatively lesser than *R. microplus* and *R. sanguineus* s.l. *Rhipicephalus pilans*. For instance, only one nucleotide result was available in the gene bank after research on the evolution and ecological niches of *Rhipicephalus* was published in the year 2021 [41].

3. Host Range of *Rhipicephalus* Species in Southeast Asia

The host specificity of *Rhipicephalus* in SEA can be narrowed down based on previous incidences and findings. They are mainly associated with several types of livestock and companion animals (**Table 1**).

Table 1. Host-tick list of *Rhipicephalus* hard tick in Southeast Asia.

Host Type	Country	Tick Species	Host	Reference
Livestock		<i>Rhipicephalus microplus</i>	Unknown	[22]
	Cambodia	<i>Rhipicephalus australis</i>	Unknown	[42]
	Indonesia	<i>Rhipicephalus australis</i>	Unknown	[42]
		<i>Rhipicephalus haemaphysaloides</i>	<i>Bos taurus</i> <i>Bubalus bubalis</i> <i>Capra aegagrus</i> <i>hircus</i>	[43]
		<i>Rhipicephalus microplus</i>	<i>Bos taurus</i> <i>Bubalus bubalis</i> <i>Capra aegagrus</i> <i>hircus</i> <i>Equus caballus</i> <i>Sus scrofa</i>	[30][43][44]
		<i>Rhipicephalus pilans</i>	<i>Bos taurus</i> <i>Bubalus bubalis</i> <i>Capra aegagrus</i> <i>hircus</i> <i>Equus caballus</i> <i>Ovis aries</i>	[30][43][44]

Host Type	Country	Tick Species	Host	Reference
		<i>Rhipicephalus sanguineus</i> s.l.	<i>Bos taurus</i> <i>Bubalus bubalis</i> <i>Gallus gallus</i> <i>domesticus</i> <i>Sus scrofa</i> <i>domesticus</i>	[44]
		<i>Rhipicephalus haemaphysaloides</i>	<i>Bos sp.</i>	[23]
	Laos	<i>Rhipicephalus microplus</i>	<i>Bos sp.</i>	[23]
	Malaysia	<i>Rhipicephalus australis</i>	Unknown	[42]
		<i>Rhipicephalus microplus</i>	<i>Bos taurus</i>	[18][45]
		<i>Rhipicephalus microplus</i>	<i>Bos sp.</i>	[19]
	Myanmar	<i>Rhipicephalus microplus</i>	<i>Bos sp.</i> <i>Sus scrofa</i>	[46]
	Singapore	<i>Rhipicephalus microplus</i>	<i>Bos sp. and Bos taurus</i>	[27][47][48]
	Thailand	<i>Rhipicephalus australis</i>	Unknown	[42]
	The Philippines	<i>Rhipicephalus microplus</i>	<i>Bos sp. and Bos indicus</i> <i>Bubalus bubalis</i> <i>Capra aegagrus hircus</i>	[28][29][49]
		<i>Rhipicephalus haemaphysaloides</i>	<i>Bos sp.</i>	[50]
		<i>Rhipicephalus microplus</i>	<i>Bos sp.</i> <i>Capra aegagrus hircus</i>	[50]
	Timor-Leste	<i>Rhipicephalus sanguineus</i> s.l.	<i>Bos taurus</i>	[50]
		<i>Rhipicephalus annulatus</i>	<i>Bos sp.</i>	[51]
	Vietnam	<i>Rhipicephalus microplus</i>	<i>Bos sp.</i>	[24]
		<i>Rhipicephalus sanguineus</i> s.l.	<i>Bos sp.</i>	[52]
		<i>Rhipicephalus</i>	<i>Canis lupus</i>	[43]

Host Type	Country	Tick Species	Host	Reference
		<i>haemaphysaloides</i>	<i>familiaris</i>	
Indonesia		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i> <i>Felis catus</i>	[53] [43] [54]
		<i>Rhipicephalus</i> <i>haemaphysaloides</i>	<i>Canis lupus</i> <i>familiaris</i>	[23]
		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[33] [55]
Laos		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[37] [56] [57] [58] [59] [60]
		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[61] [62]
Malaysia		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[34]
		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[53] [62] [63]
		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[21] [36] [62]
Companion animals	Myanmar	<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i> <i>Felis catus</i>	[53] [62] [63]
		<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[21] [36] [62]
	Thailand	<i>Rhipicephalus sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i> <i>Felis catus</i>	[53] [29] [62]
		<i>Rhipicephalus</i> <i>haemaphysaloides</i>	<i>Canis lupus</i> <i>familiaris</i>	[52]
	The Philippines	<i>Rhipicephalus</i> <i>sanguineus</i> s.l.	<i>Canis lupus</i> <i>familiaris</i>	[21] [35] [52] [62]
		<i>Rhipicephalus</i> <i>haemaphysaloides</i>	<i>Forest rats</i> *	[43]
Rodents	Vietnam	<i>Rhipicephalus</i> <i>microplus</i>	<i>Rattus exulans</i> <i>Rattus hoffmanni</i> <i>Rattus rattus</i>	[44]
		<i>Rhipicephalus</i> <i>pilans</i>	<i>Niviventer</i> <i>fulvescens</i> <i>Rattus argentiventer</i> <i>Rattus exulans</i> <i>Rattus rattus</i> <i>Rattus tiomanicus</i>	[43] [44] [64]
		<i>Rhipicephalus</i> sp.	<i>Sundamys muelleri</i>	[65]
	Indonesia			

Host Type	Country	Tick Species	Host	Reference
Wild animals	Malaysia	<i>Rhipicephalus haemaphysaloides</i>	<i>Pteropus vampirus</i>	
			<i>Rusa unicolor</i>	
			<i>Helarctos malayanus</i>	
			<i>Panthera tigris</i>	[43][66]
			<i>Varanus salvator</i>	
	Indonesia	<i>Rhipicephalus microplus</i>	<i>Sus scrofa</i>	
			<i>Hylomys suillus</i>	
			<i>Bos javanicus</i>	
			<i>Manis javanica</i>	[43][44]
			<i>Rusa timorensis</i>	
Human	Thailand	<i>Rhipicephalus pilans</i>	<i>Rusa unicolor</i>	
			<i>Crocidura nigripes</i>	
			<i>Hylomys suillus</i>	
			<i>Rusa timorensis</i>	[43][67]
			<i>Suncus murinus</i>	
	Indonesia	<i>Rhipicephalus sanguineus</i> s.l.	<i>Sus scrofa</i>	
			<i>Bos javanicus</i>	[43]
			<i>Rusa unicolor</i>	
			<i>Arctictis binturong</i>	
			<i>Cuon alpinus</i>	[68]
	Thailand	<i>Rhipicephalus haemaphysaloides</i>	<i>Martes flavigula</i>	
			<i>Neofelis nebulosa</i>	
			-	[44]
			-	
			-	

7. Steinfeld, H.; Gerber, P.; Wassenaar, T.; Castel, V.; Rosales, M.; de Haan, C. Livestock's Long Shadow: Environmental Issues and Options; Food and Agriculture Organization of the United Nations (FAO): Rome, Italy, 2006.

8. Ilea, R.C. Intensive livestock farming: Global trends, increased environmental concerns, and ethical solutions. *J. Agric. Environ. Ethics.* 2009, 22, 153–167.

4. The Impacts of Ticks and Tick-Borne Diseases

9. Food and Agriculture Organization of the United Nations (FAO). Agricultural Transformation of Middle-Income Asian Economies: Diversification, Farm Size and Mechanization; Dawe, D., Ed.; Tick-borne diseases transmitted by *Rhipicephalus* ticks affect cattle production worldwide, including SEA countries [70][71][72]. *ESR Working Paper No. 15-04*; Food and Agriculture Organization of the United Nations: Rome, Italy, 2015. Studies have shown the potentially devastating impact of *R. microplus* infestation on developing countries' livestock economies [31]. These losses are bolstered by developing countries' inability to control and monitor the diseases; hence, it impairs the livestock economy [73]. The distribution and prevalence of these

10. Dantas-Torres, F.; Baggio, B.; Gómez, O.; et al. Ticks and tick-borne diseases in Asia: Health perspective. *Trends Parasitol.* **2012**, *28*, 137–146.

11. de la Fuente, J.; Estrada-Peña, A.; Venzal, J.M.; Kocan, K.M.; Sonenshine, D.E. Overview: Ticks as vectors of pathogens that cause disease in humans and animals. *Front. Biosci.* **2008**, *13*, 6938–6946.

12. Yu, Z.; Wang, H.; Wang, T.; Sun, W.; Yang, X.; Liu, J. Tick-borne pathogens and the vector potential of ticks in China. *Parasites Vectors* **2015**, *8*, 24.

13. William, L.N.; Sonenshine, D.E.; Noden, B.H.; Brown, R.N. Ticks (Ixodida). In *Medical and Veterinary Entomology*; Mullen, G.R., Durden, L.A., Eds.; Elsevier Inc.: Amsterdam, The Netherlands, 2019, pp. 603–672.

14. Ernejoor, F.G.; Erna, G.; Mariana, A. Phenotypic and genotypic identification of hard ticks of ruminants and companion animals. *Anaplasma marginale* and *A. centrale* are the notable species in cattle and buffaloes across SEA [49], while *A. platys* occur in dogs [44–49].

15. Petney, T.N.; Sajjuntha, W.; Boulanger, N.; Chitimia-Dobler, L.; Pfeffer, M.; Eamudomkarn, C.; Andrews, R.H.; Ahamad, M.; Putthasorn, N.; Muders, S.V.; et al. Ticks (Argasidae, Ixodidae) and sensitivity of pathogens detection with the help of molecular work could logically fit this scenario. However, it remains unclear why such co-morbidities are consistently challenging to treat, and the ticks are difficult to control in the environment. Hence, an elaborate effort is required to identify the epidemiological patterns of *Rhipicephalus*, the pathogens they transmitted and the rising incidence of resistance to control drugs of this tick in SEA. Molecular detection of the presence of pathogens in squashed ticks is more direct in understanding the host-parasite dynamics for TBDs should be extended further to involve more host species of *Rhipicephalus* in the region. It remains crucial to determine the extent to which *Rhipicephalus* species act as biological, mechanical vectors or both for pathogens of interest.

16. Iain, P.J.; Jefferies, R. Arthropod-transmitted diseases of companion animals in Southeast Asia. *Trends Parasitol.* **2004**, *20*, 27–34.

17. Colella, V.; Nguyen, V.L.; Tan, D.Y.; Lu, N.; Pang, F.; ZhiJuan, Y.; Wang, J.; Liu, X.; Chen, X.; Dong, J.; et al. Zoonotic vectorborne pathogens and ectoparasites of dogs and cats in Eastern and Southeast Asia. *Emerg. Infect. Dis.* **2020**, *26*, 1221–1233.

18. Low, V.L.; Tay, S.T.; Kho, K.L.; Koh, F.X.; Tan, T.K.; Lim, Y.A.L.; Ong, B.L.; Panchadcharam, C.; Norma-Rashid, Y.; Sofian-Azirun, M. Molecular characterisation of the tick *Rhipicephalus microplus* in Malaysia: New insights into the cryptic diversity and distinct genetic assemblages throughout the world. *Parasites Vectors* **2015**, *8*, 1–10.

19. Roy, B.C.; Estrada-Peña, A.; Krucken, J.; Rehman, A.; Nijhof, A.M. Morphological and phylogenetic analyses of *Rhipicephalus microplus* ticks from Bangladesh, Pakistan and Myanmar. *Ticks Tick Borne Dis.* **2018**, *9*, 1069–1079.

5. Resistant and Susceptibility Host Responses

20. Csordas, B.G.; Garcia, M.V.; Cunha, R.C.; Giachetto, P.F.; Blecha, I.M.Z.; Andreotti, R. New insights from molecular characterization of the tick *Rhipicephalus (Boophilus) microplus* in Brazil. The complex interaction, mainly due to the host's diverse immune mechanisms and non-immune structural components, has contributed to various responses towards tick feeding [83]. Most mammals mount an early genetic diversity of *Rhipicephalus* resistance genes to the tick bite in the Old and New Worlds cattle breeds. *Parasites Vectors* **2018**, *10*, 213.

21. Dantas-Torres, F.; Lata, J.; Mg, S.; Antunes, C.; Gómez, O.; Gómez, C.; Gómez, A.; Gómez, J.; De Santis, J. The morphological and genetic diversity of *Rhipicephalus* spp. in cattle breeds from the Americas and Old World cattle breeds. *Parasites Vectors* **2018**, *10*, 213.

Rhipicephalus spp. than *B. taurus* breeds, although considerable variation in resistance exists between and within breeds [84]. The pattern of host resistance to ticks in the SEA region is not necessarily

20. Bongers, T.; Doherty, S.; Barker, S. *In silico* phylogenetic analysis of endo-chordei genomic sequences of several species indicates that the generally Rhipicephalus (Boophilus) microplus, highly asymptomatic species could be selected. *Parasitology* 2014, **137**, 241–253.

23. Vongphayloth, K.; Brey, P.T.; Robbins, R.G.; Sutherland, I.W. First survey of the hard tick (Acaria): Overall, resistance to *R. microplus* infestation in cattle has many effector mechanisms. Although some of the Ixodidae fauna of Nakai District, Khammouane Province, Laos, and an updated checklist of the mechanisms and modulating factors have been identified and quantified, much remains to be explained. Studying ticks of Laos. *Syst. Appl. Acarol.* 2016, **21**, 166–180.

the genetic resistance to ticks among different breeds of cattle can contribute to alternative control methods.

24. Hwang, N.T.; Aboeshie, M. Evaluation of the efficacy of *Camellia sinensis* methanolic extract against the cattle tick *Rhipicephalus (Boophilus) microplus* and the dog tick *Rhipicephalus sanguineus* in a host-parasite relationship. *Plant. Res.* 2011, **3**, 284–289.

25. Kolonin, G.V. Review of the Ixodid Tick Fauna (Acaria: Ixodidae) of Vietnam. *J. Med. Entomol.* 1995, **32**, 276–282.

diversification purposes.

26. Muramatsu, Y.; Usaki, N.; Thongchai, C.; Kramomtong, I.; Kriengsak, P.; Tamura, Y.

6. Controlling and Acaricides Resistance
Seroepidemiologic survey in Thailand of *Coxiella burnetii* infection in cattle and chickens and presence in ticks attached to dairy cattle. *Southeast Asian J. Trop. Med. Public Health* 2014, **45**,

1167.

Rhipicephalus ticks' control mainly depends on conventional acaricides. However, the exhaustive use of these chemicals has resulted in tick populations developing resistance to major acaricide chemical classes. [86].

27. Kaewmongkol, S.; Kaewmongkol, S.; Intromongkol, N.; Lankjaroen, N.; Sirimatumit, T.; Berry, C.M.; Jonsson, N.N.; Stich, R.W.; Sittapalapong, S. Variation among Bm86 sequences in *Rhipicephalus* in goat and sheep farms in Malaysia collected from cattle across Thailand. *Exp. Appl. Acarol.* 2015, **66**, 247–256.

28. Ybanez, A.P.; Siyakumar, T.; Ybanez, R.H.D.; Ratilla, J.C.; Perez, Z.O.; Gabotero, S.R.; Inokuma, H. First molecular characterization of *Anaplasma marginale* in cattle and *Rhipicephalus* (Boophilus) microplus ticks in Cebu, Philippines. *J. Vet. Med. Sci.* 2013, **75**, 27–36.

29. Portugálza, H.P.; Bagot, M.A. Different species of lice (Phthiraptera), fleas (Siphonaptera) and ticks (Ixodidae) collected from livestock, poultry, reptile and companion animal in Leyte Island, Philippines. *Livest. Res. Rural.* 2015, **27**, 1–10.

The success of mosquito control using genetic control methods [90] rekindled interest in using this method to control

30. Sahara, A.; Nugraheni, Y.R.; Patra, G.; Prastowo, J.; Priowidodo, D. Ticks (Acaria: Ixodidae) infestation on cattle in various regions in Indonesia. *Vet. World* 2019, **12**, 1755.

31. Gris, L.; Leite, R.C.; Martins, J.R.D.S.; Barros, A.T.M.D.; Andreotti, R.; Cancado, P.H.D.; León, A.A.P.D.; Pereira, J.B.; Villela, H.S. Reassessment of the potential economic impact of cattle

The per capita consumption of livestock products among SEA countries is projected to increase in the years to come [92] significantly. The increase in demand for livestock products has intensified the race to acquire agricultural

32. Nava, S.; Estrada-Peña, A.; Petney, T.; Beati, L.; Labruna, M.B.; Szabó, M.P.; Venzel, J.M.; Mastropaoletti, M.; Mangold, A.J.; Guglielmone, A.A. The taxonomic status of *Rhipicephalus* sanguineus (Latreille, 1806). *Vet. Parasitol.* 2015, **208**, 2–8.

33. Kernif, T.; Socolovschi, C.; Wells, K.; Lakim, M.B.; Inthalad, S.; Slesak, G.; Boudebouch, N.; Beaucournu, J.C.; Newton, P.N.; Raoult, D.; et al. *Bartonella* and *Rickettsia* in arthropods from the

probable Dermatophagoides or *Bromoides*, Malaysia. Case report. *Int J Infect Dis*. 2012; 35:5157. [94].

34. Hmoon, M.M.; Htun, L.L.; Wai, S.S.; Thu, M.J.; Aung, S.T.; Chel, H.M.; Thaw, Y.N.; Win, S.Y.; Soe, N.C.; Bawm, S. Morphological and molecular identification of ticks infested in stray dogs within Nay Pyi Taw Area, Myanmar. *South Asian J. Life Sci.* 2018, 6, 41–45.

35. Nguyen, V.E.; Coombe, V.; Ratta, P.; Buntar, S.; Dantas-Torres, F. Stray dog ticks and associated pathogens from dogs in northern Vietnam. *Parasitol. Res.* 2019; 118, 139–142.

36. Changbunjong, T.; Buddhirongawat, R.; Suwanpakdee, S.; Siengsanan, J.; Yongyuttawichai, P.; smallholder farmers' resort to using acaricide as it is the most cost-effective method to control tick infestation. Cheewajorn, K.; Jangjaras, J.; Sangloung, C.; Ratanakorn, P. A survey of ectoparasitic arthropods on domestic animals in Tak Province, Thailand. *Southeast Asian J. Trop. Med. Public Health* educated on sustainable agricultural practices and shown the impact of such practices in improving income levels 2009, 40, 435–442. [100]. Besides, there should be more university-industry-farm partnerships for the pilot-testing of newer technologies

37. Koh, F.X.; Panchadcharam, C.; Tay, S.T. Vector-borne diseases in stray dogs in peninsular Malaysia and molecular detection of *Anaplasma* and *Ehrlichia* spp. from *Rhipicephalus sanguineus* (Acarina: Ixodidae) ticks. *J. Med. Entomol.* 2015; 53, 183–187.

38. Koh, F.X.; Panchadcharam, C.; Tay, S.T. Vector-borne diseases in stray dogs in peninsular Malaysia and molecular detection of *Anaplasma* and *Ehrlichia* spp. from *Rhipicephalus sanguineus* (Acarina: Ixodidae) ticks. *J. Med. Entomol.* 2015; 53, 183–187.

39. Galay, R.L.; Manalo, A.A.L.; Dolores, S.L.D.; Aguilar, J.P.M.; Sandalo, K.A.C.; Cruz, K.B.; Divina, B.P.; Andoh, M.; Masatani, T.; Tanaka, T. Molecular detection of tick-borne pathogens in canine companion animals to rodents. The presence and host range of *Rhipicephalus* species in the wild is yet to be studied and understood. There is a realm of unknown ecodynamics for this species. Nevertheless, *Rhipicephalus* population and *Rhipicephalus sanguineus* (sensu lato) ticks from southern Metro Manila and Laguna, Philippines. *Parasites Vectors* 2018; 11, 643.

40. Haen, U.K.; Sovanna, S.; Pratono, P.R.C. Prevalence of ticks and tick-borne diseases in Indonesian dogs. *J. Vet. Sci. Technol.* 2016; 7, 530.

41. Bakkes, D.K.; Ropiquet, A.; Chitimia-Dobler, L.; Matloa, D.E.; Apanaskevich, D.A.; Horak, I.G.; Mans, B.J.; Matthee, C.A. Adaptive radiation and speciation in *Rhipicephalus* ticks: A medley of novel hosts, nested predator-prey food webs, off-host periods and dispersal along temperature variation gradients. *Mol. Phylogenetics Evol.* 2021; 162, 107178.

42. Estuado-Pérez, J.; Venza, J.M.; Nava, S.; Mangold, A.; Eugenio, A.A.; Labora, M.B.; de la Fuente, J. Reinstatement of *Rhipicephalus* (Baboonius) *australis* (Acarini: Ixodidae) with a redescription of the adult and larval stages. *J. Med. Entomol.* 2012; 49, 794–802.

43. Munaf, H.B. Keanekaragaman hospes jenis-jenis caplak marga-marga Amblyomma, *Boophilus* establish a rapporteur system for tick and TBDs in SEA. Babesiosis, anaplasmosis, and theileriosis are the most dan *Rhipicephalus* (Acarina: Ixodidae) yang tercatat memarasit kerbau dan sapi di Indonesia. Ber. reported tick-borne disease of animals in SEA. Diagnosis is usually based on clinical signs of anemia, jaundice, Biol. 1986, 3.

44. Kedarshein, S. *Salivary Exopidia Ticks of Indonesia (Acarina: Ixodidae)*. Ph.D. Dissertation, University of Maryland, College Park, MD, USA, 1971.

45. Tay, S.T.; Koh, F.X.; Kho, K.L.; Ong, B.L. Molecular survey and sequence analysis of *Anaplasma* spp. in cattle and ticks in a Malaysian farm. *Trop. Biomed.* 2014, 31, 769–776.

46. Petney, T.N. A preliminary study of the significance of ticks and tick-borne diseases in South-east Asia. *Mitt. Österr. Ges. Tropenmed. Parasitol.* 1993, 15, 33–42.

47. Jittapalapong, S.; Thanasilp, S.; Kengradomkit, C.; Sirinarukmit, T.; Kaewmongkol, G.; Stich, R.W. Molecular cloning, sequence analysis, and immune recognition of Bm95 from Thai strains of *Rhipicephalus* (*Boophilus*) *microplus*. *Ann. N. Y. Acad. Sci.* 2008, 1149, 45–48.

48. Kaewhom, P.; Stich, R.W.; Needham, G.R.; Jittapalapong, S. Molecular analysis of calreticulin expressed in salivary glands of *Rhipicephalus* (*Boophilus*) *microplus* indigenous to Thailand. *Ann. N. Y. Acad. Sci.* 2008, 1149, 53–57.

49. Swann, P.H.P.; Claveria, F.G. *Rhipicephalus* (*Boophilus*) *microplus* ticks (Family Ixodidae) in goats raised in a small private farm in San Jose del Monte, Bulacan, Central Luzon, Philippines. *Philipp. J. Sci.* 2017, 146, 493–496.

50. Silva, H.R.B.C. *Prospecção Parasitológica em Timor. Subsídios para o Estudo da Fauna Parasitológica dos Seus Animais Domésticos*; Junta de Investigações do Ultramar: Lisboa, Portugal, 1960.

51. Chien, N.T.H.; Linh, B.K.; Van Tho, N.; Hieu, D.D.; Lan, N.T. Status of cattle ticks infection in yellow and dairy cows in Ba Vi District. In Proceedings of the International Conference on Agriculture Development in the Context of International Integration: Opportunities and Challenges, Hanoi, Vietnam, 7–8 December 2016; pp. 115–119.

52. Dong, T.L.; Minh, D.B. Determine the presence of pathogens on ticks in the Mekong Delta region. In International Conference on the Development of Biomedical Engineering in Vietnam; Springer: Singapore, 2020; pp. 707–713.

53. Murrell, A.; Campbell, N.J.H.; Barker, S.C. A total evidence phylogeny of ticks provides insights into the evolution of life cycles and biogeography. *Mol. Phylogenet. Evol.* 2001, 21, 244–258.

54. Sinaga, B.V.; Hariani, N. Prevalensi dan Intensitas ektoparasit pada anjing peliharaan (*Canis familiaris*) di Kalimantan Timur, Indonesia. *J. Bioterididik* 2019, 7, 5.

55. Wilson, N. New distributional records of ticks from Southeast Asia and the Pacific (Metastigmata: Argasidae, Ixodidae). *Orient. Insects* 1970, 4, 37–46.

56. Low, V.L.; Prakash, B.K. First genetic characterization of the brown dog tick *Rhipicephalus sanguineus* sensu lato in Peninsular Malaysia. *Exp. Appl. Acarol.* 2018, 75, 299–307.

57. Macadam, I.; Gudan, D.; Timbs, D.V.; Urquhart, H.R.; Sewell, M.M.H. Metazoan parasites of dogs in Sabah, Malaysia. *Trop. Anim. Health Prod.* 1984, 16, 34–38.

58. Latrofa, M.S.; Dantas-Torres, F.; Giannelli, A.; Otranto, D. Molecular detection of tick-borne pathogens in *Rhipicephalus sanguineus* group ticks. *Ticks Tick Borne Dis.* 2014, 5, 943–946.

59. Watanabe, M.; Nakao, R.; Amin-Babjee, S.M.; Maizatul, A.M.; Youn, J.H.; Qiu, Y.; Watanabe, M. Molecular screening for *Rickettsia*, *Anaplasmataceae* and *Coxiella burnetii* in *Rhipicephalus sanguineus* ticks from Malaysia. *Trop. Biomed.* 2015, 32, 390–398.

60. Prakash, B.K.; Low, V.L.; Tan, T.K.; Vinnie-Siow, W.Y.; Lim, Y.A.L.; Morvarid, A.R.; Sofian-Azirun, M. Detection of *Hepatozoon canis* in the brown dog tick and domestic dogs in Peninsular Malaysia. *J. Med. Entomol.* 2018, 55, 1346–1348.

61. Prakash, B.K.; Low, V.L.; Vinnie-Siow, W.Y.; Tan, T.K.; Lim, Y.A.L.; Morvarid, A.R.; Sofian-Azirun, M. Detection of *Babesia* spp. in dogs and their ticks from Peninsular Malaysia: Emphasis on *Babesia gibsoni* and *Babesia vogeli* infections in *Rhipicephalus sanguineus* sensu lato (Acarı: Ixodidae). *J. Med. Entomol.* 2018, 55, 1337–1340.

62. Nguyen, V.L.; Colella, V.; Greco, G.; Fang, F.; Nurcahyo, W.; Hadi, U.K.; Venturina, V.; Tong, K.B.Y.; Tsai, Y.L.; Taweethavonsawat, P.; et al. Molecular detection of pathogens in ticks and fleas collected from companion dogs and cats in East and Southeast Asia. *Parasites Vectors* 2020, 13, 420.

63. Theis, J.H.; Franti, C.E. Changing infestation rates of *Rhipicephalus sanguineus* (Latreille) (Ixodidae) ticks on dogs on Singapore Island, 1965–1966. *J. Med. Entomol.* 1971, 8, 23–28.

64. Hasan, M.B. Tick fauna of Baluran Wildlife Reserve, Indonesia. *Hemera Zoa* 1978, 70, 37–44.

65. Adrus, M.; Ahamad, M.; Abdullah, M.T. Detection of rickettsiae in engorged ticks from small mammals in Malaysia. *Borneo J. Resour. Sci. Technol.* 2014, 4, 34–41.

66. Anastos, G. Two new species of ticks from Soembawa Island, Indonesia (Acarina: Ixodidae). *J. Parasitol. Res.* 1956, 42, 306–310.

67. Walker, J.B.; Keirans, J.E.; Horak, I.G. Genus *Rhipicephalus* (Acari, Ixodidae). A Guide to the Brown Ticks of the World; Cambridge University Press: Cambridge, UK, 2000.

68. Cornet, J.P.; Demoraes, F.; Souris, M.; Kittayapong, P.; Gonzalez, J.P. Spatial distribution of ticks in Thailand: A discussion basis for tick-borne virus spread assessment. *Int. J. Geoinformat.* 2009, 5, 57–62.

69. Kitaoka, S.; Suzuki, H. Studies on the parasite fauna of Thailand. Parasitic ticks on mammals and description of *Ixodes siamensis* sp. n. and *Rhipicephalus tetracornus* sp. n. (Acarina: Ixodidae). *Trop. Med.* 1983, 25, 205–219.

70. Rahman, W.A.; Lye, Y.P.; Chandrawathani, P. The seroprevalence of bovine babesiosis in Malaysia. *Trop. Biomed.* 2010, 27, 301–307.

71. Ochirkhuu, N.; Konnai, S.; Mingala, C.N.; Okagawa, T.; Villanueva, M.; Pilapil, F.M.I.R.; Murata, S.; Ohashi, K. Molecular epidemiological survey and genetic analysis of vector-borne infections of cattle in Luzon Island, the Philippines. *Vet. Parasitol.* 2015, 212, 161–167.

72. Jirapattharasate, C.; Moumouni, P.F.A.; Cao, S.; Iguchi, A.; Liu, M.; Wang, G.; Zhou, M.; Vudriko, P.; Efstratiou, A.; Changbunjong, T.; et al. Molecular detection and genetic diversity of bovine *Babesia* spp., *Theileria orientalis*, and *Anaplasma marginale* in beef cattle in Thailand. *Parasitol. Res.* 2017, 116, 751–762.

73. Perry, B.; Grace, D. The impacts of livestock diseases and their control on growth and development processes that are pro-poor. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 2009, 364, 2643–2655.

74. Inpankaew, T.; Hii, S.F.; Chimnoi, W.; Traub, R.J. Canine vector-borne pathogens in semi-domesticated dogs residing in northern Cambodia. *Parasites Vectors* 2016, 9, 253.

75. Dantrakool, A.; Somboon, P.; Hashimoto, T.; Saito-Ito, A. Identification of a new type of *Babesia* species in wild rats (*Bandicota indica*) in Chiang Mai Province, Thailand. *J. Clin. Microbiol.* 2004, 42, 850–854.

76. Lim, P.L.; Chavatte, J.M.; Vasoo, S.; Yang, J. Imported human Babesiosis, Singapore, 2018. *Emerg. Infect. Dis.* 2020, 26, 826–828.

77. Nugraha, A.B.; Cahyaningsih, U.; Amrozi, A.; Ridwan, Y.; Agungpriyono, S.; Taher, D.M.; Guswanto, A.; Gantuya, S.; Tayebwa, D.S.; Tuvshintulga, B.; et al. Serological and molecular prevalence of equine piroplasmosis in Western Java, Indonesia. *Vet. Parasitol. Reg. Stud. Rep.* 2018, 14, 1–6.

78. Kamyingkird, K.; Yangtara, S.; Desquesnes, M.; Cao, S.; Moumouni, A.; Jittapalapong, S.; Nimsupan, B.; Terkawi, M.A.; Masatani, T.; Nishikawa, Y.; et al. Seroprevalence of *Babesia caballi* and *Theileria equi* in horses and mules from Northern Thailand. *J. Protozool. Res.* 2014, 24, 11–17.

79. Ybañez, A.P.; Ybañez, R.H.D.; Claveria, F.G.; Cruz-Flores, M.J.; Xuenan, X.; Yokoyama, N.; Inokuma, H. High genetic diversity of *Anaplasma marginale* detected from Philippine cattle. *J. Vet. Med. Sci.* 2014, 76, 1009–1014.

80. Faizal, M.D.; Haryanto, A.; Tjahajati, I. Diagnosis and molecular characterization of *Anaplasma platys* in dog patients in Yogyakarta area, Indonesia. *Indones. J. Biotechnol.* 2019, 24, 43–50.

81. Chansiri, L. Tick-borne diseases in Thailand. *Trop. Anim. Health Prod.* 1997, 29, 52S.

82. Ghosh, S.; Azhahianambi, P.; de la Fuente, J. Control of ticks of ruminants, with special emphasis on livestock farming systems in India: Present and future possibilities for integrated control—A review. *Exp. Appl. Acarol.* 2006, 40, 49–66.

83. Jonsson, N.N.; Piper, E.K.; Constantinoiu, C.C. Host resistance in cattle to infestation with the cattle tick *Rhipicephalus microplus*. *Parasite Immunol.* 2014, 36, 551–557.

84. Seifert, G. Variations between and within breeds of cattle in resistance to field infestations of the cattle tick (*Boophilus microplus*). *Aust. J. Agric. Res.* 1971, 22, 159–168.

85. Kongsuwan, K.; Josh, P.; Colgrave, M.L.; Bagnall, N.H.; Gough, J.; Burns, B.; Pearson, R. Activation of several key components of the epidermal differentiation pathway in cattle following infestation with the cattle tick, *Rhipicephalus* (*Boophilus*) *microplus*. *Int. J. Parasitol.* 2010, 40, 499–507.

86. Rodriguez-Vivas, R.I.; Jonsson, N.N.; Bhushan, C. Strategies for the control of *Rhipicephalus microplus* ticks in a world of conventional acaricide and macrocyclic lactone resistance. *Parasitol. Res.* 2018, 117, 3–29.

87. Basripuzi, H.B.; Sani, R.A.; Ariff, O.M. Anthelmintic resistance in selected goat farms in Kelantan. *Mal. J. Anim. Sci.* 2012, 15, 47–56.

88. Puspitasari, S.; Farajallah, A.; Erni Sulistiawati, M. Effectiveness of Ivermectin and Albendazole against *Haemonchus contortus* in Sheep in West Java, Indonesia. *Trop. Life Sci. Res.* 2016, 27, 135–144.

89. Kochapakdee, S.; Pandey, V.S.; Pralomkarn, W.; Choldumrongkul, S.; Ngampongsai, W.; Lawpetchara, A. Anthelmintic resistance in goats in southern Thailand. *Vet. Rec.* 1995, 137, 124–125.

90. Bouyer, J.; Culbert, N.J.; Dicko, A.H.; Pacheco, M.G.; Virginio, J.; Pedrosa, M.C.; Garziera, L.; Pinto, A.T.M.; Klaptoz, A.; Germann, J.; et al. Field performance of sterile male mosquitoes released from an uncrewed aerial vehicle. *Sci. Robot.* 2020, 5, eaba6251.

91. Osburn, R.L.; Knippling, E.F. The potential use of sterile hybrid *Boophilus* ticks (Acari: Ixodidae) as a supplemental eradication technique. *J. Med. Entomol.* 1982, 19, 637–644.

92. Loong, S.K.; Lim, F.S.; Khoo, J.J.; Lee, H.Y.; Suntharalingam, C.; Ishak, S.N.; Mohd-Taib, F.S.; AbuBakar, S. Culturable pathogenic bacteria in ticks parasitizing farm animals and rodents in Malaysia. *Trop. Biomed.* 2020, 37, 803–811.

93. Wong, C.C.; Moog, F.; Chen, C.P. Forage and ruminant livestock integration in tree crop plantations of Southeast Asia. In *Grasslands: Developments Opportunities Perspectives*; Reynolds, S., Frame, J., Eds.; Taylor and Francis: London, UK, 2019; pp. 403–431.

94. Ndawula, C., Jr.; Tabor, A.E. Cocktail anti-tick vaccines: The unforeseen constraints and approaches toward enhanced efficacies. *Vaccines* 2020, 8, 457.

95. Rodríguez-Molano, C.E.; Torres, S.U.; Monrroy, L.N. Background on the control of the cattle tick *R. (B.) microplus* and the use of coumarin substances as an alternative. *Pharm. Pharmacol. Int. J.*

2020, 8, 215–232.

96. Kwanmuang, K.; Pongputhinan, T.; Jabri, A.; Chitchumnung, P. Small-scale farmers under Thailand's smart farming system. FFTC-AP 2020, 2647. Available online: (accessed on 4 March 2021).

97. Tran, C.T. Overview of agricultural policies in Vietnam. FFTC-AP 2014, 629. Available online: (accessed on 5 March 2021).

98. Kustiari, R. Livestock development policy in Indonesia. FFTC-AP 2014, 728. Available online: (accessed on 5 March 2021).

99. Hashim, F.A.H. Strategies to strengthen livestock industry in Malaysia. FFTC-AP 2015, 911. Available online: (accessed on 5 March 2021).

100. Altieri, M.A.; Nichols, C.I. Agroecology scaling up for food sovereignty and resiliency. In Sustainable Agriculture Reviews; Lichtenföhr, E., Ed.; Springer: Berlin, Germany, 2012; pp. 1–29.

Retrieved from <https://encyclopedia.pub/entry/history/show/28264>