

Gastroschisis

Subjects: **Others**

Contributor: Vineet Bhandari

The International Clearinghouse for Birth Defects Surveillance and Research defines gastroschisis as "a congenital malformation characterized by visceral herniation usually through a right side abdominal wall defect to an intact umbilical cord and not covered by a membrane". Approximately 10% of infants with gastroschisis have intestinal stenosis or atresia resulting from vascular insufficiency due to a volvulus or compression of vascular pedicle by a narrowing abdominal ring.

abdominal defect

infant

newborn

congenital anomaly

nutrition

1. Introduction

In recent years, the subject of gastroschisis (Figure 1) has received considerable attention due to: (1) the controversy surrounding its pathogenesis in light of our understanding of its developmental biology; (2) a temporal increase in incidence of gastroschisis for reasons that are not entirely clear; (3) improvements in our ability to diagnose the condition early in pregnancy and identify "high-risk" fetuses; (4) improvements in operative techniques; and (5) advances in parenteral and enteral nutritional support during the postoperative period.

The first case of an abdominal wall defect was reported by Conrad Wolffhart (Lycosthenes), an Alsatian humanist and theologian, who described an infant, born in 1547, with "a large extrusion of intestines from abdomen and chest, feet by the head, and a tall, pointed skull" [1]. The term "gastroschisis" (from ancient Greek, *gastro* = stomach, and *schisi* = split), first coined by Taruffi in 1894, is actually a misnomer, since it is the anterior abdominal wall, not the stomach, that is split. In 1953, Moore and Stokes first described the features differentiating gastroschisis from omphalocele [2].



Figure 1. Term infant with simple gastroschisis. The picture shows a small abdominal wall defect to the right of the site of umbilical cord insertion.

2. Pathogenesis

Several embryologic hypotheses have been proposed to elucidate the pathogenesis of gastroschisis including a failure of differentiation of the embryonic mesenchyme due to a teratogenic exposure [3], rupture of the amniotic membrane at the base of the umbilical cord [4], abnormal involution of the right umbilical vein, leading to impaired viability of surrounding mesenchyme [5], interruption of the omphalomesenteric artery leading to localized necrosis of the abdominal wall at the base of the cord [6], abnormal folding of the embryo leading to a ventral body wall defect [7], failure of the yolk sac and related vitelline structures to be incorporated into the umbilical stalk, leading to a perforation in the abdominal wall separate from the umbilicus [8], and amniotic rupture in the pars flaccida part of the umbilical cord secondary to a genetic predisposition or exogenous factors e.g., toxins, drugs, viruses, or radiation [9].

More recently, Lubinsky proposed a binary vascular–thrombotic model for gastroschisis, where the normal involution of the umbilical vein creates a potential site for thrombosis adjacent to the umbilical ring. Subsequent thrombosis, associated with factors increasing maternal estrogen levels, weakens the umbilical ring, creating a site for potential herniation. This model can explain the morphological findings (location at the umbilical ring, typical right-sidedness, and amniocyte inclusions) as well the epidemiologic risk factors (rising incidence due to increasing environmental contamination with estrogen disruptors) [10]. However, evidence for this to be occurring in humans is lacking.

Recent human evidence appears to support the theory that gastroschisis is not a defect of the abdominal wall, but an abnormality of the rudimentary umbilical ring, resulting in a separation of the fetal ectoderm from the amnion's epithelium on the right side [9][11][12].

3. Epidemiology

The epidemiology of gastroschisis has been a subject of considerable interest, given the steady increase in its prevalence over the past three decades. In the US, the prevalence of gastroschisis nearly doubled during the period from 1995 to 2005 [13]. However, this increasing trend is not universal; low and stable rates have been reported from Southern Europe (particularly Italy) and the Middle East [14]. The current prevalence rate of gastroschisis in the US is 4.5 per 10,000 live births [15].

Gastroschisis is unusual among birth defects in that it disproportionately affects younger mothers, with the highest prevalence among mothers aged <20 years (15.7 per 10,000 live births) [14][15][16][17][18][19]. Approximately 70% of infants with gastroschisis are born to women under 25 years of age, and the incidence among teenage mothers is more than seven times that among mothers aged ≥25 years [15]. The prevalence rates are higher among white and Hispanic mothers than among black mothers [14][15][18].

In a recent meta-analysis of 29 studies, maternal smoking (relative risk (RR), 1.56; 95% confidence interval (CI), 1.40–1.74), illicit drug use (RR, 2.14; 95% CI, 1.48–3.07), and alcohol consumption (RR, 1.39; 95% CI, 1.13–1.70) during early pregnancy were associated with an increased risk of gastroschisis [20]. A younger age of the father has been found to be a risk factor in some studies [21][22][23][24]. A recent ecologic analysis found that gastroschisis prevalence was higher in areas with high or medium opioid prescription rates, compared with that in areas with low rates [15]. In a Canadian cohort study involving two large databases, smoking (odds ratio (OR), 2.86; 95% CI, 2.22–3.66), a history of pregestational or gestational diabetes (OR,

2.81; 95% CI, 1.42–5.5), and use of medication to treat depression (OR, 4.4; 95% CI, 1.38–11.8) emerged as significant associations with gastroschisis pregnancies [17]. The National Birth Defects Prevention Study 1997–2011 reported genitourinary tract infections during the periconceptional period to be significantly associated with gastroschisis (OR, 1.5; 95% CI, 1.3–1.9) [25].

There appears to be a spatial variation in the rates of gastroschisis, with "case clusters" in discrete areas relative to surrounding areas [24][25][26][27], suggesting the possible role of agrichemicals as an etiologic factor [28][29].

4. Types of Gastroschisis

First proposed by Molik et al. [30], gastroschisis is classified into simple and complex types based on the condition of the bowel. In simple gastroschisis, the bowel is in good condition with no intestinal complications. Complex gastroschisis, on the other hand, is gastroschisis associated with congenital intestinal complications in the form of an atresia, perforation, ischemia, necrosis, or volvulus. Closed or closing gastroschisis is a subset of complex gastroschisis, in which the abdominal wall defect closes around the prolapsed bowel, resulting in exit and/or entry intestinal stricture, atresia, ischemia, necrosis, or resorption. These intestinal complications result from a combination of exposure to digestive compounds in the amniotic fluid and ischemia due to mesenteric constriction at the level of the defect. Infants with closing gastroschisis tend to have a high incidence of short-bowel syndrome (SBS). In extremely-rare cases, known as the "vanishing gut syndrome", the abdominal wall defect closes completely prenatally, resulting in extreme short gut [31]. In a recent meta-analysis [32], infants with complex gastroschisis, occurring in 17% of cases, have significantly higher mortality rates (16.67%) compared to those with simple gastroschisis (2.18%). Infants with complex gastroschisis also suffer from significantly-higher rates of morbidities. They are started on enteral feeds later, they take longer to reach full enteral feedings, and require a longer duration of parenteral nutrition (PN). They have a higher risk of sepsis, SBS, and necrotizing enterocolitis (NEC). They tend to stay longer in the hospital and are more likely to be sent home with enteral tube feedings and PN. Left-sided gastroschisis is a rare entity, where the abdominal wall defect occurs in the left paraumbilical region. It is more common in females, and associated with a higher incidence of extraintestinal anomalies compared with the right-sided lesions [33].

5. Surgical Management

The goals of surgical management of gastroschisis include reduction of the herniated viscera into the peritoneal cavity while avoiding direct trauma to the bowel and excessive intra-abdominal pressure, and closure of the abdominal wall defect. While the condition of the exposed bowel and the degree of abdominovisceral disproportion primarily dictate the type and timing of surgical intervention, other factors such as gestational maturity and the infant's weight and co-morbidities also need to be considered. The surgical options for closure include:

- (a) Primary reduction, with either immediate sutured closure or sutureless closure;**5.1. Primary Reduction**
- (b) Prosthetic silo placement, gradual visceral reduction followed by delayed sutured or sutureless closure.

Primary reduction of the gastroschisis is usually performed if the herniated bowel can be safely placed back into the abdominal cavity without causing excessive intra-abdominal pressure. If the herniated bowel loops appear near-normal, this can be performed in the neonatal intensive care unit (NICU) under mild sedation/analgesia. Although rare, in infants with closing gastroschisis, the narrow fascial defect can compromise intestinal blood flow, and may require urgent enlargement of the defect to preserve bowel perfusion and facilitate reduction.

"Abdominal compartment syndrome", a serious and potentially-life-threatening complication after primary closure, is characterized by respiratory compromise, and/or lower limb, renal and intestinal ischemia. Intra-gastric or intra-vesical

pressures >20 mmHg, or central venous pressure >4 mmHg have been shown to correlate with decreased perfusion to the kidneys and bowel, and potential risk of compartment syndrome [34][35]. Peak inspiratory pressures of <25 cm H₂O on the ventilator after closure also predicts low risk for abdominal compartment syndrome [36]. Following the reduction of the bowel into the abdominal cavity, the defect is closed either by sutured fascial closure technique or by sutureless closure technique.

5.2. Staged Reduction

Staged reduction is achieved by the placement of a spring-loaded silo that provides coverage to the exposed bowel, with subsequent definitive closure of the defect. This method can be used as initial therapy and also following failure of primary reduction. The spring-loaded ring of the silo is inserted through the abdominal wall defect and rests beneath the fascia inside the abdominal cavity without the need for placement of fascial sutures. The procedure can be performed at the bedside under mild sedation/analgesia, without the need for endotracheal intubation. The transparent bag covers the eviscerated bowel, which is reduced daily by tying umbilical tapes around the bag (Figure 2). Once the bowel is completely reduced into the abdomen (Figure 3), closure is performed using either the sutured fascial closure or the sutureless closure techniques.



Figure 2. Staged reduction of gastroschisis. Shown is a picture with the bowel loops placed in a silo.



Figure 3. Staged reduction of gastroschisis. Shown is a picture with the bowel loops reduced into the abdomen using a silo.

5.3. Sutureless Closure

First described by Sandler et al. [37], the "plastic" sutureless closure technique utilizes the umbilical cord, left deliberately long at the time of birth, as a biologic dressing. After careful reduction of the eviscerated bowel into the abdomen, the gastroschisis defect is covered with the umbilical cord cut and tailored to fit the opening. A clear plastic dressing (Tegaderm 3M) is placed over the defect, which is then allowed to heal by secondary intention. The umbilical-cord-covered defect contracts circumferentially resulting in a scarless abdomen and a cosmetically-acceptable appearance of the umbilicus in 2–4 weeks ([Figure 4](#)). This technique can be used after primary reduction as well as following staged reduction with silo placement. In the latter case, the cord is wrapped in Vaseline gauze and kept moist while the silo is in place or placed inside the silo to maintain viability.



Figure 4. The infant shown in [Figure 1](#) underwent a primary sutureless closure soon after birth. Shown at 19 days of age, he is on full oral feeds.

5.4. Primary Reduction vs. Staged Reduction

The optimal timing of abdominal wall closure in gastroschisis remains debatable, with general agreement that early reduction is best.

Staged reduction with silo placement has the theoretical advantage of achieving reduced intra-abdominal pressure at the time of definitive closure, leading to improved splanchnic perfusion, resulting in faster return of bowel function, reduced rates of infection and NEC and decreased risk of long-term bowel dysfunction [\[38\]\[39\]](#). Placement of a silo also allows for ongoing assessment of bowel perfusion through the transparent bag. Prolonged use of the silo, however, can lead to pressure necrosis around the silo ring.

In a meta-analysis that included studies with least selection bias, staged closure with silo was associated with better outcomes and a significant reduction in ventilator days, time to first feed, and infection rates [\[40\]](#). However, in the same meta-analysis, when all studies were included, primary closure was associated with improved outcomes.

A recent large, multicenter retrospective observational study involving 866 neonates with gastroschisis compared infants who underwent immediate closure with those who had a silo placed for ≤ 5 days [\[41\]](#). The two groups had generally-equivalent outcomes, except for a higher incidence of ventral hernias in infants who underwent immediate closure compared to those who had silo placed for a short duration. There were no significant differences between the two groups in terms of mortality, time to tolerance of full enteral feeds, duration of TPN, or hospital LOS. Infants who require a silo for longer than 5 days likely represent a subset of patients with more severe disease and less favorable outcomes, and therefore, may not be appropriate to compare to those infants who underwent immediate closure.

Two small prospective randomized trials demonstrated no difference between infants undergoing primary closure and those undergoing silo placement with delayed closure in terms of time to full enteral feeds, duration of ventilatory support, LOS, or

incidence of sepsis [42][43]. In one of these studies [43], patients in the silo group were closed at a median of 4 days after silo placement (interquartile range (IQR), 2–5.8 days).

5.5. Sutured Closure vs. Sutureless Closure

The sutureless closure method has gained popularity in recent years due to its simplicity, and purported advantages of the ability to perform the procedure at the bedside without the need for general anesthesia, lower intra-abdominal pressures after closure, shorter duration of mechanical ventilation, decreased need for pain medication, superior cosmetic appearance of the umbilicus, and lower hospital cost [37][44][45].

In a large, single-center cohort study of 97 infants with gastroschisis, sutureless repair was associated with significant reduction in duration of mechanical ventilation and pain medication requirements, but with an increased risk of umbilical hernias, compared to sutured closure [46]. In a prospective randomized controlled trial involving 39 infants with gastroschisis, sutureless repair was associated with a significant increase in time to full feeds and time to discharge [47]. A meta-analysis of studies comparing flap closure with fascial closure for gastroschisis by Yousseff et al. demonstrated no significant differences in mortality, LOS, or feeding parameters between the two groups [48]. The infants who underwent flap closure had significantly fewer wound infections, but had an increased risk of umbilical hernia.

In infants with gastroschisis with associated ileal atresia, the surgical management must be individualized based on gestational maturity, birth weight, clinical status, and the condition of the bowel. If the bowel is in good condition, primary anastomosis with abdominal wall closure can be performed. Otherwise, options include creation of stoma with closure or reduction of unprepared bowel into the abdomen with closure and delayed exploration to establish bowel continuity [36]. The musculature of the abdominal wall is normal in gastroschisis.

[Figure 5](#) provides an algorithm for the choice of optimal surgical procedure based on the type of gastroschisis and the ability to safely reduce the bowel.




Figure 5. Algorithm for reduction and closure of gastroschisis. (Adapted from Ref. [36]).

6. Postoperative Management

The postoperative course in infants with gastroschisis is primarily dependent upon the extent and severity of intestinal injury. Most infants tend to have some degree of adynamic ileus, intestinal dysmotility, and nutrient malabsorption and require nasogastric decompression via a Replogle (nasogastric) tube and TPN via a secure central venous catheter, either a peripherally-inserted central catheter (PICC) or a tunneled Broviac catheter. Feeding is initiated upon return of bowel function (resolution of abdominal distension and bilious gastric drainage). The time to initiation of feeds can range from one to two weeks, or longer, depending on the condition of the bowel. Early initiation of trophic feeds can improve peristalsis, prevent villous atrophy, and reduce bacterial overgrowth. Feeds are advanced slowly as feeding intolerance is common due to intestinal dysmotility. Feeding with maternal expressed breast milk, when available, may help to protect the infant from developing NEC [49]. Early oral stimulation is important to preserve sucking/swallowing reflexes and prevent oral aversion. Infants with gastroschisis have been shown to have abnormal esophageal motor function leading to delayed acquisition of oral feeding milestones, and the possible need for chronic tube feeding [50]. In a multicenter, randomized, double-blind, placebo-controlled trial, enterally-administered erythromycin conferred no advantage in the time taken to achieve full enteral feeding after primary repair of uncomplicated gastroschisis [51].

There appears to a wide variation in the use of prophylactic antibiotics following abdominal wall closure [52]. Based on a multi-institutional review of practice and outcomes, it is recommended that antibiotics be discontinued within 48 h of abdominal wall closure in the absence of culture-positive sepsis or clinical instability [53].

The incidence of NEC in infants with gastroschisis is approximately 5%, which is higher than that predicted based on their gestational age. However, NEC in gastroschisis tends to be clinically less severe than that seen in preterm infants, with only one-quarter of these infants requiring surgery for NEC [54].

Despite infants with gastroschisis having some degree of malrotation, the incidence of midgut volvulus is relatively low at approximately 1% [55]. This may relate to bowel inflammation and handling leading to formation of protective adhesions. Given the low risk of midgut volvulus and the risk of bowel adhesions and obstruction, Ladd's procedure is not routinely performed in gastroschisis. However, the diagnosis of volvulus must be considered in any infant with gastroschisis presenting with delayed bowel obstruction.

SBS can occur in infants with complex gastroschisis with extensive bowel injury resulting from jejunointestinal atresia, midgut volvulus, NEC, or refractory intestinal failure. These infants are dependent on PN for prolonged periods and therefore require placement of indwelling central venous catheters. As a result, they can develop intestinal bacterial overgrowth, bacteremia due to translocation of enteric bacteria, or intestinal failure-associated liver disease. The risk of liver injury can be reduced by using ursodeoxycholic acid, which acts by displacing hydrophobic, more toxic bile salts [56], use of SMOFlipid, an intravenous lipid emulsion containing soybean oil, medium-chain triglycerides, olive oil, and fish oil [57], and cycling of PN to reduce hepatic steatosis [58]. Using enteral antibiotics and probiotics can be helpful in treating bacterial overgrowth. Other pharmacologic interventions used in SBS include use of gastric-acid-suppressing agents, e.g., H₂-receptor blockers and proton pump inhibitors, prokinetic agents such as erythromycin, anti-diarrheal agents such as loperamide and codeine, absorbent agents such as pectin and guar gum, or cholestyramine [59]. The surgical approaches to SBS include intestinal lengthening procedures, e.g., longitudinal intestinal lengthening and tailoring (LILT) and serial transverse enteroplasty (STEP), and in patients with irreversible hepatic and intestinal failure, intestinal transplantation [60]. Management of SBS often requires multidisciplinary "intestinal failure" teams that include neonatologists, gastroenterologists, surgeons, nutritionists, pharmacists, nurses, and social workers.

Cryptorchidism is common in gastroschisis, occurring in approximately 35% of all male infants [61][62]. It has been postulated that increased intra-abdominal pressure is responsible for the normal descent of the testis, and in gastroschisis, forces that promote testicular descent are absent. However, there is a high rate of spontaneous migration during the first year of life, with nearly 50% of the testes relocating without intervention [61][62]. The watch-and-wait approach is recommended for management of cryptorchidism in gastroschisis due to a high rate of spontaneous migration during the first year of life and greater than 90% of testes being viable on follow-up. Laparoscopic orchidopexy is a safe and feasible option for testes that remain intraabdominal at follow-up [62]. In contrast, the rate of spontaneous testicular descent was only 10% in boys with omphalocele [61].

References

1. Opitz, J.M.; Feldkamp, M.L.; Botto, L.D. An evolutionary and developmental biology approach to gastroschisis. *Birth Defects Res.* 2019, 111, 294–311.
2. Moore, T.C.; Stokes, G.E. Gastroschisis; Report of two cases treated by a modification of the gross operation for omphalocele. *Surgery* 1953, 33, 112–120.
3. Duhamel, B. Embryology of exomphalos and allied malformations. *Arch. Dis. Child.* 1963, 38, 142–147.
4. Shaw, A. The myth of gastroschisis. *J. Pediatr. Surg.* 1975, 10, 235–244.
5. De Vries, P.A. The pathogenesis of gastroschisis and omphalocele. *J. Pediatr. Surg.* 1980, 15, 245–251.
6. Hoyme, H.E.; Higginbottom, M.C.; Jones, K.L. The vascular pathogenesis of gastroschisis; intrauterine interruption of the omphalomesenteric artery. *J. Pediatr.* 1981, 98, 228–231.
7. Feldkamp, M.L.; Carey, J.C.; Sadler, T.W. Development of gastroschisis: Review of hypotheses, a novel hypothesis, and implications for research. *Amer. J. Med. Genet.* 2007, 143A, 639–652.
8. Stevenson, R.E.; Rogers, R.C.; Chandler, J.C.; Gauderer, M.W.L.; Hunter, A.G.W. Escape of the yolk sac: A hypothesis to explain the embryogenesis of gastroschisis. *Cin. Genet.* 2009, 75, 326–333.
9. Bargy, F.; Beaudoin, S. Comprehensive developmental mechanisms in Gastroschisis. *Fetal Diagn. Ther.* 2014, 36, 223–230.
10. Lubinsky, M. A vascular and thrombotic model of gastroschisis. *Amer. J. Med. Genet.* 2014, 164A, 915–917.
11. Rittler, M.; Vauthay, L.; Mazzitelli, N. Gastroschisis is a defect of the umbilical ring: Evidence from morphological evaluation of stillborn fetuses. *Birth Defects Res. A Clin. Mol. Teratol.* 2013, 97, 198–209.
12. Beaudoin, S. Insights into the etiology and embryology of gastroschisis. *Semin. Pediatr. Surg.* 2018, 27, 283–288.
13. Kirby, R.S.; Marshall, J.; Tanner, J.P.; Salemi, J.L.; Feldkamp, M.L.; Marengo, L.; Meyer, R.E.; Druschel, C.M.; Rickard, R.; Kucik, J.E.; National Birth defects Prevention Network. Prevalence and correlates of gastroschisis in 15 states, 1995 to 2005. *Obstetr. Gynecol.* 2013, 122, 275–281. Available online: https://journals.lww.com/greenjournal/Fulltext/2013/08000/Prevalence_and_Correlates_of_Gastroschisis_in_15.14.aspx (accessed on 4 August 2020).

14. Castilla, E.E.; Mastroiacovo, P.; Orioli, I.M. *Gastroschisis: International Epidemiology and Public Health Perspectives*. *Amer. J. Med. Genet. Part C Semin. Med. Genet.* 2008, **148C**, 162–179.
15. Short, T.D.; Stallings, E.B.; Isenburg, J.; O’Leary, L.A.; Yazdy, M.M.; Bohm, M.K.; Ethen, M.; Chen, X.; Tran, T.; Fox, D.J.; et al. *Gastroschisis trends and ecologic link to opioid prescription rates—United States, 2006–2015*. *Morb. Mortal. Wkly. Rep.* 2019, **68**, 31–36.
16. Loane, M.; Dolk, H.; Bradbury, I.; EUROCAT Working Group. Increasing prevalence of gastroschisis in Europe 1980–2002; a phenomenon restricted to younger mothers? *Pediatr. Perinat. Epidemiol.* 2007, **21**, 363–369.
17. Skarsgard, E.D.; Meaney, C.; Bassil, K.; Brindle, M.; Arbour, L.; Moineddin, R.; Canadian Pediatric Surgery Network (CAPSNet). *Maternal Risk Factors for Gastroschisis in Canada*. *Birth Def. Res.* 2015, **103**, 111–118.
18. Jones, A.M.; Isenburg, J.; Salemi, J.L.; Arnold, K.E.; Mai, C.T.; Aggarwal, D.; Arias, W.; Carrino, G.E.; Ferrell, E.; Folorunso, O.; et al. *Increasing Prevalence of Gastroschisis—14 States, 1995–2012*. *Morb. Mortal. Wkly. Rep.* 2016, **65**, 23–26.
19. Stallings, E.B.; Isenburg, J.L.; Short, T.D.; Heinke, D.; Kirby, R.S.; Romitti, P.A.; Canfield, M.A.; O’Leary, L.A.; Liberman, R.F.; Forestieri, N.E.; et al. *Population-based birth defects in the United States, 2012–2016: A focus on abdominal wall defects*. *Birth Def. Res.* 2019, **111**, 1436–1447.
20. Baldacci, S.; Santoro, M.; Coi, A.; Mezzasalma, L.; Bianchi, F.; Pierini, A. *Lifestyle and sociodemographic risk factors for gastroschisis; a systematic review and meta-analysis*. *Arch. Dis. Child.* 2020, **105**, 756–764, doi:10.1136/archdischild-2019-318412.
21. Chabra, S.; Gleason, C.A.; Seidel, K.; Williams, M.A. *Rising Prevalence of Gastroschisis in Washington State*. *J. Toxicol. Environ. Health. A* 2011, **74**, 336–345.
22. Archer, N.P.; Langlois, P.H.; Suarez, L.; Brender, J.; Shanmugam, R. *Association of Paternal Age with Prevalence of Selected Birth Defects*. *Birth Defects Res. A Clin. Mol. Teratol.* 2007, **79**, 27–34.
23. Green, R.F.; Devine, O.; Crider, K.S.; Olney, R.S.; Archer, N.; Olshan, A.F.; Shapira, S.K.; National Birth Defects Prevention Study. *Association of Paternal Age and Risk for Major Congenital Anomalies from the National Birth Defects Prevention Study, 1997 to 2004*. *Ann. Epidemiol.* 2010, **20**, 241–249.
24. Kazaura, M.R.; Lie, R.T.; Irgens, L.M.; Didriksen, A.; Kapstad, M.; Egenaes, J.; Bjerkedal, T. *Increasing Risk of Gastroschisis in Norway: An Age-Period-Cohort Analysis*. *Am. J. Epidemiol.* 2004, **159**, 358–363.
25. Feldkamp, M.L.; Arnold, K.E.; Krikov, S.; Reehuis, J.; Almli, L.M.; Moore, C.A.; Botto, L.D. *Risk of gastroschisis with maternal genitourinary infections: The US National birth defects prevention study 1997–2011*. *BMJ Open* 2019, **9**, e026297, doi:10.1136/bmjopen-2018-026297.
26. Root, E.D.; Meyer, R.E.; Emch, M.E. *Evidence of Localized Clustering of Gastroschisis Births in North Carolina, 1999–2004*. *Soc. Sci. Med.* 2009, **68**, 1361–1367.
27. Bassil, K.L.; Yang, J.; Arbour, L.; Moineddin, R.; Brindle, M.E.; Hazell, E.; Skarsgard, E.D. *Spatial variability of gastroschisis in Canada, 2006–2011: An exploratory analysis*. *Can. J. Public Health* 2016, **107**, e62–e67.
28. Waller, S.A.; Paul, K.; Peterson, S.E.; Hitti, J.E. *Agricultural-related Chemical Exposures, Season of Conception, and Risk of Gastroschisis in Washington State*. *Am. J. Obstet. Gynecol.* 2010, **202**, 241.e1–e6.

29. Souther, C.; Puapong, D.P.; Woo, R.; Johnson, S.M. Possible etiologies of increased incidence of gastroschisis. *Pediatr. Surg. Int.* 2017, 33, 1209–1213.

30. Molik, K.A.; Gingalewski, C.A.; West, K.W.; Rescorla, F.J.; Scherer, L.R.; Engum, S.A.; Grosfeld, J.L. Gastroschisis: A Plea for Risk Categorization. *J. Pediatr. Surg.* 2001, 36, 51–55.

31. Perrone, E.E.; Olson, J.; Golden, J.M.; Besner, G.E.; Gayer, C.P.; Islam, S.; Gollin, G. Closing Gastroschisis: The Good, the Bad, and the Not-So Ugly. *J. Pediatr. Surg.* 2019, 54, 60–64.

32. Bergholz, R.; Boettcher, M.; Reinshagen, K.; Wenke, K. Complex Gastroschisis Is a Different Entity to Simple Gastroschisis Affecting Morbidity and Mortality—A Systematic Review and Meta-Analysis. *J. Pediatr. Surg.* 2014, 49, 1527–1532.

33. Suver, D.; Lee, S.L.; Shekherdimian, S.; Kim, S.S. Left-sided Gastroschisis: Higher Incidence of Extraintestinal Congenital Anomalies. *Am. J. Surg.* 2008, 195, 663–666.

34. Yaster, M.; Scherer, T.L.; Stone, M.M.; Schleien, C.L.; Wetzel, R.C.; Buck, J.R.; Nichols, D.; Colombani, P.M.; Dudgeon, D.L. Prediction of successful primary closure of congenital abdominal wall defects using intraoperative measurements. *J. Pediatr. Surg.* 1989, 24, 1217–1220.

35. Olesevich, M.; Alexander, F.; Khan, M.; Cotman, K. Gastroschisis revisited: Role of intraoperative measurement of abdominal pressure. *J. Pediatr. Surg.* 2005, 40, 789–792.

36. Petrosyan, M.; Sandler, A.D. Closure methods in gastroschisis. *Semin. Pediatr. Surg.* 2018, 27, 304–308.

37. Sandler, A.; Lawrence, J.; Meehan, J.; Phearman, L.; Soper, R. A “plastic” sutureless abdominal wall closure in gastroschisis. *J. Pediatr. Surg.* 2004, 39, 738–741.

38. Kidd, J.N. Jr.; Jackson, R.J.; Smith, S.D.; Wagner, C.W. Evolution of staged versus primary closure of gastroschisis. *Ann. Surg.* 2003, 237, 759–765.

39. Schlatter, M.; Norris, K.; Uitvlugt, N.; DeCou, J.; Connors, R. Improved outcomes in the treatment of gastroschisis using a preformed silo and delayed repair approach. *J. Pediatr. Surg.* 2003, 38, 459–464.

40. Kunz, S.N.; Tieder, J.S.; Whitlock, K.; Jackson, J.C.; Avansino, J.R. Primary fascial closure versus staged closure with silo in patients with gastroschisis: A meta-analysis. *J. Pediatr. Surg.* 2013, 48, 845–857.

41. Hawkins, R.B.; Raymond, S.L.; St Peter, S.D.; Downard, C.D.; Qureshi, F.G.; Renaud, E.; Danielson, P.D.; Islam, S. Immediate Versus Silo Closure for Gastroschisis: Results of a Large Multicenter Study. *J. Pediatr. Surg.* 2020, 55, 1280–1285, doi:10.1016/j.jpedsurg.2019.08.002.

42. Pastor, A.C.; Phillips, J.D.; Fenton, S.J.; Meyers, R.L.; Lamm, A.W.; Raval, M.V.; Lehman, E.; Karp, T.B.; Wales, P.W.; Langer, J.C. Routine use of a SILASTIC spring-loaded silo for infants with gastroschisis: A multicenter randomized controlled trial. *J. Pediatr. Surg.* 2008, 43, 1807–1812.

43. Poola, A.S.; Aguayo, P.; Fraser, J.D.; Hendrickson, R.J.; Weaver, K.L.; Gonzalez, K.W.; Peter, S.D.S. Primary Closure versus Bedside Silo and Delayed Closure for Gastroschisis: A Truncated Prospective Randomized Trial. *Eur. J. Pediatr. Surg.* 2019, 29, 203–208.

44. Emami, C.N.; Youssef, F.; Baird, R.J.; Laberge, J.-M.; Skarsgard, E.D.; Puligandla, P.S. A risk-stratified comparison of fascial versus flap closure techniques on the early outcomes of infants with gastroschisis. *J. Pediatr. Surg.* 2015, 50, 102–106.

45. Pet, G.E.; Stark, R.A.; Meehan, J.J.; Javid, P.J. Outcomes of bedside sutureless umbilical closure without endotracheal intubation for gastroschisis repair in surgical infants. *Am. J. Surg.* 2017, 213, 958–962.

46. Witt, R.G.; Zobel, M.; Padilla, B.; Lee, H.; MacKenzie, T.C.; Vu, L. Evaluation of Clinical Outcomes of Sutureless vs Sutured Closure Techniques in Gastroschisis Repair. *JAMA Surg.* 2019, 154, 33–39.

47. Bruzoni, M.; Jaramillo, J.D.; Dunlap, J.L.; Abrajano, C.; Stack, S.W.; Hintz, S.R.; Hernandez-Boussard, T.; Dutta, S. Sutureless vs Sutured Gastroschisis Closure: A Prospective Randomized Controlled Trial. *J. Am. Coll. Surg.* 2017, 224, 1091–1096.e1.

48. Youssef, F.; Gorgy, A.; Arbash, G.; Puligandla, P.S.; Baird, R.J. Flap versus fascial closure for gastroschisis: A systematic review and meta-analysis. *J. Pediatr. Surg.* 2016, 51, 718–725.

49. Jadcherla, S.R.; Gupta, A.; Stoner, E.; Fernandez, S.; Caniano, D.; Rudolph, C.D. Neuromotor markers of esophageal motility in feeding intolerant infants with gastroschisis. *J. Pediatr. Gastroenterol. Nutr.* 2008, 47, 158–164.

50. Jayanthi, S.; Seymour, P.; Puntis, J.W.; Stringer, M.D. Necrotizing enterocolitis after gastroschisis repair: A preventable complication? *J. Pediatr. Surg.* 1998, 33, 705–707.

51. Curry, J.I.; Lander, A.D.; Stringer, M.D.; BAPS Multicentre Research Committee. A multicenter, randomized, double-blind, placebo-controlled trial of the prokinetic agent erythromycin in the postoperative recovery of infants with gastroschisis. *J. Pediatr. Surg.* 2004, 39, 565–569.

52. Lusk, L.A.; Brown, E.G.; Overcash, R.T.; Grogan, T.R.; Keller, R.L.; Kim, J.H.; Poulain, F.R.; Shew, S.B.; Uy, C.; DeUgarte, D.A. University of California Fetal Consortium. Multi-institutional practice patterns and outcomes in uncomplicated gastroschisis: A report from the University of California Fetal Consortium (UCfC). *J. Pediatr. Surg.* 2014, 49, 1782–1786.

53. DeUgarte, D.A.; Calkins, K.L.; Guner, Y.; Kim, J.; Kling, K.; Kramer, K.; Lee, H.; Lusk, L.; Saadai, P.; Uy, C.; et al. University of California Fetal Consortium. Adherence to and outcomes of a University-Consortium gastroschisis pathway. *J. Pediatr. Surg.* 2020, 55, 45–48.

54. Fullerton, B.S.; Velazco, C.S.; Sparks, E.A.; Morrow, K.A.; Edwards, E.M.; Soll, R.F.; Modi, B.P.; Horbar, J.D.; Jaksic, T. Contemporary Outcomes of Infants with Gastroschisis in North America: A Multicenter Cohort Study. *J. Pediatr.* 2017, 188, 192–197.e6.

55. Fawley, J.A.; Abdelhafeez, A.H.; Schultz, J.A.; Ertl, A.; Cassidy, L.D.; St Peter, S.; Wagner, A.J. The risk of midgut volvulus in patients with abdominal wall defects: A multi-institutional study. *J. Pediatr. Surg.* 2017, 52, 26–29.

56. De Marco, G.; Sordino, D.; Bruzzese, E.; Di Caro, S.; Mambretti, D.; Tramontano, A.; Colombo, C.; Simoni, P.; Guarino, A. Early treatment with ursodeoxycholic acid for cholestasis in children on parenteral nutrition because of primary intestinal failure. *Aliment. Pharmacol. Ther.* 2006, 24, 387–394.

57. Muhammed, R.; Bremner, R.; Protheroe, S.; Johnson, T.; Holden, C.; Murphy, M.S. Resolution of parenteral nutrition-associated jaundice on changing from a soybean oil emulsion to a complex mixed-lipid emulsion. *J. Pediatr. Gastroenterol. Nutr.* 2012, 54, 797–802.

58. Jensen, A.R.; Goldin, A.B.; Koopmeiners, J.S.; Stevens, J.; Waldhausen, J.H.; Kim, S.S. The association of cyclic parenteral nutrition and decreased incidence of cholestatic liver disease in patients with gastroschisis. *J. Pediatr. Surg.* 2009, 44, 183–189.

59. Amin, S.C.; Pappas, C.; Iyengar, H.; Maheshwari, A. Short bowel syndrome in the NICU. *Clin. Perinatol.* 2013, 40, 53–68.
60. Iyer, K.R. Surgical management of short bowel syndrome. *JPEN J Parenter. Enteral Nutr.* 2014, 38 (Suppl. 1), 53S–59S.
61. Yardley, I.E.; Bostock, E.; Jones, M.O.; Turnock, R.R.; Corbett, H.J.; Losty, P.D. Congenital abdominal wall defects and testicular maldescent—A 10-year single-center experience. *J. Pediatr. Surg.* 2012, 47, 1118–1122.
62. Hill, S.J.; Durham, M.M. Management of cryptorchidism and gastroschisis. *J. Pediatr. Surg.* 2011, 46, 1798–1803.

Retrieved from <https://encyclopedia.pub/entry/history/show/16295>