

# Infections&Immunotherapy in Lung Cancer

Subjects: **Others**

Contributor: Sara Pilotto

Infectious diseases represent a relevant issue in lung cancer patients. Bacterial and viral infections might influence the patients' prognosis, both directly affecting the immune system and indirectly impairing the outcome of anticancer treatments, mainly immunotherapy. In this analysis, we aimed to review the current evidence in order to clarify the complex correlation between infections and lung cancer. In detail, we mainly explored the potential impact on immunotherapy outcome/safety of (1) bacterial infections, with a detailed focus on antibiotics; and (2) viral infections, discriminating among (a) human immune-deficiency virus (HIV), (b) hepatitis B/C virus (HBV-HCV), and (c) Sars-Cov-2. A series of studies suggested the prognostic impact of antibiotic therapy administration, timing, and exposure ratio in patients treated with immune checkpoint inhibitors, probably through an antibiotic-related microbiota dysbiosis. Although cancer patients with HIV, HBV, and HCV were usually excluded from clinical trials evaluating immunotherapy, some retrospective and prospective trials performed in these patient subgroups reported similar results compared to those described in not-infected patients, with a favorable safety profile. Moreover, patients with thoracic cancers are particularly at risk of COVID-19 severe outcomes and mortality. Few reports speculated about the prognostic implications of anticancer therapy, including immunotherapy, in lung cancer patients with concomitant Sars-Cov-2 infection, showing, to date, inconsistent results. The correlation between infectious diseases and immunotherapy remains to be further explored and clarified in the context of dedicated trials. In clinical practice, the accurate and prompt multidisciplinary management of lung cancer patients with infections should be encouraged in order to select the best treatment options for these patients, avoiding unexpected toxicities, while maintaining the anticancer effect.

non-small cell lung cancer

antibiotic therapy

microbiota

HIV

antiretroviral therapy

HBV

HCV

COVID-19

## 1. Introduction

The use of immune checkpoint inhibitors (ICIs), both alone or in combination with chemotherapy, has radically changed the treatment algorithm of locally advanced and metastatic non-small cell lung cancer (NSCLC), improving patients' survival with a favorable safety profile [1].

Nevertheless, ICIs' efficacy and safety in less-investigated special subgroups of lung cancer patients are still far from being clarified. Indeed, lung cancer patients could suffer from concomitant infections.

With regard to bacterial infections, to which lung cancer patients are particularly susceptible, several studies reported the prognostic implications of antibiotic therapy administration, timing, and cumulative exposure ratio (proportion between days of antibiotic therapy and days of immunotherapy) [2][3].

Patients concurrently affected by lung cancer and viral infections, such as the human immune-deficiency virus (HIV), hepatitis B virus (HBV), and hepatitis C virus (HCV), have usually been excluded from those clinical trials that led to immunotherapy regulatory approval, mainly due to concerns about tolerance, efficacy, and risk of viral reactivation. However, a series of case reports, retrospective, and recent prospective studies are available, suggesting that ICIs seem to be safe and active in people living with HIV (PLWH) and in patients affected by chronic HBV/HCV, without reliable risks of viral reactivation [4][5].

Among viral infections, since the end of 2019, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which causes the coronavirus disease 2019 (COVID-19), is progressively spreading worldwide [6]. Overall, patients with cancer appeared to be at higher risk of severe events and death compared with non-cancer patients [7][8][9]. Among oncological patients, those affected by lung cancer seem to be particularly vulnerable, with an estimated mortality rate of around 32% [10]. Nevertheless, data exploring the potential correlation between the type and timing of anticancer therapies, including immunotherapy, and COVID-19 outcome are still preliminary.

## 2. Bacterial Infections in Lung Cancer

Patients affected by NSCLC have a higher risk of experiencing recurrent infections related to the oncological disease itself, to frequently co-occurring factors, such as old age, smoking history, presence of chronic obstructive pulmonary disease (COPD) [11][12], as well as to the immunosuppression related to oncological therapies [13]. The majority of infections involve the respiratory tract, mainly bacterial pneumonia, typically diagnosed with productive cough with fever and/or radiological infiltrates [13][14]. Other frequent infections may involve blood, urinary, or the gastrointestinal tract [15].

According to the etiology, the most commonly administered antibiotics are broad spectrum, mainly  $\beta$ -lactams (+/- inhibitors), fluoroquinolones, and macrolides [13][14].

### 2.1. Antibiotics and Immune Checkpoint Inhibitors in Lung Cancer

Despite previously unseen results in non-oncogene addicted NSCLC, only 20–40% of patients experience a durable benefit from immunotherapy [16], suggesting the importance of appropriately identifying those clinical, pathological, and/or laboratory parameters able to predict an optimized application of ICIs. Tumor-related factors alone, such as programmed death-ligand 1 (PD-L1) expression, tumor mutational burden, tumor-infiltrating lymphocytes, and high microsatellite instability, do not completely explain the existing differences in immunotherapy efficacy [17].

Considering this, many studies have focused on host factors. Among them, gut microbiota, clearly emerged as a crucial player in modulating the activity and toxicity of anticancer treatments, including immunotherapy [18]. In this regard, the development of an altered systemic immune response, mediated by gut microbiota dysbiosis [19], might contribute to explaining the reported worse prognosis observed with the concomitant administration of antibiotic therapy and ICIs [20].

Therefore, in the last few years, several studies and meta-analyses suggested the potential negative prognostic impact of antibiotics administration in immunotherapy-treated patients ([Table 1](#)).

**Table 1.** Meta-analyses evaluating the impact of antibiotics administration on clinical outcomes of advanced cancer patients treated with immune-checkpoint inhibitors, focusing on results obtained in non-small-cell lung cancer.

Meta-Analysis.	N. of Included Studies	Pooled HR PFS		pooled HR OS	NSCLC-HR	NSCLC-HR OS
		[95% CI]	p-Value	[95% CI]	PFS [95% CI]	[95% CI]
[21]	19	1.84 [1.49–2.26]	p < 0.001	2.37 [2.05–2.75]	p < 0.001	1.79 [1.29–2.49] 2.68 [2.19–3.28]
[22]	18	1.65 [1.3–2.1]	p < 0.0001	1.92 [1.37–2.68]	p < 0.001	1.64 [1.07–2.52] 2.00 [1.23–3.24]
[23]	20	1.53 [1.30–1.79]	p < 0.01	1.90 [1.55–2.34]	p < 0.01	1.39 [1.16–1.67] 1.73 [1.26–2.38]
[24]	15	1.53 [1.22–1.93]	p < 0.01	2.07 [1.51–2.84]	N.A.	N.A.

[25] *	23	N.A.	N.A.	1.47 [1.13–1.90]	1.69 [1.25–2.29]
				<i>p</i> < 0.01	<i>p</i> < 0.01
[2]	33	1.76 [1.47–2.12]	1.76 [1.41–2.19]	1.70 [1.21–2.27]	1.80 [1.28–2.55]
		<i>p</i> < 0.00001	<i>p</i> < 0.00001	<i>p</i> = 0.0004	<i>p</i> = 0.0008

HR, hazard ratio; OS, overall survival; PFS, progression-free survival; NSCLC, non-small-cell lung cancer; N.A., not available. \* Only NSCLC patients were included.

The first meta-analysis that systematically evaluated the association between antibiotic therapy and clinical efficacy of ICIs in solid cancers was conducted in 2019 [21]. A statistically significant reduction in both overall survival (OS) and progression-free survival (PFS) was globally observed and confirmed for NSCLC in the subgroup analysis stratified by cancer type (OS: HR 2.68, 95% CI 2.19–3.28, *p* < 0.001; PFS: HR 1.79, 95% CI 1.29–2.49, *p* < 0.001). Regarding the time window of exposure, the survival impact of antibiotics when used within 60 days from immunotherapy initiation was weaker (HR 1.97, 95% CI 1.49–2.59, *p* < 0.001) if compared with their use within 30 days (HR 2.23, 95% CI 1.82–2.74, *p* < 0.001) [21].

## 2.2. Antibiotics Timing and/or Cumulative Exposure: What Matters?

The time window of antibiotics exposure in relation to immunotherapy initiation seems to represent a crucial factor in determining the different outcomes of patients affected by NSCLC, as well as other solid tumors. While a statistically significant detrimental prognosis was consistently documented when antibiotics were administered shortly before and/or after ICIs initiation, this impact vanishes when the timeframe included immunotherapy course and later times.

In this sense, a monocentric analysis performed in cancer patients enrolled in ICIs phase I clinical trials reported a statistically significant negative effect on OS only in case of antibiotic use within 30 days prior to immunotherapy initiation (HR 2.00, 95% CI 1.2–3.3, *p* = 0.011) [26]. This finding was further confirmed in a prospective multicenter study, suggesting that prior antibiotic therapy (within 30 days before immunotherapy initiation), but not concurrent, was associated with worse ICIs activity and efficacy in patients treated in routine clinical practice (OS: HR 7.4, 95% CI 4.3–12.8, *p* < 0.001) [27]. In NSCLC patients, OS was 2.5 months in case of antibiotics administration within 30 days before ICIs therapy versus 26 months without any antibiotic exposure (*p* < 0.001) [27].

On the other hand, the detrimental prognostic effect of the antibiotic administration in ICIs-treated NSCLC patients might be associated with the antibiotic-immunotherapy exposure ratio (AIER), a candidate new variable defined as the proportion between days of antibiotic therapy and days of immunotherapy [3]. During the entire immunotherapy

course, a higher AIER was reported to be related with a worse outcome, when considered as both a continuous variable (PFS: HR 1.053,  $p = 0.0029$ , OS: HR 1.064,  $p < 0.0001$ ), as well as using the cutoff value of 4.2% (the median AIER in the whole immunotherapy treatment). In detail, when comparing patients with AIER < 4.2% and  $\geq 4.2\%$ , the median PFS was 3.5 versus 1.9 months ( $p < 0.0001$ ) and the median OS was 13.2 versus 5.1 months ( $p = 0.0004$ ). In contrast with the previously described meta-analyses, the antibiotic administration in the early ICIs period did not significantly influence patients' prognosis [3]. Similarly, a retrospective analysis performed in ICIs-treated patients affected by advanced solid tumors, supporting the potential stronger prognostic impact of the cumulative antibiotic exposure variable compared with the administration timeframe [28].

## 2.3. Host Microbiome, Immunotherapy, and Antibiotics in Lung Cancer

The host microbiome, and gut microbiota in particular, has been widely recognized as a fundamental modulator of cancer development and progression, as well as of anticancer treatments' efficacy and toxicity [29]. Among them, several chemotherapeutic drugs, such as 5-fluorouracil, gemcitabine, cyclophosphamide, irinotecan, cisplatin, and tyrosine kinase inhibitors, acting as targeted agents are included [30].

Regarding immunotherapy, commensal microorganisms might influence systemic immune functions and, therefore, treatment response through different mechanisms. A high diversity of the gut microbiome triggers Th1 lymphocyte and M1 macrophage differentiation, activation of helper and cytotoxic T cells, and upregulation of PD-1 expression [31]. In terms of composition, preclinical data showed that an overrepresentation of *Bifidobacterium* species in the gut microbiota of melanoma mouse models increased the response to anti-PD-L1 agents, with the possibility of restoring the antitumor activity through the oral administration of commensal *Bifidobacteria* in those mice harboring an unfavorable microbiota [32]. Moreover, the antitumor activity of cytotoxic T-lymphocyte antigen-4 (CTLA-4) inhibitors changed according to distinct *Bacteroides* species. Indeed, fecal transplantation with peculiar *Bacteroides* species (such as *B. fragilis*) from human to mice allowed remarkable responses to CTLA-4 blockade to be achieved [33]. The modification of gut microbiota with fecal transplantation or oral supplementation was also studied in a series of patients, including NSCLC, demonstrating that the clinical response to PD-1 blockade was correlated to the abundance in patients' stool samples of *Akkermansia muciniphila*, probably potentiating the T lymphocytes-mediated immune response through an interleukin-12-dependent mechanism [34]. Focusing on NSCLC, responders to nivolumab had a higher gut microbiota diversity at treatment initiation, remaining stable during therapy [35].

In this light, antibiotics lead to an abnormal gut microbiota composition, potentially impairing ICIs efficacy, through different mechanisms. First, antibiotics administration might decrease the diversity of intestinal microbiota, eliminating the most immunogenic bacteria [36]. Second, early antibiotic use might influence the plasma levels of citrulline, a recognized biomarker of intestinal barrier and enterocytes integrity and functioning, that correlate with the clinical outcome in nivolumab-treated NSCLC patients [37]. Third, when commensal microbes decrease and pathogenic bacteria increase, toxins (such as the cytolethal distending toxin, the cytotoxic necrotizing factor-1, and the *B. fragilis* toxin), hydrogen sulfide, superoxide radicals and carcinogens, such as acetaldehyde, are abnormally released, potentially triggering DNA damages. Moreover, dysbiosis induces the lack of those anti-inflammatory

actions mediated by the short-chain fatty acids normally fermented by the healthy microbiota, thus inducing an aberrant proliferation of the epithelial cells and, potentially, the carcinogenesis process [38]. Finally, another candidate mechanism is represented by the alteration of the  $\beta$ -catenin pathway, leading to loss of cell polarity, dysregulation of cellular growth, and acquisition of stem cell-like characteristics, due to the direct bond of microbial proteins to E-cadherin and/or to the translocation of some effectors into the host cells' cytoplasm. Similar mechanisms are speculated to also be involved in the carcinogenesis process [39].

Although most of the available data in the oncological setting explored gut microbiota, a less-investigated but very promising microbial community is represented by the respiratory microbiota. The composition of lung microbiota is different from that characterizing other body districts and might be altered by smoking and environmental exposures [40]. Specifically, it seems to be determined by the balance of three main factors: microbial immigration, microbial elimination, and relative reproduction rates of its members [41]. Although specific communities are particularly enriched in lung cancer patients [42][43][44], the underlying mechanisms supporting this correlation are still far from being clarified. Nevertheless, the inflammatory response induced by quantitative and qualitative changes in lung microbiota is likely to be a determinant, as for other chronic respiratory diseases [45][46][47]. Interleukin-17C induced by aberrant bacteria (e.g., *Haemophilus influenzae*) increased the level of neutrophilic inflammation in COPD patients, thus mediating tumor proliferative effects [43]. Moreover, T helper cells induced by bacteria could promote angiogenesis and lung cancer cell proliferation [41]. In this light, epidemiological data suggested a correlation between prolonged antibiotic exposure and lung cancer development [48]. Finally, a complex bidirectional crosstalk between gut and lung microbiota has been demonstrated, in terms of both microbes transfer through bloodstream and lymphatic system and systemic immune response modulation. For example, short-chain fatty acids (SCFAs), metabolized by gut microbes and released in the systemic circulation, modulate several immune and epithelial cells functions through the regulation of G-protein-coupled receptors and histone deacetylase [49].

## 2.4. Open Issues and Future Perspectives in NSCLC Patients Treated with ICIs and Antibiotics

Although most of the available data are consistent in suggesting the negative prognostic impact of antibiotics administration in immunotherapy-treated patients, particularly if used shortly before and/or after ICIs initiation, other studies did not show this difference [3][50] or, even, they observed an opposite correlation [51][52].

Lack of definitive data in this setting might be related to different limitations. First, most of the available studies were retrospective, heterogenous in terms of patients' selection, comorbidities, tumor burden, and treatment line. Second, the concurrent use of other drugs potentially modulating the immune response and host microbiome, such as corticosteroids and proton pump inhibitors (PPI), should be considered. In this regard, no detrimental effects were observed in terms of ICIs outcome when corticosteroids were administered as premedication, but their prognostic impact according to different dosages, timeframes, treatment durations, and administration routes is still debated [53]. Regarding PPI, in a pooled analysis of the OAK and POPLAR trials, Chalabi et al. reported a shorter PFS (1.9 versus 2.8 months; HR 1.30, 95% CI 1.10–1.53,  $p = 0.001$ ) and OS (9.6 versus 14.5 months; HR 1.45,

95% CI 1.20–1.75,  $p = 0.0001$ ) for PPI-treated versus not-treated NSCLC patients undergoing atezolizumab [54]. In addition, a strong heterogeneity was also observed in terms of antibiotic type, single or multiple antimicrobial drugs combinations, treatment durations, and administration routes. The retrospective study of *Mielgo-Rubio* et al. suggested that intravenous administration might be associated with a worse prognosis than oral (OS: 2.9 versus 14.2 months,  $p = 0.0001$ ; PFS: 2.2 versus 5.9 months,  $p = 0.001$ ) in advanced immunotherapy-treated NSCLC patients [55]. An additional crucial and still-unanswered question is whether the suspected detrimental effect on ICIs outcomes is really due to the antibiotics, rather than to the infection itself, as a direct cause or consequence of a systemic immune system impairment.

Finally, although microbiome dysbiosis is the most reliable mechanism justifying the harmful effect of antibiotics, a systematic evaluation of gut (and lung) microbiota composition and perturbation before, during, and after antibiotics and ICIs administration should be performed to provide concrete data. Moreover, when dealing with the microbiome, the existence of other potential modulators, such as the dietary pattern and the geographical origin, has to be considered within dedicated studies.

## 3. Viral Infections in Lung Cancer

Chronic viral infections represent frequent comorbidities in lung cancer patients. Furthermore, viral infections, such as HIV, HBV, and HCV, can increase the risk of developing malignancies, including lung cancer [56][57]. Specific antiviral therapies, such as antiretroviral therapy (ART) for HIV and antiviral drugs for HBV/HCV, are now the standard of care in chronic viral infections [58]. To date, most of the clinical trials testing immunotherapy in solid cancers have excluded patients affected by chronic viral infections, such as HIV, HBV, and HCV, due to concerns about viral reactivation, as well as ICIs safety and efficacy in these peculiar populations. Nowadays, dealing with viral infections, a specific mention should be made about the current Sars-Cov-2 pandemic that is spreading worldwide and features high aggressiveness and mortality in oncological patients, particularly in thoracic malignancies.

### 3.1. Human Immune-Deficiency Virus (HIV)

#### 3.1.1. HIV and Risk of Lung Cancer

Due to active antiretroviral therapy (ART), in the last years, the mortality and morbidity of PLWH has progressively decreased [59]. The HIV infection was historically associated with an increased risk of solid and hematologic cancers, including aggressive non-Hodgkin lymphoma, Kaposi sarcoma, cervical, and anal cancer, globally representing the acquired immune deficiency syndrome (AIDS)-defining cancers (ADCs) [60]. Nevertheless, due to the recent improved control of HIV infections, HIV-positive patients have a higher risk of developing other types of solid tumors, as lung cancer, gastrointestinal carcinoma, and skin cancer (known as non-AIDS-defining cancers, NADCs) [57]. Overall, cancer represents a leading cause of death for more than 37 million people worldwide living with HIV [61][62].


Regarding the risk of developing lung cancer, several factors might be implicated, such as the increased smoking habit in HIV-positive patients [63], the immunosenescence phenomenon due to their longer life expectancy [64], as well as the negative modulation of the immune system caused by the virus-related chronic inflammation [65]. Of interest, in the United States, HIV-positive patients, smokers, with a good disease control with ART are more likely to die for NSCLC than for AIDS [66].

### 3.1.2. HIV and Immunotherapy in Lung Cancer

For several HIV-associated cancers, including lung cancer, treatment outcomes seem to be similar to those observed in the overall oncological population [67].

Regarding immunotherapy, although the use of ICIs might be theoretically beneficial for treating HIV-associated cancers, PLWH has usually been excluded from pivotal clinical trials testing these drugs, due to concerns about ICIs safety/efficacy and risk of viral reactivation.

Preclinical data suggested that PD-1/PD-L1 blockade might induce an increased functioning of HIV-specific T cells. In detail, during chronic HIV infection, virus-specific CD8+ T cells might undergo a PD-1-mediated functional exhaustion, reducing proliferation and cytokine production. Indeed, PD-1 expression has been correlated with T cell exhaustion and disease progression in HIV-infected patients [68]. Furthermore, an effective ART has been shown to downregulate the expression of PD-1 on CD4+ and CD8+ T cells [69]. With these premises, PD-1 inhibition could play a crucial role in immune reconstitution of HIV-infected patients, highlighting the potential therapeutic benefit of blocking PD-1/PD-L1 interactions for enhancing CD8+ and CD4+ T cells [70]. In vivo studies confirmed that PD-1 blockade induced both an improvement in antitumor immune response and a restoration of the functional quality of virus-specific CD8+ T cells [68] (Figure 1).



**Figure 1.** (a) During chronic HIV infection, high expression of PD-1 leads to a decrease in cytokines production and in HIV-specific CD8+ T cells proliferation. (b) Indeed, PD-1 expression is correlated with T cell exhaustion and disease progression in HIV-infected patients. The antiretroviral therapy (ART) seems to be able to inhibit PD-1 expression on CD8+ and CD4+ T cells. (c) The PD-1 blockade could provide an enhancement in CD8+ and CD4+

T cells and a restoration of the functional quality of virus-specific CD8+ T cells, improving immune response. Furthermore, immune checkpoint inhibitors could reverse HIV latency.

Interestingly, the PD-L1 expression in NSCLC tissues was similar between HIV- and non-HIV-infected patients, but this biomarker was associated with poor prognosis particularly in patients with HIV infection, suggesting a more potent systemic immune suppression through the PD-1/PD-L1 axis in NSCLC patients with HIV than in those not infected [71].

A series of case reports [72][73], retrospective, and recent prospective studies (Table 2) investigated the safety and efficacy of ICIs in HIV patients affected by solid tumors, including lung cancer. In seven HIV-positive advanced NSCLC patients, nivolumab and pembrolizumab demonstrated a good safety profile with a similar efficacy compared to HIV-negative populations [74]. Similar results were obtained in another relatively small population of lung cancer patients [75]. Among 12 HIV-infected patients with advanced NSCLC treated with second-line nivolumab, a favorable clinical outcome (7/12 disease control with 3 partial responses and 4 disease stabilizations), without clinically meaningful side effects (except one case of neurosyphilis) and neither a relevant impact on HIV viral load nor CD4+/CD8+ cell count were reported. Furthermore, nivolumab was able to enhance the capacities of HIV-specific CD8+ cells to proliferate and secrete cytokines, expanding the PD-1 low T cell subset [76].

**Table 2.** Clinical studies investigating safety and efficacy of ICIs in HIV patients affected by solid tumors, including lung cancer.

Trial	Phase	Type of Cancer	N. of Patients	Type of ICIs	Primary Endpoints	Secondary Endpoints	Main Findings	Status
[4] [NCT02595866]	1	Advanced or metastatic solid and hematological cancers	60 HIV-positive *	Pembrolizumab	Safety	ORR, PFS, DoR, OS	Results on 30 patients:  - most common irAEs: fatigue, hypothyroidism, anemia;  - signals of activity in several cancer types;	Recruiting

								- - the only one NSCLC patient obtained CR.
[77]	3b/4 [NCT02869789]	Advanced or metastatic NSCLC	1036 [4 HIV-positive in cohort 1A] §	Nivolumab plus Ipilimumab	Safety	PFS, ORR, DoR, FACT-L, OS	- good safety profile; - - promising OS outcomes.	Active, not recruiting patients described for
[78]	2 [NCT03094286]	Advanced or metastatic solid and hematological cancers	20 HIV-positive [14 NSCLC, 2 melanoma, 1 SCLC, 2 anal carcinoma, 1 bladder carcinoma] §	Durvalumab	Feasibility (ability to receive at least a median number of 4 cycles)	ORR, PFS, OS	[85] - durvalumab was feasible and safe; [86] - DCR (on 16 patients) 50%; [87] - one NSCLC patient with CD4+ cell count<200 had no side effect with long-lasting PR.	Active, not recruiting transplant, treatment owed an ocellular observed latent viral
[79]	1 [NCT02408861]	Advanced or metastatic solid and hematological cancers	96 HIV-positive *	Nivolumab plus Ipilimumab	MTD	ORR, immune function, change in immune	Results on 37 patients: - 11% of patients	BV/HCV patients no severe abilities or the ORR

was 35%, with a median PFS of 4.5 months, including deep and prolonged responses among several patients [5].

### 3.3. Open Issues and Future Perspectives in HIV, HBV, and HCV Lung Cancer Patients Treated with ICIs

The above-mentioned studies consistently supported that lung cancer patients with chronic viral infections, such as HIV, HBV, and HCV, are likely to achieve a clinically meaningful benefit from immunotherapy, comparable to that reported in not-infected oncological patients, without unexpected toxicities.

45,100,000 cases and 1,255,157 deaths [6]. Although we should consider that the currently high COVID-19-related pressure on the healthcare systems might lead to confounding findings, a growing amount of data suggests that oncological patients are particularly vulnerable in terms of risk of both severe events (hospitalization up to 40%, severe respiratory illness 20%) and death (13–35%) compared with patients without a history of cancer [7][8][9][89]. Beyond patient-related parameters, such as age, sex, smoking status, and comorbidities, some cancer-specific factors have been associated with an increased mortality, mainly the PS according to ECOG and the presence of an active oncological disease [89].

Thoracic cancers are at a higher risk of death compared to the general population but also to other cancer types, with an estimated mortality rate of around 32% [10]. Among patients affected by thoracic malignancies, older age

[90]

least a median number of 4 cycles)  
HIV reservoir

erged as

outcome

[7][9][89], the real correlation between the type and timing of anticancer therapies, including immunotherapy, and COVID-19 complications and mortality still needs clarification.

ICIs, immune checkpoint inhibitors; NSCLC, non-small cell lung cancer; HCC, hepatocellular carcinoma; ORR, objective response rate; PFS, progression-free survival; DoR, duration of response; OS, overall survival; irAEs, immune related adverse events; CR, complete response; FACT-L, functional assessment of cancer therapy-lung;

SCLC, small-cell lung cancer; DCR, disease control rate; PR, partial response; MTD, maximum tolerated dose; SOC, standard of care; BOR, best overall response; NA, not available. \*Estimated enrollment, §actual enrollment.

The concomitant diagnosis of bacterial or viral infections in patients affected by lung cancer represents a relevant issue in clinical practice, particularly nowadays that the main therapeutic strategy includes immunotherapy and, therefore, the modulation of the systemic immune response.

An observational French study by the CANCERVIH network, evaluated 23 HIV-positive cancer patients (including 21 advanced NSCLC) treated with nivolumab or pembrolizumab, reporting a good tolerability, a disease control rate (DCR) of 41% and a median OS of 10.7 months. Of note, no significant impact on CD4+ count or HIV viremia was described [82].

antibiotics type, route and duration of administration, type of bacterial infections, concurrent use of other drugs, and ICIs efficacy. Moreover, the role of gut microbiota in response to ICIs and the crosstalk with lung microbiota should be explored in order to prevent wrong use of antibiotics, as well as to identify potential therapeutical approaches to exploit this crosstalk.

objective response rate (ORR) was 13% among 8 NSCLC patients treated with ICIs monotherapy and 75% among

4 NSCLC patients treated with chemoimmunotherapy combination. Again, viral reactivations were not observed [83].

efficacy and safety of immunotherapy for infected NSCLC patients, in a similar way with the general oncological population. A multidisciplinary approach is required for the optimal clinical management of these patients.

A dedicated phase I clinical trial evaluated as the primary endpoint the safety of pembrolizumab in 30 PLWH, 11

with ADCs and 19 with NADCs, including one NSCLC patient (sarcomatoid carcinoma). The most common immune-related adverse events were fatigue, hypothyroidism, and anemia. HIV was controlled in all participants, without any statistically significant increase in CD4+ cells count. In addition, promising signals of activity were observed among patients affected by Kaposi sarcoma, lung cancer, primary effusion lymphoma, and diffuse large

B-cell lymphoma. Of note, the NSCLC patient obtained a complete objective response due to pembrolizumab

Regarding [4] immunotherapy, some reports initially described the occurrence of explosive deteriorations of the patient's clinical condition related to the concomitant ICIs administration and SARS-CoV-2 infection, probably due to a pathological hyperactivation of CD8+ T-cells, inducing an abnormally excessive immune response (cytokine storm), finally leading to the severe acute respiratory distress syndrome [9]. Nevertheless, although the occurrence of a negative synergy cannot always be excluded, other evidence did not find a significant association between

ipilimumab and nivolumab plus ipilimumab. NSCLC was the most prevalent cancer type (34.2%), followed by melanoma (21.9%) and Kaposi sarcoma (12.3%). Overall, 95% of patients were receiving ART at the time of immunotherapy initiation (mainly in monotherapy, but 5 patients received them in combination with anti-PD-1/anti-

CTLA-4 inhibitors). Among 37 patients with a known baseline viral load, 31 (83.8%) had an undetectable viral load at death, differently from chemotherapy, both as a unique modality or in combination with ICIs [90]. A recent update of the COVID-19 and Cancer Consortium (CCCI9) analysis on 3000 patients suggested that the 30-day mortality

was higher among those undergoing an active treatment (with the exception of endocrine therapy), particularly HBV, none of whom showed any treatment-related change in liver function. Among 34 patients with known paired

patients and 3 post-treatment COVID-19 diagnosis. Of suspected 2nd-day mortality was the highest for patients with a 25% post-treatment CD4+ cells values, the count significantly increased with a mean change of 12.3/ $\mu$ L. In terms of antitumor activity, ORR was overall promising: 30% for NSCLC, 27% for melanoma, and 63% for Kaposi's sarcoma.

## Open Issues and Future Perspectives of Sars-Cov-2 Infection in Lung Cancer Patients

The Checkmate 817 trial is a phase 3b/4 study that aimed to evaluate the efficacy and safety of the nivolumab plus ipilimumab combination in metastatic previously untreated NSCLC patients, including a special cohort of ECOG Performance Status (PS) 2 or ECOG PS 0-1 and 1 among asymptomatic untreated brain metastases, hepatic or renal impairment, HIV. Four HIV-positive patients (7%) were enrolled. In 2019, the first results of this trial showed a consistent safety profile and promising OS outcomes in the special advanced NSCLC population.

Beyond safety issues, mainly related to the combination of immunotherapy with cytotoxic chemotherapy, diagnostic difficulties might emerge in discriminating COVID-19 from disease-related symptoms, lung cancer progression, as well as from immune-related adverse events.

Recently, results from the DURVAST trial, a phase 2 study evaluating durvalumab in HIV-infected patients with solid tumors, were published. Among 20 enrolled patients, 14 had NSCLC, 2 melanoma, 1 small-cell lung cancer (SCLC), 2 anal carcinoma, and 1 bladder carcinoma. All patients, receiving ART, showed undetectable HIV plasma viremia. Durvalumab demonstrated to be feasible (the primary endpoint, defined as the ability to receive at least a median number of 4 cycles) and safe. The most common adverse events were diarrhea, arthromyalgia, and asthenia. There were no safety concerns related to HIV reactivation. HIV viremia remained undetectable, and the CD4+/CD8+ T cells count was stable throughout durvalumab treatment. Of note, 1 NSCLC patient with a CD4+ T-cell count of less than 200 cells/mm<sup>3</sup> had no side effects and experienced a long-lasting partial response, suggesting that treating patients with low basal CD4+ T cell counts might also be safe.

In terms of antitumor activity, DCR, evaluated in 16 patients, was 50%, including long-lasting responses. The results of this trial, although limited by the small sample size, suggested a longer duration of clinical benefit in patients treated with INSTIs, a newer class of antiretroviral drugs, allowing the authors to speculate that INSTIs could contribute to the antitumoral immune response of durvalumab.

The results of a phase 1 trial of nivolumab in advanced HIV-associated solid tumors, including NSCLC and SCLC,

## References

the secondary endpoints were the evaluation of nivolumab impact on immune function (HIV viral load, CD4+/CD8+ cells) and objective response. A total of 11% of patients experienced treatment-related serious adverse events, mainly fatigue and rash. An ORR of 24% was observed in immunotherapy-responsive cancers. Moreover, there were no significant changes in HIV viral load during the study period. In light of these findings, the authors concluded that HIV-infected patients treated with ART, having a CD4+ T-cell count greater than 100 cells/ $\mu$ L and undetectable viral load might safely receive nivolumab in clinical trials.

**Administration shortly before or after immunotherapy initiation is correlated with poor prognosis in solid cancer patients: An up-to-date systematic review and meta-analysis.** *Int. Immunopharmacol.* 2020, **88**, 106876.

The French Cooperative Thoracic Intergroup reported the preliminary results of the CHIVA-2 trial, a phase II study of nivolumab after prior chemotherapy for HIV-infected advanced NSCLC patients. A total of 16 patients were enrolled. The majority of them (69%) received nivolumab as second-line treatment, with a median treatment duration of 3.5 months. Similarly to what was observed in the general NSCLC population, a DCR (primary endpoint) of 62.5%, with a median PFS and OS (secondary endpoints) of 3.4 and 14.1 months, respectively, were reported. Nivolumab was well tolerated without serious adverse events nor opportunistic infections reported.

Of little risk, T-Cell Checkpoint Inhibitors and the MacCarty's OWEI Using B. trachealis to Improve Survival in Therapy, open to the Kaiser Permanente. *Assessment of the safety of pembrolizumab in patients with HIV and advanced cancer—A phase 1 study*. *JAMA Oncol*. 2019.

Finally, another early phase trial is currently ongoing in order to specifically evaluate the efficacy and safety of nivolumab in combination with cabozantinib (NCT04514484) in HIV-positive patients, including lung cancer [[Table 2](#)]. Costa, D.B.; Awad, M.M. Safety and efficacy of immune checkpoint inhibitors in patients with non-small cell lung cancer and hepatitis B or hepatitis C infection. *Lung Cancer* 2020, 145, 181–185.

6. World Health Organization (WHO). Available online:

<https://www.who.int/emergencies/diseases/novel-coronavirus-2019> (accessed on 8 November 2020).

7. Liang, W.; Guan, W.; Chen, R.; Wang, W.; Li, J.; Xu, K.; Li, C.; Ai, Q.; Lu, W.; Liang, H.; et al. Cancer patients in SARS-CoV-2 infection: A nationwide analysis in China. *Lancet Oncol*. 2020, 21, 335–337.

8. Mehta, V.; Goel, S.; Kabarriti, R.; Cole, D.; Goldfinger, M.; Acuna-Villaorduna, A.; Pradhan, K.; Thota, R.; Reissman, S.; Sparano, J.A.; et al. Case fatality rate of cancer patients with COVID-19 in a New York hospital system. *Cancer Discov*. 2020, 10, 935–941.

9. Zhang, L.; Zhu, F.; Xie, L.; Wang, C.; Wang, J.; Chen, R.; Jia, P.; Guan, H.Q.; Peng, L.; Chen, Y.; et al. Clinical characteristics of COVID-19-infected cancer patients: A retrospective case study in three hospitals within Wuhan, China. *Ann. Oncol.* 2020, 31, 894–901.

10. Garassino, M.C.; Whisenant, J.G.; Huang, L.-C.; Trama, A.; Torri, V.; Agustoni, F.; Baena, J.; Banna, G.; Berardi, R.; Bettini, A.C.; et al. COVID-19 in patients with thoracic malignancies (TERAVOLT): First results of an international, registry-based, cohort study. *Lancet Oncol*. 2020, 21, 914–922.

11. Rabe, K.F.; Watz, H. Chronic obstructive pulmonary disease. *Lancet* 2017, 389, 1931–1940.

12. Akinosoglou, K.S.; Karkoulias, K.; Marangos, M. Infectious complications in patients with lung cancer. *Eur. Rev. Med. Pharmacol. Sci.* 2013, 17, 8–18.

13. Perlin, E.; Bang, K.M.; Shah, A.; Hursey, P.D.; Whittingham, W.L.; Hashmi, K.; Campbell, L.; Kassim, O.O. The impact of pulmonary infections on the survival of lung cancer patients. *Cancer* 1990, 66, 593–596.

14. Kohno, S.; Koga, H.; Oka, M.; Kadota, J.; Kaku, M.; Soda, H.; Tomono, K.; Hara, K. The pattern of respiratory infection in patients with lung cancer. *Tohoku J. Exp. Med.* 1994, 173, 405–411.

15. Berghmans, T.; Sculier, J.P.; Klastersky, J. A prospective study of infections in lung cancer patients admitted to the hospital. *Chest* 2003, 124, 114–120.

16. Assi, H.I.; Kamphorst, A.O.; Moukalled, N.M.; Ramalingam, S.S. Immune checkpoint inhibitors in advanced non-small cell lung cancer. *Cancer* 2018, 124, 248–261.

17. Prelaj, A.; Tay, R.; Ferrara, R.; Chaput, N.; Besse, B.; Califano, R. Predictive biomarkers of response for immune checkpoint inhibitors in non-small-cell lung cancer. *Eur. J. Cancer* 2019, 106, 144–159.
18. Shui, L.; Yang, X.; Li, J.; Yi, C.; Sun, Q.; Zhu, H. Gut microbiome as a potential factor for modulating resistance to cancer immunotherapy. *Front. Immunol.* 2019, 10, 2989.
19. Jernberg, C.; Lofmark, S.; Edlund, C.; Jansson, J.K. Long-term impacts of antibiotic exposure on the human intestinal microbiota. *Microbiology* 2010, 156, 3216–3223.
20. Banna, G.L.; Passiglia, F.; Colonese, F.; Canova, S.; Menis, J.; Addeo, A.; Russo, A.; Cortinovis, D.L. Immune-checkpoint inhibitors in non-small cell lung cancer: A tool to improve patients' selection. *Crit. Rev. Oncol. Hematol.* 2018, 129, 27–39.
21. Huang, X.Z.; Gao, P.; Song, Y.X.; Xu, Y.; Sun, J.X.; Chen, X.W.; Zhao, J.H.; Wang, Z.N. Antibiotic use and the efficacy of immune checkpoint inhibitors in cancer patients: A pooled analysis of 2740 cancer patients. *Oncoimmunology* 2019, 8, e1665973.
22. Wilson, B.E.; Routy, B.; Nagrial, A.; Chin, V.T. The effect of antibiotics on clinical outcomes in immune-checkpoint blockade: A systematic review and meta-analysis of observational studies. *Cancer Immunol. Immunother.* 2020, 69, 343–354.
23. Xu, H.; Xu, X.; Wang, H.; Ge, W.; Cao, D. The association between antibiotics use and outcome of cancer patients treated with immune checkpoint inhibitors: A systematic review and meta-analysis. *Crit. Rev. Oncol. Hematol.* 2020, 149, 102909.
24. Petrelli, F.; Iaculli, A.; Signorelli, D.; Ghidini, A.; Dottorini, L.; Perego, G.; Ghidini, M.; Zaniboni, A.; Gori, S.; Inno, A. Survival of patients treated with antibiotics and immunotherapy for cancer: A systematic review and meta-analysis. *J. Clin. Med.* 2020, 9, 1458.
25. Lurienne, L.; Cervesi, J.; Duhalde, L.; de Gunzburg, J.; Andremont, A.; Zalcman, G.; Buffet, R.; Bandinelli, P.A. NSCLC Immunotherapy efficacy and antibiotic use: A systematic review and meta-analysis. *J. Thorac. Oncol.* 2020, 15, 1147–1159.
26. Sen, S.; Pestana, R.C.; Hess, K.; Viola, G.M.; Subbiah, V. Impact of antibiotic use on survival in patients with advanced cancers treated on immune checkpoint inhibitor phase I clinical trials. *Ann. Oncol.* 2018, 29, 2396–2398.
27. Pinato, D.J.; Howlett, S.; Ottaviani, D.; Urus, H.; Patel, A.; Mineo, T.; Brock, C.; Power, D.; Hatcher, O.; Falconer, A.; et al. Association of prior antibiotic treatment with survival and response to immune checkpoint inhibitor therapy in patients with cancer. *JAMA Oncol.* 2019, 5, 1774–1778.
28. Iglesias-Santamaria, A. Impact of antibiotic use and other concomitant medications on the efficacy of immune checkpoint inhibitors in patients with advanced cancer. *Clin. Transl. Oncol.* 2020, 22, 1481–1490.

29. Helmink, B.A.; Khan, M.A.W.; Hermann, A.; Gopalakrishnan, V.; Wargo, J.A. The microbiome, cancer, and cancer therapy. *Nat. Med.* 2019, 25, 377–388.

30. Gori, S.; Inno, A.; Belluomini, L.; Bocus, P.; Bisoffi, Z.; Russo, A.; Arcaro, G. Gut microbiota and cancer: How gut microbiota modulates activity, efficacy and toxicity of antitumoral therapy. *Crit. Rev. Oncol. Hematol.* 2019, 143, 139–147.

31. Li, W.; Deng, Y.; Chu, Q.; Zhang, P. Gut microbiome and cancer immunotherapy. *Cancer Lett.* 2019, 447, 41–47.

32. Sivan, A.; Corrales, L.; Hubert, N.; Williams, J.B.; Aquino-Michaels, K.; Earley, Z.M.; Benyamin, F.W.; Lei, Y.M.; Jabri, B.; Alegre, M.L.; et al. Commensal *Bifidobacterium* promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science* 2015, 350, 1084–1089.

33. Vetizou, M.; Pitt, J.M.; Daillere, R.; Lepage, P.; Waldschmitt, N.; Flament, C.; Rusakiewicz, S.; Routy, B.; Roberti, M.P.; Duong, C.P.; et al. Anticancer immunotherapy by CTLA-4 blockade relies on the gut microbiota. *Science* 2015, 350, 1079–1084.

34. Routy, B.; le Chatelier, E.; Derosa, L.; Duong, C.P.M.; Alou, M.T.; Daillere, R.; Fluckiger, A.; Messaoudene, M.; Rauber, C.; Roberti, M.P.; et al. Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* 2018, 359, 91–97.

35. Jin, Y.; Dong, H.; Xia, L.; Yang, Y.; Zhu, Y.; Shen, Y.; Zheng, H.; Yao, C.; Wang, Y.; Lu, S. The Diversity of gut microbiome is associated with favorable responses to anti-programmed death 1 immunotherapy in Chinese patients with NSCLC. *J. Thorac. Oncol.* 2019, 14, 1378–1389.

36. Derosa, L.; Routy, B.; Kroemer, G.; Zitvogel, L. The intestinal microbiota determines the clinical efficacy of immune checkpoint blockers targeting PD-1/PD-L1. *Oncoimmunology* 2018, 7, e1434468.

37. Krief, J.O.; de Tauriers, P.H.; Dumenil, C.; Neveux, N.; Dumoulin, J.; Giraud, V.; Labrune, S.; Tisserand, J.; Julie, C.; Emile, J.-F.; et al. Role of antibiotic use, plasma citrulline and blood microbiome in advanced non-small cell lung cancer patients treated with nivolumab. *J. Immuno Ther. Cancer* 2019, 7, 176.

38. Nagasaka, M.; Sexton, R.; Alhasan, R.; Rahman, S.; Azmi, A.S.; Sukari, A. Gut microbiome and response to checkpoint inhibitors in non-small cell lung cancer—A review. *Crit. Rev. Oncol. Hematol.* 2020, 145, 102841.

39. Abreu, M.T.; Peek, R.M., Jr. Gastrointestinal malignancy and the microbiome. *Gastroenterology* 2014, 146, 1534–1546.

40. Yu, G.; Gail, M.H.; Consonni, D.; Carugno, M.; Humphrys, M.; Pesatori, A.C.; Caporaso, N.E.; Goedert, J.J.; Ravel, J.; Landi, M.T. Characterizing human lung tissue microbiota and its relationship to epidemiological and clinical features. *Genome Biol.* 2016, 17, 163.

41. Dickson, R.P.; Erb-Downward, J.R.; Huffnagle, G.B. Towards an ecology of the lung: New conceptual models of pulmonary microbiology and pneumonia pathogenesis. *Lancet Respir. Med.* 2014, 2, 238–246.
42. Jin, J.; Gan, Y.; Liu, H.; Wang, Z.; Yuan, J.; Deng, T.; Zhou, Y.; Zhu, Y.; Zhu, H.; Yang, S.; et al. Diminishing microbiome richness and distinction in the lower respiratory tract of lung cancer patients: A multiple comparative study design with independent validation. *Lung Cancer* 2019, 136, 129–135.
43. Laroumagne, S.; Salinas-Pineda, A.; Hermant, C.; Murris, M.; Gourraud, P.A.; Do, C.; Segonds, C.; Didier, A.; Mazieres, J. Incidence and characteristics of bronchial colonisation in patient with lung cancer: A retrospective study of 388 cases. *Rev. Mal. Respir.* 2011, 28, 328–335.
44. Liu, Y.; O'Brien, J.L.; Ajami, N.J.; Scheurer, M.E.; Amirian, E.S.; Armstrong, G.; Tsavachidis, S.; Thrift, A.P.; Jiao, L.; Wong, M.C.; et al. Lung tissue microbial profile in lung cancer is distinct from emphysema. *Am. J. Cancer Res.* 2018, 8, 1775–1787.
45. Chang, S.H.; Mirabolfathinejad, S.G.; Katta, H.; Cumpian, A.M.; Gong, L.; Caetano, M.S.; Moghaddam, S.J.; Dong, C. T helper 17 cells play a critical pathogenic role in lung cancer. *Proc. Natl. Acad. Sci. USA* 2014, 111, 5664–5669.
46. Dickson, R.P.; Martinez, F.J.; Huffnagle, G.B. The role of the microbiome in exacerbations of chronic lung diseases. *Lancet* 2014, 384, 691–702.
47. Jungnickel, C.; Schmidt, L.H.; Bittigkoffer, L.; Wolf, L.; Wolf, A.; Ritzmann, F.; Kamyschnikow, A.; Herr, C.; Menger, M.D.; Spieker, T.; et al. IL-17C mediates the recruitment of tumor-associated neutrophils and lung tumor growth. *Oncogene* 2017, 36, 4182–4190.
48. Boursi, B.; Mamiani, R.; Haynes, K.; Yang, Y.X. Recurrent antibiotic exposure may promote cancer formation—Another step in understanding the role of the human microbiota? *Eur. J. Cancer* 2015, 51, 2655–2664.
49. Carbone, C.; Piro, G.; di Noia, V.; D'Argento, E.; Vita, E.; Ferrara, M.G.; Pilotto, S.; Milella, M.; Cammarota, G.; Gasbarrini, A.; et al. Lung and gut microbiota as potential hidden driver of immunotherapy efficacy in lung cancer. *Mediators Inflamm.* 2019, 2019, 7652014.
50. Kaderbhai, C.; Richard, C.; Fumet, J.D.; Aarnink, A.; Foucher, P.; Coudert, B.; Favier, L.; Lagrange, A.; Limagne, E.; Boidot, R.; et al. Antibiotic Use Does Not Appear to Influence Response to Nivolumab. *Anticancer Res.* 2017, 37, 3195–3200.
51. Huemer, F.; Rinnerthaler, G.; Lang, D.; Hackl, H.; Lamprecht, B.; Greil, R. Association between antibiotics use and outcome in patients with NSCLC treated with immunotherapeutics. *Ann. Oncol.* 2019, 30, 652–653.
52. Metges, J.-P.; Michaud, E.; Lagadec, D.D.; Marhuenda, F.; Chaslerie, A.; Grude, F. Impact of anti-infectious and corticosteroids on immunotherapy: Nivolumab and pembrolizumab follow-up in a

French study. *J. Clin. Oncol.* 2018, 36, 15157.

53. Rossi, G.; Pezzuto, A.; Sini, C.; Tuzi, A.; Citarella, F.; McCusker, M.G.; Nigro, O.; Tanda, E.; Russo, A. Concomitant medications during immune checkpoint blockage in cancer patients: Novel insights in this emerging clinical scenario. *Crit. Rev. Oncol. Hematol.* 2019, 142, 26–34.

54. Chalabi, M.; Cardona, A.; Nagarkar, D.R.; Scala, A.D.; Gandara, D.R.; Rittmeyer, A.; Albert, M.L.; Powles, T.; Kok, M.; Herrera, F.G.; et al. Efficacy of chemotherapy and atezolizumab in patients with non-small-cell lung cancer receiving antibiotics and proton pump inhibitors: Pooled post hoc analyses of the OAK and POPLAR trials. *Ann. Oncol.* 2020, 31, 525–531.

55. Mielgo-Rubio, X.; Chara, L.; Sotelo-Lezama, M.; Castro, R.L.; Rubio-Martínez, J.; Velastegui, A.; Olier-Garate, C.; Falagan, S.; Gómez-Barreda, I.; Bautista-Sanz, P.; et al. MA10.01 antibiotic use and PD-1 inhibitors: Shorter survival in lung cancer, especially when given intravenously. Type of infection also matters. *J. Thorac. Oncol.* 2018, 13.

56. Mahajan, R.; Xing, J.; Liu, S.J.; Ly, K.N.; Moorman, A.C.; Rupp, L.; Xu, F.; Holmberg, S.D. Chronic hepatitis cohort study, I. Mortality among persons in care with hepatitis C virus infection: The chronic hepatitis cohort study (CHeCS), 2006–2010. *Clin. Infect. Dis.* 2014, 58, 1055–1061.

57. Yarchoan, R.; Uldrick, T.S. HIV-associated cancers and related diseases. *N. Engl. J. Med.* 2018, 378, 2145.

58. Novakova, L.; Pavlik, J.; Chrenkova, L.; Martinec, O.; Cerveny, L. Current antiviral drugs and their analysis in biological materials—Part II: Antivirals against hepatitis and HIV viruses. *J. Pharm. Biomed. Anal.* 2018, 147, 378–399.

59. UNAIDS. Global HIV & AIDS Statistics—2020 Fact Sheet. Available online: <https://www.unaids.org/en/resources/fact-sheet> (accessed on 30 October 2020).

60. Grulich, A.E.; van Leeuwen, M.T.; Falster, M.O.; Vajdic, C.M. Incidence of cancers in people with HIV/AIDS compared with immunosuppressed transplant recipients: A meta-analysis. *Lancet* 2007, 370, 59–67.

61. Shiels, M.S.; Pfeiffer, R.M.; Gail, M.H.; Hall, H.I.; Li, J.; Chaturvedi, A.K.; Bhatia, K.; Uldrick, T.S.; Yarchoan, R.; Goedert, J.J.; et al. Cancer burden in the HIV-infected population in the United States. *J. Natl. Cancer Inst.* 2011, 103, 753–762.

62. Vandenhende, M.A.; Roussillon, C.; Henard, S.; Morlat, P.; Oksenhendler, E.; Aumaitre, H.; Georget, A.; May, T.; Rosenthal, E.; Salmon, D.; et al. Cancer-related causes of death among HIV-infected patients in France in 2010: Evolution since 2000. *PLoS ONE* 2015, 10, e0129550.

63. Mdodo, R.; Frazier, E.L.; Dube, S.R.; Mattson, C.L.; Sutton, M.Y.; Brooks, J.T.; Skarbinski, J. Cigarette smoking prevalence among adults with HIV compared with the general adult population in the United States: Cross-sectional surveys. *Ann. Intern. Med.* 2015, 162, 335–344.

64. Serrao, R.; Pinero, C.; Velez, J.; Coutinho, D.; Maltez, F.; Lino, S.; Sarmento, E.C.R.; Tavares, A.P.; Pacheco, P.; Lopes, M.J.; et al. Non-AIDS-related comorbidities in people living with HIV-1 aged 50 years and older: The AGING POSITIVE study. *Int. J. Infect. Dis.* 2019, **79**, 94–100.

65. Sigel, K.; Wisnivesky, J.; Gordon, K.; Dubrow, R.; Justice, A.; Brown, S.T.; Goulet, J.; Butt, A.A.; Crystal, S.; Rimland, D.; et al. HIV as an independent risk factor for incident lung cancer. *AIDS* 2012, **26**, 1017–1025.

66. Reddy, K.P.; Kong, C.Y.; Hyle, E.P.; Baggett, T.P.; Huang, M.; Parker, R.A.; Paltiel, A.D.; Losina, E.; Weinstein, M.C.; Freedberg, K.A.; et al. Lung cancer mortality associated with smoking and smoking cessation among people living with HIV in the United States. *JAMA Intern. Med.* 2017, **177**, 1613–1621.

67. Uldrick, T.S.; Ison, G.; Rudek, M.A.; Noy, A.; Schwartz, K.; Bruinooge, S.; Schenkel, C.; Miller, B.; Dunleavy, K.; Wang, J.; et al. Modernizing clinical trial eligibility criteria: Recommendations of the American Society of clinical oncology-friends of cancer research HIV working group. *J. Clin. Oncol* 2017, **35**, 3774–3780.

68. Velu, V.; Shetty, R.D.; Larsson, M.; Shankar, E.M. Role of PD-1 co-inhibitory pathway in HIV infection and potential therapeutic options. *Retrovirology* 2015, **12**, 14.

69. Grabmeier-Pfistershammer, K.; Steinberger, P.; Rieger, A.; Leitner, J.; Kohrgruber, N. Identification of PD-1 as a unique marker for failing immune reconstitution in HIV-1-infected patients on treatment. *J. Acquir. Immune Defic. Syndr.* 2011, **56**, 118–124.

70. Porichis, F.; Hart, M.G.; Massa, A.; Everett, H.L.; Morou, A.; Richard, J.; Brassard, N.; Veillette, M.; Hassan, M.; Ly, N.L.; et al. Immune checkpoint blockade restores HIV-specific CD4 T cell help for NK cells. *J. Immunol.* 2018, **201**, 971–981.

71. Okuma, Y.; Hishima, T.; Kashima, J.; Homma, S. High PD-L1 expression indicates poor prognosis of HIV-infected patients with non-small cell lung cancer. *Cancer Immunol. Immunother.* 2018, **67**, 495–505.

72. Guihot, A.; Marcellin, A.G.; Massiani, M.A.; Samri, A.; Soulie, C.; Autran, B.; Spano, J.P. Drastic decrease of the HIV reservoir in a patient treated with nivolumab for lung cancer. *Ann. Oncol.* 2018, **29**, 517–518.

73. McCullar, B.; Alloway, T.; Martin, M. Durable complete response to nivolumab in a patient with HIV and metastatic non-small cell lung cancer. *J. Thorac. Dis.* 2017, **9**, 540–542.

74. Ostios-Garcia, L.; Faig, J.; Leonardi, G.C.; Adeni, A.E.; Subegdjo, S.J.; Lydon, C.A.; Rangachari, D.; Huberman, M.S.; Sehgal, K.; Shea, M.; et al. Safety and efficacy of PD-1 inhibitors among HIV-positive patients with non-small cell lung cancer. *J. Thorac. Oncol.* 2018, **13**, 1037–1042.

75. Lavole, A.; Guihot, A.; Veyri, M.; Lambotte, O.; Autran, B.; Cloarec, N.; le Garff, G.; Flament, T.; Cadranel, J.; Spano, J.P. PD-1 blockade in HIV-infected patients with lung cancer: A new

challenge or already a strategy? *Ann. Oncol.* 2018, 29, 1065–1066.

76. Samri, A.; Lavolé, A.; Even, S.; Lambert-Niclot, S.; le Garff, G.; Cadranel, J.; Spano, J.-P.; Autran, B.; Marcelin, A.-G.; Guihot, A. Immunovirological evolution in HIV-infected patients treated with anti-PD1 therapy. In Proceedings of the 9th IAS Conference on HIV Science (IAS 2017), Paris, France, 23–26 July 2017.

77. Barlesi, F.; Audigier-Valette, C.; Felip, E.; Ciuleanu, T.E.; Jao, K.; Rijavec, E.; Urban, L.; Aucoin, J.S.; Zannori, C.; Vermaelen, K.; et al. Nivolumab plus low-dose IPILIMUMAB as first-line treatment of advanced NSCLC: Overall survival analysis of checkmate 817. *Ann. Oncol.* 2019, 30, 33–34.

78. Gonzalez-Cao, M.; Moran, T.; Dalmau, J.; Garcia-Corbacho, J.; Bracht, J.W.P.; Bernabe, R.; Juan, O.; de Castro, J.; Blanco, R.; Drozdowskyj, A.; et al. Assessment of the feasibility and safety of durvalumab for treatment of solid tumors in patients with HIV-1 infection: The phase 2 DURVAST study. *JAMA Oncol.* 2020, 6, 1063–1067.

79. Rajdev, L.; Lensing, S.; Ramos, J.C.; Baiocchi, R.; Wang, C.C.J.; Ratner, L.; Rubinstein, P.G.; Ambinder, R.; Henry, D.; Streicher, H.; et al. 1023MO AMC 095: A report of nivolumab (nivo) in advanced HIV associated solid tumours (ST). *Ann. Oncol.* 2020, 31, 706.

80. Lavole, A.; Mazieres, J.; Schneider, S.; Brosseau, S.; Kiakouama, L.M.; Greillier, L.; Guihot, A.; Abbar, B.; Baron, M.; Makinson, A.; et al. 1389P IFCT-1602 CHIVA2 phase II trial: Nivolumab in previously treated HIV-patients with advanced non-small cell lung cancer (NSCLC). *Ann. Oncol.* 2020, 31, 882–883.

81. Sezer, A.; Kilickap, S.; Gümüş, M.; Bondarenko, I.; Özgüroğlu, M.; Gogishvili, M.; Turk, H.M.; Çiçin, İ.; Bentsion, D.; Gladkov, O.; et al. LBA52 EMPOWER-Lung 1: Phase III first-line (1L) cemiplimab monotherapy vs. platinum-doublet chemotherapy (chemo) in advanced non-small cell lung cancer (NSCLC) with programmed cell death-ligand 1 (PD-L1)  $\geq 50\%$ . *Ann. Oncol.* 2020, 31, 1182–1183.

82. Spano, J.P.; Veyri, M.; Gobert, A.; Guihot, A.; Perre, P.; Kerjouan, M.; Brosseau, S.; Cloarec, N.; Montaudie, H.; Helissey, C.; et al. Immunotherapy for cancer in people living with HIV: Safety with an efficacy signal from the series in real life experience. *AIDS* 2019, 33, 13–19.

83. Shah, N.J.; Al-Shbool, G.; Blackburn, M.; Cook, M.; Belouali, A.; Liu, S.V.; Madhavan, S.; He, A.R.; Atkins, M.B.; Gibney, G.T.; et al. Safety and efficacy of immune checkpoint inhibitors (ICIs) in cancer patients with HIV, hepatitis B, or hepatitis C viral infection. *J. Immunother. Cancer* 2019, 7, 353.

84. Cook, M.R.; Kim, C. Safety and efficacy of immune checkpoint inhibitor therapy in patients with HIV infection and advanced-stage cancer: A systematic review. *JAMA Oncol.* 2019, 5, 1049–1054.

85. Kothapalli, A.; Khattak, M.A. Safety and efficacy of anti-PD-1 therapy for metastatic melanoma and non-small-cell lung cancer in patients with viral hepatitis: A case series. *Melanoma Res.* 2018, 28, 155–158.

86. Tio, M.; Rai, R.; Ezeoke, O.M.; McQuade, J.L.; Zimmer, L.; Khoo, C.; Park, J.J.; Spain, L.; Turajlic, S.; Ardolino, L.; et al. Anti-PD-1/PD-L1 immunotherapy in patients with solid organ transplant, HIV or hepatitis B/C infection. *Eur. J. Cancer* 2018, 104, 137–144.

87. Galanina, N.; Goodman, A.M.; Cohen, P.R.; Frampton, G.M.; Kurzrock, R. Successful treatment of HIV-associated kaposi sarcoma with immune checkpoint blockade. *Cancer Immunol. Res.* 2018, 6, 1129–1135.

88. Sahin, I.H.; Kane, S.R.; Brutcher, E.; Guadagno, J.; Smith, K.E.; Wu, C.; Lesinski, G.B.; Gunthel, C.J.; El-Rayes, B.F. Safety and efficacy of immune checkpoint inhibitors in patients with cancer living with HIV: A perspective on recent progress and future needs. *JCO Oncol. Pract.* 2020, 16, 319–325.

89. Kuderer, N.M.; Choueiri, T.K.; Shah, D.P.; Shyr, Y.; Rubinstein, S.M.; Rivera, D.R.; Shete, S.; Hsu, C.Y.; Desai, A.; Lopes, G.D.L., Jr.; et al. Clinical impact of COVID-19 on patients with cancer (CCC19): A cohort study. *Lancet* 2020, 395, 1907–1918.

90. Espinar, J.B.; Torri, V.; Whisenant, J.; Hirsch, F.R.; Rogado, J.; de Castro Carpeño, J.; Halmos, B.; Ceresoli, G.L.; Rueda, A.G.; Tiseo, M.; et al. LBA75 Defining COVID-19 outcomes in thoracic cancer patients: TERAVOLT (Thoracic cancERs international coVid 19 cOllaboraTion). *Ann. Oncol.* 2020, 31, 1204–1205.

91. Di Noia, V.; D'Aveni, A.; Squadroni, M.; Beretta, G.D.; Ceresoli, G.L. Immune checkpoint inhibitors in SARS-CoV-2 infected cancer patients: The spark that ignites the fire? *Lung Cancer* 2020, 145, 208–210.

92. Luo, J.; Rizvi, H.; Egger, J.V.; Preeshagul, I.R.; Wolchok, J.D.; Hellmann, M.D. Impact of PD-1 blockade on severity of COVID-19 in patients with lung cancers. *Cancer Discov.* 2020, 10, 1121–1128.

93. Wise-Draper, T.M.; Desai, A.; Elkrief, A.; Rini, B.I.; Flora, D.B.; Bowles, D.W.; Shah, D.; Rivera, D.; Johnson, D.B.; Lopes, G.; et al. LBA71 Systemic cancer treatment-related outcomes in patients with SARS-CoV-2 infection: A CCC19 registry analysis. *Ann. Oncol.* 2020, 31, 1201–1202.

94. Davis, A.P.; Boyer, M.; Lee, J.H.; Kao, S.C. COVID-19: The use of immunotherapy in metastatic lung cancer. *Immunotherapy* 2020, 12, 545–548.

95. Passaro, A.; Addeo, A.; von Garnier, C.; Blackhall, F.; Planchard, D.; Felip, E.; Dziadziszko, R.; de Marinis, F.; Reck, M.; Bouchaab, H.; et al. ESMO Management and treatment adapted recommendations in the COVID-19 era: Lung cancer. *ESMO Open* 2020, 5.

Retrieved from <https://encyclopedia.pub/entry/history/show/15047>