

TAF1 Gene

Subjects: Genetics & Heredity

Contributor: Rui Liu

TATA-box binding protein associated factor 1: The TAF1 gene provides instructions for making part of a protein called transcription factor IID (TFIID).

Keywords: genes

1. Normal Function

The *TAF1* gene provides instructions for making part of a protein called transcription factor IID (TFIID). This protein is active in cells and tissues throughout the body, where it attaches (binds) to DNA. Transcription factor IID plays an essential role in regulating the activity of most genes.

The *TAF1* gene is part of a complex region of DNA known as the *TAF1/DYT3* multiple transcript system. This region consists of short stretches of DNA from the *TAF1* gene plus some extra segments of genetic material near the gene. These stretches of DNA can be combined in different ways to create various sets of instructions for making proteins. Researchers believe that some of these variations are critical for the normal function of nerve cells (neurons) in the brain.

2. Health Conditions Related to Genetic Changes

2.1. X-linked dystonia-parkinsonism

Several changes in the *TAF1/DYT3* multiple transcript system have been identified in people with X-linked dystonia-parkinsonism. Some alter single DNA building blocks (nucleotides) in the gene; these changes are described as disease-specific single-nucleotide changes (DSCs). Another genetic change deletes a small number of nucleotides from the gene. Researchers are uncertain how these changes are related to the movement abnormalities characteristic of the disease.

X-linked dystonia-parkinsonism may also be related to an extra segment of DNA in the *TAF1/DYT3* multiple transcript system. The extra segment results from the insertion of a retrotransposon, which is a small piece of DNA that can move around to different positions in a cell's genetic material. When a retrotransposon inserts itself in or near a gene, it can disrupt the gene's function. In this case, the retrotransposon insertion probably interferes with the normal function of the *TAF1/DYT3* multiple transcript system.

Researchers suspect that changes in the *TAF1/DYT3* multiple transcript system disrupt the regulation of critical genes in neurons. This defect leads to the eventual death of these cells, particularly in areas of the brain called the caudate nucleus and putamen. These regions are critical for normal movement, learning, and memory. It is unclear why the effects of changes in the *TAF1/DYT3* multiple transcript system appear to be limited to dystonia and parkinsonism.

3. Other Names for This Gene

- BA2R
- CCG1
- CCGS
- Cell cycle gene 1 protein
- cell cycle, G1 phase defect
- complementation of cell cycle block, G1-to-S
- DYT3

- DYT3/TAF1
- KAT4
- N-TAF1
- NSCL2
- P250
- TAF(II)250
- TAF1 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 250kDa
- TAF1_HUMAN
- TAF2A
- TAFII-250
- TAFII250
- TATA box binding protein associated factor 1
- TATA box-binding protein-associated factor 1
- TATA box-binding protein-associated factor 2A
- TBP-associated factor 1
- TBP-associated factor 250 kDa
- TBP-associated factor, RNA polymerase II, 250-kD
- transcription factor TFIID p250 polypeptide
- Transcription initiation factor TFIID 250 kDa subunit
- Transcription initiation factor TFIID subunit 1

References

1. Herzfeld T, Nolte D, Müller U. Structural and functional analysis of the humanTAF1/DYT3 multiple transcript system. Mamm Genome. 2007 Nov;18(11):787-95.
 2. Maile T, Kwoczyński S, Katzenberger RJ, Wassarman DA, Sauer F. TAF1 activates transcription by phosphorylation of serine 33 in histone H2B. Science. 2004 May14;304(5673):1010-4. Retraction in: McNutt M. Science. 2014 May 30;344(6187):981.
 3. Makino S, Kaji R, Ando S, Tomizawa M, Yasuno K, Goto S, Matsumoto S, TabuenaMD, Maranon E, Dantes M, Lee LV, Ogasawara K, Tooyama I, Akatsu H, Nishimura M,Tamiya G. Reduced neuron-specific expression of the TAF1 gene is associated with X-linked dystonia-parkinsonism. Am J Hum Genet. 2007 Mar;80(3):393-406.
 4. Muller U, Herzfeld T, Nolte D. The TAF1/DYT3 multiple transcript system inX-linked dystonia-parkinsonism. Am J Hum Genet. 2007 Aug;81(2):415-7; authorreply 417-8.
 5. Nolte D, Niemann S, Müller U. Specific sequence changes in multiple transcripts system DYT3 are associated with X-linked dystonia parkinsonism. Proc Natl AcadSci U S A. 2003 Sep 2;100(18):10347-52.
-