

Otopalatodigital Syndrome Type 1

Subjects: **Genetics & Heredity**

Contributor: Rita Xu

Otopalatodigital syndrome type 1 is a disorder primarily involving abnormalities in skeletal development.

genetic conditions

1. Introduction

It is a member of a group of related conditions called otopalatodigital spectrum disorders, which also includes otopalatodigital syndrome type 2, frontometaphyseal dysplasia, Melnick-Needles syndrome, and terminal osseous dysplasia. In general, these disorders involve hearing loss caused by malformations in the tiny bones in the ears (ossicles), problems in the development of the roof of the mouth (palate), and skeletal abnormalities involving the fingers or toes (digits).

Otopalatodigital syndrome type 1 is usually the mildest of the otopalatodigital spectrum disorders. People with this condition usually have characteristic facial features including wide-set and downward-slanting eyes; prominent brow ridges; and a broad, flat nose. Affected individuals have abnormalities of the fingers and toes, such as blunt, square-shaped (spatulate) fingertips; shortened thumbs and big toes; unusually long second toes; and a wide gap between the first and second toes (known as a sandal gap). Affected individuals also have hearing loss.

Infants with otopalatodigital syndrome type 1 may be born with an opening in the roof of the mouth (a cleft palate). Individuals with this condition often have fewer teeth than normal (hypodontia). They may have mild abnormal curvature (bowing) of their limbs, and limited range of motion in some joints. People with otopalatodigital syndrome type 1 may be somewhat shorter than other members of their family.

Females with otopalatodigital syndrome type 1 often have more variable signs and symptoms compared to affected males, with females typically having fewer signs and symptoms.

2. Frequency

Otopalatodigital syndrome type 1 is a rare disorder, affecting fewer than 1 in every 100,000 individuals. Its specific incidence is unknown.

3. Causes

Otopalatodigital syndrome type 1 is caused by mutations in the *FLNA* gene. The *FLNA* gene provides instructions for producing the protein filamin A, which helps build the network of protein filaments (cytoskeleton) that gives structure to cells and allows them to change shape and move. Filamin A attaches (binds) to another protein called actin, and helps the actin to form the branching network of filaments that make up the cytoskeleton. Filamin A also links actin to many other proteins to perform various functions within the cell.

The *FLNA* gene mutations that cause otopalatodigital syndrome type 1 result in changes to the filamin A protein in the region that binds to actin. The mutations are described as "gain-of-function" because they appear to lead to a protein with an increased ability to bind to actin. Researchers believe that the mutations impair the stability of the cytoskeleton and disrupt cellular processes involved in skeletal development, but it is not known how changes in the protein relate to the specific signs and symptoms of otopalatodigital syndrome type 1.

The Gene Associated with Otopalatodigital Syndrome Type 1

- *FLNA*

4. Inheritance

This condition is inherited in an X-linked pattern. A condition is considered X-linked if the mutated gene that causes the disorder is located on the X chromosome, one of the two sex chromosomes in each cell. In males, who have only one X chromosome, a mutation in the only copy of the gene in each cell is sufficient to cause the condition. In females, who have two copies of the X chromosome, one altered copy of the gene in each cell can lead to less severe features of the condition or may cause no signs or symptoms at all. A characteristic of X-linked inheritance is that fathers cannot pass X-linked traits to their sons.

5. Other Names for This Condition

- cranioorodigital syndrome
- faciopalatoosseous syndrome
- FPO
- OPD syndrome, type 1
- oto-palato-digital syndrome, type I
- Taybi syndrome

References

1. Hidalgo-Bravo A, Pompa-Mera EN, Kofman-Alfaro S, Gonzalez-Bonilla CR, Zenteno JC. A novel filamin A D203Y mutation in a female patient with otopalatodigital type 1 syndrome and extremely skewed X chromosome inactivation. *Am J Med Genet A*.2005 Jul 15;136(2):190-3.

2. Joksic I, Cuturilo G, Jurisic A, Djuricic S, Peterlin B, Mijovic M, KaradzovON, Egic A, Milovanovic Z. Otopalatodigital Syndrome Type I: Novel Characteristics and Prenatal Manifestations in two Siblings. *Balkan J Med Genet.* 2019 Dec 21;22(2):83-88. doi: 10.2478/bjmg-2019-0024.
3. Moutton S, Fergelot P, Naudion S, Cordier MP, Solé G, Guerineau E, Hubert C, Rooryck C, Vuillaume ML, Houcinat N, Deforges J, Bouron J, Devès S, Le Merrer M, David A, Geneviève D, Giuliano F, Journel H, Megarbane A, Faivre L, Chassaing N, Francannet C, Sarrazin E, Stattin EL, Vigneron J, Leclair D, Abadie C, Sarda P, Baumann C, Delrue MA, Arveiler B, Lacombe D, Goizet C, Coupry I. Otopalatodigital spectrum disorders: refinement of the phenotypic and mutational spectrum. *J Hum Genet.* 2016 Aug;61(8):693-9. doi: 10.1038/jhg.2016.37.
4. Robertson S. X-Linked Otopalatodigital Spectrum Disorders. 2005 Nov 30 [updated 2019 Oct 3]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. *GeneReviews® [Internet].* Seattle (WA): University of Washington, Seattle; 1993-2020. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1393/>
5. Robertson SP, Twigg SR, Sutherland-Smith AJ, Biancalana V, Gorlin RJ, Horn D, Kenwick SJ, Kim CA, Morava E, Newbury-Ecob R, Orstavik KH, Quarrell OW, Schwartz CE, Shears DJ, Suri M, Kendrick-Jones J, Wilkie AO; OPD-spectrum Disorders Clinical Collaborative Group. Localized mutations in the gene encoding the cytoskeletal protein filamin A cause diverse malformations in humans. *Nat Genet.* 2003 Apr;33(4):487-91.
6. Robertson SP. Otopalatodigital syndrome spectrum disorders: otopalatodigital syndrome types 1 and 2, frontometaphyseal dysplasia and Melnick-Needles syndrome. *Eur J Hum Genet.* 2007 Jan;15(1):3-9.

Retrieved from <https://encyclopedia.pub/entry/history/show/11839>