

Oxidative Stress, NAD⁺ in ALS

Subjects: Neurosciences

Contributor: ELENA OBRADOR

ALS, often referred to as “Lou Gehrig’s disease”, is a progressive neurodegenerative disease that damages motor neurons (MNs) in the brain and the spinal cord. This loss of MNs is responsible for progressive weakness and paralysis. There is mounting evidence involving oxidative stress as a main pathophysiological mechanism leading to MN damage and death. Conceptually, oxidative stress is an imbalance of oxygen-derived free radicals and antioxidants in the body, which can lead to cell and organ damage. Reactive oxygen species (ROS) contain an uneven number of electrons and, at high non-physiological levels, can cause oxidative damages to nucleic acids, lipids, and proteins. Moreover, the generation of NO and H₂O₂ (induced by proinflammatory cytokines in e.g., endothelial cells) can lead to the formation of highly damaging –OONO radicals. On the other hand, NAD⁺ is involved in cell bioenergetics, redox regulation, signaling, homeostasis, adaptive response to stress, and survival. Specifically, different NAD⁺-dependent enzymes are implicated in mechanisms regulating synaptic plasticity and neuronal resilience to stress.

Keywords: amyotrophic lateral sclerosis ; motor neurons ; oxidative stress ; NAD⁺ ; antioxidants

1. NAD⁺ and Physiological Antioxidant Levels in ALS

NAD⁺ is a coenzyme in redox reactions, a donor of ADP-ribose for ADP-ribosylation reactions, a precursor of cyclic ADP-ribose, and a substrate for sirtuins (SIRT) that use the cofactor to remove acetyl groups from proteins. SIRT link NAD⁺ levels to mitochondrial function, dynamics and biogenesis, and to cellular antioxidant defenses ^{[1][2]}. Interestingly, under the conditions of reduced cellular energy (as it occurs in the MNs during the progression of ALS), SIRT may not consume sufficient NAD⁺ to preclude any cell survival-promoting effects of its deacetylase action on protein substrates ^[3]. For instance, poly(ADP-ribose) polymerase 1 (PARP1) is involved in DNA replication, transcription, DNA repair, apoptosis, and genome stability. However, DNA damage may overactivate PARP1 and lead to cell death and inflammation ^[4]. SIRT1 function (nucleus and cytoplasm) can facilitate cell survival under stress conditions by the deacetylation-dependent deactivation of PARP1 ^[5]. However, SIRT1 also consumes NAD⁺. Thus, inducing an increase in SIRT levels and activity (at least above an undefined threshold) could be counterproductive to neuronal survival, particularly to those energetically compromised. As an example, the nicotinamide-induced inhibition of SIRT1 was shown to protect neurons from death under acute anoxic injury ^[6]. Moreover, in SIRT1 knockout mice, their brains showed low levels of oxidative stress-related markers (i.e., carbonylated proteins and isoprostanes) ^[7]. These facts suggest a delicate balance between SIRT1 activity and the survival of neurons subjected to stress-associated insults. Despite the controversial facts regarding the activity of SIRT1, it is widely accepted that the maintenance of high NAD⁺ levels promotes cell homeostasis and survival ^[8]. Consequently, the consumption of NAD⁺ without an adequate method of replenishment is deleterious for the cell physiology.

In mammalian cells, NAD⁺ can be synthesized de novo from tryptophan (a multi-step enzymatic process which is fairly inefficient), or from nicotinic acid, nicotinamide (NAM), nicotinamide mononucleotide (NMN), or nicotinamide riboside (NR) ^{[9][10]}. For instance, it has been shown that NAM prevents NAD⁺ depletion and protects neurons against excitotoxicity and cerebral ischemia ^[11]. NR and NMN were shown to confer axonal protection in a Wallerian degeneration model of neuronal explant cultures ^[12]. Interestingly, the mechanism that neurons undergoing axonal degeneration use for protection is to upregulate the expression of nicotinamide riboside kinase (NRK) and nicotinamide mononucleotide adenylyl transferase 1 (NNMAT1), the enzymes required to convert NR (or NMN) to NAD⁺ ^[12]. Further, NR rescues mitochondrial defects and neuronal loss in models of Parkinson’s diseases ^[13], or increasing cytosolic and mitochondrial NAD⁺ content in ALS astrocytes increases the oxidative stress resistance and reverts their toxicity towards MNs ^[14]. Furthermore, it has been shown that SARM1 (sterile α and TIR motif-containing 1) is a key factor in triggering axon degeneration after an injury. Once activated in neurons, SARM1, which metabolizes NAD⁺ to NAM, depletes NAD⁺, leading to a massive loss of the energy supply within the axons ^[15]. These researchers were capable of reversing this detrimental effect by supplementing the neurons in which SARM1 was activated with NR ^[15].

NAD⁺ levels in mouse cortical neurons are of approximately 10 nmol/mg of protein [16], which is similar to those found in different types of neurons. The altered expression of enzymes involved in NAD⁺ synthesis (nicotinamide phosphoribosyltransferase and nicotinamide nucleotide adenyltransferase 2) and the decreased SIRT6 expression found in the spinal cord of ALS patients suggest deficits of this neuroprotective pathway in the human pathology [17]. Although precise in vivo data on NAD⁺ levels in the MNs of ALS models and patients are lacking, taking into account all available evidence, a significant NAD⁺ depletion is expected (see e.g., Park 2016) [18].

SIRT3, the main SIRT deacetylase in mitochondria, deacetylates and increases the activity of both superoxide dismutase 2 (SOD2) (thereby controlling the excess of toxic O₂ [19]) and isocitrate dehydrogenase (which increases NADPH and, that way, favors the glutathione (GSH) reductase reaction) [20]. GSH is a main physiological antioxidant and the GSH/glutathione disulfide (GSSG) ratio is a measure of cellular oxidative stress [21]. Oxidation of this tripeptide, which cannot be synthesized inside the mitochondria and must be transported from the cytosol [22], facilitates the opening of the mitochondrial permeability transition pore and the release of proapoptotic death signals [23]. Importantly, early studies also demonstrated that GSH deficiency leads to mitochondrial damage in the brain [24].

There is evidence to suggest that the response to oxidative stress is dampened in ALS. GSH depletion promotes neurological deficits, mitochondrial dysfunction, and MN degeneration in mutant SOD1 ALS mice [25]. Levels of GSH are lower in the motor cortex of ALS patients as compared to healthy volunteers [26].

Consequently, this background strongly supports the notion of a close relationship between oxidative stress, NAD⁺ levels, cellular redox status, and mitochondrial function, all of which are involved in the pathophysiology of ALS [1].

2. NAD⁺ Promoters and Antioxidants to Protect Motor Neurons

2.1. NAD⁺ Promoters

NAD⁺ is a coenzyme that facilitates redox reactions and is found in all living cells. Potential NAD⁺ promoters now under study include (but are not limited to) niacin (NA), NAM, NMN, and NR [27][28][29][30].

NA is a water-soluble vitamin B3 that has been shown to increase tissue NAD⁺ in humans. In a recent clinical trial, patients with mitochondrial myopathy or a healthy age-matched control group were given a steadily increasing dose of NA, starting at 250 mg/day to 750–1000 mg/day over a 4-month period, and then a 10-month follow-up treatment period. NA treatment increased muscle NAD⁺ levels 1.3-fold at 4 months and 2.3-fold after 10 months in the study group. The control group saw no increase in NAD⁺ (NCT03973203, www.clinicaltrials.gov (accessed on 30 July 2021)) [31]. Nevertheless, the side effects and risks of taking high doses of NA are well known and include flushing, an upset stomach, diarrhea, liver damage, stomach ulcers, changes to glucose levels, muscle damage, low blood pressure, and changes in the heart rhythm [32].

NAM, which is also very soluble in water (approximately 1 g/mL), has the advantage of not causing skin flushing. At effective doses, its side effects are minimal, although at high doses it may also cause liver toxicity, nausea, vomiting, headache, fatigue, dizziness, and low platelets in the blood [33]. Moreover, a chronic and excessive intake of NAM may exacerbate motor symptoms of Parkinson's disease [34].

A clinical trial, designed to assess the efficacy of a single-dose supplementation of NAM on NAD⁺ levels, showed that NAM intake within the daily tolerable upper level (200 mg) significantly increased the whole blood NAD⁺ 2–2.5-fold [35]. Olson et al. [36], using radiometric methods to characterize the uptake of [¹⁴C] NAM in human leukemic K-562 cells, demonstrated the binding of NAM to the plasma membrane, followed by its intracellular uptake and immediate synthesis to NAD⁺. However, it is uncertain how many cell types can take up significant amounts of NAM. Moreover, its efficacy decreases with age and stress [33], and inhibits different SIRT activities [37].

NMN is the phosphorylated form of NR. However, its NAD⁺-promoting activity mainly depends on the extracellular conversion of NMN to NR [38]. Then, NR enters the cells through nucleoside transporters and can be re-phosphorylated back to NMN through the reaction catalyzed by NRK [39]. Nevertheless, in rodents, it has been found that orally administered NMN quickly enters the cells of the small intestine through the Slc12a8 NMN transporter, and then is used to generate NAD⁺ [40]. This Slc12a8 carrier system appears approximately 100-fold more active in the small intestine than in other tissues, e.g., adipose cells and the brain [40]. Nevertheless, the preponderance of the evidence supports the idea that the main mechanism, whereby NMN increases intracellular NAD⁺ levels, relies on the extracellular metabolism of NMN to NR, which is then taken up by the cell and converted into NAD⁺. Considering all of this, it is critical to carefully take into account that findings in animal studies do not necessarily translate to humans. In practice, the first-in-humans

clinical trial using NMN was recently published [41]. Specifically, 10 weeks of daily NMN supplementation (250 mg/day) in prediabetic women was shown to increase both circulating NAD⁺ levels and muscle insulin sensitivity [41]. NMN levels in the body were not reported.

Nicotinic acid has also been shown to both increase NAD⁺ levels in human cells through the nicotinic acid phosphoribosyltransferase-catalyzed reaction and to exert protection against oxidative stress [42].

Clinical research on the bioavailability of NR supplementation began in 2016 with a pilot study where a few people were given three single doses of NR. Each person completed each dose (100, 300, or 1000 mg) with a 7-day washout period between each one. The results showed that NR was bioavailable in supplement form and increased blood NAD⁺ when taken in aggregate (no single dose increased NAD⁺ levels significantly) [43]. In 2017, a randomized double-blind trial included 120 healthy subjects (60–80 years old) who were assigned to one of three groups in which each person completed eight weeks of daily supplementation: (1) a placebo group, (2) a dose of 250 mg NR + 50 mg pterostilbene (PT, a natural antioxidant found e.g., in blueberries), or (3) a dose of 500 mg NR + 100 mg PT [44]. The low-dose group showed a 40% increase in blood NAD⁺ compared to the placebo, while NAD⁺ levels in the higher-dose group increased by ~90%. No adverse side effects were observed, and it was concluded that the sustained use of NR safely increases NAD⁺ levels.

In 2019, a small group of 70 to 80-year-old men were given 1000 mg of NR supplementation per day in a randomized control crossover trial. The results showed that NR increased NAD⁺ in the skeletal muscle and also exerted anti-inflammatory properties [45]. In a work published that same year, Conze et al. reported the safety of the long-term (8 weeks) administration of NR (up to 1 g per day) to healthy overweight humans [46]. This study employed a dietary restriction designed to limit the amount of NAD⁺ precursors that could be obtained from the diet. The success of the dietary restriction is shown by the continued decline in NAD⁺ levels of the placebo group during the study. NR administration was also associated with an increase in blood NAD⁺ levels under these conditions [46].

Other approaches that could also be useful to increase NAD⁺ levels in vivo may include, analogs of NMN [47] or inhibitors of CD38 [48], PARP1 [49], SARM1 [50], and α -amino- β -carboxymuconate- ϵ -semialdehyde decarboxylase (ACMSD, a critical enzyme in de novo NAD⁺ biosynthesis; www.tespharma.com (accessed on 30 July 2021)).

2.2. Antioxidants

At present, only edaravone, a pyrazolone free-radical scavenger that may decrease oxidative damage, has been approved for ALS therapy. Its antioxidant effect has been known since 1994 and its impact on other diseases, such as ischemic stroke, was also investigated. However, it was not until 2006–2008 that its effectiveness was demonstrated in the SOD1^{G93A} mouse model [51][52]. In 2017, the FDA's approval was based on the results of a double-blind, randomized, and placebo-controlled phase 3 clinical trial (NCT01492686) that evaluated its safety and effectiveness in treating ALS. After a 12-week observation period, participants were distributed into two groups. One group received 60 mg of edaravone (i.v.) per day for six months, while participants in the other group received a placebo. After the six-month-period of treatment, those who received edaravone experienced a 33% lower decline in the Revised Amyotrophic Lateral Sclerosis Functional Rating Scale (ALSFRS-R) score than those in the control group. Nevertheless, edaravone can cause hives, swelling, and shortness of breath in some people, due to anaphylactic reactions to sulfite-containing infusion components. Otherwise, it is rather well tolerated, although common side effects include walking problems, bruises, and headaches [53]. In a meta-analysis of published post-marketing clinical data, edaravone appeared effective in Asian countries, where its reported benefits on ALSFRS-R scores and lung capacity were similar to those seen in the clinical trials. The drug seems to have little clinical benefit in European countries, and the reason for this difference remains unclear [54].

In addition, other antioxidant molecules have also been considered for ALS therapy, i.e., (but not limited to) vitamin E, coenzyme Q10, melatonin, β -carotene, thiol donors, and natural polyphenols. However, data obtained in regular clinical trials are scarce and discouraging. The most promising of these antioxidant treatments, in the opinion of the authors of this review, are discussed below in more detail.

Vitamin E, in combination with the antiparkinsonian drug selegiline, was first assayed in an 18-month randomized treatment trial, but this long-term antioxidative treatment did not benefit patients with ALS [55]. However, in another double-blind placebo-controlled randomized trial, though vitamin E did not appear to affect the survival and neuromotor functions, patients receiving riluzole plus vitamin E remained in the milder clinical state of the disease for a longer period of time [56]. Nevertheless, at present, there is no evidence that vitamin E may benefit ALS patients once the disease has been diagnosed.

Coenzyme Q10, a mitochondrial cofactor known for its antioxidant properties, has a prolonged survival in mouse models of ALS and has a slowed functional decline in Parkinson's disease. However, a phase II trial of Coenzyme Q10 in ALS did not render sufficient evidence to justify a phase III [57]. Nevertheless, a combined treatment of edaravone and coenzyme Q10 has been proposed [58].

In a small clinical safety study, chronic high-dose (300 mg/day) rectal melatonin was well tolerated during an observation period of up to 2 years. Circulating serum protein carbonyls (a marker of protein oxidation) were elevated in ALS patients, but were normalized to control values by melatonin treatment [59]. Moreover, two ALS cases, in which cocktails of supplements including melatonin were associated with a partial recovery of lost motor function, support further studies with melatonin, at least in a pilot trial [60].

High dietary intakes of β -carotene and lutein have been inversely associated with ALS risk [61]. Thus, the authors of this study suggest that the consumption of foods high in carotenoids may help to prevent or delay the onset of ALS, though we do not yet have evidence of the efficacy of this suggestion.

N-acetyl-L-cysteine (NAC), a direct thiol donor, was shown to improve the survival and neuromotor functions in the SOD1^{G93A} mouse model [62]. However, 50 mg of NAC/kg per day (S.C.) failed to significantly increase the survival or slow disease progression in a randomized, double-blind, and placebo-controlled clinical trial [63].

An ongoing 6-month open-label pilot trial is testing the effect of curcumin (a natural polyphenol responsible for turmeric's yellow color) in ALS patients. This study is expected to be completed in 2021 (NCT04499963).

References

1. Obrador, E.; Salvador-Palmer, R.; López-Blanch, R.; Jihad-Jebbar, A.; Vallés, S.L.; Estrela, J.M. The Link between Oxidative Stress, Redox Status, Bioenergetics and Mitochondria in the Pathophysiology of ALS. *Int. J. Mol. Sci.* 2021, 22, 6352.
2. Pehar, M.; Harlan, B.A.; Killoy, K.M.; Vargas, M.R. Nicotinamide Adenine Dinucleotide Metabolism and Neurodegeneration. *Antioxid. Redox Signal.* 2018, 28, 1652–1668.
3. Liu, D.; Gharavi, R.; Pitta, M.; Gleichmann, M.; Mattson, M.P. Nicotinamide Prevents NAD⁺ Depletion and Protects Neurons against Excitotoxicity and Cerebral Ischemia: NAD⁺ Consumption by SIRT1 May Endanger Energetically Compromised Neurons. *Neuromol. Med.* 2009, 11, 28–42.
4. Bouchard, V.J.; Rouleau, M.; Poirier, G.G. PARP-1, a Determinant of Cell Survival in Response to DNA Damage. *Exp. Hematol.* 2003, 31, 446–454.
5. Rajamohan, S.B.; Pillai, V.B.; Gupta, M.; Sundaresan, N.R.; Birukov, K.G.; Samant, S.; Hottiger, M.O.; Gupta, M.P. SIRT1 Promotes Cell Survival under Stress by Deacetylation-Dependent Deactivation of Poly(ADP-Ribose) Polymerase 1. *Mol. Cell. Biol.* 2009, 29, 4116–4129.
6. Chong, Z.-Z.; Lin, S.-H.; Li, F.; Maiese, K. The Sirtuin Inhibitor Nicotinamide Enhances Neuronal Cell Survival during Acute Anoxic Injury through AKT, BAD, PARP, and Mitochondrial Associated “Anti-Apoptotic” Pathways. *Curr. Neurovasc. Res.* 2005, 2, 271–285.
7. Li, Y.; Xu, W.; McBurney, M.W.; Longo, V.D. SirT1 Inhibition Reduces IGF-I/IRS-2/Ras/ERK1/2 Signaling and Protects Neurons. *Cell Metab.* 2008, 8, 38–48.
8. Katsyuba, E.; Romani, M.; Hofer, D.; Auwerx, J. NAD⁺ Homeostasis in Health and Disease. *Nat. Metab.* 2020, 2, 9–31.
9. Belenky, P.; Bogan, K.L.; Brenner, C. NAD⁺ Metabolism in Health and Disease. *Trends. Biochem. Sci.* 2007, 32, 12–19.
10. Strømmland, Ø.; Niere, M.; Nikiforov, A.A.; VanLinden, M.R.; Heiland, I.; Ziegler, M. Keeping the Balance in NAD Metabolism. *Biochem. Soc. Trans.* 2019, 47, 119–130.
11. Slomka, M.; Zieminska, E.; Salinska, E.; Lazarewicz, J.W. Neuroprotective Effects of Nicotinamide and 1-Methylnicotinamide in Acute Excitotoxicity in Vitro. *Folia Neuropathol.* 2008, 46, 69–80.
12. Sasaki, Y.; Araki, T.; Milbrandt, J. Stimulation of Nicotinamide Adenine Dinucleotide Biosynthetic Pathways Delays Axonal Degeneration after Axotomy. *J. Neurosci.* 2006, 26, 8484–8491.
13. Schöndorf, D.C.; Ivanyuk, D.; Baden, P.; Sanchez-Martinez, A.; De Cicco, S.; Yu, C.; Giunta, I.; Schwarz, L.K.; Di Napoli, G.; Panagiotakopoulou, V.; et al. The NAD⁺ Precursor Nicotinamide Riboside Rescues Mitochondrial Defects and Neuronal Loss in iPSC and Fly Models of Parkinson's Disease. *Cell Rep.* 2018, 23, 2976–2988.
14. Harlan, B.A.; Pehar, M.; Sharma, D.R.; Beeson, G.; Beeson, C.C.; Vargas, M.R. Enhancing NAD⁺ Salvage Pathway Reverts the Toxicity of Primary Astrocytes Expressing Amyotrophic Lateral Sclerosis-Linked Mutant Superoxide Dismutase

15. Gerdts, J.; Brace, E.J.; Sasaki, Y.; DiAntonio, A.; Milbrandt, J. SARM1 Activation Triggers Axon Degeneration Locally via NAD⁺ Destruction. *Science* 2015, 348, 453–457.
16. Alano, C.C.; Tran, A.; Tao, R.; Ying, W.; Karliner, J.S.; Swanson, R.A. Differences among Cell Types in NAD(+) Compartmentalization: A Comparison of Neurons, Astrocytes, and Cardiac Myocytes. *J. Neurosci. Res.* 2007, 85, 3378–3385.
17. Harlan, B.A.; Killoy, K.M.; Pehar, M.; Liu, L.; Auwerx, J.; Vargas, M.R. Evaluation of the NAD⁺ Biosynthetic Pathway in ALS Patients and Effect of Modulating NAD⁺ Levels in HSOD1-Linked ALS Mouse Models. *Exp. Neurol.* 2020, 327, 113219.
18. Park, J.H.; Long, A.; Owens, K.; Kristian, T. Nicotinamide Mononucleotide Inhibits Post-Ischemic NAD(+) Degradation and Dramatically Ameliorates Brain Damage Following Global Cerebral Ischemia. *Neurobiol. Dis.* 2016, 95, 102–110.
19. Karnati, S.; Lüers, G.; Pfreimer, S.; Baumgart-Vogt, E. Mammalian SOD2 Is Exclusively Located in Mitochondria and Not Present in Peroxisomes. *Histochem. Cell. Biol.* 2013, 140, 105–117.
20. Jo, S.H.; Son, M.K.; Koh, H.J.; Lee, S.M.; Song, I.H.; Kim, Y.O.; Lee, Y.S.; Jeong, K.S.; Kim, W.B.; Park, J.W.; et al. Control of Mitochondrial Redox Balance and Cellular Defense against Oxidative Damage by Mitochondrial NADP⁺-Dependent Isocitrate Dehydrogenase. *J. Biol. Chem.* 2001, 276, 16168–16176.
21. Sies, H.; Berndt, C.; Jones, D.P. Oxidative Stress. *Annu. Rev. Biochem.* 2017, 86, 715–748.
22. Mårtensson, J.; Lai, J.C.; Meister, A. High-Affinity Transport of Glutathione Is Part of a Multicomponent System Essential for Mitochondrial Function. *Proc. Natl. Acad. Sci. USA* 1990, 87, 7185–7189.
23. Berman, S.B.; Watkins, S.C.; Hastings, T.G. Quantitative Biochemical and Ultrastructural Comparison of Mitochondrial Permeability Transition in Isolated Brain and Liver Mitochondria: Evidence for Reduced Sensitivity of Brain Mitochondria. *Exp. Neurol.* 2000, 164, 415–425.
24. Jain, A.; Mårtensson, J.; Stole, E.; Auld, P.A.; Meister, A. Glutathione Deficiency Leads to Mitochondrial Damage in Brain. *Proc. Natl. Acad. Sci. USA* 1991, 88, 1913–1917.
25. Killoy, K.M.; Harlan, B.A.; Pehar, M.; Helke, K.L.; Johnson, J.A.; Vargas, M.R. Decreased Glutathione Levels Cause Overt Motor Neuron Degeneration in HSOD1WT Over-Expressing Mice. *Exp. Neurol.* 2018, 302, 129–135.
26. Weiduschat, N.; Mao, X.; Hupf, J.; Armstrong, N.; Kang, G.; Lange, D.J.; Mitumoto, H.; Shungu, D.C. Motor Cortex Glutathione Deficit in ALS Measured in Vivo with the J-Editing Technique. *Neurosci. Lett.* 2014, 570, 102–107.
27. Magni, G.; Amici, A.; Emanuelli, M.; Raffaelli, N.; Ruggieri, S. Enzymology of NAD⁺ Synthesis. *Adv. Enzymol. Relat. Areas Mol. Biol.* 1999, 73, 135–182, xi.
28. Bogan, K.L.; Brenner, C. Nicotinic Acid, Nicotinamide, and Nicotinamide Riboside: A Molecular Evaluation of NAD⁺ Precursor Vitamins in Human Nutrition. *Annu. Rev. Nutr.* 2008, 28, 115–130.
29. Grozio, A.; Sociali, G.; Sturla, L.; Caffa, I.; Soncini, D.; Salis, A.; Raffaelli, N.; De Flora, A.; Nencioni, A.; Bruzzone, S. CD73 Protein as a Source of Extracellular Precursors for Sustained NAD⁺ Biosynthesis in FK866-Treated Tumor Cells. *J. Biol. Chem.* 2013, 288, 25938–25949.
30. Braidy, N.; Berg, J.; Clement, J.; Khorshidi, F.; Poljak, A.; Jayasena, T.; Grant, R.; Sachdev, P. Role of Nicotinamide Adenine Dinucleotide and Related Precursors as Therapeutic Targets for Age-Related Degenerative Diseases: Rationale, Biochemistry, Pharmacokinetics, and Outcomes. *Antioxid. Redox Signal.* 2019, 30, 251–294.
31. Pirinen, E.; Auranen, M.; Khan, N.A.; Brilhante, V.; Urho, N.; Pessia, A.; Hakkarainen, A.; Kuula, J.; Heinonen, U.; Schmidt, M.S.; et al. Niacin Cures Systemic NAD⁺ Deficiency and Improves Muscle Performance in Adult-Onset Mitochondrial Myopathy. *Cell Metab.* 2020, 31, 1078–1090.e5.
32. Habibe, M.N.; Kellar, J.Z. Niacin Toxicity. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2021.
33. Hwang, E.S.; Song, S.B. Possible Adverse Effects of High-Dose Nicotinamide: Mechanisms and Safety Assessment. *Biomolecules* 2020, 10, 687.
34. Harrison, I.F.; Powell, N.M.; Dexter, D.T. The Histone Deacetylase Inhibitor Nicotinamide Exacerbates Neurodegeneration in the Lactacystin Rat Model of Parkinson's Disease. *J. Neurochem.* 2019, 148, 136–156.
35. Ito, T.K.; Sato, T.; Hakamata, A.; Onoda, Y.; Sato, S.; Yamazaki, F.; Horikawa, M.; Takahashi, Y.; Kitamoto, T.; Suzuki, M.; et al. A Nonrandomized Study of Single Oral Supplementation within the Daily Tolerable Upper Level of Nicotinamide Affects Blood Nicotinamide and NAD⁺ Levels in Healthy Subjects. *Transl. Med. Aging* 2020, 4, 45–54.
36. Olsson, A.; Olofsson, T.; Pero, R.W. Specific Binding and Uptake of Extracellular Nicotinamide in Human Leukemic K-562 Cells. *Biochem. Pharmacol.* 1993, 45, 1191–1200.

37. Avalos, J.L.; Bever, K.M.; Wolberger, C. Mechanism of Sirtuin Inhibition by Nicotinamide: Altering the NAD(+) Cosubstrate Specificity of a Sir2 Enzyme. *Mol. Cell* 2005, 17, 855–868.
38. Campagna, R.; Mateuszuk, Ł.; Wojnar-Lason, K.; Kaczara, P.; Tworzydło, A.; Kij, A.; Bujok, R.; Mlynarski, J.; Wang, Y.; Sartini, D.; et al. Nicotinamide N-Methyltransferase in Endothelium Protects against Oxidant Stress-Induced Endothelial Injury. *Biochim. Biophys. Acta Mol. Cell Res.* 2021, 1868, 119082.
39. Ratajczak, J.; Joffraud, M.; Trammell, S.A.J.; Ras, R.; Canela, N.; Boutant, M.; Kulkarni, S.S.; Rodrigues, M.; Redpath, P.; Migaud, M.E.; et al. NRK1 Controls Nicotinamide Mononucleotide and Nicotinamide Riboside Metabolism in Mammalian Cells. *Nat. Commun.* 2016, 7, 13103.
40. Grozio, A.; Mills, K.F.; Yoshino, J.; Bruzzone, S.; Sociali, G.; Tokizane, K.; Lei, H.C.; Cunningham, R.; Sasaki, Y.; Migaud, M.E.; et al. Slc12a8 Is a Nicotinamide Mononucleotide Transporter. *Nat. Metab.* 2019, 1, 47–57.
41. Yoshino, M.; Yoshino, J.; Kayser, B.D.; Patti, G.J.; Franczyk, M.P.; Mills, K.F.; Sindelar, M.; Pietka, T.; Patterson, B.W.; Imai, S.-I.; et al. Nicotinamide Mononucleotide Increases Muscle Insulin Sensitivity in Prediabetic Women. *Science* 2021, 372, 1224–1229.
42. Hara, N.; Yamada, K.; Shibata, T.; Osago, H.; Hashimoto, T.; Tsuchiya, M. Elevation of Cellular NAD Levels by Nicotinic Acid and Involvement of Nicotinic Acid Phosphoribosyltransferase in Human Cells. *J. Biol. Chem.* 2007, 282, 24574–24582.
43. Trammell, S.A.J.; Schmidt, M.S.; Weidemann, B.J.; Redpath, P.; Jaksch, F.; Dellinger, R.W.; Li, Z.; Abel, E.D.; Migaud, M.E.; Brenner, C. Nicotinamide Riboside Is Uniquely and Orally Bioavailable in Mice and Humans. *Nat. Commun.* 2016, 7, 1–14.
44. Dellinger, R.W.; Santos, S.R.; Morris, M.; Evans, M.; Alminana, D.; Guarente, L.; Marcotulli, E. Repeat Dose NRPT (Nicotinamide Riboside and Pterostilbene) Increases NAD⁺ Levels in Humans Safely and Sustainably: A Randomized, Double-Blind, Placebo-Controlled Study. *NPJ Aging Mech. Dis.* 2017, 3, 17.
45. Elhassan, Y.S.; Kluckova, K.; Fletcher, R.S.; Schmidt, M.S.; Garten, A.; Doig, C.L.; Cartwright, D.M.; Oakey, L.; Burley, C.V.; Jenkinson, N.; et al. Nicotinamide Riboside Augments the Aged Human Skeletal Muscle NAD⁺ Metabolome and Induces Transcriptomic and Anti-Inflammatory Signatures. *Cell Rep.* 2019, 28, 1717–1728.e6.
46. Conze, D.; Brenner, C.; Kruger, C.L. Safety and Metabolism of Long-Term Administration of NIAGEN (Nicotinamide Riboside Chloride) in a Randomized, Double-Blind, Placebo-Controlled Clinical Trial of Healthy Overweight Adults. *Sci. Rep.* 2019, 9, 9772.
47. Christ, W.; Coper, H. Preparation and Purification of Nicotinamide Mononucleotide Analogs. *Methods Enzymol.* 1980, 66, 71–81.
48. Tarragó, M.G.; Chini, C.C.S.; Kanamori, K.S.; Warner, G.M.; Caride, A.; de Oliveira, G.C.; Rud, M.; Samani, A.; Hein, K. Z.; Huang, R.; et al. A Potent and Specific CD38 Inhibitor Ameliorates Age-Related Metabolic Dysfunction by Reversing Tissue NAD⁺ Decline. *Cell Metab.* 2018, 27, 1081–1095.e10.
49. Bian, C.; Zhang, C.; Luo, T.; Vyas, A.; Chen, S.-H.; Liu, C.; Kassab, M.A.; Yang, Y.; Kong, M.; Yu, X. NADP⁺ Is an Endogenous PARP Inhibitor in DNA Damage Response and Tumor Suppression. *Nat. Commun.* 2019, 10, 693.
50. Hughes, R.O.; Bosanac, T.; Mao, X.; Engber, T.M.; DiAntonio, A.; Milbrandt, J.; Devraj, R.; Krauss, R. Small Molecule SARM1 Inhibitors Recapitulate the SARM1^{-/-} Phenotype and Allow Recovery of a Metastable Pool of Axons Fated to Degenerate. *Cell Rep.* 2021, 34, 108588.
51. Ito, H.; Wate, R.; Zhang, J.; Ohnishi, S.; Kaneko, S.; Ito, H.; Nakano, S.; Kusaka, H. Treatment with Edaravone, Initiated at Symptom Onset, Slows Motor Decline and Decreases SOD1 Deposition in ALS Mice. *Exp. Neurol.* 2008, 213, 448–455.
52. Zona, C.; Pieri, M.; Carunchio, I. Voltage-Dependent Sodium Channels in Spinal Cord Motor Neurons Display Rapid Recovery from Fast Inactivation in a Mouse Model of Amyotrophic Lateral Sclerosis. *J. Neurophysiol.* 2006, 96, 3314–3322.
53. Luo, L.; Song, Z.; Li, X.; Huiwang, Zeng, Y.; Qinwang, Meiqi, He, J. Efficacy and Safety of Edaravone in Treatment of Amyotrophic Lateral Sclerosis—a Systematic Review and Meta-Analysis. *Neurol. Sci.* 2019, 40, 235–241.
54. Ortiz, J.F.; Khan, S.A.; Salem, A.; Lin, Z.; Iqbal, Z.; Jahan, N. Post-Marketing Experience of Edaravone in Amyotrophic Lateral Sclerosis: A Clinical Perspective and Comparison With the Clinical Trials of the Drug. *Cureus* 2020, 12, e10818.
55. Kweciński, H.; Janik, P.; Jamrozik, Z.; Opuchlik, A. The effect of selegiline and vitamin E in the treatment of ALS: An open randomized clinical trials. *Neurol. Neurochir. Pol.* 2001, 35, 101–106.
56. Desnuelle, C.; Dib, M.; Garrel, C.; Favier, A. A Double-Blind, Placebo-Controlled Randomized Clinical Trial of Alpha-Tocopherol (Vitamin E) in the Treatment of Amyotrophic Lateral Sclerosis. *Amyotroph. Lateral Scler. Other Motor. Neuron Disord.* 2001, 2, 9–18.

57. Kaufmann, P.; Thompson, J.L.P.; Levy, G.; Buchsbaum, R.; Shefner, J.; Krivickas, L.S.; Katz, J.; Rollins, Y.; Barohn, R. J.; Jackson, C.E.; et al. Phase II Trial of CoQ10 for ALS Finds Insufficient Evidence to Justify Phase III. *Ann. Neurol.* 2009, 66, 235–244.
58. Nagase, M.; Yamamoto, Y.; Miyazaki, Y.; Yoshino, H. Increased Oxidative Stress in Patients with Amyotrophic Lateral Sclerosis and the Effect of Edaravone Administration. *Redox Rep.* 2016, 21, 104–112.
59. Weishaupt, J.H.; Bartels, C.; Pölking, E.; Dietrich, J.; Rohde, G.; Poeggeler, B.; Mertens, N.; Sperling, S.; Bohn, M.; Hüt her, G.; et al. Reduced Oxidative Damage in ALS by High-Dose Enteral Melatonin Treatment. *J. Pineal. Res.* 2006, 41, 313–323.
60. ALSUNTANGLED GROUP. ALSUntangled #61: Melatonin. *Amyotroph. Lateral Scler. Frontotemporal. Degener.* 2021, 1–4.
61. Fitzgerald, K.C.; O'Reilly, É.J.; Fondell, E.; Falcone, G.J.; McCullough, M.L.; Park, Y.; Kolonel, L.N.; Ascherio, A. Intake s of Vitamin C and Carotenoids and Risk of Amyotrophic Lateral Sclerosis: Pooled Results from 5 Cohort Studies. *Ann. Neurol.* 2013, 73, 236–245.
62. Andreassen, O.A.; Dedeoglu, A.; Klivenyi, P.; Beal, M.F.; Bush, A.I. N-Acetyl-L-Cysteine Improves Survival and Preserv es Motor Performance in an Animal Model of Familial Amyotrophic Lateral Sclerosis. *Neuroreport* 2000, 11, 2491–2493.
63. Louwerse, E.S.; Weverling, G.J.; Bossuyt, P.M.M.; Meyjes, F.E.P.; Jong, J.M.B.V. de Randomized, Double-Blind, Contr olled Trial of Acetylcysteine in Amyotrophic Lateral Sclerosis. *Arch. Neurol.* 1995, 52, 559–564.

Retrieved from <https://encyclopedia.pub/entry/history/show/32468>