

Congenital Afibrinogenemia

Subjects: Genetics & Heredity

Contributor: Nicole Yin

Congenital afibrinogenemia is a bleeding disorder caused by impairment of the blood clotting process.

Keywords: genetic conditions

1. Introduction

Normally, blood clots protect the body after an injury by sealing off damaged blood vessels and preventing further blood loss. However, bleeding is uncontrolled in people with congenital afibrinogenemia. Newborns with this condition often experience prolonged bleeding from the umbilical cord stump after birth. Nosebleeds (epistaxis) and bleeding from the gums or tongue are common and can occur after minor trauma or in the absence of injury (spontaneous bleeding). Some affected individuals experience bleeding into the spaces between joints (hemarthrosis) or the muscles (hematoma). Rarely, bleeding in the brain or other internal organs occurs, which can be fatal. Women with congenital afibrinogenemia can have abnormally heavy menstrual bleeding (menorrhagia). Without proper treatment, women with this disorder may have difficulty carrying a pregnancy to term, resulting in repeated miscarriages.

2. Frequency

Congenital afibrinogenemia is a rare condition that occurs in approximately 1 in 1 million newborns.

3. Causes

Congenital afibrinogenemia results from mutations in one of three genes, *FGA*, *FGB*, or *FGG*. Each of these genes provides instructions for making one part (subunit) of a protein called fibrinogen. This protein is important for blood clot formation (coagulation), which is needed to stop excessive bleeding after injury. In response to injury, fibrinogen is converted to fibrin, the main protein in blood clots. Fibrin proteins attach to each other, forming a stable network that makes up the blood clot.

Congenital afibrinogenemia is caused by a complete absence of fibrinogen protein. Most *FGA*, *FGB*, and *FGG* gene mutations that cause this condition result in a premature stop signal in the instructions for making the respective protein. If any protein is made, it is nonfunctional. When any one subunit is missing, the fibrinogen protein is not assembled, which results in the absence of fibrin. Consequently, blood clots do not form in response to injury, leading to the excessive bleeding seen in people with congenital afibrinogenemia.

3.1. The Genes Associated with Congenital Afibrinogenemia

- *FGA*
- *FGB*
- *FGG*

4. Inheritance

Congenital afibrinogenemia is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene. The parents have about half the normal level of fibrinogen in their blood but typically do not show signs and symptoms of the condition.

5. Other Names for This Condition

- afibrinogenemia
- familial afibrinogenemia

References

1. Casini A, Lukowski S, Quintard VL, Crutu A, Zak M, Regazzoni S, de Moerloose P, Neerman-Arbez M. FGB mutations leading to congenital quantitative fibrinogen deficiencies: an update and report of four novel mutations. *Thromb Res.* 2014 May;133(5):868-74. doi: 10.1016/j.thromres.2014.01.022.
2. de Moerloose P, Casini A, Neerman-Arbez M. Congenital fibrinogen disorders: an update. *Semin Thromb Hemost.* 2013 Sep;39(6):585-95. doi: 10.1055/s-0033-1349222.
3. Lak M, Keihani M, Elahi F, Peyvandi F, Mannucci PM. Bleeding and thrombosis in 55 patients with inherited afibrinogenemia. *Br J Haematol.* 1999 Oct;107(1):204-6.
4. Neerman-Arbez M, de Moerloose P, Bridel C, Honsberger A, Schönborn A, Rossier C, Peerlinck K, Claeysens S, Di Michele D, d'Oiron R, Dreyfus M, Laubriat-Bianchin M, Dieval J, Antonarakis SE, Morris MA. Mutations in the fibrinogen alpha gene account for the majority of cases of congenital afibrinogenemia. *Blood.* 2000 Jul 1;96(1):149-52.
5. Neerman-Arbez M, de Moerloose P, Honsberger A, Parlier G, Arnuti B, Biron C, Borg JY, Eber S, Meili E, Peter-Salonen K, Ripoll L, Vervel C, d'Oiron R, Staeger P, Antonarakis SE, Morris MA. Molecular analysis of the fibrinogen gene cluster in 16 patients with congenital afibrinogenemia: novel truncating mutations in the FGA and FGG genes. *Hum Genet.* 2001 Mar;108(3):237-40.
6. Neerman-Arbez M, Honsberger A, Antonarakis SE, Morris MA. Deletion of the fibrinogen [correction of fibrogen] alpha-chain gene (FGA) causes congenital afibrinogenemia. *J Clin Invest.* 1999 Jan;103(2):215-8. Erratum in: *J Clin Invest* 1999 Mar;103(5):759.
7. Neerman-Arbez M. Molecular basis of fibrinogen deficiency. *Pathophysiol Haemost Thromb.* 2006;35(1-2):187-98. Review.
8. Weisel JW. Fibrinogen and fibrin. *Adv Protein Chem.* 2005;70:247-99. Review.

Retrieved from <https://encyclopedia.pub/entry/history/show/11269>