

Predefined Diets of IBD

Subjects: Nutrition & Dietetics

Contributor: Pablo Caballero

Inflammatory bowel disease (IBD) is a chronic disease mediated by the immune system and characterized by the importance of diet in pathological development.

Keywords: Inflammatory bowel diseases ; nutrition therapy

1. Introduction

Inflammatory bowel disease (IBD) is characterized by chronic inflammation of the gastrointestinal tract with periods of remission or recurrence and includes both Crohn's disease (CD) and ulcerative colitis (UC) ^[1].

In CD, inflammation can be present in any area of the entire gastrointestinal tract, whereas in UC, the inflammatory process affects only the colon ^[2].

The symptoms of this type of disease are diverse, including the appearance of diarrhea, bloody stools, abdominal pain, fatigue, weight loss, etc. ^[3]. The prevalence exceeds 0.3% in North America, Oceania, and many European countries, and the incidence of these types of pathologies has increased rapidly in recently industrialized countries, producing a high burden on health systems ^{[4][5]}.

IBD, apart from being treated with expensive medical treatments to alleviate its activity, leads to a decrease in the patient's quality of life that can affect the degree of disability and work productivity and is associated with more symptoms of anxiety and depression ^{[6][7][8]}. In addition, malnutrition and specific nutritional deficiencies are frequent among these types of patients, depending on the state and/or progress of the disease, and most patients impose dietary restrictions based on their own beliefs ^{[9][10]}.

The etiology of IBD is unknown; in fact, it is a multifactorial disease. However, Westernised lifestyles and diets are one of the main drivers of the increasing incidence ^{[11][12]}. Diet plays an important role in the gut's microbial composition and functioning, intestinal barrier, host immunity, and intestinal physiology ^{[12][13]}.

Regarding nutritional treatment, for artificial nutrition, such as enteral and parenteral nutrition, there is evidence of efficacy ^{[14][15]}, but for natural nutrition, it seems that a diet that is low in fiber, high in fat, and high in carbohydrates can lead to severe dysbiosis, while one richer in fruits, vegetables, and olive oil could prevent it. However, together with the role of certain foods such as meat, fish, and dairy products, their role in the development of the disease is controversial and uncertain in the absence of studies ^{[16][17][18]}.

On the other hand, while for pediatric patients it seems that the implementation of predefined diets, such as the specific carbohydrate diet (SCD) or the Crohn's disease exclusion diet (CDED), is positive ^[19], for the adult population, the studies performed to date have been inconclusive and the findings are inconsistent in addition to being mostly concerned with symptomatology of the inactive disease ^{[20][21][22][23]}.

Therefore, there is a need for more and clearer evidence that allows health professionals to increase their knowledge in order to advise their patients on what type of specific dietary formulas or nutrients they will need that will allow them not only to comply with nutritional requirements but also to improve the outcome of their symptoms, with positive repercussions for a better quality of life ^{[24][25]}.

2. Discussion

Our systematic review included a total of 31 studies, which compiled information from 5331 individuals with IBD and who had an intervention with different predefined diets. All the studies had a broad reach, and within the diverse effects found, CDAI, FC, CRP, and ALB were the most common, allowing us to conduct a meta-analysis to arrive at more complete conclusions.

The main premise of these types of diets was based on the reduction of some types of pro-inflammatory foods and the increase of others, which are believed to promote a favorable intestinal microbiota [26]. In combination with the high prevalence of malnutrition, the importance of diets that can modify the intestinal barrier and host immunity must be increased [27][28]. In fact, although we did not observe an amelioration in terms of ALB, CRP, and CF levels, an improvement in CDAI levels was observed through interventions with predefined diets, more specifically of microparticles diet, semi-vegetarian diet, and immunoglobulin exclusion diet, in patients with CD.

The low FODMAP diet (LFD) reduces fermented oligosaccharides, disaccharides, monosaccharides, and polyols because they are poorly absorbed in the small intestine and are fermented by bacteria in the colon, triggering intestinal discomfort and gas in sensitive individuals [29][30][31]. This diet has been used mainly with patients with irritable bowel syndrome; however, it has been transferred to patients with IBD due to the similarity of functional gut symptoms such as bloating, abdominal pain, wind, and diarrhea [32][31]. As for the results obtained in our systematic review, most of the individuals improved their symptoms of the disease [32][33][34][35]. This coincides with other studies, in which an improvement was reported due to the use of an LFD for the treatment of gastrointestinal symptoms [23][36]. Furthermore, according to Pedersen et al., Testa et al., Bodini et al., and Cox et al., the LFD reported a better quality of life, although it was measured with different questionnaires [37][38][39][34]. Results of good adherence to this type of diet have also been reported [32][38][34][40], but in terms of disease activity, the results have been controversial; while for some authors no improvements were found for biomarkers or indices such as CRP, FC, HBI, or IBS-SSS, others did obtain improvements [32][37][38][39][34].

All of this, together with the concern of several authors who expressed the possibility that this type of diet may alter the microbiome by increasing the colonic pH, thereby allowing enteropathogenic colonization and causing an increase in dysbiosis [26][41][42], indicate that the use of supplementation should be considered to avoid deficiencies that could be caused by an LFD for long periods of time. Furthermore, it is of great importance that it be considered in the "induction" phase of prescription of diet modification, and if patients do not respond to the modification, the FODMAP restriction should be discontinued [41], as it can compromise the nutritional status of the patient and, to some extent, can affect intestinal inflammation [42].

The Specific Carbohydrate Diet (SCD) is based on the hypothesis that IBD patients have a dysfunction of disaccharidases, which are necessary to digest and absorb disaccharides and amylopectin. Therefore, high amounts of these compounds could cause an overgrowth of bacteria and intestinal lesions which can increase the intestinal permeability, and this is why this type of diet allows foods with carbohydrates that consist only of monosaccharides and excludes disaccharides and most polysaccharides [20]. An improvement in the symptomatology and an increase in clinical remissions are the most important results reported by Suskind et al. [39].

Both SCD and LFD have the potential to contribute to vitamin D deficiency. Therefore, their follow-up and clinical evaluation is very important due to the association of this deficiency with an increased risk of surgery and hospitalization [43][44][45][46].

The Immunoglobulin Exclusion Diet (IGED) is a dietary strategy associated with the identification of foods that cause a certain degree of intolerance, meaning an IgG-mediated reaction that acts as a delayed-type hypersensitivity response to antigen exposure, all of which result in excessive protective immune responses that could lead to increased disease activity [47][48][49]. The researchers Rajendran et al., Gunasekeera et al., and Uzunismail obtained improvements in the activity of the pathology through various tools. However, contradictory results were found for symptomatology, quality of life, and certain biochemical parameters such as CRP and ALB [50][51][52][53][54].

Several authors state that vegetarian dietary patterns are associated with a decrease in serum CRP, fibrinogen, and total leukocyte concentrations [55]. This coincides with the results obtained by Chiba et al., in which an improvement in the CRP, symptoms, and certain laboratory data could be observed [56][57]. However, it can cause an increase in posttraumatic stress and poorer mental health [58].

With respect to the Mediterranean diet (MED), characterized by the consumption of important sources of fiber (cereals, legumes, vegetables, fruits, and nuts) and with a high content of chemical compounds with antioxidant properties such as flavonoids, phytosterols, vitamins, terpenes, and polyphenols [43][59], we have obtained positive results with quality of life, HBI, FC, and cholesterol [60][61][62]. Currently, there is some controversy regarding the role of this diet in IBD, as several authors indicated that a healthy diet pattern, which includes the MED, is associated with significant reductions in inflammation-related CRP [63], and other researchers concluded that this type of diet does not have significant effects on inflammatory substances [64].

Also, there is the gluten-free diet (GFD), which eliminates the gliadin protein located in wheat, barley, rye, and other grains. This diet has been traditionally used for patients with celiac disease and more recently in people with sensitivity to non-celiac gluten [65]. However, the nutrient responsible for improvement is controversial, since these cereals have more than one possible symptom inducer such as gluten, fructans, trypsin amylase inhibitors, and lectins [66][67][68]. The results from our systematic review are controversial. On the one hand, the use of this type of diet improved the symptoms of pathology; however, it could also lead to an increase in anxiety and depression, possibly due to the difficulty of adherence [69][58]. These findings coincide with the results from some authors, who state that GFD, despite the existence of data indicating low adherence, suggests a potential benefit and great utility in the management of IBD [26][44].

Despite being the first systematic review that deals with the general effects of predefined diets on adult patients with IBD, this article is not exempt from limitations. It is possible that the CONSORT questionnaire was not the best for evaluating the Non-randomized controlled clinical trials (NRCCT) and Uncontrolled and non-randomized clinical trial (UNRCT) reviewed; however, we tried to avoid this limitation by adjusting the items of this tool to the type of study, as no questionnaire was found that evaluated the Randomized controlled clinical trial (RCCT), the NRCCT, and the UNRCT [70][71]. Also, some studies were somewhat old, which could have reduced the score of this tool on the methodological quality due to the lack of standard criteria at the time the clinical trials were conducted. The UC and CD data were combined to perform a meta-analysis for the variables CDAI, FC, CRP, and ALB due to the low number of studies that separated these diseases to elaborate on their results and the great variability, not only of the tools used but also of the unit of measurement employed. However, these clinical entities have different clinical courses. The results derived from this work could help in clinical practice to help health professionals, through the creation of a guide oriented towards evaluating the addition of predefined diets within the set of medical therapies for an adult patient diagnosed with IBD. Both clinical trials and observational studies have been used within this systematic review, a parameter that has allowed us to have a more global view of the effect of intervention.

As future lines of research, the use of other types of predefined diets should be considered, which have been observed to show positive results in such patients and for which little evidence is found [72][73][60][74][75][69][76][57][77][78].

References

1. Chen, X.-L.; Zhong, L.; Wen, Y.; Liu, T.-W.; Li, X.-Y.; Hou, Z.-K.; Hu, Y.; Mo, C.; Liu, F.-B. Inflammatory bowel disease-specific health-related quality of life instruments: a systematic review of measurement properties. *Health Qual. Life Outcomes* 2017, 15, 177, doi:10.1186/s12955-017-0753-2.
2. Tárrago, C.P.; Maestu, A.P.; Miján De La Torre, A. Tratamiento nutricional en la enfermedad inflamatoria intestinal. *Nutr. Hosp.* 2008, 23, 418–428.
3. Cortés, R.L.; Fernández, B.M.; Montoro, C.H.; Hernández, P.E.; Aznarez, A.C.S.; Gutiérrez, C.R. Calidad de vida relacionada con la salud en pacientes con enfermedad inflamatoria intestinal. *An. Sist. Sanit. Navar.* 2016, 39, 123–131, doi:10.23938/S1137-6627/2016000100014.
4. Ng, S.C.; Shi, H.Y.; Hamidi, N.; Underwood, F.E.; Tang, W.; Benchimol, E.I.; Panaccione, R.; Ghosh, S.; Wu, J.C.Y.; Chan, F.K.L.; et al. Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet (London, England)* 2018, 390, 2769–2778, doi:10.1016/S0140-6736(17)32448-0.
5. Molodecky, N.A.; Soon, I.S.; Rabi, D.M.; Ghali, W.A.; Ferris, M.; Chernoff, G.; Benchimol, E.I.; Panaccione, R.; Ghosh, S.; Barkema, H.W.; et al. Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. *Gastroenterology* 2012, 142, 46-54.e42, doi:10.1053/j.gastro.2011.10.001.
6. De Boer, A.G.E.M.; Bennebroek Evertsz, F.; Stokkers, P.C.; Bockting, C.L.; Sanderman, R.; Hommes, D.W.; Sprangers, M.A.G.; Frings-Dresen, M.H.W. Employment status, difficulties at work and quality of life in inflammatory bowel disease patients. *Eur. J. Gastroenterol. Hepatol.* 2016, 28, 1130–1136, doi:10.1097/MEG.0000000000000685.
7. Lo, B.; Prosberg, M. V; Gluud, L.L.; Chan, W.; Leong, R.W.; van der List, E.; van der Have, M.; Sarter, H.; Gower-Rousseau, C.; Peyrin-Biroulet, L.; et al. Systematic review and meta-analysis: assessment of factors affecting disability in inflammatory bowel disease and the reliability of the inflammatory bowel disease disability index. *Aliment. Pharmacol. Ther.* 2018, 47, 6–15, doi:10.1111/apt.14373.
8. Everhov, A.H.; Khalili, H.; Askling, J.; Myrelid, P.; Ludvigsson, J.F.; Halfvarson, J.; Nordenvall, C.; Soderling, J.; Olen, O.; Neovius, M. Sick Leave and Disability Pension in Prevalent Patients With Crohn's Disease. *J. Crohn's Colitis* 2018, 12, 1418–1428, doi:10.1093/ecco-jcc/jjy123.

9. Casanova, M.J.; Chaparro, M.; Molina, B.; Merino, O.; Batanero, R.; Dueñas-Sadornil, C.; Robledo, P.; Garcia-Albert, A.M.; Gómez-Sánchez, M.B.; Calvet, X.; et al. Prevalence of malnutrition and nutritional characteristics of patients with inflammatory bowel disease. *J. Crohn's Colitis* 2017, 11, 1430–1439, doi:10.1093/ecco-jcc/jjx102.
10. Bergeron, F.; Bouin, M.; D'Aoust, L.; Lemoyne, M.; Presse, N. Food avoidance in patients with inflammatory bowel disease: What, when and who? *Clin. Nutr.* 2018, 37, 884–889, doi:10.1016/j.clnu.2017.03.010.
11. Chiba, M.; Nakane, K.; Komatsu, M. Westernized Diet is the Most Ubiquitous Environmental Factor in Inflammatory Bowel Disease. *Perm. J.* 2019, 23, 18–107, doi:10.7812/TPP/18-107.
12. Reddavid, R.; Rotolo, O.; Caruso, M.G.; Stasi, E.; Notarnicola, M.; Miraglia, C.; Nouvenne, A.; Meschi, T.; De' Angelis, G.L.; Di Mario, F.; et al. The role of diet in the prevention and treatment of Inflammatory Bowel Diseases. *Acta Biomed.* 2018, 89, 60–75, doi:10.23750/abm.v89i9-S.7952.
13. Rizzello, F.; Spisni, E.; Giovanardi, E.; Imbesi, V.; Salice, M.; Alvisi, P.; Valerii, M.C.; Gionchetti, P. Implications of the Westernized Diet in the Onset and Progression of IBD. *Nutrients* 2019, 11, doi:10.3390/nu11051033.
14. Comeche, J.M.; Caballero, P.; Gutierrez-Hervas, A.; Garcia-Sanjuan, S.; Comino, I.; Altavilla, C.; Tuells, J. Enteral Nutrition in Patients with Inflammatory Bowel Disease. Systematic Review, Meta-Analysis, and Meta-Regression. *Nutrients* 2019, 11, doi:10.3390/nu11112657.
15. Comeche, J.M.; Comino, I.; Altavilla, C.; Tuells, J.; Gutierrez-Hervas, A.; Caballero, P. Parenteral Nutrition in Patients with Inflammatory Bowel Disease Systematic Review, Meta-Analysis and Meta-Regression. *Nutrients* 2019, 11, doi:10.3390/nu11122865.
16. Tomasello, G.; Mazzola, M.; Leone, A.; Sinagra, E.; Zummo, G.; Farina, F.; Damiani, P.; Cappello, F.; Geagea, A.G.; Jurjus, A.; et al. Nutrition, oxidative stress and intestinal dysbiosis: Influence of diet on gut microbiota in inflammatory bowel diseases. *Biomed. Pap.* 2016, 160, 461–466, doi:10.5507/bp.2016.052.
17. Baños Madrid, R.; Salama Benarroch, H.; Morán Sánchez, S.; Gallardo Sanchez, F.; Albaladejo Meroño, A.; Mercader Martínez, J. Lactose malabsorption in patients with inflammatory bowel disease without activity: It would be necessary to exclude the lactose products in their diet to all the patients? [Malabsorción de lactosa en pacientes con enfermedad inflamatoria intestinal inactiva. *An. Med. Interna* 2004, 21, 212–214.
18. Vidarsdottir, J.B.; Johannsdottir, S.E.; Thorsdottir, I.; Björnsson, E.; Ramel, A. A cross-sectional study on nutrient intake and -status in inflammatory bowel disease patients. *Nutr. J.* 2016, 15, 61, doi:10.1186/s12937-016-0178-5.
19. Penagini, F.; Dilillo, D.; Borsani, B.; Cococcioni, L.; Galli, E.; Bedogni, G.; Zuin, G.; Zuccotti, G.V. Nutrition in Pediatric Inflammatory Bowel Disease: From Etiology to Treatment. A Systematic Review. *Nutrients* 2016, 8, doi:10.3390/nu8060334.
20. Aleksandrova, K.; Romero-Mosquera, B.; Hernandez, V. Diet, Gut Microbiome and Epigenetics: Emerging Links with Inflammatory Bowel Diseases and Prospects for Management and Prevention. *Nutrients* 2017, 9, doi:10.3390/nu9090962.
21. Gu, P.; Feagins, L.A. Dining With Inflammatory Bowel Disease: A Review of the Literature on Diet in the Pathogenesis and Management of IBD. *Inflamm. Bowel Dis.* 2020, 26, 181–191, doi:10.1093/ibd/izz268.
22. Limketkai, B.N.; Iheozor-Ejiofor, Z.; Gjuladin-Hellon, T.; Parian, A.; Matarese, L.E.; Bracewell, K.; MacDonald, J.K.; Gordon, M.; Mullin, G.E. Dietary interventions for induction and maintenance of remission in inflammatory bowel disease. *Cochrane Database Syst. Rev.* 2019, 2019, doi:10.1002/14651858.CD012839.pub2.
23. Zhan, Y.; Zhan, Y.-A.; Dai, S.-X. Is a low FODMAP diet beneficial for patients with inflammatory bowel disease? A meta-analysis and systematic review. *Clin. Nutr.* 2018, 37, 123–129, doi:10.1016/j.clnu.2017.05.019.
24. Guagnozzi, D.; González-Castillo, S.; Oliveira, A.; Lucendo, A.J. Nutritional treatment in inflammatory bowel disease. An update. *Rev Esp Enferm Dig* 2012, 104, 479–488.
25. Charlebois, A.; Rosenfeld, G.; Bressler, B. The Impact of Dietary Interventions on the Symptoms of Inflammatory Bowel Disease: A Systematic Review. *Crit. Rev. Food Sci. Nutr.* 2016, 56, 1370–1378, doi:10.1080/10408398.2012.760515.
26. Best, W.R.; Beckett, J.M.; Singleton, J.W.; Kern, F.J. Development of a Crohn's disease activity index. National Cooperative Crohn's Disease Study. *Gastroenterology* 1976, 70, 439–444.
27. Limketkai, B.N.; Wolf, A.; Parian, A.M. Nutritional Interventions in the Patient with Inflammatory Bowel Disease. *Gastroenterol. Clin. North Am.* 2018, 47, 155–177, doi:10.1016/j.gtc.2017.09.007.
28. Li, S.; Ney, M.; Eslamparast, T.; Vandermeer, B.; Ismond, K.P.; Kroeker, K.; Halloran, B.; Raman, M.; Tandon, P. Systematic review of nutrition screening and assessment in inflammatory bowel disease. *World J. Gastroenterol.* 2019, 25, 3823–3837, doi:10.3748/wjg.v25.i28.3823.

29. Levine, A.; Sigall Boneh, R.; Wine, E. Evolving role of diet in the pathogenesis and treatment of inflammatory bowel diseases. *Gut* 2018, 67, 1726–1738, doi:10.1136/gutjnl-2017-315866.
30. Ankersen, D.V.; Carlsen, K.; Marker, D.; Munkholm, P.; Burisch, J. Using eHealth strategies in delivering dietary and other therapies in patients with irritable bowel syndrome and inflammatory bowel disease. *J. Gastroenterol. Hepatol.* 2017, 32 Suppl 1, 27–31, doi:10.1111/jgh.13691.
31. Barrett, J.S. How to institute the low-FODMAP diet. *J. Gastroenterol. Hepatol.* 2017, 32 Suppl 1, 8–10, doi:10.1111/jgh.13686.
32. Croagh, C.; Shepherd, S.J.; Berryman, M.; Muir, J.G.; Gibson, P.R. Pilot study on the effect of reducing dietary FODMAP intake on bowel function in patients without a colon. *Inflamm. Bowel Dis.* 2007, 13, 1522–1528, doi:10.1002/ibd.20249.
33. Gearry, R.B.; Irving, P.M.; Barrett, J.S.; Nathan, D.M.; Shepherd, S.J.; Gibson, P.R. Reduction of dietary poorly absorbed short-chain carbohydrates (FODMAPs) improves abdominal symptoms in patients with inflammatory bowel disease—a pilot study. *J. CROHNS COLITIS* 2009, 3, 8–14, doi:10.1016/j.crohns.2008.09.004.
34. Konijeti, G.G.; Kim, N.; Lewis, J.D.; Groven, S.; Chandrasekaran, A.; Grandhe, S.; Diamant, C.; Singh, E.; Oliveira, G.; Wang, X.; et al. Efficacy of the Autoimmune Protocol Diet for Inflammatory Bowel Disease. *Inflamm. Bowel Dis.* 2017, 23, 2054–2060, doi:10.1097/MIB.0000000000001221.
35. Godny, L.; Reshef, L.; Pfeffer-Gik, T.; Goren, I.; Yanai, H.; Tulchinsky, H.; Gophna, U.; Dotan, I. Adherence to the Mediterranean diet is associated with decreased fecal calprotectin in patients with ulcerative colitis after pouch surgery. *Eur. J. Nutr.* 2020, 59, 3183–3190, doi:10.1007/s00394-019-02158-3.
36. Lewis, J.D.; Abreu, M.T. Diet as a Trigger or Therapy for Inflammatory Bowel Diseases. *Gastroenterology* 2017, 152, 398+, doi:10.1053/j.gastro.2016.10.019.
37. Herfarth, H.H.; Martin, C.F.; Sandler, R.S.; Kappelman, M.D.; Long, M.D. Prevalence of a gluten-free diet and improvement of clinical symptoms in patients with inflammatory bowel diseases. *Inflamm. Bowel Dis.* 2014, 20, 1194–1197, doi:10.1097/MIB.000000000000077.
38. Kyaw, M.H.; Moshkovska, T.; Mayberry, J. A prospective, randomized, controlled, exploratory study of comprehensive dietary advice in ulcerative colitis: impact on disease activity and quality of life. *Eur. J. Gastroenterol. Hepatol.* 2014, 26, 910–917, doi:10.1097/MEG.0000000000000127.
39. Prince, A.C.; Myers, C.E.; Joyce, T.; Irving, P.; Lomer, M.; Whelan, K. Fermentable Carbohydrate Restriction (Low FODMAP Diet) in Clinical Practice Improves Functional Gastrointestinal Symptoms in Patients with Inflammatory Bowel Disease. *Inflamm. Bowel Dis.* 2016, 22, 1129–1136, doi:10.1097/MIB.0000000000000708.
40. Bodini, G.; Zanella, C.; Crespi, M.; Lo Pumo, S.; Demarzo, M.G.; Savarino, E.; Savarino, V.; Giannini, E.G. A randomized, 6-wk trial of a low FODMAP diet in patients with inflammatory bowel disease. *NUTRITION* 2019, 67–68, doi:10.1016/j.nut.2019.06.023.
41. Marsh, A.; Eslick, E.M.; Eslick, G.D. Does a diet low in FODMAPs reduce symptoms associated with functional gastrointestinal disorders? A comprehensive systematic review and meta-analysis. *Eur. J. Nutr.* 2016, 55, 897–906, doi:10.1007/s00394-015-0922-1.
42. Barbalho, S.M.; Goulart, R. de A.; Aranao, A.L. de C.; de Oliveira, P.G.C. Inflammatory Bowel Diseases and Fermentable Oligosaccharides, Disaccharides, Monosaccharides, and Polyols: An Overview. *J. Med. Food* 2018, 21, 633–640, doi:10.1089/jmf.2017.0120.
43. Gibson, P.R. Use of the low-FODMAP diet in inflammatory bowel disease. *J. Gastroenterol. Hepatol.* 2017, 32 Suppl 1, 40–42, doi:10.1111/jgh.13695.
44. Hou, J.K.; Lee, D.; Lewis, J. Diet and inflammatory bowel disease: review of patient-targeted recommendations. *Clin. Gastroenterol. Hepatol.* 2014, 12, 1592–1600, doi:10.1016/j.cgh.2013.09.063.
45. Ruemmele, F.M. Role of Diet in Inflammatory Bowel Disease. *Ann. Nutr. Metab.* 2016, 68, 33–41, doi:10.1159/000445392.
46. Kakodkar, S.; Mutlu, E.A. Diet as a Therapeutic Option for Adult Inflammatory Bowel Disease. *Gastroenterol. Clin. North Am.* 2017, 46, 745–767, doi:10.1016/j.gtc.2017.08.016.
47. Mouli, V.P.; Ananthakrishnan, A.N. Review article: vitamin D and inflammatory bowel diseases. *Aliment. Pharmacol. Ther.* 2014, 39, 125–136, doi:10.1111/apt.12553.
48. Crowe, S.E.; Perdue, M.H. Gastrointestinal food hypersensitivity: basic mechanisms of pathophysiology. *Gastroenterology* 1992, 103, 1075–1095, doi:10.1016/0016-5085(92)90047-3.

49. Ortolani, C.; Pastorello, E.A. Food allergies and food intolerances. *Best Pract. Res. Clin. Gastroenterol.* 2006, 20, 467–483, doi:10.1016/j.bpg.2005.11.010.
50. Berghouse, L.; Hori, S.; Hill, M.; Hudson, M.; Lennard-Jones, J.E.; Rogers, E. Comparison between the bacterial and oligosaccharide content of ileostomy effluent in subjects taking diets rich in refined or unrefined carbohydrate. *Gut* 1984, 25, 1071–1077, doi:10.1136/gut.25.10.1071.
51. Jones, V.A.; Dickinson, R.J.; Workman, E.; Wilson, A.J.; Freeman, A.H.; Hunter, J.O. Crohn's disease: maintenance of remission by diet. *Lancet (London, England)* 1985, 2, 177–180, doi:10.1016/s0140-6736(85)91497-7.
52. Rajendran, N.; Kumar, D. Food-specific IgG4-guided exclusion diets improve symptoms in Crohn's disease: a pilot study. *Colorectal Dis.* 2011, 13, 1009–1013, doi:10.1111/j.1463-1318.2010.02373.x.
53. Halmos, E.P.; Christophersen, C.T.; Bird, A.R.; Shepherd, S.J.; Muir, J.G.; Gibson, P.R. Consistent Prebiotic Effect on Gut Microbiota With Altered FODMAP Intake in Patients with Crohn's Disease: A Randomised, Controlled Cross-Over Trial of Well-Defined Diets. *Clin. Transl. Gastroenterol.* 2016, 7, e164, doi:10.1038/ctg.2016.22.
54. Schreiner, P.; Yilmaz, B.; Rossel, J.-B.; Franc, Y.; Misselwitz, B.; Scharl, M.; Zeitz, J.; Frei, P.; Greuter, T.; Vavricka, S.R.; et al. Vegetarian or gluten-free diets in patients with inflammatory bowel disease are associated with lower psychological well-being and a different gut microbiota, but no beneficial effects on the course of the disease. *UNITED Eur. Gastroenterol. J.* 2019, 7, 767–781, doi:10.1177/2050640619841249.
55. Cai, C.; Shen, J.; Zhao, D.; Qiao, Y.; Xu, A.; Jin, S.; Ran, Z.; Zheng, Q. Serological investigation of food specific immunoglobulin G antibodies in patients with inflammatory bowel diseases. *PLoS One* 2014, 9, doi:10.1371/journal.pone.0112154.
56. Bentz, S.; Hausmann, M.; Piberger, H.; Kellermeier, S.; Paul, S.; Held, L.; Falk, W.; Obermeier, F.; Fried, M.; Scholmerich, J.; et al. Clinical relevance of IgG antibodies against food antigens in Crohn's disease: a double-blind cross-over diet intervention study. *Digestion* 2010, 81, 252–264, doi:10.1159/000264649.
57. Pedersen, N.; Ankersen, D.V.; Felding, M.; Wachmann, H.; Vegh, Z.; Molzen, L.; Burisch, J.; Andersen, J.R.; Munkholm, P. Low-FODMAP diet reduces irritable bowel symptoms in patients with inflammatory bowel disease. *World J. Gastroenterol.* 2017, 23, 3356–3366, doi:10.3748/wjg.v23.i18.3356.
58. Papada, E.; Amerikanou, C.; Forbes, A.; Kaliora, A.C. Adherence to Mediterranean diet in Crohn's disease. *Eur. J. Nutr.* 2020, 59, 1115–1121, doi:10.1007/s00394-019-01972-z.
59. Craddock, J.C.; Neale, E.P.; Peoples, G.E.; Probst, Y.C. Vegetarian-Based Dietary Patterns and their Relation with Inflammatory and Immune Biomarkers: A Systematic Review and Meta-Analysis. *Adv. Nutr.* 2019, 10, 433–451, doi:10.1093/advances/nmy103.
60. Ritchie, J.K.; Wadsworth, J.; Lennard-Jones, J.E.; Rogers, E. Controlled multicentre therapeutic trial of an unrefined carbohydrate, fibre rich diet in Crohn's disease. *Br. Med. J. (Clin. Res. Ed)*. 1987, 295, 517–520, doi:10.1136/bmj.295.6597.517.
61. Komperød, M.J.; Sommer, C.; Mellin-Olsen, T.; Iversen, P.O.; Røseth, A.G.; Valeur, J. Persistent symptoms in patients with Crohn's disease in remission: An exploratory study on the role of diet. *Scand. J. Gastroenterol.* 2018, 53, 573–578, doi:10.1080/00365521.2017.1397736.
62. Chiba, M.; Nakane, K.; Tsuji, T.; Tsuda, S.; Ishii, H.; Ohno, H.; Watanabe, K.; Ito, M.; Komatsu, M.; Yamada, K.; et al. Relapse Prevention in Ulcerative Colitis by Plant-Based Diet Through Educational Hospitalization: A Single-Group Trial. *Perm. J.* 2018, 22, 17–167, doi:10.7812/TPP/17-167.
63. Bifulco, M. Mediterranean diet: the missing link between gut microbiota and inflammatory diseases. *Eur. J. Clin. Nutr.* 2015, 69, 1078.
64. Neale, E.P.; Batterham, M.J.; Tapsell, L.C. Consumption of a healthy dietary pattern results in significant reductions in C-reactive protein levels in adults: a meta-analysis. *Nutr. Res.* 2016, 36, 391–401, doi:10.1016/j.nutres.2016.02.009.
65. Mayr, H.L.; Tierney, A.C.; Thomas, C.J.; Ruiz-Canela, M.; Radcliffe, J.; Itsiopoulos, C. Mediterranean-type diets and inflammatory markers in patients with coronary heart disease: a systematic review and meta-analysis. *Nutr. Res.* 2018, 50, 10–24, doi:10.1016/j.nutres.2017.10.014.
66. Elli, L.; Tomba, C.; Branchi, F.; Roncoroni, L.; Lombardo, V.; Bardella, M.T.; Ferretti, F.; Conte, D.; Valiante, F.; Fini, L.; et al. Evidence for the Presence of Non-Celiac Gluten Sensitivity in Patients with Functional Gastrointestinal Symptoms: Results from a Multicenter Randomized Double-Blind Placebo-Controlled Gluten Challenge. *Nutrients* 2016, 8, 84, doi:10.3390/nu8020084.
67. Skodje, G.I.; Sarna, V.K.; Minelle, I.H.; Rolfsen, K.L.; Muir, J.G.; Gibson, P.R.; Veierød, M.B.; Henriksen, C.; Lundin, K.E.A. Fructan, Rather Than Gluten, Induces Symptoms in Patients With Self-Reported Non-Celiac Gluten Sensitivity. *Gastroenterology* 2018, 154, 529–539.e2, doi:10.1053/j.gastro.2017.10.040.

68. Skodje, G.I.; Minelle, I.H.; Rolfsen, K.L.; Iacovou, M.; Lundin, K.E.A.; Veierød, M.B.; Henriksen, C. Dietary and symptom assessment in adults with self-reported non-coeliac gluten sensitivity. *Clin. Nutr. ESPEN* 2019, 31, 88–94, doi:10.1016/j.clnesp.2019.02.012.
69. Marlow, G.; Ellett, S.; Ferguson, I.R.; Zhu, S.; Karunasinghe, N.; Jesuthasan, A.C.; Han, D.Y.; Fraser, A.G.; Ferguson, L.R. Transcriptomics to study the effect of a Mediterranean-inspired diet on inflammation in Crohn's disease patients. *Hum. Genomics* 2013, 7, 24, doi:10.1186/1479-7364-7-24.
70. Campos Cañuelo, D.; Sanz-Valero, J.; Wanden-Berghe, C. Consecuencias de la nutrición parenteral domiciliaria en adultos con síndrome de intestino corto: revisión exploratoria. *Hosp. a Domic.* 2019, 3, 149, doi:10.22585/hospdomic.v3i2.60.
71. Gibson, P.R.; Skodje, G.I.; Lundin, K.E.A. Non-coeliac gluten sensitivity. *J. Gastroenterol. Hepatol.* 2017, 32 Suppl 1, 86–89, doi:10.1111/jgh.13705.
72. Guido Schwarzer, James R. Carpenter and Gerta Rücker (2020). *metasens: Advanced Statistical Methods to Model and Adjust for Bias in Meta-Analysis*. R package version 0.5-0. <https://CRAN.R-project.org/package=metasens>;
73. Heaton, K.W.; Thornton, J.R.; Emmett, P.M. Treatment of Crohn's disease with an unrefined-carbohydrate, fibre-rich diet. *Br. Med. J.* 1979, 2, 764–766, doi:10.1136/bmj.2.6193.764.
74. Lomer, M.C.; Harvey, R.S.; Evans, S.M.; Thompson, R.P.; Powell, J.J. Efficacy and tolerability of a low microparticle diet in a double blind, randomized, pilot study in Crohn's disease. *Eur. J. Gastroenterol. Hepatol.* 2001, 13, 101–106, doi:10.1097/00042737-200102000-00003.
75. Uzunışmail, H.; Cengiz, M.; Uzun, H.; Ozbakir, F.; Göksel, S.; Demirdağ, F.; Can, G.; Balci, H. The effects of provocation by foods with raised IgG antibodies and additives on the course of Crohn's disease: a pilot study. *Turkish J. Gastroenterol. Off. J. Turkish Soc. Gastroenterol.* 2012, 23, 19–27, doi:10.4318/tjg.2012.0332.
76. Gunasekera, V.; Mendall, M.A.; Chan, D.; Kumar, D. Treatment of Crohn's Disease with an IgG4-Guided Exclusion Diet: A Randomized Controlled Trial. *Dig. Dis. Sci.* 2016, 61, 1148–1157, doi:10.1007/s10620-015-3987-z.
77. Jian, L.; Anqi, H.; Gang, L.; Litian, W.; Yanyan, X.; Mengdi, W.; Tong, L. Food Exclusion Based on IgG Antibodies Alleviates Symptoms in Ulcerative Colitis: A Prospective Study. *Inflamm. Bowel Dis.* 2018, 24, 1918–1925, doi:10.1093/ibd/izy110.
78. Testa, A.; Imperatore, N.; Rispo, A.; Rea, M.; Tortora, R.; Nardone, O.M.; Lucci, L.; Accarino, G.; Caporaso, N.; Castiglione, F. Beyond Irritable Bowel Syndrome: The Efficacy of the Low Fodmap Diet for Improving Symptoms in Inflammatory Bowel Diseases and Celiac Disease. *Dig. Dis.* 2018, 36, 271–280, doi:10.1159/000489487.

Retrieved from <https://encyclopedia.pub/entry/history/show/16630>