

Ethylene

Subjects: Biology

Contributor: Monika Keisham

Salinity stress is one of the major threats to agricultural productivity across the globe. Research in the past three decades, therefore, has focused on analyzing the effects of salinity stress on the plants. Evidence gathered over the years supports the role of ethylene as a key regulator of salinity stress tolerance in plants. This gaseous plant hormone regulates many vital cellular processes starting from seed germination to photosynthesis for maintaining the plants' growth and yield under salinity stress. Ethylene modulates salinity stress responses largely via maintaining the homeostasis of Na^+/K^+ , nutrients, and reactive oxygen species (ROS) by inducing antioxidant defense in addition to elevating the assimilation of nitrates and sulfates. Moreover, a cross-talk of ethylene signaling with other phytohormones has also been observed, which collectively regulate the salinity stress responses in plants.

Keywords: ROS ; ethylene ; antioxidants ; salinity stress ; photosynthesis ; programmed cell death ; seed germination ; hormone cross-talk

1. Introduction

Ethylene, the first gaseous plant hormone to be identified, is a key regulator of plant growth and development. Although growth-regulating effects of ethylene were first observed in 1901 by a Russian physiologist Dimitry K. Neljubov, ethylene was established as a plant hormone almost 60 years later in 1965 ^[1]. The biosynthetic pathway of ethylene production was elucidated in 1970–80, while its signaling components were identified in the 1990s after the development of *Arabidopsis thaliana* as a model system that facilitated the development and screening of genetic mutants ^[2]. Subsequent studies on ethylene led to the identification of an array of genes and transcriptional factors working downstream that widen our understanding of how this colorless and odorless plant hormone functions ^[3]. Because of the gaseous nature of ethylene, it can easily diffuse to nearby cells and, therefore, ethylene production predominantly takes place locally at the site of its action. Earlier work showed that ethylene biosynthesis is dramatically induced during fruit ripening and leaf senescence, among others ^[4]. Ethylene is also well known for its triple response which includes (i) inhibition of hypocotyl and root elongation, (ii) swelling of the hypocotyl, and (iii) exaggerated tightening of the apical hook ^[5].

In addition to regulating plant growth and development, research in the past two decades has also highlighted the involvement of ethylene in regulating plant responses to various biotic and abiotic stresses ^{[6][7][8][9]}. Among different abiotic stresses, ethylene has emerged as one of the important positive mediators for salinity stress tolerance in the model plant *A. thaliana* as well as in many crop plants including grapevines, maize, and tomato ^{[10][11][12][13]}. Salinity stress is one of the major abiotic stresses, posing a major threat to agricultural productivity ^{[14][15]}. Globally, more than 20% of irrigated land is affected by salinity stress, resulting in an average yield fall of more than 50% for major crops ^{[16][17]}. As per the Food and Agriculture Organization (FAO), 13,003 million hectares (Mha) of the global land area is devoted to agriculture ^[18] in which soil salinity has adversely affected about 30% of the irrigated land and 6% of the total land area ^{[19][20]}, resulting in a monetary loss of approximately 27.3 billion USD per year ^[21]. Plants growing in geographically over-salted soils in areas where hydraulic lifting of saline underground water occurs and in coastal areas are frequently exposed to salinity stress ^[22]. The decline of farmable land due to salinity is a major concern for food security in the current and future scenario of a growing population which has been estimated to reach 8.5 billion over the next 25 years ^[23]. Salinity stress is exerted because of excess of one or more salt ions in the soil including sodium (Na^+), bicarbonate (HCO_3^-), magnesium (Mg^{2+}), sulfate (SO_4^{2-}), potassium (K^+), chloride (Cl^-), calcium (Ca^{2+}), and carbonate (CO_3^{2-}) ^[24]. Salinity stress negatively influences seed germination, growth, physiology, productivity, and reproduction and sometimes even results in death under severe conditions ^[25]. At the onset of salinity stress, the capacity for water absorption by roots decreases and the transpiration rate increases due to osmotic imbalance, thereby generating hyperosmotic stress ^{[26][27]}. Osmotic stress, in turn, induces closure of stomata, which restricts CO_2 uptake, resulting in reduced carbon fixation and assimilation in leaf tissues ^{[28][29]}. Rates of photosynthesis and carbohydrate production are therefore reduced, which impacts plant growth and yield. Other consequences of stomatal closure include reduced evapotranspiration of water and

accumulation of reactive oxygen species (ROS) [30]. Salinity stress-induced ROS accumulation can lead to uncontrolled oxidation of membranes, proteins, and DNA, ultimately resulting in cell death [31]. In order to maintain ROS homeostasis [32], plants are equipped with a powerful and multifaceted antioxidant system, consisting of enzymatic and nonenzymatic components. While enzymatic components include superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), glutathione reductase (GR), guaiacol peroxidase (POD), glutathione transferase (GST), glutathione peroxidase-like (GPXLs), and thioredoxin peroxidase (TPOX), non-enzymatic antioxidants consist of ascorbic acid (AsA), glutathione (GSH), α -tocopherol (TOC), flavonoids, and carotenoids [33][34]. Successful detoxification of the stress-induced ROS is one of the crucial factors in salinity stress adaptation, and ethylene seems to play a pivotal role in ROS detoxification, thereby providing adaptation to salinity stress [35][36].

2. Ethylene is a Key Modulator of Salinity Stress Responses in Plants

Ethylene biosynthesis and signaling are implicated in salinity stress tolerance in plants (Table S1). Overproduction of endogenous ethylene or exogenous treatment of ethylene-releasing compounds such as ethephon or ethylene precursors such as 1-aminocyclopropane-1-carboxylic acid (ACC) increase salinity stress tolerance in various plants including *Arabidopsis* [37] and maize [12]. Moreover, ethylene has also been found as an essential positive mediator of salinity stress tolerance in grapevine [13], maize [12], and tomato [38]. Promising evidence of the involvement of melatonin in enhancing salinity stress tolerance by promoting *MYB108A*-mediated ethylene biosynthesis was also reported in grapevines [13]. It has been reported that inhibition of ethylene biosynthesis and/or signaling leads to increased sensitivity of plants to salinity stress. Gharbi et al. [38] hypothesize that ethylene exhibits a positive effect in adaptation to salinity stress probably by maintaining stomatal conductance, water use efficiency, and osmotic adjustment in *Solanum chilense* [38]. Calcium carbide (CaC_2), a precursor of acetylene exhibiting similar effects to ethylene, is commonly used to improve seed germination rates and ethylene concentration in germinating seeds under salinity stress. It is proposed that CaC_2 is involved in osmotic adjustments and management of oxidative stress by increasing solute concentrations and activities of antioxidant enzymes in germinating seeds. CaC_2 considerably enhances the activity of SOD and CAT besides reducing the H_2O_2 and malondialdehyde (MDA) concentrations for improving seed germination in *Cucumis sativus* under salinity stress conditions [39]. These reports collectively suggest a positive regulatory role of ethylene in salt stress tolerance in plants; however, negative regulation of ethylene in salinity stress response has also been reported in many plants, including rice. Transgenic plants with reduced ethylene biosynthesis showed elevated salinity tolerance in tobacco [40]. Similarly, exogenous treatment with ethylene in rice resulted in salinity hypersensitivity [41][42]. Elevation of ethylene production under salinity stress significantly reduced growth, grain filling, and development of spikelets in rice [43]. Exogenous application of 1-MCP, an ethylene action inhibitor, to the rice spikelets resulted in improved physiological, agronomical, and biochemical characteristics under salinity stress, further suggesting a negative role of ethylene in salinity stress tolerance in rice [44].

Gene mutation and transgenic analyses have shown that almost all the components of ethylene biosynthesis and signal transduction pathway respond to salinity stress either positively or negatively. In cotton, short- and long-term salinity stress resulted in upregulation of different sets of genes of ethylene signaling involved in the regulation of salinity stress [45]. These genes include (a) ethylene biosynthesis genes (homologs of *ACS1*, *ACS12*, *ACO1*, and *ACO3*), (b) ethylene receptor genes (*ETR1*, *ETR2*, and *EIN4*), (c) ethylene signaling pathway (*ERF1*, *ERF2*, *EIN3*, and *MEKK1-MKK2-MPK4/6* kinases), and (d) feedback mechanism gene (*CTR1*) [45].

2.1. Salinity Stress and Ethylene Receptors

Ethylene receptors are negative regulators of ethylene signaling, and interestingly, an inhibition of ethylene receptors has been observed during salinity stress in several plant species. In *Arabidopsis*, salinity stress has been shown to suppress *ETR1* expression [46]. In addition, *etr* loss-of-function mutants showed enhanced tolerance while *etr-1* gain-of-function mutants showed increased sensitivity to salinity stress in *Arabidopsis* [45][47][48]. Wilson et al. [49] reported that *ETR1* and *ETR2* function differently in *Arabidopsis* during seed germination under salinity stress. Loss-of-function mutants of *etr1* germinated earlier than the wild type (WT), while that of *etr2* germinated after WT [49]. Moreover, it has also been shown that *ETR1* and *ETR2* regulate abscisic acid (ABA) signaling independently of ethylene signaling and lead to contrasting germination during salinity stress [49]. In tobacco, salinity stress increases *NTHK1* mRNA levels dramatically [47], suggesting a negative regulation of ethylene in salinity stress tolerance. However, overexpression of *NTHK1* in tobacco resulted in early inductions of the ACC oxidase (*NtACO3*) and ERF (*NtERF1* and *NtERF4*) genes during salinity stress [50]. In contrast, the expression level of a salinity-inducible ACC synthase gene (*NtACS1*) was greatly suppressed in the overexpression lines. Further overexpression of *NTHK1* in *Arabidopsis* resulted in enhanced sensitivity to salinity stress but reduced sensitivity to ethylene [50][51]. Recently, it was reported that *NTHK1* interacts with an ankyrin domain-containing protein NEIP2 (*NTHK1* ethylene receptor-interacting protein 2) to improve the salinity and oxidative stress

tolerance in tobacco [52]. Overexpression of *NTHK1* resulted in the accumulation of NEIP2 in the presence of both ethylene and salinity stress. On the other hand, overexpression of *NEIP2* inhibited ethylene responses, similar to the functions of *NTHK1* [52]. All of these results suggest a negative regulation of ethylene receptors in salinity stress tolerance, indicating ethylene as a positive mediator of salinity stress tolerance in plants.

In addition to the ethylene receptors, *CTR1* is regulated by salinity stress. It has been reported that *ctr1* loss-of-function mutants showed enhanced tolerance to salinity stress, possibly by modulation of shoot Na^+/K^+ ratio, which is dependent on *ETR1*-*CTR1*-regulated signaling [45][53]. Moreover, high survival rates of *CTR1* mutant *ctr1-1* was observed under salinity and osmotic stress conditions in comparison to loss-of-function mutants *ein2* and *ein3-1 eil1-1* (double mutations of *EIN3* and *EIN3*-Like1 (*EIL1*)) which showed remarkably reduced tolerance to salinity [48][54][55]. Ge et al. [56] showed that the heterotrimeric G-protein $\text{G}\alpha$ subunit *GPA1* is involved in ethylene-induced stomatal closure via NADPH oxidase-dependent H_2O_2 synthesis. Interestingly, *GPA1* also functions downstream of *RAN1*, *ETR1*, *ERS1*, *EIN4*, and *CTR1* and upstream of *EIN2*, *EIN3*, and *ARR2*. In guard cells of *Arabidopsis* leaves, *ETR1* and *ERS1* mediate both ethylene and H_2O_2 signaling, highlighting ethylene-mediated regulation of H_2O_2 concentrations during salinity stress [56].

2.2. Salinity Stress and EIN Proteins

EIN proteins are the transcription factors that function downstream of *CTR1* in the ethylene signaling cascade. It has been observed that salinity stress-induced stabilization of *EIN3/EIL1* promotes tolerance to salinity stress by averting ROS accumulation [45]. In *Arabidopsis*, loss-of-function of *ein2* resulted in enhanced sensitivity to salinity stress while its overexpression lines showed reduced sensitivity [45][55]. Lei et al. [55] identified a MA3 domain-containing protein *EIN2* C-Terminus Interacting Protein 1 (*ECIP1*), which interacts with *EIN2*. *ecip1* loss-of-function mutants confer sensitivity to salinity stress during seed germination in *Arabidopsis*. However, these mutants show insignificant changes in ethylene responses and salinity stress tolerance. In *Arabidopsis*, several studies based on gene mutations showed that the ethylene signal from *EIN2* to the nucleus is transduced by *EIN3/EILs*. Overexpression of *EIN3* remarkably enhanced tolerance to salinity stress. Interestingly, both loss-of-function mutants *ein3-1* and double mutant *ein3eil1* showed severe sensitivity to salinity stress [45][54][55]. Mao huzi (*MHZ7*) and *MHZ6* of rice are homologs of *Arabidopsis* *EIN2* and *EIN3* [57]. *OsEIL2*, *OsEIN2* (*MHZ7*), and *OsEIL1* (*MHZ6*) were found to play a negative role in salinity stress tolerance in rice, unlike *EIN2* and *EIN3*, which play positive roles in *Arabidopsis* [42]. Knockout mutants of *OsEIL2*, *MHZ7/OsEIN2*, or *MHZ6/OsEIL1* enhanced salinity stress tolerance, while their overexpression mutants exhibited increased sensitivity to salinity stress in rice [42]. Downstream of *EIN3/EIL1* is *ESE1*, which is positively regulated by ethylene signaling. *ESE1* enhances plant tolerance to salinity stress by binding to promoters of salinity stress-responsive genes such as *RD29A* and *COR15A* [58]. Quan et al. [59] proposed that *EIN3*-*SOS2* modulate salinity stress tolerance possibly by linking the ethylene signaling and salinity overly sensitive (*SOS*) pathways. However, further characterization showed that both *EIN2* and *EIN3* failed to change the expression of *SOS* genes in *Arabidopsis*. Interestingly, *SOS2* phosphorylates *EIN3* and activates salinity-inducible *ESE1* [59]. In rice, *OsDOF15* was identified as a gene involved in the coordination between salinity and ethylene biosynthesis in rice to inhibit primary root development by affecting cell proliferation in the root apical meristem [60]. In addition, the expression levels of *OsEIL2* were found to be upregulated during salinity stress in WT plants; however, *OsEIL2* overexpression plants showed growth retardation with shortened roots and shoots than control plants [61]. In addition, *OsEIL2* overexpression plants showed increased ethylene sensitivity and accelerated leaf senescence [61]. Further, it has been observed that *OsEIL2* negatively regulates the expression of *BURP* genes *OsBURP14* and *OsBURP16* to reduce the pectin content during salinity stress in rice [61]. In the case of mulberry, increased expression of *MnEIL3* was observed during salinity stress both in the root and shoot [62]. Transgenic plants overexpressing *MnEIL3* showed an upregulation of ethylene biosynthetic genes in *Arabidopsis* to enhance salinity tolerance. Moreover, *MnEIL3* may enhance the activities of *MnACO1* and *MnACS1* promoters, indicating functioning of an ethylene–*EIN3/EILs*–1-aminocyclopropane-1-carboxylic acid (*ACC*) oxidase (*ACO*)/ *ACC* synthase (*ACS*) regulatory loop under salinity stress [62].

Cortical microtubule reorganization is crucial for survival under salinity stress [63][64], and ethylene has been found to modulate the same in roots cells and etiolated hypocotyls [65][66][67][68]. In *Arabidopsis*, microtubule-stabilizing protein Wave-Dampened2-Like5 (*WDL5*) takes part in ethylene signaling to inhibit etiolated hypocotyl elongation [67][69]. Dou et al. [69] found that ethylene signaling has a positive role in the regulation of microtubule reassembly and that *WDL5* functions as a downstream effector of signaling involved in ethylene-mediated microtubule reassembly in salinity stress. Knockout of *WDL5* partly suppressed ethylene-induced microtubule reassembly, whereas its upregulation partially protects from the salinity stress. Microtubule reassembly under salinity stress is insensitive to the effect of *ACC* in *ein3eil1* cells; therefore, transcriptional regulation by *EIN3/EIL1* is essential for ethylene signaling-mediated reassembly of microtubules in response to salinity stress [69].

2.3. Effects of Salinity Stress on ERFs and other Ethylene-Responsive Transcription Factors

Ethylene-responsive element binding factors (ERFs) are transcription factors functioning downstream of EIN3 in ethylene signaling. Cheng et al. [70] reported that overexpression of *ERF1* enhances tolerance of plants to salinity, drought, and high-temperature stress conditions in an ethylene-independent manner. Moreover, enhanced expressions of three *ERF* genes including *ESE1*, *ESE2*, and *ESE3* were observed in response to salinity stress and ethylene in *Arabidopsis* [58]. Recently, it was shown that *ERF1/2* was upregulated whereas *CTR1* and EIN3-binding F-box protein 1/2 (*EBF1/2*) were downregulated during salinity stress in *Cynanchum auriculatum* [14]. Li et al. [71] cloned an APETALA2/ethylene responsive factor (*AP2/ERF*) gene from salinity-tolerant sweet potato line ND98 and named it *IbRAP2-12*. Based on the transient expression in tobacco epidermal cells and transcriptional activation analysis, the protein of *IbRAP2-12* was found to be localized in nucleus and it was observed that *IbRAP2-12* exhibits transcriptional activation because of a domain located at the C-terminus of the protein. As compared to the WT, *IbRAP2-12*-overexpression lines showed higher accumulation of proline and lower concentration of H_2O_2 under salinity stress in *Arabidopsis*. Moreover, multiple genes involved in the ROS-detoxification including *SAPX*, *GPX7*, and *CAT5* were found to be upregulated in the transgenic plants under salinity stress [71].

Similarly, overexpression of an ethylene-responsive transcription factor (*TdSHN1*) from durum wheat resulted in the development of a cuticle and lower stomatal density than WT. *TdSHN1* overexpression lines, therefore, showed enhanced tolerance to salinity stress because of a reduced water loss from the leaf lamina [72]. On exposure to salinity stress, *MdERF4* from apple is induced to decrease the salinity stress tolerance by binding and inhibiting the expression of *MdERF3*, suggesting that the *MdERF4*–*MdERF3* interaction may be a feedback regulation mechanism in salinity stress to maintain ethylene homeostasis in plants [73]. Similarly, the expression of the *ERF38* gene from poplar 84K (*Populus alba* × *Populus glandulosa*) was induced under salinity stress. Overexpression of *ERF38* in poplar reduced membrane lipid peroxidation, decreased ROS accumulation, accumulated proline and soluble proteins, and exhibited higher POD and SOD activities in transgenic plants than WT, suggesting a multifaceted role of *ERF38* in improving salinity and osmotic stress tolerance in poplar [74]. Acireductone dioxygenase (*ARD*) is an active metal-binding metalloenzyme and is involved in the formation of 2-keto-4-methylthiobutyrate (*KMTB*) to further produce methionine (*Met*) on the methionine salvage pathway, which is an initial substrate in ethylene synthesis pathway [75][76][77]. Overexpression of *OsARD1* increases the water-holding capacity and relative water content in the leaves of *OsARD1* overexpression lines to reduce the sensitivity to salinity and osmotic stresses at germination stage [78]. It was speculated that the drop in sensitivity to salinity and osmotic stresses results in an increased ethylene concentration in *OsARD1* overexpression plants [78].

3. Seed Germination Regulated by Ethylene under Salinity Stress

Successful seed germination is the most crucial phase in the initiation of the life cycle of plants and is regulated by many external and internal factors including phytohormones, light, temperature, drought, and salinity [79][80][81][82]. Seed germination is severely affected in saline soil, which negatively influences plant growth and crop yield [10] (Figure 1). Different components of ethylene signaling participate either positively or negatively during seed germination and seedling growth under salinity stress [83]. For example, seed germination in *Arabidopsis* was inhibited by *ETR1* and *EIN4*, whereas *ETR2* was found to be a positive regulator involved in stimulating seed germination during salinity stress conditions [49].




Figure 1. Role of ethylene signaling in seed germination under salinity stress. Ethylene induced by salinity stress activates the signaling pathway by inhibiting the active receptors and by releasing the Constitutive Triple Response1 (CTR1). Salinity-induced ethylene signal is transduced mainly through the classical receptors–CTR1–Ethylene-Insensitive2 (EIN2)–EIN3 pathway to regulate many effectors involved in plant growth and salinity response. EIN3/EIN3-Like (EIL) promotes the entry of Constitutive Photomorphogenesis1 (COP1) into the nucleus and degrades the Elongated Hypocotyl5 (HY5) protein, which inhibits seed germination by upregulating *ABI5* gene expression. Degradation of HY5 inhibits the expression of *ABI5* and ultimately induced seed germination under salinity stress. On the other hand, EIN3/EIL also directly induces seed germination by scavenging reactive oxygen species (ROS) through upregulating peroxidase activities.

Ethylene antagonistically modulates seed germination in *Arabidopsis* under salinity stress via the Constitutive Photomorphogenesis1 (COP1)-mediated downregulation of Elongated Hypocotyl5 (*HY5*) and ABA Insensitive5 (*ABI5*) in the nucleus [84] (Figure 1). Salinity stress inhibits seed germination through elevation of the H_2O_2 and exogenous treatment of ethylene precursor (ACC) has been shown to regulate the ROS homeostasis to induce the seed germination [85]. Interaction between ethylene and nitric oxide has also been shown to regulate seed germination by decreasing the H_2O_2 level induced by salinity stress, further suggesting that ethylene promotes the seed germination rate by modulation of ROS production in salinity stress. On the other hand, ethylene can also inhibit the seed germination induced by salinity stress in many plant species. For example, Chang et al. [86] reported that ethylene is involved in the suppression of seed germination in cucumber (*Cucumis sativus* L.) and that l-Glu interacts with ethylene in the regulation of seed germination under salinity stress. Ethylene produced during salinity stress helps to maintain the Na^+/K^+ homeostasis to provide salinity stress tolerance [53].

Overexpression of the *Malus hupehensis* SHINE clade protein (*MhSHN1*) gene, a member of the AP2/ERF transcription factor, does not regulate the seed germination in transgenic tobacco plants yet it enhances salinity and osmotic stress tolerance during seed germination [87]. In *Stylosanthes humilis*, a forage legume naturally growing in the saline soils, ethylene production in the seeds provides tolerance to salinity stress [88][89][90]. Salinity induced production of ABA, and ethylene forms a point of union between the two and enables the regulation of energy metabolism and embryo growth in *S. humilis* seeds within a given pH condition [90]. Ectopic expression of a zinc finger transcription factor *Gossypium hirsutum* plant AT-rich sequence and zinc-binding (*GhPLATZ1*) in *Arabidopsis* regulated seed germination and seedling establishment under salinity and mannitol stress conditions. Further experimentations revealed that the inhibition of *ABI4* and *ETO1* expressions suppressed ACS gene expression to alter the ABA, GA, and ethylene pathways in transgenic lines [91]. Ectopic expression of a homolog of *AtERF38* (*GhERF38* from *G. hirsutum*) in *Arabidopsis* resulted in ABA sensitivity in transgenic lines; therefore, reduced seed germination under salinity and drought stress was observed in the transgenic plants as compared to WT [92]. Plant growth-promoting *Pseudomonas fluorescens* strains improve salinity tolerance of plants due to its ability to produce ACC deaminase and, consequently, to stimulate seed germination in wheat under salinity stress [93]. With the higher activity of ACC deaminase, the *Enterobacter cloacae* HSNJ4 strain could effectively promote seed germination and could provide the salinity tolerance by degrading ACC, thus inhibiting ethylene

synthesis [94]. Moreover, seeds of the transgenic line overexpressing ethylene response factors (ERF95 and ERF96) showed better germination and seedling establishment as compared to the WT during salinity stress conditions [95]. A novel ethylene-responsive transcription factor from *Lycium chinense* *LchERF* provides salinity tolerance to transgenic tobacco during seed germination and vegetative growth [96]. Notwithstanding, Chang et al. [86] reported that salinity interferes with the ethylene signaling pathway and decreases ethylene production in seeds of *C. sativus*, which was associated with the inhibition of germination. In Faba beans (*Vicia faba*), seed germination in salinity-tolerant Y134 is not inhibited during salinity stress as compared to the salinity-sensitive Y078 probably because of the downregulation of genes related to ABA and ethylene signaling pathways and upregulation of late embryogenesis abundant (LEA) genes [97]. Seeds of *Capsicum annuum* primed with SA showed higher a germination rate due to suppression of the ethylene level as well as elevation of total soluble sugar contents and SOD activity [98].

4. Fine-Tuning of Photosynthetic Machinery by Ethylene during Salinity Stress

Homeostasis of essential elements like N, P, K, S, and Ca is altered during salinity stress, which in turn affects the photosynthetic efficiency of plants [99][100][101]. Salinity stress induces oxidative stress through increased production of ROS, which can disrupt chloroplast functions (Figure 2). Salts at higher concentrations induce both osmotic and ionic stresses, which affect photosynthetic activity either by closing the stomata or by reducing the activity of CO₂-fixing enzymes and availability of water in the plant cells [102] (Figure 2). Activities of CO₂-fixing enzymes are reduced at higher concentrations of Na⁺, and tolerance of these enzymes to Na⁺ concentrations varies from species to species [103]. Na⁺ ions imbalance the proton motive force and thus influence photosynthetic machinery and chloroplastic functions [104]. Salinity stress influences the photosynthetic parameters including chlorophyll, photosystems, net photosynthesis rate (Pn), chlorophyll fluorescence parameters, soluble sugar contents, and ribulose biphosphate carboxylase/oxygenase (RuBisCO) activity [105]. Recently, it was reported that tomato plants showed improved photosynthesis, metabolic homeostasis, and growth rate as a result of elevated CO₂ under salinity stress by decreasing the amount of ABA hormone and ACC [106]. Among all the photosynthetic parameters, photosystem II (PSII) is the most susceptible to various abiotic stresses including salinity [107][108]. Homeostasis of Na⁺ ions maintain membrane integrity, relative water content, net photosynthesis, and yield. Ethylene has been shown to promote the homeostasis of Na⁺/K⁺, nutrients, and ROS to enhance plant tolerance to salinity [57]. The *ctr1-1* mutants maintain relatively higher concentrations of K⁺ and lower concentrations of Na⁺ in contrast to *ein2-5* or *ein3* plants, where an opposite trend of K⁺ and Na⁺ concentration was observed compared with the WT undertreated and optimum conditions [53]. Because of this altered K⁺ and Na⁺ homeostasis, *ctr1-1* plants displayed a slight reduction in leaf area and root elongation, while *ein2-5* or *ein3-1* mutants showed magnified retardation in plant growth compared to the WT under salinity stress [53]. In pomegranate, salinity decreased the net photosynthetic rate, chlorophyll content, stomatal conductance, relative water content, and electrical conductivity [105][109][110]. Further heat map analysis showed that antiapoptotic genes *BAG6* and *BAG7* were clustered together with *ERS2*, *EIN3*, and *ACS* and that the transcripts levels of *BAG6*, *BAG7*, *ERS2*, and *ACS2* were significantly suppressed in the response to salinity. The inclusion of ACC or ethylene source in the saline solution restored the expression levels of *BAG6* and *BAG7*, suggesting the involvement of ethylene in the regulation of these antiapoptotic genes under salinity stress [111].




Figure 2. Functions of ethylene in the regulation of photosynthesis under salinity stress. In the absence of ethylene, salinity stress results in an imbalance of Na⁺/K⁺ homeostasis, which leads to the production of ROS. This salinity-induced ROS production, in turn, exerts oxidative stress on plants, resulting in stomatal closure and reduced activity of CO₂-fixing enzymes, resulting in a decrease in photosynthesis. In the presence of ethylene, Na⁺/K⁺ homeostasis and nutrients homeostasis are maintained, and the antioxidant defense mechanism is activated, which limits ROS production, thereby

preventing ROS-induced oxidative stress. In the absence of oxidative stress, the rate of photosynthesis is maintained even during salinity stress.

In *Arabidopsis* and alfalfa, *Enterobacter* sp. SA187 mediates salinity tolerance by producing 2-keto-4-methylthiobutyric acid (KMBA), which is converted into ethylene *in planta* ^{[112][113]}. This *Enterobacter*-produced KMBA is involved in the maintenance of photosynthesis and primary metabolism together with the reduction of ABA-mediated stress responses in plants. Gene expression analysis revealed that, after SA187 inoculation, genes related to photosynthesis and primary metabolism remain unaltered under salinity stress conditions as compared to the mock plants ^{[113][114]}. Similarly, in rice, inoculation of *Glutamicibacter* sp. YD01 facilitated rice plants to combat stress by ethylene-mediated regulation of ROS accumulation, ion homeostasis, and photosynthetic capacity and by enhancing stress-responsive gene expression ^[115]. Tight regulation of ROS homeostasis also accelerates photosynthesis and growth by abating lipid peroxidation in chloroplasts ^{[116][117][118]}.

It is well established that salinity stress also affects nitrogen and sulfur assimilation in plants. Plants grown on low nitrate (5 mM) showed lower photosynthesis and growth compared to the plants grown in sufficient nitrate concentrations during non-saline conditions ^[119]. When excess nitrate (20 mM) was applied under non-saline conditions, an inhibitory effect on photosynthesis was observed, which was related to higher ethylene production. However, under salinity stress conditions, as the demand for N increased, the excess N optimized ethylene, enhanced proline production, and promoted photosynthesis and growth ^[119]. Recently, it was reported that cadmium and sodium stress conditions induce ethylene and Jasmonic acid (JA) signaling. Both of these signaling pathways converged at *EIN3/EIL1* and resulted in an enhanced expression of a nitrate transporter *NRT1.8* and reduced expression of *NRT1.5*. Although it resulted in decreased plant growth, it promoted plant tolerance to stress in a nitrate reductase-dependent manner by mediating the stress-initiated nitrate allocation to roots, which decoupled nitrate assimilation and photosynthesis ^[120]. Taken together, these studies clearly highlight that ethylene plays a major role in stabilizing photosynthesis under salinity stress conditions by maintaining the ROS accumulation, ion homeostasis, and mineral homeostasis and by elevating the antioxidant defense mechanism.

References

1. Neljubow, D. Über die horizontale Nutation der Stengel von *Pisum Sativum* und einiger anderen Pflanzen. Bot. Cent. Beih. 1901, 10, 128–139.
2. Hua, J. Isolation of Components Involved in Ethylene Signaling. In *Ethylene in Plants*; Springer: Dordrecht, The Netherlands, 2015; pp. 27–44. ISBN 9401794839.
3. Abeles, F.B.; Morgan, P.W.; Salinityveit, M.E., Jr. *Ethylene in Plant Biology*; Academic Press: San Diego, CA, USA, 2012; ISBN 0080916287.
4. Burg, S.P.; Burg, E.A. Molecular requirements for the biological activity of ethylene. *Plant Physiol.* 1967, 42, 144–152.
5. Guzman, P.; Ecker, J.R. Exploiting the triple response of *Arabidopsis* to identify ethylene-related mutants. *Plant Cell* 1990, 2, 513–523.
6. Gómez-Cadenas, A.; Tadeo, F.R.; Primo-Millo, E.; Talón, M. Involvement of abscisic acid and ethylene in the responses of citrus seedlings to salinity shock. *Physiol. Plant.* 1998, 103, 475–484.
7. Arraes, F.B.M.; Beneventi, M.A.; de Sa, M.E.L.; Paixao, J.F.R.; Albuquerque, E.V.S.; Marin, S.R.R.; Purgatto, E.; Nepomuceno, A.L.; Grossi-de-Sa, M.F. Implications of ethylene biosynthesis and signaling in soybean drought stress tolerance. *BMC Plant Biol.* 2015, 15, 213.
8. Gupta, R.; Min, C.W.; Kim, S.W.; Yoo, J.S.; Moon, A.R.; Shin, A.Y.; Kwon, S.Y.; Kim, S.T. A TMT-Based Quantitative Proteome Analysis to Elucidate the TSWV Induced Signaling Cascade in Susceptible and Resistant Cultivars of *Solanum lycopersicum*. *Plants* 2020, 9, 290.
9. Gupta, R.; Min, C.W.; Kim, Y.J.; Kim, S.T. Identification of Msp1-Induced Signaling Components in Rice Leaves by Integrated Proteomic and Phosphoproteomic Analysis. *Int. J. Mol. Sci.* 2019, 20, 4135.
10. Munns, R.; Tester, M. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* 2008, 59, 651–681.
11. Siddikee, M.A.; Chauhan, P.S.; Sa, T. Regulation of ethylene biosynthesis under salinity stress in red pepper (*Capsicum annuum* L.) by 1-aminocyclopropane-1-carboxylic acid (ACC) deaminase-producing halotolerant bacteria. *J. Plant Growth Regul.* 2012, 31, 265–272.
12. Freitas, V.S.; de Souza Miranda, R.; Costa, J.H.; de Oliveira, D.F.; de Oliveira Paula, S.; de Castro Miguel, E.; Freire, R.S.; Prisco, J.T.; Gomes-Filho, E. Ethylene triggers salinity tolerance in maize genotypes by modulating polyamine

catabolism enzymes associated with H₂O₂ production. *Environ. Exp. Bot.* 2018, 145, 75–86.

13. Xu, L.; Xiang, G.; Sun, Q.; Ni, Y.; Jin, Z.; Gao, Z.; Yao, Y. Melatonin enhances salinity tolerance by promoting MYB108A-mediated ethylene biosynthesis in grapevines. *Hortic. Res.* 2019, 6, 114.
14. Zhang, M.; Hong, L.Z.; Gu, M.F.; Wu, C.D.; Zhang, G. Transcriptome analyses revealed molecular responses of *Cynanchum auriculatum* leaves to saline stress. *Sci. Rep.* 2020, 10, 449.
15. Kim, S.W.; Min, C.W.; Gupta, R.; Jo, I.H.; Bang, K.H.; Kim, Y.C.; Kim, K.H.; Kim, S.T. Proteomics analysis of early salinity-responsive proteins in ginseng (*Panax ginseng* CA Meyer) leaves. *Korean J. Med. Crop. Sci.* 2014, 22, 398–404.
16. Bray, E.A.; Bailey-Serres, J.; Weretilnyk, E. Responses to abiotic stress. In *Biochemistry and Molecular Biology of Plants*; Buchanan, B., Gruissem, W., Jones, R., Eds.; American Society of Plant Physiology: Rockville, MD, USA, 2000; pp. 1158–1203.
17. Pitman, M.G.; Läuchli, A. Global Impact of Salinity and Agricultural Ecosystems. In *Salinity: Environment-Plants-Molecules*; Springer: Dordrecht, The Netherlands, 2002; pp. 3–20.
18. Polle, A.; Chen, S. On the salinity side of life: Molecular, physiological and anatomical adaptation and acclimation of trees to extreme habitats. *Plant Cell Environ.* 2015, 38, 1794–1816.
19. Ghatak, A.; Chaturvedi, P.; Paul, P.; Agrawal, G.K.; Rakwal, R.; Kim, S.T.; Weckwerth, W.; Gupta, R. Proteomics survey of Solanaceae family: Current status and challenges ahead. *J. Proteom.* 2017, 169, 41–57.
20. Chaves, M.M.; Flexas, J.; Pinheiro, C. Photosynthesis under drought and salinity stress: Regulation mechanisms from whole plant to cell. *Ann. Bot.* 2009, 103, 551–560.
21. Qadir, M.; Quillérou, E.; Nangia, V.; Murtaza, G.; Singh, M.; Thomas, R.J.; Drechsel, P.; Noble, A.D. Economics of salinity-induced land degradation and restoration. *Nat. Resour. Forum* 2014, 38, 282–295.
22. Neumann, P.M. Inhibition of root growth by salinity stress: Toxicity or an adaptive biophysical response? In *Structure and Function of Roots*; Springer: Dordrecht, The Netherlands, 1995; pp. 299–304.
23. Rasool, S.; Hameed, A.; Azooz, M.M.; Siddiqi, T.O.; Ahmad, P. Salinity stress: Causes, types and responses of plants. In *Ecophysiology and Responses of Plants under Salinity Stress*; Springer: New York, NY, USA, 2013; pp. 1–24.
24. Muhammad, N.; Samina, B.; Yasmin, K.; Roqayya, M.; Zabta, K.S.; Abdul, L.K.; Ajmal, K.; Ahmed, A. Plant growth promoting bacteria as an alternative strategy for salinity tolerance in plants: A review. *Microbiol. Res.* 2018, 209, 21–32.
25. Mahajan, S.; Tuteja, N. Cold, salinity and drought stresses: An overview. *Arch. Biochem. Biophys.* 2005, 444, 139–158.
26. Munns, R. Genes and salinity tolerance: Bringing them together. *New Phytol.* 2005, 167, 645–663.
27. Yang, Y.; Guo, Y. Elucidating the molecular mechanisms mediating plant salinity-stress responses. *New Phytol.* 2018, 217, 523–539.
28. Fricke, W. Rapid and tissue-specific accumulation of solutes in the growth zone of barley leaves in response to salinity. *Planta* 2004, 219, 515–525.
29. James, R.A.; von Caemmerer, S.; Condon, A.T.; Zwart, A.B.; Munns, R. Genetic variation in tolerance to the osmotic stress component of salinity stress in durum wheat. *Funct. Plant Biol.* 2008, 35, 111–123.
30. Møller, I.M.; Jensen, P.E.; Hansson, A. Oxidative modifications to cellular components in plants. *Annu. Rev. Plant Biol.* 2007, 58, 459–481.
31. Fichman, Y.; Miller, G.; Mittler, R. Whole-plant live imaging of reactive oxygen species. *Mol. Plant* 2019, 12, 1203–1210.
32. Mittler, R. ROS are good. *Trends Plant Sci.* 2017, 22, 11–19.
33. Parida, A.K.; Das, A.B. Salinity tolerance and salinity effect on plants: A review. *Ecotoxicol. Environ. Safe* 2005, 60, 324–349.
34. Younis, M.E.; Hasaneen, M.N.A.; Kazamel, A.M.S. Exogenously applied ascorbic acid ameliorates detrimental effects of NaCl and mannitol stress in *Vicia faba* seedlings. *Protoplasma* 2010, 239, 39–48.
35. Wang, Y.; Wu, J.; Choi, Y.W.; Jun, T.H.; Kwon, S.W.; Choi, I.S.; Kim, Y.C.; Gupta, R.; Kim, S.T. Expression Analysis of *Oryza sativa* Ascorbate Peroxidase 1 (OsAPx1) in Response to Different Phytohormones and Pathogens. *J. Life Sci.* 2015, 25, 1091–1097.
36. Wang, Y.; Wu, J.; Kim, S.G.; Tsuda, K.; Gupta, R.; Park, S.Y.; Kim, S.T.; Kang, K.Y. Magnaporthe oryzae-secreted protein MSP1 induces cell death and elicits defense responses in rice. *Mol. Plant-Microbe Interact.* 2016, 29, 299–312.
37. Yang, L.; Zu, Y.G.; Tang, Z.H. Ethylene improves Arabidopsis salinity tolerance mainly via retaining K⁺ in shoots and roots rather than decreasing tissue Na⁺ content. *Environ. Exp. Bot.* 2013, 86, 60–69.

38. Gharbi, E.; Martínez, J.P.; Benahmed, H.; Lepoint, G.; Vanpee, B.; Quinet, M.; Lutts, S. Inhibition of ethylene synthesis reduces salinity-tolerance in tomato wild relative species *Solanum chilense*. *J. Plant Physiol.* 2017, 210, 24–37.
39. Shakar, M.; Yaseen, M.; Mahmood, R.; Ahmad, I. Calcium carbide induced ethylene modulate biochemical profile of *Cucumis sativus* at seed germination stage to alleviate salinity stress. *Sci. Hortic.* 2016, 213, 179–185.
40. Tavladoraki, P.; Cona, A.; Federico, R.; Tempera, G.; Viceconte, N.; Saccoccio, S.; Battaglia, V.; Toninello, A.; Agostinelli, E. Polyamine catabolism: Target for antiproliferative therapies in animals and stress tolerance strategies in plants. *Amino Acids* 2012, 42, 411–426.
41. Müller, M.; Munnébosch, S. Ethylene response factors. A key regulatory hub in hormone and stress signaling. *Plant Physiol.* 2015, 169, 32–41.
42. Yang, C.; Ma, B.; He, S.J.; Xiong, Q.; Duan, K.X.; Yin, C.C.; Chen, H.; Lu, X.; Chen, S.H.; Zhang, J.H. MAOHUZI6/ETHYLENE INSENSITIVE3-LIKE1 and ETHYLENE INSENSITIVE3-LIKE2 regulate ethylene response of roots and coleoptiles and negatively affect salinity tolerance in rice. *Plant Physiol.* 2015, 169, 148–165.
43. Sajid, H.; Chu, Z.; Zhigang, B.; Xiaochuang, C.; Lianfeng, Z.; Azhar, H.; Zhu, C.; Fahad, S.; James, A.B.; Zhang, J.; et al. Effects of 1-Methylcyclopropene on rice growth characteristics and superior and inferior spikelet development under salinity stress. *J. Plant Growth Regul.* 2018, 37, 1368–1384.
44. Hussain, S.; Bai, Z.; Huang, J.; Cao, X.; Zhu, L.; Zhu, C.; Zhang, J. 1-Methylcyclopropene Modulates Physiological, Biochemical, and Antioxidant Responses of Rice to Different Salinity Stress Levels. *Front. Plant Sci.* 2019, 10, 124.
45. Peng, Z.; He, S.; Gong, W.; Sun, J.; Pan, Z.; Xu, F.; Lu, Y.; Du, X. Comprehensive analysis of differentially expressed genes and transcriptional regulation induced by salt stress in two contrasting cotton genotypes. *BMC Genom.* 2014, 15, 760.
46. Zhao, X.C.; Schaller, G.E. Effect of salinity and osmotic stress upon expression of the ethylene receptor ETR1 in *Arabidopsis thaliana*. *FEBS Lett.* 2004, 562, 189–192.
47. Zhou, H.L.; Cao, W.H.; Cao, Y.R.; Liu, J.; Hao, Y.J.; Zhang, J.S.; Chen, S.Y. Roles of ethylene receptor NTHK1 domains in plant growth, stress response and protein phosphorylation. *FEBS Lett.* 2006, 580, 1239–1250.
48. Cao, W.H.; Liu, J.; He, X.J.; Mu, R.L.; Zhou, H.L.; Chen, S.Y.; Zhang, J.S. Modulation of ethylene responses affects plant salinity-stress responses. *Plant Physiol.* 2007, 143, 707–719.
49. Wilson, R.L.; Kim, H.; Bakshi, A.; Binder, B.M. The ethylene receptors ETHYLENE RESPONSE1 and ETHYLENE RESPONSE2 have contrasting roles in seed germination of *Arabidopsis* during salinity stress. *Plant Physiol.* 2014, 165, 1353–1366.
50. Cao, W.H.; Liu, J.; Zhou, Q.Y.; Cao, Y.R.; Zheng, S.F.; Du, B.X.; Zang, J.S.; Chen, S.Y. Expression of tobacco ethylene receptor NTHK1 alters plant responses to salinity stress. *Plant Cell Environ.* 2006, 29, 1210–1219.
51. Chen, T.; Liu, J.; Lei, G.; Liu, Y.F.; Li, Z.G.; Tao, J.J.; Hao, Y.J.; Cao, Y.R.; Lin, Q.; Zhang, W.K.; et al. Effects of tobacco Ethylene receptor mutations on receptor kinase activity, plant growth and stress responses. *Plant Cell Physiol.* 2009, 50, 1636–1650.
52. Cao, Y.R.; Chen, H.W.; Li, Z.G.; Tao, J.J.; Ma, B.; Zhang, W.K.; Chen, S.Y.; Zhang, J.S. Tobacco ankyrin protein NEIP2 interacts with ethylene receptor NTHK1 and regulates plant growth and stress responses. *Plant Cell Physiol.* 2015, 56, 803–818.
53. Jiang, C.; Belfield, E.J.; Cao, Y.; Smith, J.A.; Harberd, N.P. An *Arabidopsis* soil-salinity-tolerance mutation confers ethylene-mediated enhancement of sodium/potassium homeostasis. *Plant Cell.* 2013, 25, 3535–3552.
54. Achard, P.; Cheng, H.; De Grauwe, L.; Decat, J.; Schoutteten, H.; Moritz, T.; Van Der Straeten, D.; Peng, J.; Harberd, N.P. Integration of plant responses to environmentally activated phytohormonal signals. *Science* 2006, 311, 91–94.
55. Lei, G.; Shen, M.; Li, Z.G.; Zhang, B.; Duan, K.X.; Wang, N.; Cao, Y.R.; Zhang, W.K.; Ma, B.; Ling, H.Q.; et al. EIN2 regulates salinity stress response and interacts with a MA3 domain-containing protein ECIP1 in *Arabidopsis*. *Plant Cell Environ.* 2011, 34, 1678–1692.
56. Ge, X.M.; Cai, H.L.; Lei, X.; Zhou, X.; Yue, M.; He, J.M. Heterotrimeric G protein mediates ethylene-induced stomatal closure via hydrogen peroxide synthesis in *Arabidopsis*. *Plant J.* 2015, 82, 138–150.
57. Tao, J.J.; Chen, H.W.; Ma, B.; Zhang, W.K.; Chen, S.Y.; Zhang, J.S. The role of ethylene in plants under salinity stress. *Front. Plant Sci.* 2015, 6, 1059.
58. Zhang, L.X.; Li, Z.F.; Quan, R.D.; Li, G.J.; Wang, R.G.; Huang, R.F. An AP2 domain-containing gene, ESE1, targeted by the ethylene signaling component EIN3 is important for the salinity response in *Arabidopsis*. *Plant Physiol.* 2011, 157, 854–865.

59. Quan, R.; Wang, J.; Yang, D.; Zhang, H.; Zhang, Z.; Huang, R. EIN3 and SOS2 synergistically modulate plant salinity tolerance. *Sci. Rep.* 2017, 7, 44637–44647.
60. Qin, H.; Wang, J.; Chen, X.; Wang, F.; Peng, P.; Zhou, Y.; Miao, Y.; Zhang, Y.; Gao, Y.; Qi, Y.; et al. Rice Os DOF 15 contributes to ethylene-inhibited primary root elongation under salinity stress. *New Phytol.* 2019, 223, 798–813.
61. Jin, J.; Duan, J.; Shan, C.; Mei, Z.; Chen, H.; Feng, H.; Zhu, J.; Cai, W. Ethylene insensitive3-like2 (OsEIL2) confers stress sensitivity by regulating OsBURP16, the β subunit of polygalacturonase (PG1 β -like) subfamily gene in rice. *Plant Sci.* 2020, 292, 110353.
62. Liu, C.; Li, J.; Zhu, P.; Yu, J.; Hou, J.; Wang, C.; Long, D.; Yu, M.; Zhao, A. Mulberry EIL3 confers salinity and drought tolerances and modulates ethylene biosynthetic gene expression. *PeerJ* 2019, 7, e6391.
63. Wang, C.; Li, J.; Yuan, M. Salinity tolerance requires cortical microtubule reorganization in Arabidopsis. *Plant Cell Physiol.* 2007, 48, 1534–1547.
64. Zhang, Q.; Lin, F.; Mao, T.; Nie, J.; Yan, M.; Yuan, M.; Zhang, W. Phosphatidic acid regulates microtubule organization by interacting with MAP65-1 in response to salinity stress in Arabidopsis. *Plant Cell* 2012, 24, 4555–4576.
65. Le, J.; Vandenbusscheb, F.; Straetenb, D.V.D.; Verbelen, J.P. Position and cell 786 type-dependent microtubule reorientation characterizes the early response of the 787 Arabidopsis root epidermis to ethylene. *Physiol. Plant.* 2004, 121, 513–519.
66. Verbelen, J.P.; Jie, L.; Vissenberg, K.; Cnodder, T.D.; Vandenbussche, F.; Sugimoto, K.; Straeten, D.V.D. Microtubules and the control of cell elongation in Arabidopsis roots. In *The Plant Cytoskeleton: A Key Tool for Agro-Biotechnology*; Springer: Dordrecht, The Netherlands, 2008; pp. 73–90.
67. Sun, J.; Ma, Q.; Mao, T. Ethylene regulates the Arabidopsis microtubule-associated protein WAVE-DAMPENED2-LIKE5 in etiolated hypocotyl elongation. *Plant Physiol.* 2015, 169, 325–337.
68. Ma, Q.; Sun, J.; Mao, T. Microtubule bundling plays a role in ethylene-mediated cortical microtubule reorientation in etiolated Arabidopsis hypocotyls. *J. Cell Sci.* 2016, 129, 2043–2051.
69. Dou, L.; He, K.; Higaki, T.; Wang, X.; Mao, T. Ethylene signaling modulates cortical microtubule reassembly in response to salinity stress. *Plant Physiol.* 2018, 176, 2071–2081.
70. Cheng, M.C.; Liao, P.M.; Kuo, W.W.; Lin, T.P. The Arabidopsis ETHYLENE RESPONSE FACTOR1 regulates abiotic stress-responsive gene expression by binding to different cis-acting elements in response to different stress signals. *Plant Physiol.* 2013, 162, 1566–1582.
71. Li, Y.; Zhang, H.; Zhang, Q.; Liu, Q.; Zhai, H.; Zhao, N.; He, S. An AP2/ERF gene, IbRAP2-12, from sweetpotato is involved in salinity and drought tolerance in transgenic Arabidopsis. *J. Plant Sci.* 2019, 281, 19–30.
72. Djemal, R.; Khoudi, H. TdSHN1, a WIN1/SHN1-type transcription factor, imparts multiple abiotic stress tolerance in transgenic tobacco. *Environ. Exp. Bot.* 2016, 131, 89–100.
73. An, J.P.; Zhang, X.W.; Xu, R.R.; You, C.X.; Wang, X.F.; Hao, Y.J. Apple MdERF4 negatively regulates salinity tolerance by inhibiting MdERF3 transcription. *Plant Sci.* 2018, 276, 181–188.
74. Cheng, Z.; Zhang, X.; Zhao, K.; Yao, W.; Li, R.; Zhou, B.; Jiang, T. Over-expression of ERF38 gene enhances salt and osmotic tolerance in transgenic poplar. *Front. Plant Sci.* 2019, 10, 1375.
75. Yang, S.F.; Hoffman, N.E. Ethylene biosynthesis and its regulation in higher plants. *Annu. Rev. Plant Physiol.* 1984, 35, 155–189.
76. Wang, K.; Li, H.; Ecker, J.R. Ethylene biosynthesis and signaling networks. *Plant Cell* 2002, 14, 131–151.
77. Sauter, M.; Lorbiecke, R.; OuYang, B.; Pochapsky, T.C.; Rzewuski, G. The immediate-early ethylene response gene OsARD1 encodes an acireductone dioxygenase involved in recycling of the ethylene precursor S-adenosylmethionine. *Plant J.* 2005, 44, 718–729.
78. Liang, S.; Xiong, W.; Yin, C.; Xie, X.; Jin, Y.J.; Zhang, S.; Yang, B.; Ye, G.; Chen, S.; Luan, W.J. Overexpression of OsARD1 Improves Submergence, Drought, and Salinity Tolerances of Seedling Through the Enhancement of Ethylene Synthesis in Rice. *Front. Plant Sci.* 2019, 10, 1088.
79. Finch-Savage, W.E.; Leubner-Metzger, G. Seed dormancy and the control of germination. *New Phytol.* 2006, 171, 501–523.
80. Holdsworth, M.J.; Bentsink, L.; Soppe, W.J. Molecular networks regulating Arabidopsis seed maturation, after-ripening, dormancy and germination. *New Phytol.* 2008, 179, 33–54.
81. Weitbrecht, K.; Müller, K.; Leubner-Metzger, G. First off the mark: Early seed germination. *J. Exp. Bot.* 2011, 62, 3289–3309.

82. Ghassemian, M.; Nambara, E.; Cutler, S.; Kawaide, H.; Kamiya, Y.; McCourt, P. Regulation of abscisic acid signaling by the ethylene response pathway in Arabidopsis. *Plant Cell* 2000, 12, 1117–1126.
83. Cao, Y.R.; Chen, S.Y.; Zhang, J.S. Ethylene signaling regulates salinity stress response: An overview. *Plant Signal. Behav.* 2008, 3, 761–763.
84. Yu, Y.; Wang, J.; Shi, H.; Gu, J.; Dong, J.; Deng, X.W.; Huang, R. Salinity stress and ethylene antagonistically regulate nucleocytoplasmic partitioning of COP1 to control seed germination. *Plant Physiol.* 2016, 170, 2340–2350.
85. Lin, Y.; Yang, L.; Paul, M.; Zu, Y.; Tang, Z. Ethylene promotes germination of Arabidopsis seed under salinity by decreasing reactive oxygen species: Evidence for the involvement of nitric oxide simulated by sodium nitroprusside. *Plant Physiol. Biochem.* 2013, 73, 211–218.
86. Chang, C.; Wang, B.; Shi, L.; Li, Y.; Duo, L.; Zhang, W. Alleviation of salinity stress-induced inhibition of seed germination in cucumber (*Cucumis sativus* L.) by ethylene and glutamate. *J. Plant Physiol.* 2010, 167, 1152–1156.
87. Zhang, J.Y.; Luo, H.T.; Guo, Z.R. Overexpression of *Malus hupehensis* MhSHN1 Gene Enhances Salinity and Osmotic Stress Tolerance in Transgenic Tobacco Plants. *Russ. J. Plant Physiol.* 2018, 65, 857–864.
88. Lovato, M.B.; de Lemos Filho, J.P.; Martins, P.S. Growth responses of *Stylosanthes humilis* (Fabaceae) populations to saline stress. *Environ. Exp. Bot.* 1999, 41, 145–153.
89. Silva, P.O.; Medina, E.F.; Barros, R.S.; Ribeiro, D.M. Germination of salt-stressed seeds as related to the ethylene biosynthesis ability in three *Stylosanthes* species. *J. Plant Physiol.* 2014, 171, 14–22.
90. Silva, N.C.; de Souza, G.A.; Pimenta, T.M.; Brito, F.A.; Picoli, E.A.; Zsögön, A.; Ribeiro, D.M. Salinity stress inhibits germination of *Stylosanthes humilis* seeds through abscisic acid accumulation and associated changes in ethylene production. *Plant Physiol. Biochem.* 2018, 130, 399–407.
91. Zhang, S.; Yang, R.; Huo, Y.; Liu, S.; Yang, G.; Huang, J.; Zheng, C.; Wu, C. Expression of cotton PLATZ1 in transgenic Arabidopsis reduces sensitivity to osmotic and salinity stress for germination and seedling establishment associated with modification of the abscisic acid, gibberellin, and ethylene signalling pathways. *BMC Plant Biol.* 2018, 18, 1–11.
92. Ma, L.; Hu, L.; Fan, J.; Amombo, E.; Khaldun, A.B.M.; Zheng, Y.; Chen, L. Cotton GhERF38 gene is involved in plant response to salinity/drought and ABA. *Ecotoxicology* 2017, 26, 841–854.
93. Safari, D.; Jamali, F.; Nooryazdan, H.R.; Bayat, F. Evaluation of ACC deaminase producing '*Pseudomonas fluorescens*' strains for their effects on seed germination and early growth of wheat under salinity stress. *Aust. J. Crop. Sci.* 2018, 12, 413.
94. Li, H.; Lei, P.; Pang, X.; Li, S.; Xu, H.; Xu, Z.; Feng, X. Enhanced tolerance to salinity stress in canola (*Brassica napus* L.) seedlings inoculated with the halotolerant *Enterobacter cloacae* HSNJ4. *Appl. Soil Ecol.* 2017, 119, 26–34.
95. Wang, X.; Hou, C.; Zheng, K.; Li, Q.; Chen, S.; Wang, S. Overexpression of ERF96, a small ethylene response factor gene, enhances salinity tolerance in Arabidopsis. *Biol. Plant* 2017, 61, 693–701.
96. Wu, D.; Ji, J.; Wang, G.; Guan, C.; Jin, C. LchERF, a novel ethylene-responsive 76transcription factor from *Lycium chinense*, confers salinity tolerance in transgenic tobacco. *Plant Cell Rep.* 2014, 33, 2033–2045.
97. Yang, F.; Chen, H.; Liu, C.; Li, L.; Liu, L.; Han, X.; Wan, Z.; Sha, A. Transcriptome profile analysis of two *Vicia faba* cultivars with contrasting salinity tolerance during seed germination. *Sci. Rep.* 2020, 10, 7250.
98. Ahmed, W.; Imran, M.; Yaseen, M.; ul Haq, T.; Jamshaid, M.U.; Rukh, S.; Ikram, R.M.; Ali, M.; Ali, A.; Maqbool, M.; et al. Role of salicylic acid in regulating ethylene and physiological characteristics for alleviating salinity stress on germination, growth and yield of sweet pepper. *PeerJ* 2020, 8, e8475.
99. Khan, M.I.R.; Asgher, M.; Khan, N.A. Alleviation of salinity-induced photosynthesis and growth inhibition by salicylic acid involves glycinebetaine and ethylene in mungbean (*Vigna radiata* L.). *Plant Physiol. Biochem.* 2014, 80, 67–74.
100. Ahanger, M.A.; Agarwal, R.M. Salinity stress induced alterations in antioxidant metabolism and nitrogen assimilation in wheat (*Triticum aestivum* L.) as influenced by potassium supplementation. *Plant Physiol. Biochem.* 2017, 115, 449–460.
101. Ahanger, M.A.; Qin, C.; Begum, N.; Maodong, Q.; Dong, X.X.; El-Esawi, M.; El-Sheikh, M.A.; Alatar, A.A.; Zhang, L. Nitrogen availability prevents oxidative effects of salinity on wheat growth and photosynthesis by up-regulating the antioxidants and osmolytes metabolism, and secondary metabolite accumulation. *BMC Plant Biol.* 2019, 19, 1–12.
102. Cheeseman, J.M. The integration of activity in saline environments: Problems and perspectives. *Funct. Plant Biol.* 2013, 40, 759–774.
103. Bose, J.; Munns, R.; Shabala, S.; Gilliham, M.; Pogson, B.; Tyerman, S.D. Chloroplast function and ion regulation in plants growing on saline soils: Lessons from halophytes. *J. Exp. Bot.* 2017, 68, 3129–3143.

104. Van Zelm, E.; Zhang, Y.; Testerink, C. Salinity tolerance mechanisms of plants. *Annu. Rev. Plant Biol.* 2020, 71, 403–433.
105. Liu, C.; Zhao, X.; Yan, J.; Yuan, Z.; Gu, M. Effects of Salinity Stress on Growth, Photosynthesis, and Mineral Nutrients of 18 Pomegranate (*Punica granatum*) Cultivars. *Agronomy* 2020, 10, 27.
106. Brito, F.A.; Pimenta, T.M.; Henschel, J.M.; Martins, S.C.; Zsögön, A.; Ribeiro, D.M. Elevated CO₂ improves assimilation rate and growth of tomato plants under progressively higher soil salinity by decreasing abscisic acid and ethylene levels. *Environ. Exp. Bot.* 2020, 176, 104050.
107. Murata, N.; Takahashi, S.; Nishiyama, Y.; Allakhverdiev, S.I. Photoinhibition of photosystem II under environmental stress. *Biochim. Biophys. Acta* 2007, 1767, 414–421.
108. Liu, C.; Li, L.L.; Li, G.Z.; Hao, L. Ethylene insensitive mutation improves Arabidopsis plant tolerance to NO₂ exposure. *Ecotoxicol. Environ. Saf.* 2020, 189, 110043.
109. Sarafi, E.; Chatzissavvidis, C.; Therios, I. Effect of calcium and boron on the ion status, carbohydrate and proline content, gas exchange parameters and growth performance of pomegranate cv. Wonderful plants grown under NaCl stress. *Türk Tarım ve Doğa Bilimleri Dergisi* 2014, 1, 1606–1617.
110. Sun, Y.; Niu, G.; Masabni, J.G.; Ganjegunte, G. Relative salinity tolerance of 22 pomegranate (*Punica granatum*) cultivars. *HortScience* 2018, 53, 1513–1519.
111. Pan, Y.J.; Liu, L.; Lin, Y.C.; Zu, Y.G.; Li, L.P.; Tang, Z.H. Ethylene antagonizes salinity-induced growth retardation and cell death process via transcriptional controlling of ethylene-, BAG-and senescence-associated genes in Arabidopsis. *Front. Plant Sci.* 2016, 7, 696.
112. Splivallo, R.; Fischer, U.; Göbel, C.; Feussner, I.; Karlovsky, P. Truffles regulate plant root morphogenesis via the production of auxin and ethylene. *Plant Physiol.* 2009, 150, 2018–2029.
113. Nouri, M.Z.; Moumeni, A.; Komatsu, S. Abiotic stresses: Insight into gene regulation and protein expression in photosynthetic pathways of plants. *Int. J. Mol. Sci.* 2015, 16, 20392–20416.
114. De Zélicourt, A.; Synek, L.; Saad, M.M.; Alzubaidy, H.; Jalal, R.; Xie, Y.; Andres-Barrao, C.; Rolli, E.; Guerard, F.; Mariappan, K.G.; et al. Ethylene induced plant stress tolerance by *Enterobacter* sp. SA187 is mediated by 2-keto-4-methylthiobutyric acid production. *PLoS Genet.* 2018, 14, e1007273.
115. Ji, J.; Yuan, D.; Jin, C.; Wang, G.; Li, X.; Guan, C. Enhancement of growth and salinity tolerance of rice seedlings (*Oryza sativa* L.) by regulating ethylene production with a novel halotolerant PGPR strain *Glutamicibacter* sp. YD01 containing ACC deaminase activity. *Acta Physiol. Plant.* 2020, 42, 1–17.
116. Foyer, C.H.; Shigeoka, S. Understanding oxidative stress and antioxidant functions to enhance photosynthesis. *Plant Physiol.* 2011, 155, 93–100.
117. Considine, M.J.; Foyer, C.H. Redox regulation of plant development. *Antioxid. Redox Signal.* 2014, 21, 1305–1326.
118. Noctor, G.; Mhamdi, A.; Foyer, C.H. The roles of reactive oxygen metabolism in drought: Not so cut and dried. *Plant Physiol.* 2014, 164, 1636–1648.
119. Iqbal, N.; Umar, S.; Khan, N.A. Nitrogen availability regulates proline and ethylene production and alleviates salinity stress in mustard (*Brassica juncea*). *J. Plant Physiol.* 2015, 178, 84–91.
120. Zhang, F.; Wan, X.; Zhong, Y. Nitrogen as an important detoxification factor to cadmium stress in poplar plants. *J. Plant Interact.* 2014, 9, 249–258.