

Milroy Disease

Subjects: Genetics & Heredity

Contributor: Rita Xu

Milroy disease is a condition that affects the normal function of the lymphatic system.

Keywords: genetic conditions

1. Introduction

The lymphatic system produces and transports fluids and immune cells throughout the body. Impaired transport with accumulation of lymph fluid can cause swelling (lymphedema). Individuals with Milroy disease typically have lymphedema in their lower legs and feet at birth or develop it in infancy. The lymphedema typically occurs on both sides of the body and may worsen over time.

Milroy disease is associated with other features in addition to lymphedema. Males with Milroy disease are sometimes born with an accumulation of fluid in the scrotum (hydrocele). Males and females may have upslanting toenails, deep creases in the toes, wart-like growths (papillomas), and prominent leg veins. Some individuals develop non-contagious skin infections called cellulitis that can damage the thin tubes that carry lymph fluid (lymphatic vessels). Episodes of cellulitis can cause further swelling in the lower limbs.

2. Frequency

Milroy disease is a rare disorder; its incidence is unknown.

3. Causes

Mutations in the *FLT4* gene cause some cases of Milroy disease. The *FLT4* gene provides instructions for producing a protein called vascular endothelial growth factor receptor 3 (VEGFR-3), which regulates the development and maintenance of the lymphatic system. Mutations in the *FLT4* gene interfere with the growth, movement, and survival of cells that line the lymphatic vessels (lymphatic endothelial cells). These mutations lead to the development of small or absent lymphatic vessels. If lymph fluid is not properly transported, it builds up in the body's tissues and causes lymphedema. It is not known how mutations in the *FLT4* gene lead to the other features of this disorder.

Many individuals with Milroy disease do not have a mutation in the *FLT4* gene. In these individuals, the cause of the disorder is unknown.

3.1. The Gene Associated with Milroy Disease

- *FLT4*

4. Inheritance

Milroy disease is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder. In many cases, an affected person inherits the mutation from one affected parent. Other cases may result from new mutations in the *FLT4* gene. These cases occur in people with no history of the disorder in their family. About 10 percent to 15 percent of people with a mutation in the *FLT4* gene do not develop the features of Milroy disease.

5. Other Names for This Condition

- congenital familial lymphedema
- hereditary lymphedema type I

- Milroy's disease
- Nonne-Milroy lymphedema

References

1. Brice G, Child AH, Evans A, Bell R, Mansour S, Burnand K, Sarfarazi M, Jeffery S, Mortimer P. Milroy disease and the VEGFR-3 mutation phenotype. *J Med Genet*. 2005 Feb;42(2):98-102. Review.
 2. Brice GW, Mansour S, Ostergaard P, Connell F, Jeffery S, Mortimer P. Milroy Disease. 2006 Apr 27 [updated 2014 Sep 25]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1239/>
 3. Butler MG, Dagenais SL, Rockson SG, Glover TW. A novel VEGFR3 mutation causes Milroy disease. *Am J Med Genet A*. 2007 Jun 1;143A(11):1212-7.
 4. Evans AL, Bell R, Brice G, Comeglio P, Lipede C, Jeffery S, Mortimer P, Sarfarazi M, Child AH. Identification of eight novel VEGFR-3 mutations in families with primary congenital lymphoedema. *J Med Genet*. 2003 Sep;40(9):697-703.
 5. Ghalamkarpour A, Morlot S, Raas-Rothschild A, Utkus A, Mulliken JB, Boon LM, Vikkula M. Hereditary lymphedema type I associated with VEGFR3 mutation: the first de novo case and atypical presentations. *Clin Genet*. 2006 Oct;70(4):330-5.
 6. Irrthum A, Karkkainen MJ, Devriendt K, Alitalo K, Vikkula M. Congenital hereditary lymphedema caused by a mutation that inactivates VEGFR3 tyrosine kinase. *Am J Hum Genet*. 2000 Aug;67(2):295-301.
 7. Karkkainen MJ, Ferrell RE, Lawrence EC, Kimak MA, Levinson KL, McTigue MA, Alitalo K, Finegold DN. Missense mutations interfere with VEGFR-3 signalling in primary lymphoedema. *Nat Genet*. 2000 Jun;25(2):153-9.
 8. Levinson KL, Feingold E, Ferrell RE, Glover TW, Traboulsi EI, Finegold DN. Age of onset in hereditary lymphedema. *J Pediatr*. 2003 Jun;142(6):704-8.
-

Retrieved from <https://encyclopedia.pub/entry/history/show/11745>