

17-Beta Hydroxysteroid Dehydrogenase 3 Deficiency

Subjects: Genetics & Heredity

Contributor: Catherine Yang

17-beta hydroxysteroid dehydrogenase 3 deficiency is a condition that affects male sexual development. People with this condition are genetically male, with one X and one Y chromosome in each cell, and they have male gonads (testes). Their bodies, however, do not produce enough of a male sex hormone (androgen) called testosterone. Testosterone has a critical role in male sexual development, and a shortage of this hormone disrupts the formation of the external sex organs before birth.

Keywords: genetic conditions

1. Introduction

Most people with 17-beta hydroxysteroid dehydrogenase 3 deficiency are born with external genitalia that appear female. In some cases, the external genitalia do not look clearly male or clearly female (sometimes called ambiguous genitalia). Still other affected infants have genitalia that appear predominantly male, often with an unusually small penis (micropenis) or the urethra opening on the underside of the penis (hypospadias).

During puberty, people with this condition develop some male secondary sex characteristics, such as increased muscle mass, deepening of the voice, and development of male pattern facial and body hair. In addition to these changes typical of adolescent boys, some affected individuals may also experience breast enlargement (gynecomastia). Despite having testes, people with this disorder are generally unable to father children (infertile).

Children with 17-beta hydroxysteroid dehydrogenase 3 deficiency are often raised as girls. About half of these individuals adopt a male gender role in adolescence or early adulthood.

2. Frequency

17-beta hydroxysteroid dehydrogenase 3 deficiency is a rare disorder. Researchers have estimated that this condition occurs in approximately 1 in 147,000 newborns. It is more common in the Arab population of Gaza, where it affects 1 in 200 to 300 people.

3. Causes

Mutations in the *HSD17B3* gene cause 17-beta hydroxysteroid dehydrogenase 3 deficiency. The *HSD17B3* gene provides instructions for making an enzyme called 17-beta hydroxysteroid dehydrogenase 3. This enzyme is active in the testes, where it helps to produce testosterone from a weaker precursor androgen called androstenedione.

Mutations in the *HSD17B3* gene result in a 17-beta hydroxysteroid dehydrogenase 3 enzyme with little or no activity, reducing production of testosterone from androstenedione. The shortage of the stronger androgen affects the development of the reproductive tract in the male fetus, resulting in the abnormalities in the external sex organs that occur in 17-beta hydroxysteroid dehydrogenase 3 deficiency.

At puberty, conversion of androstenedione to testosterone increases in various tissues of the body through processes involving other enzymes. The additional testosterone results in the development of male secondary sex characteristics in adolescents, including those with 17-beta hydroxysteroid dehydrogenase 3 deficiency.

A portion of the androstenedione is also converted to the female sex hormone estrogen. Since impairment of the conversion to testosterone in this disorder results in excess androstenedione in the body, a corresponding excess of estrogen may be produced, leading to breast enlargement in some affected individuals.

3.1. The gene associated with 17-beta hydroxysteroid dehydrogenase 3 deficiency

- HSD17B3

4. Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition. Individuals who are genetically male and have two copies of a mutated gene in each cell are affected by 17-beta hydroxysteroid dehydrogenase 3 deficiency. People with two mutations who are genetically female do not usually experience any signs and symptoms of this disorder.

5. Other Names for This Condition

- 17-beta hydroxysteroid dehydrogenase III deficiency
- 17-ketosteroid reductase deficiency of testis
- 17-KSR deficiency
- neutral 17-beta-hydroxysteroid oxidoreductase deficiency
- pseudohermaphroditism, male, with gynecomastia
- testosterone 17-beta-dehydrogenase deficiency

References

1. Ben Rhouma B, Kallabi F, Mahfoudh N, Ben Mahmoud A, Engeli RT, Kamoun H, Keskes L, Odermatt A, Belguith N. Novel cases of Tunisian patients with mutations in the gene encoding 17 β -hydroxysteroid dehydrogenase type 3 and a founder effect. *J Steroid Biochem Mol Biol*. 2017 Jan;165(Pt A):86-94. doi:10.1016/j.jsbmb.2016.03.007.
 2. Bertelloni S, Balsamo A, Giordani L, Fischetto R, Russo G, Delvecchio M, Gennari M, Nicoletti A, Maggio MC, Concolino D, Cavallo L, Cicognani A, Chiumello G, Hiort O, Baroncelli GI, Faienza MF. 17beta-Hydroxysteroid dehydrogenase-3 deficiency: from pregnancy to adolescence. *J Endocrinol Invest*. 2009 Sep;32(8):666-70. doi: 10.3275/6281.
 3. Castro CC, Guaragna-Filho G, Calais FL, Coeli FB, Leal IR, Cavalcante-Junior EF, Monlleó IL, Pereira SR, Silva RB, Gabiatti JR, Marques-de-Faria AP, Maciel-Guerra AT, Mello MP, Guerra-Junior G. Clinical and molecular spectrum of patients with 17 β -hydroxysteroid dehydrogenase type 3 (17- β -HSD3) deficiency. *Arq Bras Endocrinol Metabol*. 2012 Nov;56(8):533-9.
 4. Engeli RT, Tsachaki M, Hassan HA, Sager CP, Essawi ML, Gad YZ, Kamel AK, Mazen I, Odermatt A. Biochemical Analysis of Four Missense Mutations in the HSD17B3 Gene Associated With 46,XY Disorders of Sex Development in Egyptian Patients. *J Sex Med*. 2017 Sep;14(9):1165-1174. doi: 10.1016/j.jsxm.2017.07.006.
 5. Faienza MF, Giordani L, Delvecchio M, Cavallo L. Clinical, endocrine, and molecular findings in 17beta-hydroxysteroid dehydrogenase type 3 deficiency. *J Endocrinol Invest*. 2008 Jan;31(1):85-91. Review.
 6. Galdiero M, Vitale P, Simeoli C, Afeltra L, Melis D, Alviggi C, Cariati F, Lo Calzo F, Di Somma C, Colao A, Pivonello R. The 17 β -hydroxysteroid dehydrogenase type 3 deficiency: a case report of an 18-year patient and review of the literature. *Minerva Endocrinol*. 2013 Mar;38(1):113-22. Review.
 7. Hassan HA, Mazen I, Gad YZ, Ali OS, Mekawy M, Essawi ML. Mutational Profile of 10 Afflicted Egyptian Families with 17- β -HSD-3 Deficiency. *Sex Dev*. 2016;10(2):66-73. doi: 10.1159/000445311.
 8. Mendonca BB, Gomes NL, Costa EM, Inacio M, Martin RM, Nishi MY, Carvalho FM, Tibor FD, Domenice S. 46,XY disorder of sex development (DSD) due to 17 β -hydroxysteroid dehydrogenase type 3 deficiency. *J Steroid Biochem Mol Biol*. 2017 Jan;165(Pt A):79-85. doi: 10.1016/j.jsbmb.2016.05.002. Review.
 9. Yu B, Liu Z, Mao J, Wang X, Zheng J, Xiong S, Cui M, Ma W, Huang Q, Xu H, Huang B, Nie M, Wu X. Novel mutations of HSD17B3 in three Chinese patients with 46,XY Disorders of Sex Development. *Steroids*. 2017 Oct;126:1-6. doi:10.1016/j.steroids.2017.07.009.
-

