

# L-Arginine Supplementation during Late Gestation

Subjects: **Zoology**

Contributor: Jinsu Hong , Lin Hu Fang , Jae Hark Jeong , Yoo Yong Kim

Arginine is one of the functional amino acids that enhances the growth of fetus and placenta development. Since the fetal growth and the nutrient requirement for fetuses are increased hugely during the late gestation period in high-prolific sows, supplementation of L-arginine could have a positive influence on the reproductive performance of sows and piglet uniformity.

arginine

late gestation

sow

reproductive performance

piglet uniformity

## 1. Introduction

The high prolific sows are developed to improve the number of piglets and sow productivity in the swine industry [1]. However, as they have greater litter size, some problems were reported that the proportion of small piglets at birth, within-litter variation of piglet birth weight, and mortality of suckling piglets were also increased [2][3][4]. These problems are caused by several maternal factors, such as inadequate nutrient intake of late gestation, high number of fetus, insufficient cervical space, or insufficient reproductive tract for fetus development [5][6][7]. Therefore, various studies were conducted to minimize these problems and improve the piglet uniformity of high prolific sows with nutritional strategies including sow body condition, nutrients type or ratio, and functional amino acids [8][9][10].

Fetal growth is rapid during the late gestation period and nutrient requirement for the fetus also increases greatly [8]. The low birth weight of piglets was reported to increase the proportion of stillborn piglets considered to be at a greater risk of mortality and morbidity [9][10]. Since supplying adequate nutrient to sows during the late gestation period is important, an increased feeding method was suggested in order to meet their nutrient requirement. Although increased feeding during late gestation is likely to meet the nutrient requirement of sows and improve piglet birth weight [11][12], it also showed a negative effect on sow body condition and postpartum agalactia [13][14], resulting in poor milk production and lactation feed intake [15][16]. With the effects of arginine, if the nutrient delivery efficiency from the dam to the fetus in late gestation increased by additional arginine supplementation, it could partly be attributed to increase in the birth weight of small fetuses and piglet uniformity at birth.

Most of the previous studies for evaluating the effect of arginine in gestating sows were investigated in the early-gestation period [17][18][19][20] or whole gestation period [21][22][23][24]. Additionally, most previous studies for the effect of arginine supplementation in sows, investigated the effect of 1% arginine supplementation in sow diet. Thus, there is a need to investigate the effects of dietary arginine below 1% of supplementation level keeping in mind the market cost of L-arginine, in order to apply this in the field.

### 3. Results

The arginine supplementation levels in the late-gestating sows' diet did not affect body weight, backfat thickness, and the lactation feed intake of sows (Table 1).

**Table 1.** Effects of arginine supplementation levels on body weight and back-fat thickness in late-gestating sows.

|                               | Treatment <sup>1</sup> |       |       | SEM <sup>2</sup> | p-Value |      |       |
|-------------------------------|------------------------|-------|-------|------------------|---------|------|-------|
|                               | CON                    | ARG10 | ARG15 |                  | Diet    | Lin. | Quad. |
| <b>Body Weight, kg</b>        |                        |       |       |                  |         |      |       |
| Day 70                        | 237.1                  | 239.1 | 238.4 | 3.58             | 0.98    | 0.90 | 0.85  |
| Day 110                       | 258.2                  | 256.3 | 252.7 | 3.33             | 0.79    | 0.50 | 0.99  |
| BW gain (70–110 days)         | 21.1                   | 17.3  | 14.3  | 2.46             | 0.53    | 0.27 | 0.79  |
| 24 h postpartum               | 233.7                  | 230.8 | 227.9 | 3.73             | 0.82    | 0.54 | 0.92  |
| Day 21 of lactation           | 229.4                  | 224.9 | 218.5 | 3.97             | 0.53    | 0.26 | 0.95  |
| BW loss (0–21 days)           | -4.3                   | -5.9  | -9.4  | 1.67             | 0.44    | 0.20 | 0.96  |
| <b>Backfat Thickness, mm</b>  |                        |       |       |                  |         |      |       |
| Day 70                        | 20.7                   | 21.4  | 19.1  | 0.77             | 0.49    | 0.35 | 0.46  |
| Day 110                       | 22.4                   | 22.6  | 20.7  | 0.84             | 0.59    | 0.37 | 0.66  |
| BF gain (70–110 days)         | 2.1                    | 1.3   | 1.6   | 0.3              | 0.51    | 0.57 | 0.30  |
| 24 h postpartum               | 21.0                   | 21.1  | 20.3  | 0.85             | 0.91    | 0.72 | 0.84  |
| Day 21 of lactation           | 20.1                   | 19.8  | 18.6  | 0.79             | 0.71    | 0.43 | 0.90  |
| BF loss (0–21 days)           | -0.9                   | -1.3  | -1.7  | 0.38             | 0.69    | 0.40 | 0.85  |
| Lactation feed intake, kg/day | 4.82                   | 4.95  | 4.97  | 0.114            | 0.85    | 0.62 | 0.77  |

In reproductive performance, dietary arginine levels during late gestation had no influence on the number of piglets for total born, stillborn, mummy, and born alive (Table 2). However, piglet birth weight showed a quadratic response ( $p < 0.03$ ) such that the piglet birth weight was decreased by the arginine levels from 0.72% to 1.0%, whereas it was increased by the arginine level from 1.0% to 1.5%. Additionally, increasing the inclusion level of dietary arginine tended to increase ( $p < 0.10$ ) the alive litter weight linearly. Increasing the dietary arginine levels in late gestation did not affect the farrowing time, helping frequency, and placenta weight.

**Table 2.** Effects of arginine supplementation levels on reproductive performance in late-gestating sows.

|                         | Treatment <sup>1</sup> |           |           | SEM <sup>2</sup> | p-Value |      |       |
|-------------------------|------------------------|-----------|-----------|------------------|---------|------|-------|
|                         | CON                    | ARG10     | ARG15     |                  | Diet    | Lin. | Quad. |
| <b>No. of Pigs</b>      |                        |           |           |                  |         |      |       |
| Total born              | 13.7                   | 15.0      | 14.8      | 0.50             | 0.53    | 0.46 | 0.41  |
| Stillborn               | 0.7                    | 0.7       | 0.8       | 0.12             | 0.93    | 0.73 | 0.89  |
| Mummy                   | 0.2                    | 0.3       | 0.1       | 0.08             | 0.49    | 0.31 | 0.53  |
| Born alive              | 12.8                   | 14.0      | 13.9      | 0.45             | 0.35    | 0.31 | 0.31  |
| Total litter weight, kg | 19.61                  | 19.91     | 21.49     | 0.521            | 0.30    | 0.13 | 0.73  |
| Alive litter weight, kg | 18.47                  | 19.01     | 20.69     | 0.506            | 0.19    | 0.07 | 0.81  |
| Piglet birth weight, kg | 1.57<br>A              | 1.38<br>B | 1.52<br>A | 0.036            | 0.08    | 0.85 | 0.03  |
| Farrowing time, min     | 182.4                  | 186.2     | 205.7     | 9.33             | 0.58    | 0.31 | 0.82  |
| Helping frequency       | 0.80                   | 0.81      | 1.00      | 0.141            | 0.82    | 0.55 | 0.85  |
| Placenta weight, kg     | 3.74                   | 3.74      | 3.90      | 0.224            | 0.95    | 0.76 | 0.90  |
| Placenta efficiency     | 5.80                   | 5.89      | 6.02      | 0.246            | 0.94    | 0.73 | 0.98  |


Increasing dietary arginine levels in late gestation linearly increased ( $p < 0.05$ ) the litter weight at day 21, and the litter weight gain, such that litter weight and litter weight gain for ARG15 tended to be greater than those for CON (Table 3). The piglet weight at day 21 of lactation was linearly increased ( $p < 0.05$ ) as the dietary arginine level increased in late gestation. Additionally, increasing the dietary arginine level in late gestation linearly increased ( $p < 0.05$ ) piglet weight gain, such that the piglet weight gain for ARG15 tended to be greater than that for CON.

**Table 3.** Effects of arginine supplementation levels on litter performance in late-gestating sows.

|                       | Treatment <sup>1</sup> |       |       | SEM <sup>2</sup> | p-Value |      |       |
|-----------------------|------------------------|-------|-------|------------------|---------|------|-------|
|                       | CON                    | ARG10 | ARG15 |                  | Diet    | Lin. | Quad. |
| <b>No. of Piglets</b> |                        |       |       |                  |         |      |       |
| After-fostering       | 11.5                   | 11.6  | 11.6  | 0.10             | 0.90    | 0.77 | 0.72  |

|                          | Treatment <sup>1</sup> |                     |                    | SEM <sup>2</sup> | p-Value |      |       |
|--------------------------|------------------------|---------------------|--------------------|------------------|---------|------|-------|
|                          | CON                    | ARG10               | ARG15              |                  | Diet    | Lin. | Quad. |
| Day 21 of lactation      | 10.7                   | 10.4                | 10.9               | 0.16             | 0.52    | 0.49 | 0.36  |
| <b>Litter Weight, kg</b> |                        |                     |                    |                  |         |      |       |
| After-fostering          | 17.67 <sup>a</sup>     | 16.01 <sup>b</sup>  | 17.73 <sup>a</sup> | 0.326            | 0.04    | 0.63 | 0.01  |
| Day 21 of lactation      | 53.74 <sup>B</sup>     | 53.31 <sup>B</sup>  | 60.27 <sup>A</sup> | 1.332            | 0.05    | 0.02 | 0.32  |
| Weight gain (0–21 days)  | 36.06 <sup>B</sup>     | 37.17 <sup>AB</sup> | 42.58 <sup>A</sup> | 1.262            | 0.07    | 0.02 | 0.64  |
| <b>Piglet Weight, kg</b> |                        |                     |                    |                  |         |      |       |
| After-fostering          | 1.55 <sup>a</sup>      | 1.38 <sup>b</sup>   | 1.53 <sup>a</sup>  | 0.031            | 0.04    | 0.82 | 0.01  |
| Day 21 of lactation      | 5.03                   | 5.14                | 5.51               | 0.103            | 0.13    | 0.04 | 0.78  |
| Weight gain (0–21 days)  | 3.48 <sup>B</sup>      | 3.74 <sup>AB</sup>  | 3.98 <sup>A</sup>  | 0.093            | 0.08    | 0.03 | 0.66  |

The effect of dietary arginine levels in late gestation on the piglet uniformity is presented in Table 4. The SD for piglet birth weight was linearly increased ( $p < 0.05$ ), and the CV for piglet birth weight had a tendency of linear increase ( $p < 0.10$ ) as the dietary arginine level increased in late gestation. The piglet BW distribution at birth showed a significant difference ( $p < 0.01$ ) among dietary treatment (Figure 1). However, dietary arginine levels in late gestation did not affect the piglet uniformity and BW distribution at day 21 of lactation.



**Figure 1.** Effect of arginine supplementation level in late gestating sows on piglet distribution into body weight classes on day 0 and day 21 of lactation by the FREQ procedure, with Chi-square test. CON: corn-SBM-based diet with Arg 0.72%, Arg10: corn-SBM-based diet with Arg 1.0%, Arg15: corn-SBM-based diet with Arg1.5%.

**Table 4.** Effects of arginine supplementation levels on piglet uniformity in late-gestating sows.

| Treatment <sup>1</sup>                               |       |       | SEM <sup>2</sup> | p-Value |      |       |
|--|-------|-------|------------------|---------|------|-------|
| CON  | ARG10 | ARG15 |                  | Diet    | Lin. | Quad. |
| <b>Piglet Uniformity at Birth</b>                    |       |       |                  |         |      |       |
| Avg. BW, kg  | 1.57  | 1.38  | 1.52             | -       | -    | -     |
| SD   | 274.0 | 293.5 | 334.3            | 12.72   | 0.13 | 0.04  |
| CV   | 18.4  | 21.8  | 22.9             | 1.01    | 0.15 | 0.07  |
| <b>Piglet Distribution into Birth Weight Classes</b> |       |       |                  |         |      |       |

|  | Treatment <sup>1</sup> |                   |                   | SEM <sup>2</sup> | p-Value |      |       |
|--|------------------------|-------------------|-------------------|------------------|---------|------|-------|
|  | CON                    | ARG10             | ARG15             |                  | Diet    | Lin. | Quad. |
| <0.5 kg, %   | 0                      | 1.0               | 1.2               | 0.30             | 0.24    | 0.15 | 0.38  |
| 0.5–0.8 kg, %  | 3.1                    | 5.6               | 4.2               | 0.79             | 0.44    | 0.59 | 0.25  |
| 0.8–1.2 kg, %  | 15.8                   | 25.8              | 18.2              | 2.36             | 0.21    | 0.87 | 0.07  |
| 1.2–1.6 kg, %  | 27.9 <sup>b</sup>      | 41.5 <sup>a</sup> | 26.3 <sup>b</sup> | 4.39             | 0.03    | 0.49 | 0.01  |
| 1.6–2.0 kg, %  | 39.0 <sup>ab</sup>     | 23.0 <sup>b</sup> | 39.7 <sup>a</sup> | 5.55             | 0.08    | 0.64 | 0.03  |
| >2.0 kg, %   | 14.5                   | 2.9               | 10.4              | 5.67             | 0.15    | 0.70 | 0.06  |
| <b>Piglet Uniformity at day 21</b>                   |                        |                   |                   |                  |         |      |       |
| Avg. BW, kg  | 5.03                   | 5.14              | 5.51              | -                | -       | -    | -     |
| SD   | 1186.8                 | 1198.1            | 1277.4            | 47.64            | 0.69    | 0.41 | 0.84  |
| CV   | 24.6                   | 24.4              | 25.1              | 1.17             | 0.97    | 0.85 | 0.89  |
| <b>Piglet Distribution into BW Classes at day 21</b> |                        |                   |                   |                  |         |      |       |
| <3 kg, %   | 9.5                    | 8.1               | 9.3               | 1.41             | 0.91    | 0.99 | 0.67  |
| 3–4 kg, %  | 11.5                   | 14.0              | 11.9              | 1.46             | 0.77    | 0.99 | 0.48  |
| 4–5 kg,  | 25.8                   | 22.1              | 23.3              | 2.39             | 0.83    | 0.72 | 0.62  |

|           | Treatment <sup>1</sup> |       |       | SEM <sup>2</sup> | p-Value |      |       |
|-----------|------------------------|-------|-------|------------------|---------|------|-------|
|           | CON                    | ARG10 | ARG15 |                  | Diet    | Lin. | Quad. |
| %         |                        |       |       |                  |         |      |       |
| 5–6 kg, % | 26.3                   | 29.9  | 19.4  | 2.17             | 0.14    | 0.12 | 0.19  |
| 6–7 kg, % | 20.2                   | 17.9  | 23.9  | 2.01             | 0.48    | 0.36 | 0.42  |
| >7 kg, %  | 6.7                    | 8.0   | 12.2  | 1.80             | 0.41    | 0.19 | 0.86  |

The blood concentration of AAs in the gestating sows is presented in Table 5. Increasing arginine supplementation to late-gestating sows linearly increased ( $p < 0.05$ ) the blood concentrations of arginine and ornithine at day 90 and day 110 of gestation, respectively. On the other hand, the blood concentrations of alanine at day 110 of gestation for sows fed a diet with higher dietary arginine were linearly decreased ( $p < 0.03$ ) compared to those for sows fed the isonitrogenous control diet. Additionally, glutamine concentration for sows at day 110 of gestation was quadratically increased ( $p < 0.05$ ), as the dietary arginine level increased. There were no significant differences in blood concentrations of other AA, among dietary treatments. The plasma concentrations of arginine, lysine, and methionine for day 90 of gestation were greater ( $p < 0.05$ ) than those for day 70 of gestation. The plasma concentrations of alanine and glutamine for day 110 of gestation were greater ( $p < 0.05$ ) than those for day 70 of gestation, whereas the plasma concentrations of citrulline, glycine, leucine, ornithine, taurine, and valine for day 110 of gestation were less ( $p < 0.05$ ) than those for day 70 of gestation. There were interactions ( $p < 0.05$ ) between arginine and the date effect in the plasma concentration for ornithine such that there was no significant difference in the plasma ornithine concentration for day 70 of gestation, whereas plasma ornithine concentrations at day 90 of gestation for ARG10 and ARG15 treatments were greater ( $p < 0.05$ ) than that for CON treatment, and plasma ornithine concentration at day 110 of gestation for ARG15 treatment was greater ( $p < 0.05$ ) than those for CON and ARG10 treatments.

**Table 5.** Effects of arginine supplementation levels on plasma amino acid profile in late-gestating sows.

|                        | Treatment <sup>1</sup> |       |       | SEM <sup>2</sup> | p-Value |      |       |       |       |              |
|------------------------|------------------------|-------|-------|------------------|---------|------|-------|-------|-------|--------------|
|                        | CON                    | ARG10 | ARG15 |                  | Diet    | Lin. | Quad. | Group | Date  | Group × Date |
| <b>Alanine, μmol/L</b> |                        |       |       |                  |         |      |       |       |       |              |
| Day 70                 | 609.5                  | 487.0 | 454.0 | 46.75            | 0.49    | 0.27 | 0.56  | 0.09  | <0.01 | 0.51         |
| Day 90                 | 709.1                  | 710.5 | 652.0 | 37.24            | 0.79    | 0.53 | 0.79  |       |       |              |

|  | Treatment <sup>1</sup> |                    |                    | SEM <sup>2</sup> | p-Value |       |       |       |       | Group x Date |
|--|------------------------|--------------------|--------------------|------------------|---------|-------|-------|-------|-------|--------------|
|  | CON                    | ARG10              | ARG15              |                  | Diet    | Lin.  | Quad. | Group | Date  |              |
| Day 110  | 878.4 <sup>A</sup>     | 916.3 <sup>A</sup> | 628.7 <sup>B</sup> | 53.05            | 0.06    | 0.03  | 0.26  |       |       |              |
| <b>Arginine, <math>\mu\text{mol/L}</math></b>      |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70   | 154.5                  | 255.0              | 195.8              | 23.65            | 0.31    | 0.66  | 0.14  | <0.01 | <0.01 | 0.27         |
| Day 90   | 262.9 <sup>b</sup>     | 349.1 <sup>b</sup> | 482.7 <sup>a</sup> | 29.44            | <0.01   | <0.01 | 0.88  |       |       |              |
| Day 110  | 196.4 <sup>b</sup>     | 301.2 <sup>a</sup> | 366.9 <sup>a</sup> | 24.34            | <0.01   | <0.01 | 0.34  |       |       |              |
| <b>Aspartic Acid, <math>\mu\text{mol/L}</math></b> |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70   | 14.0                   | 14.3               | 13.5               | 1.33             | 0.97    | 0.88  | 0.89  | 0.69  | 0.15  | 0.74         |
| Day 90   | 18.8                   | 31.4               | 18.3               | 4.79             | 0.46    | 0.82  | 0.23  |       |       |              |
| Day 110  | 14.4                   | 14.0               | 14.4               | 0.87             | 0.98    | 0.96  | 0.85  |       |       |              |
| <b>Citrulline, <math>\mu\text{mol/L}</math></b>    |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70   | 128.0                  | 109.3              | 114.5              | 5.64             | 0.54    | 0.48  | 0.34  | 0.96  | <0.01 | 0.58         |
| Day 90   | 114.8                  | 133.8              | 127.3              | 6.99             | 0.54    | 0.58  | 0.35  |       |       |              |
| Day 110  | 97.8                   | 89.7               | 98.9               | 4.19             | 0.68    | 0.80  | 0.40  |       |       |              |
| <b>Cystine, <math>\mu\text{mol/L}</math></b>       |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70   | 2.00                   | 0.33               | 1.00               | 0.408            | 0.38    | 0.47  | 0.20  | 0.70  | 0.50  | 0.75         |
| Day 90   | 0.88                   | 1.50               | 1.29               | 0.308            | 0.71    | 0.68  | 0.49  |       |       |              |
| Day 110  | 3.40                   | 2.33               | 1.00               | 0.928            | 0.57    | 0.31  | 0.93  |       |       |              |
| <b>Glutamic Acid, <math>\mu\text{mol/L}</math></b> |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70   | 240.5                  | 260.3              | 198.0              | 30.72            | 0.72    | 0.58  | 0.66  | 0.60  | 0.07  | 0.81         |
| Day 90   | 173.3                  | 211.6              | 189.3              | 16.13            | 0.63    | 0.81  | 0.36  |       |       |              |
| Day 110  | 170.6                  | 153.8              | 152.6              | 13.82            | 0.84    | 0.64  | 0.76  |       |       |              |

|   | Treatment <sup>1</sup> |                    |                    | SEM <sup>2</sup> | p-Value |      |       |       |       | Group x Date |
|---|------------------------|--------------------|--------------------|------------------|---------|------|-------|-------|-------|--------------|
|   | CON                    | ARG10              | ARG15              |                  | Diet    | Lin. | Quad. | Group | Date  |              |
| <b>Glutamine, <math>\mu\text{mol/L}</math></b>  |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70  | 260.5                  | 220.7              | 249.5              | 23.86            | 0.84    | 0.95 | 0.58  | 0.27  | 0.01  | 0.48         |
| Day 90  | 343.8                  | 328.9              | 289.4              | 14.64            | 0.32    | 0.14 | 0.88  |       |       |              |
| Day 110   | 299.8 <sup>a</sup>     | 330.8 <sup>a</sup> | 259.9 <sup>b</sup> | 10.39            | 0.03    | 0.04 | 0.04  |       |       |              |
| <b>Glycine, <math>\mu\text{mol/L}</math></b>    |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70  | 1224.5                 | 1186.7             | 1073.5             | 38.59            | 0.27    | 0.13 | 0.85  | 0.58  | 0.03  | 0.76         |
| Day 90  | 1206.0                 | 1291.5             | 1204.9             | 30.62            | 0.43    | 0.83 | 0.20  |       |       |              |
| Day 110   | 1070.9                 | 1105.3             | 1117.9             | 39.67            | 0.88    | 0.66 | 0.85  |       |       |              |
| <b>Histidine, <math>\mu\text{mol/L}</math></b>  |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70  | 93.0                   | 103.3              | 93.0               | 5.31             | 0.72    | 0.90 | 0.46  | 0.36  | 0.21  | 0.98         |
| Day 90  | 103.9                  | 116.5              | 96.6               | 6.80             | 0.51    | 0.56 | 0.31  |       |       |              |
| Day 110   | 90.8                   | 97.0               | 88.7               | 3.60             | 0.69    | 0.71 | 0.42  |       |       |              |
| <b>Isoleucine, <math>\mu\text{mol/L}</math></b> |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70  | 107.0                  | 121.3              | 104.0              | 6.11             | 0.51    | 0.72 | 0.32  | 0.65  | 0.81  | 0.92         |
| Day 90  | 120.4                  | 125.8              | 115.7              | 3.46             | 0.52    | 0.50 | 0.35  |       |       |              |
| Day 110   | 134.2                  | 119.5              | 109.4              | 11.82            | 0.69    | 0.42 | 0.84  |       |       |              |
| <b>Leucine, <math>\mu\text{mol/L}</math></b>    |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70  | 210.5                  | 220.3              | 202.8              | 10.62            | 0.82    | 0.74 | 0.66  | 0.66  | 0.02  | 0.84         |
| Day 90  | 217.6                  | 228.4              | 205.7              | 6.21             | 0.36    | 0.35 | 0.26  |       |       |              |
| Day 110   | 194.8                  | 165.5              | 168.0              | 13.38            | 0.61    | 0.48 | 0.54  |       |       |              |
| <b>Lysine, <math>\mu\text{mol/L}</math></b>     |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70  | 281.5                  | 321.0              | 239.5              | 18.85            | 0.17    | 0.24 | 0.19  | 0.44  | <0.01 | 0.71         |

|                              | Treatment <sup>1</sup> |                    |                    | SEM <sup>2</sup> | p-Value |       |       |       |       | Group x Date |
|------------------------------|------------------------|--------------------|--------------------|------------------|---------|-------|-------|-------|-------|--------------|
|                              | CON                    | ARG10              | ARG15              |                  | Diet    | Lin.  | Quad. | Group | Date  |              |
| Day 90                       | 409.4                  | 423.6              | 438.9              | 18.45            | 0.83    | 0.55  | 0.93  |       |       |              |
| Day 110                      | 282.4                  | 326.5              | 296.3              | 14.23            | 0.48    | 0.85  | 0.25  |       |       |              |
| <b>Methionine, µmol/L</b>    |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70                       | 48.5                   | 57.7               | 50.8               | 2.80             | 0.47    | 0.92  | 0.24  | 0.48  | <0.01 | 0.92         |
| Day 90                       | 73.0                   | 72.4               | 68.6               | 2.48             | 0.76    | 0.47  | 0.86  |       |       |              |
| Day 110                      | 58.6                   | 62.3               | 59.0               | 2.20             | 0.79    | 0.96  | 0.50  |       |       |              |
| <b>Ornithine, µmol/L</b>     |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70                       | 208.0                  | 163.3              | 146.0              | 17.04            | 0.43    | 0.22  | 0.58  | 0.12  | <0.01 | <0.01        |
| Day 90                       | 149.6 <sup>b</sup>     | 194.1 <sup>a</sup> | 228.1 <sup>a</sup> | 9.85             | <0.01   | <0.01 | 0.32  |       |       |              |
| Day 110                      | 114.4 <sup>b</sup>     | 143.8 <sup>b</sup> | 184.6 <sup>a</sup> | 9.38             | <0.01   | <0.01 | 0.80  |       |       |              |
| <b>Phenylalanine, µmol/L</b> |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70                       | 93.0                   | 98.0               | 84.5               | 13.60            | 0.53    | 0.42  | 0.52  | 0.24  | 0.07  | 0.77         |
| Day 90                       | 85.5                   | 104.8              | 85.6               | 4.51             | 0.13    | 0.75  | 0.05  |       |       |              |
| Day 110                      | 79.3                   | 81.7               | 76.4               | 3.83             | 0.89    | 0.72  | 0.71  |       |       |              |
| <b>Proline, µmol/L</b>       |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70                       | 408.0                  | 422.0              | 379.5              | 21.51            | 0.73    | 0.59  | 0.67  | 0.75  | 0.21  | 0.75         |
| Day 90                       | 389.5                  | 400.6              | 434.3              | 13.71            | 0.42    | 0.19  | 0.87  |       |       |              |
| Day 110                      | 349.8                  | 386.5              | 377.7              | 15.44            | 0.60    | 0.55  | 0.47  |       |       |              |
| <b>Serine, µmol/L</b>        |                        |                    |                    |                  |         |       |       |       |       |              |
| Day 70                       | 186.5                  | 179.7              | 153.8              | 7.66             | 0.17    | 0.09  | 0.75  | 0.26  | 0.08  | 0.83         |
| Day 90                       | 167.6                  | 215.6              | 158.0              | 16.59            | 0.33    | 0.64  | 0.16  |       |       |              |

|                           | Treatment <sup>1</sup> |                    |                    | SEM <sup>2</sup> | p-Value |      |       |       |       | Group x Date |
|---------------------------|------------------------|--------------------|--------------------|------------------|---------|------|-------|-------|-------|--------------|
|                           | CON                    | ARG10              | ARG15              |                  | Diet    | Lin. | Quad. | Group | Date  |              |
| Day 110                   | 139.5                  | 153.3              | 134.6              | 7.01             | 0.61    | 0.65 | 0.35  |       |       |              |
| <b>Taurine, μmol/L</b>    |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70                    | 141.5                  | 119.0              | 83.0               | 13.35            | 0.22    | 0.10 | 0.96  | 0.13  | <0.01 | 0.14         |
| Day 90                    | 83.0                   | 92.9               | 88.4               | 4.40             | 0.67    | 0.72 | 0.42  |       |       |              |
| Day 110                   | 67.8                   | 66.2               | 72.9               | 2.96             | 0.68    | 0.44 | 0.62  |       |       |              |
| <b>Threonine, μmol/L</b>  |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70                    | 177.0                  | 191.7              | 161.3              | 13.47            | 0.68    | 0.61 | 0.56  | 0.05  | 0.05  | 0.82         |
| Day 90                    | 138.1                  | 149.1              | 126.0              | 6.86             | 0.42    | 0.39 | 0.31  |       |       |              |
| Day 110                   | 133.2                  | 167.2              | 105.6              | 11.06            | 0.11    | 0.16 | 0.08  |       |       |              |
| <b>Tryptophan, μmol/L</b> |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70                    | 61.0                   | 76.0               | 62.5               | 4.37             | 0.36    | 0.92 | 0.19  | 0.10  | 0.40  | 0.85         |
| Day 90                    | 62.8                   | 67.1               | 64.1               | 2.35             | 0.75    | 0.91 | 0.46  |       |       |              |
| Day 110                   | 54.6                   | 66.5               | 60.0               | 3.00             | 0.29    | 0.62 | 0.15  |       |       |              |
| <b>Tyrosine, μmol/L</b>   |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70                    | 98.0                   | 104.7              | 91.5               | 7.10             | 0.77    | 0.69 | 0.63  | 0.17  | 0.23  | 0.96         |
| Day 90                    | 98.1                   | 120.9              | 97.7               | 6.37             | 0.24    | 0.77 | 0.10  |       |       |              |
| Day 110                   | 87.1                   | 101.3              | 89.7               | 4.03             | 0.36    | 0.99 | 0.16  |       |       |              |
| <b>Valine, μmol/L</b>     |                        |                    |                    |                  |         |      |       |       |       |              |
| Day 70                    | 306.5                  | 330.3              | 284.0              | 15.04            | 0.46    | 0.48 | 0.39  | 0.25  | 0.01  | 0.93         |
| Day 90                    | 276.9 <sup>AB</sup>    | 293.6 <sup>A</sup> | 253.4 <sup>B</sup> | 7.49             | 0.09    | 0.12 | 0.10  |       |       |              |
| Day 110                   | 252.0                  | 230.3              | 202.7              | 19.14            | 0.57    | 0.31 | 0.93  |       |       |              |

The dietary arginine levels for the late-gestating sows had no significant influence on the blood concentrations of BUN, creatinine, total protein, and urea for sows in the late gestation period (Table 6). The blood concentrations of BUN, creatinine, and urea for day 110 of gestation were greater ( $p < 0.05$ ) than those for day 70 of gestation. There were no interactions between arginine and the date effect in the blood concentrations for BUN, creatinine, total protein, and urea. Increasing the dietary arginine levels for sows during late gestation did not affect the milk composition of colostrum and milk (day 21 of lactation) in lactating sows (Table 7).

**Table 6.** Effects of arginine supplementation levels on blood profiles in late-gestating sows.

|                                   | Treatment <sup>1</sup> |       |       | SEM <sup>2</sup> | p-Value |      |       |       |       |              |
|-----------------------------------|------------------------|-------|-------|------------------|---------|------|-------|-------|-------|--------------|
|                                   | CON                    | ARG10 | ARG15 |                  | Diet    | Lin. | Quad. | Group | Date  | Group x Date |
| <b>Blood urea nitrogen, mg/dL</b> |                        |       |       |                  |         |      |       |       |       |              |
| Day 70                            | 12.5                   | 11.7  | 11.9  | 0.45             | 0.85    | 0.72 | 0.63  | 0.99  | 0.01  | 0.93         |
| Day 90                            | 11.3                   | 12.0  | 12.5  | 0.57             | 0.69    | 0.41 | 0.83  |       |       |              |
| Day 110                           | 14.3                   | 14.5  | 14.1  | 0.57             | 0.56    | 0.84 | 0.83  |       |       |              |
| <b>Creatinine, mg/dL</b>          |                        |       |       |                  |         |      |       |       |       |              |
| Day 70                            | 1.76                   | 1.74  | 1.95  | 0.091            | 0.59    | 0.41 | 0.70  | 0.48  | <0.01 | 0.89         |
| Day 90                            | 2.23                   | 2.43  | 2.39  | 0.104            | 0.74    | 0.63 | 0.55  |       |       |              |
| Day 110                           | 2.97                   | 2.90  | 3.18  | 0.097            | 0.48    | 0.32 | 0.50  |       |       |              |
| <b>Total protein, g/dL</b>        |                        |       |       |                  |         |      |       |       |       |              |
| Day 70                            | 6.85                   | 6.67  | 7.15  | 0.256            | 0.76    | 0.63 | 0.67  | 0.95  | 0.98  | 0.65         |
| Day 90                            | 6.86                   | 7.23  | 6.71  | 0.170            | 0.47    | 0.60 | 0.27  |       |       |              |
| Day 110                           | 6.93                   | 6.73  | 7.00  | 0.175            | 0.82    | 0.80 | 0.57  |       |       |              |
| <b>Urea, mg/dL</b>                |                        |       |       |                  |         |      |       |       |       |              |
| Day 70                            | 26.8                   | 25.0  | 25.5  | 0.97             | 0.85    | 0.73 | 0.62  | 0.99  | 0.01  | 0.93         |
| Day 90                            | 24.1                   | 25.6  | 26.7  | 1.22             | 0.69    | 0.42 | 0.83  |       |       |              |
| Day 110                           | 30.6                   | 31.1  | 30.1  | 1.22             | 0.96    | 0.84 | 0.83  |       |       |              |

**Table 7.** Effects of arginine supplementation levels during late gestation on milk composition in lactating sows.

|                         | Treatment <sup>1</sup> |       |       | SEM <sup>2</sup> | p-Value |      |       |
|-------------------------|------------------------|-------|-------|------------------|---------|------|-------|
|                         | CON                    | ARG10 | ARG15 |                  | Diet    | Lin. | Quad. |
| <b>Casein, %</b>        |                        |       |       |                  |         |      |       |
| Colostrum               | 6.57                   | 6.08  | 5.62  | 0.571            | 0.81    | 0.52 | 0.91  |
| Milk (21 days)          | 4.05                   | 4.32  | 4.09  | 0.119            | 0.63    | 0.98 | 0.34  |
| <b>Fat, %</b>           |                        |       |       |                  |         |      |       |
| Colostrum               | 6.68                   | 5.43  | 6.91  | 0.485            | 0.43    | 0.69 | 0.23  |
| Milk (21 days)          | 4.04                   | 6.82  | 6.32  | 0.368            | 0.13    | 0.23 | 0.10  |
| <b>Protein, %</b>       |                        |       |       |                  |         |      |       |
| Colostrum               | 8.28                   | 7.91  | 7.29  | 0.804            | 0.89    | 0.63 | 0.99  |
| Milk (21 days)          | 4.72                   | 4.77  | 4.61  | 0.115            | 0.86    | 0.68 | 0.72  |
| <b>Lactose, %</b>       |                        |       |       |                  |         |      |       |
| Colostrum               | 4.72                   | 4.35  | 4.31  | 0.187            | 0.65    | 0.44 | 0.60  |
| Milk (21 days)          | 5.77                   | 6.26  | 5.91  | 0.134            | 0.32    | 0.86 | 0.14  |
| <b>Total Solid, %</b>   |                        |       |       |                  |         |      |       |
| Colostrum               | 21.9                   | 20.1  | 20.9  | 0.76             | 0.67    | 0.67 | 0.43  |
| Milk (21 days)          | 16.9                   | 19.2  | 18.1  | 0.54             | 0.23    | 0.49 | 0.12  |
| <b>Solid not Fat, %</b> |                        |       |       |                  |         |      |       |
| Colostrum               | 13.3                   | 12.7  | 11.9  | 0.68             | 0.74    | 0.44 | 0.93  |
| Milk (21 days)          | 10.7                   | 11.0  | 10.6  | 0.17             | 0.64    | 0.66 | 0.41  |

## 4. Discussion

Supplementing sow diet with L-arginine during the late gestation did not influence the gain of BW and BF in sows for the late gestation period and loss of BW and BF in sows for the lactation period. This was in agreement with the result of Quesnel et al. [25], who reported that dietary supplementation with 25.5 g/day L-arginine from day 77 to farrowing did not affect the BW after farrowing and BF before farrowing of sows. Additionally, Bass et al. [26] reported that dietary supplementation with 1% L-arginine for gestating sows from day 93 to day 110 had no influence on BW loss, following farrowing or during lactation. However, they observed that improved late gestation BW gain of sows fed a diet with 1% L-arginine was revealed in parity 0 and parity 1, whereas, there was no

difference in late gestation BW gain compared to the control animals and 1% L-arginine-supplemented sows in parity 2+, implying that supplementation with 1% L-arginine partially met the arginine requirement in gilts or sows with parity 1 sows, not in sows with parity 2+. The difference between the current study and that of Bass et al. [26] with regard to the effect of dietary arginine in late gestation on the changes of BW or BF for sows could have been due to the difference in the parities of sows in the treatment group. Additionally, the lack of difference in the changes of BW and BF for sows during the whole experiment period could partly have been due to the same nitrogen content of diets among the dietary treatments, with the addition of L-Alanine as the isonitrogenous control.

Dietary effects of L-arginine was reported such that supplementation of L-arginine was found to increase nitric oxide, enhancing the delivery of essential nutrients from maternal to fetal blood [27][28][29], and increasing the polyamines necessary for embryogenesis and placental growth [30][31][32]. Since the nitric oxide and polyamines are important for angiogenesis and embryogenesis, arginine enhanced the growth of fetus and placenta development [32][33]. The litter size, including the number of total born or born alive was not affected by the dietary arginine effect in the present study, which was in agreement with the results from the studies of Quesnel et al. [25] and Nantapaitoon et al. [34], who reported that 1% L-arginine supplementation for late-gestating sows had no influence on the number of piglets at birth. However, previous studies reported that arginine supplementation enhanced conceptus survival or sow litter size in the period of early-gestation [19][35], or whole gestation [21][22][24]. The number of developing embryos is decided in early gestation, because most of the embryonic losses occurred during early gestation [36], and the dead embryos were not reabsorbed by the uterus after day 40 of gestation [37]. Thus, the differences among studies with regard to the effects of dietary arginine on the litter size of sows could have been due to differences in the supplemented period for the diet with L-arginine.

## References

1. Rooney, H.B.; O'driscoll, K.; O'doherty, J.V.; Lawlor, P.G. Effect of increasing dietary energy density during late gestation and lactation on sow performance, piglet vitality, and lifetime growth of offspring. *J. Anim. Sci.* 2020, 98, skz379, doi:10.1093/jas/skz379.
2. Quesnel, H.; Brossard, L.; Valancogne, A.; Quiniou, N. Influence of some sow characteristics on within-litter variation of piglet birth weight. *Animal* 2008, 2, 1842–1849.
3. De Vos, M.; Che, L.; Huygelen, V.; Willemen, S.; Michiels, J.; Van Cruchten, S.; Van Ginneken, C. Nutritional interventions to prevent and rear low-birthweight piglets. *J. Anim. Physiol. Anim. Nutr.* 2014, 98, 609–619.
4. Wang, J.; Feng, C.; Liu, T.; Shi, M.; Wu, G.; Bazer, F.W. Physiological alterations associated with intrauterine growth restriction in fetal pigs: Causes and insights for nutritional optimization. *Mol. Reprod. Dev.* 2017, 84, 897–904.
5. Tse, W.Y.; Town, S.C.; Murdoch, G.K.; Novak, S.; Dyck, M.K.; Putman, C.T.; Foxcroft, G.R.; Dixon, W.T. Uterine crowding in the sow affects litter sex ratio, placental development and embryonic

myogenin expression in early gestation. *Reprod. Feril. Dev.* 2008, 20, 497–504.

6. Rekiel, A.; Więcek, J.; Batorska, M.; Kulisiewicz, J. Effect of sow prolificacy and nutrition on pre and postnatal growth of progeny—a review. *Ann. Anim. Sci.* 2014, 14, 3–15.

7. Matheson, S.M.; Walling, G.A.; Edwards, S.A. Genetic selection against intrauterine growth retardation in piglets: A problem at the piglet level with a solution at the sow level. *Genet. Sel. Evol.* 2018, 50, 46, doi:10.1186/s12711-018-0417-7.

8. Kim, S.W.; Hurley, W.L.; Wu, G.; Ji, F. Ideal amino acid balance for sows during gestation and lactation. *J. Anim. Sci.* 2009, 87, E123–E132.

9. Le Cozler, Y.; Guyomarc'h, C.; Pichodo, X.; Quinio, P.Y.; Pellois, H. Factors associated with stillborn and mummified piglets in high-prolific sows. *Anim. Res.* 2002, 51, 261–268.

10. Ashworth, C.J. Late pregnancy: The effects of intra-uterine life on production traits in offspring. *Anim. Front.* 2013, 3, 62–67.

11. Noblet, J.; Close, W.H.; Heavens, R.P.; Brown, D. Studies on the energy metabolism of the pregnant sow: 1. Uterus and mammary tissue development. *Br. J. Nutr.* 1985, 53, 251–265.

12. Walker, B.; Young, B.A. Modelling the development of uterine components and sow body composition in response to nutrient intake during pregnancy. *Livest. Prod. Sci.* 1992, 30, 251–264.

13. Johnston, L.J.; Fogwell, R.L.; Weldon, W.C.; Ames, N.K.; Ullrey, D.E.; Miller, E.R. Relationship between body fat and postweaning interval to estrus in primiparous sows. *J. Anim. Sci.* 1989, 67, 943–950.

14. Dourmad, J.Y.; Etienne, M.; Prunier, A.; Noblet, J. The effect of energy and protein intake of sows on their longevity: A review. *Livest. Prod. Sci.* 1994, 40, 87–97.

15. Mullan, B.P.; Williams, I.H. The effect of body reserves at farrowing on the reproductive performance of first-litter sows. *Anim. Sci.* 1989, 48, 449–457.

16. Weldon, W.C.; Lewis, A.J.; Louis, G.F.; Kovar, J.L.; Giesemann, M.A.; Miller, P.S. Postpartum hypophagia in primiparous sows: I. Effects of gestation feeding level on feed intake, feeding behavior, and plasma metabolite concentrations during lactation. *J. Anim. Sci.* 1994, 72, 387–394.

17. Li, X.; Bazer, F.W.; Johnson, G.A.; Burghardt, R.C.; Erikson, D.W.; Frank, J.W.; Spencer, T.E.; Shinzato, I.; Wu, G. Dietary supplementation with 0.8% L-arginine between days 0 and 25 of gestation reduces litter size in gilts. *J. Nutr.* 2010, 140, 1111–1116.

18. Novak, S.; Moore, H.S.; Paradis, F.; Murdoch, G.; Dyck, M.K.; Dixon, W.T.; Foxcroft, G.R. Temporal candidate gene expression patterns in the sow placenta during early gestation and the effect of maternal L-arginine supplementation. *Soc. Reprod. Fertil. Suppl.* 2010, 66, 201–202.

19. Li, J.; Xia, H.; Yao, W.; Wang, T.; Li, J.; Piao, X.; Thacker, P.; Wu, G.; Wang, F. Effects of arginine supplementation during early gestation (day 1 to 30) on litter size and plasma metabolites in gilts and sows. *J. Anim. Sci.* 2015, **93**, 5291–5303.
20. Garbossa, C.A.P.; Júnior, F.C.; Silveira, H.; Faria, P.B.; Schinckel, A.P.; Abreu, M.L.T.; Cantarelli, V.S. Effects of ractopamine and arginine dietary supplementation for sows on growth performance and carcass quality of their progenies. *J. Anim. Sci.* 2015, **93**, 2872–2884.
21. Mateo, R.D.; Wu, G.; Bazer, F.W.; Park, J.C.; Shinzato, I.; Kim, S.W. Dietary L-arginine supplementation enhances the reproductive performance of gilts. *J. Nutr.* 2007, **137**, 652–656.
22. Che, L.; Yang, P.; Fang, Z.; Lin, Y.; Wu, D. Effects of dietary arginine supplementation on reproductive performance and immunity of sows. *Czech J. Anim. Sci.* 2013, **58**, 167–175.
23. Mateo, R.D.; Wu, G.; Moon, H.K.; Carroll, J.A.; Kim, S.W. Effects of dietary arginine supplementation during gestation and lactation on the performance of lactating primiparous sows and nursing piglets. *J. Anim. Sci.* 2008, **86**, 827–835.
24. Gao, K.; Jiang, Z.; Lin, Y.; Zheng, C.; Zhou, G.; Chen, F.; Yang, L.; Wu, G. Dietary L-arginine supplementation enhances placental growth and reproductive performance in sows. *Amino Acids* 2012, **42**, 2207–2214.
25. Quesnel, H.; Quiniou, N.; Roy, H.; Lottin, A.; Boulot, S.; Gondret, F. Supplying dextrose before insemination and arginine during the last third of pregnancy in sow diets: Effects on within-litter variation of piglet birth weight. *J. Anim. Sci.* 2014, **92**, 1445–1450.
26. Bass, B.E.; Bradley, C.L.; Johnson, Z.B.; Boyd, R.D.; Usry, J.L.; Maxwell, C.V.; Frank, J.W. Influence of dietary L-arginine supplementation to sows during late gestation on sow and litter performance during lactation. *Res. Ser.* 2011, **597**, 151–155.
27. Sladek, S.M.; Magness, R.R.; Conrad, K.P. Nitric oxide and pregnancy. *Am. J. Physiol.* 1997, **272**, R441–R463.
28. Fan, W.Q.; Smolich, J.J.; Wild, J.; Victor, Y.H.; Walker, A.M. Major vasodilator role for nitric oxide in the gastrointestinal circulation of the mid-gestation fetal lamb. *Pediatr. Res.* 1998, **44**, 344–350.
29. Gardner, P.R.; Martin, L.A.; Hall, D.; Gardner, A.M. Dioxygen-dependent metabolism of nitric oxide in mammalian cells. *Free Radical Biol. Med.* 2001, **31**, 191–204.
30. Wu, G.; Morris, S.M. Arginine metabolism: Nitric oxide and beyond. *Biochem. J.* 1998, **336**, 1–17.
31. Kim, S.W.; Mateo, R.D.; Yin, Y.L.; Wu, G. Functional amino acids and fatty acids for enhancing production performance of sows and piglets. *Asian-Australas. J. Anim. Sci.* 2007, **20**, 295–306.
32. Wu, G.; Jaeger, L.A.; Bazer, F.W.; Rhoads, J.M. Arginine deficiency in preterm infants: Biochemical mechanisms and nutritional implications. *J. Nutr. Biochem.* 2004, **15**, 442–451.

33. Reynolds, L.P.; Redmer, D.A. Angiogenesis in the placenta. *Biol. Reprod.* 2001, 64, 1033–1040.
34. Nuntapaitoon, M.; Muns, R.; Theil, P.K.; Tummaruk, P. L-arginine supplementation in sow diet during late gestation decrease stillborn piglet, increase piglet birth weight and increase immunoglobulin G concentration in colostrum. *Theriogenology* 2018, 121, 27–34.
35. Bérard, J.; Bee, G. Effects of dietary L-arginine supplementation to gilts during early gestation on foetal survival, growth and myofiber formation. *Animal* 2010, 4, 1680–1687.
36. Foxcroft, G.R.; Dixon, W.T.; Novak, S.; Putman, C.T.; Town, S.C.; Vinsky, M.D.A. The biological basis for prenatal programming of postnatal performance in pigs. *J. Anim. Sci.* 2006, 84, E105–E112.
37. Town, S.C.; Patterson, J.L.; Pereira, C.Z.; Gourley, G.; Foxcroft, G.R. Embryonic and fetal development in a commercial dam-line genotype. *Anim. Reprod. Sci.* 2005, 85, 301–316.

Retrieved from <https://encyclopedia.pub/entry/history/show/3662>