

EDN3 Gene

Subjects: **Genetics & Heredity**

Contributor: Vivi Li

Endothelin 3: The EDN3 gene provides instructions for making a protein called endothelin 3.

genes

1. Normal Function

Proteins in the endothelin family are produced in various cells and tissues, where they are involved in the development and function of blood vessels, the production of certain hormones, and the stimulation of cell growth and division.

Endothelin 3 functions by interacting with another protein, endothelin receptor type B (produced from the *EDNRB* gene), on the surface of cells. During early development before birth, endothelin 3 and endothelin receptor type B together play an important role in neural crest cells. These cells migrate from the developing spinal cord to specific regions in the embryo, where they give rise to many different types of cells. In particular, endothelin 3 and its receptor are essential for the formation of nerves in the intestine (enteric nerves) and for the production of specialized cells called melanocytes. Melanocytes produce melanin, a pigment that contributes to skin, hair, and eye color. Melanin is also involved in the normal function of the inner ear.

2. Health Conditions Related to Genetic Changes

2.1 Hirschsprung Disease

About 10 mutations in the *EDN3* gene have been found to cause Hirschsprung disease, a disorder that causes severe constipation or blockage of the intestine. Although Hirschsprung disease is a feature of another disorder called Waardenburg syndrome type IV (described below), *EDN3* gene mutations can also cause Hirschsprung disease in people without Waardenburg syndrome. These mutations change one DNA building block (nucleotide) or insert an additional nucleotide in the gene. Changes in the *EDN3* gene disrupt the normal function of endothelin 3, preventing it from playing its usual role in the development of enteric nerves. As a result, these cells do not form normally during embryonic development. A lack of enteric nerves prevents stool from being moved through the intestine normally, leading to severe constipation or intestinal blockage.

2.2 Waardenburg Syndrome

At least five mutations in the *EDN3* gene have been identified in people with Waardenburg syndrome type IV (also known as Waardenburg-Shah syndrome). This type of Waardenburg syndrome is characterized by changes in skin, hair, and eye coloring; hearing loss; and Hirschsprung disease. *EDN3* mutations change single nucleotides in the gene, preventing the production of a functional endothelin 3 protein. Because active endothelin 3 is necessary for the formation of enteric nerves and melanocytes, these cell types do not form normally during embryonic development. Missing enteric nerves in certain parts of the intestine cause the signs and symptoms of Hirschsprung disease. A lack of melanocytes affects the coloring of skin, hair, and eyes and causes the hearing loss characteristic of Waardenburg syndrome.

3. Other Names for This Gene

- EDN3_HUMAN
- endothelin 3 precursor
- ET3
- HSCR4
- PPET3
- Preproendothelin-3
- RP4-614C15.1
- WS4B

References

1. Baynash AG, Hosoda K, Giard A, Richardson JA, Emoto N, Hammer RE, Yanagisawa M. Interaction of endothelin-3 with endothelin-B receptor is essential for development of epidermal melanocytes and enteric neurons. *Cell*. 1994 Dec 30;79(7):1277-85.
2. Duan XL, Zhang XS, Li GW. Clinical relationship between EDN-3 gene, EDNRB gene and Hirschsprung's disease. *World J Gastroenterol*. 2003 Dec;9(12):2839-42.
3. Gazquez E, Watanabe Y, Broders-Bondon F, Paul-Gilloteaux P, Heysch J, Baral V, Bondurand N, Dufour S. Endothelin-3 stimulates cell adhesion and cooperates with β 1-integrins during enteric nervous system ontogenesis. *Sci Rep*. 2016 Dec 1;6:37877. doi: 10.1038/srep37877.

4. Heuckeroth RO. Finding your way to the end: a tale of GDNF and endothelin-3. *Neuron*. 2003 Dec 4;40(5):871-3. Review.
5. Kapur RP, Gershon MD, Milla PJ, Pachnis V. The influence of Hox genes and three intercellular signalling pathways on enteric neuromuscular development. *Neurogastroenterol Motil*. 2004 Apr;16 Suppl 1:8-13. Review.
6. Pingault V, Bondurand N, Lemort N, Sancandi M, Ceccherini I, Hugot JP, JoukPS, Goossens M. A heterozygous endothelin 3 mutation in Waardenburg-Hirschsprung disease: is there a dosage effect of EDN3/EDNRB gene mutations on neurocristopathy phenotypes? *J Med Genet*. 2001 Mar;38(3):205-9.
7. Shin MK, Levorse JM, Ingram RS, Tilghman SM. The temporal requirement for endothelin receptor-B signalling during neural crest development. *Nature*. 1999 Dec 2;402(6761):496-501.
8. Svensson PJ, Von Tell D, Molander ML, Anvret M, Nordenskjöld A. A heterozygous frameshift mutation in the endothelin-3 (EDN-3) gene in isolated Hirschsprung's disease. *Pediatr Res*. 1999 May;45(5 Pt 1):714-7.
9. Sánchez-Mejías A, Fernández RM, López-Alonso M, Antiñolo G, Borrego S. New roles of EDNRB and EDN3 in the pathogenesis of Hirschsprung disease. *Genet Med*. 2010 Jan;12(1):39-43. doi: 10.1097/GIM.0b013e3181c371b0.

Retrieved from <https://encyclopedia.pub/entry/history/show/12372>