

EIF2B4 Gene

Subjects: Genetics & Heredity

Contributor: Vivi Li

Eukaryotic translation initiation factor 2B subunit delta

Keywords: genes

1. Normal Function

The *EIF2B4* gene provides instructions for making one of five parts of a protein called eIF2B, specifically the delta subunit of this protein. The eIF2B protein helps regulate overall protein production (synthesis) in the cell by interacting with another protein, eIF2. The eIF2 protein is called an initiation factor because it is involved in starting (initiating) protein synthesis.

Under some conditions, eIF2B increases protein synthesis by helping to recycle molecules called GTP, which carry energy to the initiation factor. Under other conditions, it slows protein synthesis by binding tightly to the initiation factor, which converts the eIF2B protein into an inactive form and prevents recycling of GTP.

Proper regulation of protein synthesis is vital for ensuring that the correct levels of protein are available for the cell to cope with changing conditions. For example, cells must synthesize protein much faster if they are multiplying than if they are in a resting state.

2. Health Conditions Related to Genetic Changes

2.1 Leukoencephalopathy with Vanishing White Matter

Mutations in the *EIF2B4* gene have been identified in a few people with leukoencephalopathy with vanishing white matter, including some affected females with a variant of the disorder in which the neurological features are accompanied by ovarian failure (ovarioleukodystrophy). These mutations cause partial loss of eIF2B function. Impairment of eIF2B function makes it more difficult for the body's cells to regulate protein synthesis and deal with changing conditions and stress. Researchers believe that cells in the white matter (nerve fibers covered by a fatty substance called myelin that insulates and protects nerves) may be particularly affected by an abnormal response to stress, resulting in the signs and symptoms of leukoencephalopathy with vanishing white matter.

3. Other Names for This Gene

- DKFZp586J0119
 - EI2BD_HUMAN
 - EIF-2B
 - EIF2B
 - EIF2Bdelta
 - eukaryotic translation initiation factor 2B, subunit 4 (delta, 67kD)
 - eukaryotic translation initiation factor 2B, subunit 4 delta
 - eukaryotic translation initiation factor 2B, subunit 4 delta, 67kDa
 - translation initiation factor eIF-2b delta subunit
-

References

1. Dietrich J, Lacagnina M, Gass D, Richfield E, Mayer-Pröschel M, Noble M, Torres C, Pröschel C. EIF2B5 mutations compromise GFAP+ astrocyte generation in vanishing white matter leukodystrophy. *Nat Med*. 2005 Mar;11(3):277-83.
2. Eureka Bioscience: Mechanism of Translation Initiation in Eukaryotes
3. Fogli A, Boespflug-Tanguy O. The large spectrum of eIF2B-related diseases. *Biochem Soc Trans*. 2006 Feb;34(Pt 1):22-9. Review.
4. Fogli A, Schiffmann R, Hugendubler L, Combes P, Bertini E, Rodriguez D, Kimball SR, Boespflug-Tanguy O. Decreased guanine nucleotide exchange factor activity in eIF2B-mutated patients. *Eur J Hum Genet*. 2004 Jul;12(7):561-6.
5. Li W, Wang X, Van Der Knaap MS, Proud CG. Mutations linked to leukoencephalopathy with vanishing white matter impair the function of the eukaryotic initiation factor 2B complex in diverse ways. *Mol Cell Biol*. 2004 Apr;24(8):3295-306.
6. *Molecular Biology of the Cell* (fourth edition, 2002): The Phosphorylation of an Initiation Factor Globally Regulates Protein Synthesis
7. Pavitt GD. eIF2B, a mediator of general and gene-specific translational control. *Biochem Soc Trans*. 2005 Dec;33(Pt 6):1487-92. Review.
8. Pronk JC, van Kollenburg B, Scheper GC, van der Knaap MS. Vanishing white matter disease: a review with focus on its genetics. *Ment Retard Dev Disabil Res Rev*. 2006;12(2):123-8. Review.
9. Scali O, Di Perri C, Federico A. The spectrum of mutations for the diagnosis of vanishing white matter disease. *Neurol Sci*. 2006 Sep;27(4):271-7. Review.
10. Scheper GC, Proud CG, van der Knaap MS. Defective translation initiation causes vanishing of cerebral white matter. *Trends Mol Med*. 2006 Apr;12(4):159-66.
11. van der Voorn JP, van Kollenburg B, Bertrand G, Van Haren K, Scheper GC, Powers JM, van der Knaap MS. The unfolded protein response in vanishing white matter disease. *J Neuropathol Exp Neurol*. 2005 Sep;64(9):770-5.
12. van Kollenburg B, van Dijk J, Garbern J, Thomas AA, Scheper GC, Powers JM, van der Knaap MS. Glia-specific activation of all pathways of the unfolded protein response in vanishing white matter disease. *J Neuropathol Exp Neurol*. 2006 Jul;65(7):707-15.

Retrieved from <https://encyclopedia.pub/entry/history/show/12377>