

Older Adult Fall Prevention

Subjects: Nursing

Contributor: María del Carmen Miranda-Duro

Falls are the second leading cause of accidental or non-intentional deaths worldwide and are the most common problem as people age. The primary purpose of addressing falls is to detect, prevent, treat, and reduce their incidence and consequences. Previous studies identified that multifactorial programs, an interprofessional team, and assistive technology are required to address falls in older adults effectively. Accordingly, the research question is as follows: what are the scope, type of studies, and approaches and strategies to fall risk using technology in the existing occupational therapy literature regarding interventions to address the effects of falls in older adults on daily living?

Keywords: older adults ; falls ; occupation ; occupational therapy ; scoping review ; smart home technology ; telehealth ; exergames ; 3D-application tool

1. Introduction

Fall prevention is focused on injuries or complications that occur because of falling^{[1][2]}. Numerous studies ^[3] have demonstrated that many falls can be prevented through adequate assessment and intervention. Some of the most common and effective interventions include^[3] gait stabilizing footwear, vitamin D, dietary supplements, medication adaptation, multiple interventions, multifactorial interventions, assistive device training, cognitive monitoring and intervention, environmental modification, and family and caregiver training^{[4][5][6][7][8][9]}.

Different authors have also suggested that multifactorial programs are useful for preventing and reducing falls because of the complexity in the types of falls ^[10]. According to the World Health Organization (WHO), the risk of falls increases when multiple risk factors are present^[11]. These multifactorial programs^[12] involve a combination of exercise (focus on strength, balance and mobility), and other options such as individualized and comprehensive fall risk assessment about the person and, their environment and education on fall prevention. The GeriaTIC project is an example of using multifactorial interventions for fall prevention. A study by Close and colleagues highlighted that an interprofessional approach to this high-risk population can significantly decrease the risk of further falls and limit functional impairment ^[13].

The effectiveness of these multifactorial programs is also based on the need to have an interprofessional team for fall prevention and treatment and geriatric approaches^[14]. This includes a physician, a neurologist, a nurse, a psychiatrist, a physical therapist, and an occupational therapist. The latter is part of the non-pharmacological intervention ^{[15][16]}. Elliot and colleagues ^[17], in a systematic review, classified occupational therapy interventions on older adults' falls into three types of intervention that are shown in [Table 1](#).

Table 1. Summary of occupational therapy interventions for falls in older adults.

Summary of Occupational Therapy Interventions for Falls in Older Adults

Single component

(includes only one of the following components)

- Exercise
- Home safety assessment
- Education about falls' prevention

Example: Lifestyle Integrated Functional Exercise study [37]

1st option: educational components as:

- Feet or footwear risk
- Energy conservation strategies
- Safe assistive device use, home modification

Multicomponent intervention

(includes exercise and one of the following options)

- Fall recovery
- Medication routines
- Nutrition and hydration
- Relaxation stress management

2nd option: home modification with other fall prevention intervention

Example: Minimally Supervised Multimodal Exercise to Reduce Falls Risk in Economically and Educationally Disadvantaged Older Adults [38]

- Fall risk
- Environment, education, and group activities
- Activities of daily living

Multifactorial intervention

(include the complex assessment of different components)

- Assistive devices
- Self-efficacy or fear of falling

Example: A single home visit by an occupational therapist reduces the risk of falling after hip fracture in elderly women: A quasi-randomized controlled trial [39]

Population-based fall prevention

(includes strategies implemented across whole communities, two different types)

· Existing effective population-based fall prevention programs

· Other population-based multicomponent interventions

Example: Stepping On -Translating a Fall Prevention Intervention Into Practice: A Randomized Community Trial [40]

2. Methodology

The authors conducted a scoping review in January 2020. The research questions that we aimed to answer were the following: what are the scope, type of studies, and approaches and strategies to fall risk using technology in the existing occupational therapy literature regarding interventions to address the effects of falls in older adults on daily living? Two approaches guided the present scoping review. On the one hand, the Arksey and O'Malley^[18] five-stage framework was used, which includes stage 1 establishment of the research question; stage 2 identification of pertinent studies and choice of studies, stage 3 study selection as explained in [Figure 1](#); and, as shown in the Results section, the stage 4 charting the data and stage 5 mapping the data and collating, summarizing, and reporting the findings. On the other hand, this scoping review also follows the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) (see [Table S1](#) ^[19]). In accordance with the aim of a scoping review, a quality appraisal is not required, as opposed to Systematic reviews and Meta-Analyses^[20].




Figure 1. Flow diagram of the Scoping Review process.

3. Bibliometric Characteristics

In total, we screened 74 documents, with 12 studies meeting the eligibility criteria (see [Figure 1](#)). The selected literature was published between 2012 and 2020. The first study to report the use of technology in an occupational therapy falls approach was published in 2012^[21].

The studies were conducted in the USA ($n = 4$)^{[21][22][23][24]}; Europe ($n = 6$)^{[25][26][27][28][29][30]}, specifically in Scotland ($n = 1$)^[25], Sweden ($n = 1$)^[26], the UK ($n = 3$)^{[27][29][30]}, and Belgium ($n = 1$)^[28]; and other countries such as Tunisia ($n = 1$)^[31] and Australia ($n = 1$)^[32].

The journals in which the articles were published were mainly those focusing on occupational therapy ($n = 6$)^{[21][22][23][24][25]}, particularly for the first publications of studies on the topic, and informatics and technological journals ($n = 6$)^{[27][28][29][30][31][32]}, which contained more of the recent studies.

The types of study were reviews—a systematic review ($n = 1$)^[21] and a critical review ($n = 1$)^[22] and original articles ($n = 10$)^{[22][23][24][30][31]}. Within the original articles, there were different research designs. Quantitative approaches were used, including a descriptive study ($n = 1$)^[29], a case study ($n = 2$)^{[23][24]}, and an experimental study ($n = 1$)^[31]. The qualitative approaches used were qualitative research ($n = 2$)^{[26][32]} and a mixed-methods study ($n = 2$)^{[28][29]}.

Table 2. Summary of data extracted from the 12 selected studies.

Author(s), Year, [Reference]	Authors' Affiliation	Journal of Publication	Type of Study and Purpose	Sample Characteristics (Size, Age, % of Female, Setting, and Others)	Technology Used	Main Findings
Chase, C.A.; Mann, K.; Wasek, S. and Arberman, M. 2012 [58]	Western Michigan University; Rehabilitation Hospital of Indiana; Ingham County Medical Center and Rehabilitation; University of Buffalo. U.S.A.	Am. J. Occup. Ther.	This systematic review aims to synthesize existing literature about the effect of home modification as both a separate intervention and a component of several fall prevention programs.	n = 33 studies, including a total of 31 randomized controlled trials Age = is focused on older adults (not specify ages) Setting = community- dwelling older adults Female = Number of females not specified	Commercially available smart home technology: operate lights, appliances, door, and windows for frail older adults living alone.	The results contribute to evidence-based practice for occupational therapy practitioners working with older adults in community- based settings and reinforce the importance of the role of occupational therapy in the home and community.

Stewart, L. and McKinstry, B. 2012 [62]	Astley Ainslie Hospital; Edinburgh University, Edinburgh	Br. J. Occup. Ther.	<p>This critical review aimed to evaluate the association between older people's fear of falling and the use of telecare and whether telecare could reduce this fear.</p>	<p>n = 10 studies, including randomized controlled trials, a cohort study, two qualitative studies, a case study, and surveys</p> <p>Age = older adults over 60 years old</p> <p>Setting = older adults from public-assisted housing, day centers, and community living</p> <p>Female = Number of females not specified</p>	<p>Telecare: this is understood as the remote or enhanced delivery of health and social services to people in their own homes using telecommunications and computerized systems.</p>	<p>Telecare's contribution to supporting an aging population at home for longer is becoming increasingly recognized by health and social care services worldwide. However, this critical review identified that few studies are investigating older people's views and the use of telecare in the domain of occupational therapy.</p>
Horowitz, B.; Nochajski, S.M. and Schweitzer, H.A. 2013 [59]	York College-CUNY; University of Buffalo; State University of New York. U.S.A.	Occup. Ther. Health Care	<p>This case study focused on the development and pilot-testing of the Home Safety Self-Assessment Tool (H.S.S.A.T.), a new home assessment tool designed for use by older adults to promote home safety and aging.</p>	<p>n= 28 older adults</p> <p>Age = between 69 to 87 years old</p> <p>Female = 68%</p> <p>Setting = community dwelling</p>	<p>Videos: developed with instructions to install home modifications to prevent falls. Several tools were included to analyze the risks associated with older adults' homes in the project.</p>	<p>The results suggest the tool may assist older adults in identifying environmental factors related to falls and facilitate their ability to age in place.</p>

Charness, n. 2014 [60]	Florida State University. U.S.A.	Occup. Ther. Health Care	This case study was used to illustrate telehealth as one important tool to improve the efficiency of healthcare delivery.	n = 9 older adults Age = older than 75 years Female = Number of females not specified Setting = community- dwelling	Telehealth system: consisting of a watch for a factor sensor system that monitors temperature with an analog display, an emergency button, and an accelerometer to provide information about activity and to monitor falls.	Telehealth systems can provide potentially important support for persons to live independently longer through automated monitoring. The purpose of this study was to find a cost-effective telehealth technology.
Bianco, M.L.; Pedell, S. and Renda, G. 2016 [69]	Swinburne University. Australia	A.C.M.	This qualitative research study assessed the perceptions of ten older adults on an augmented reality tool.	n= 10 older adults Age = between 69 and 92 years old with a mean age of 79.1 years Female = 60% Setting = not specified	An augmented reality application prototype on an iPad. The application is a 3D model library bank of typical and novel home modifications. Professionals can access this modification bank to superimpose a proposed recommendation into their client's home environment for evaluation and discussion.	The findings indicate that many older adults welcome augmented reality as a design and communication medium. It can be used as a bridging mechanism to increase the person- centeredness of fall prevention services.

Glannfjord, F.; Hemmingsson, H. and Larsson, A. 2016 [63]	Linköping University, Sweden	Scand. J. Occup. Ther.	This qualitative (phenomenology) study examined how older adults perceive the Wii, namely the Wii sports bowling game, in an activity group.	n= between 10 and 12		The Nintendo Wii sports bowling game, with the Wii controller, was compared with real-life bowling.	The Wii was found to be an enjoyable and social activity. The interviewers looked forward to participating in the activity each week. Participants felt like they were really bowling in this virtual activity, and positive differences between regular bowling and virtual bowling were identified; virtual options were identified as being easier.
				Age = mean age of 78, between 64 and 98 years old	Female = Number of females not specified		
Hamm, J.; Money, G.A. 2017 [64]	Brunel University, London South Bank University, U.K.	Health Informatics J.	This mixed-study explored occupational therapists' perceptions of an early-stage, three-dimensional measurement aid prototype, which provides enhanced assistive equipment provision process guidance to clinicians.	Setting = Activity center for elderly people	n= 10 occupational therapists	A 3D measurement aid prototype (3D-MAP) application, using 3D visualization technology was deployed on a tablet, mobile phone, or laptop. This was based on the five most commonly measured items with the Assistive Equipment Provision Process tool (bed, bath, toilet, chair, and stairs).	The results show that occupational therapists considered that the 3D-MAP application could effectively augment existing 2D diagrams and deliver numerous benefits.
					Female = 100% A ten pounds voucher was offered 2 to 31 years of experience. The occupational therapists' experience was in adults, social services, surgical rehabilitation, neurology, rehabilitation, and social services.		

Lemmens, R.; Gielen, C. and Spooren, A. 2017 [65]	P.X.L. University College. Belgium	Stud. Health. Technol. Inform.	This was a qualitative study about developing a screening tool to enable occupational therapists to assess people's home environments to facilitate independent living.	1st phase:	The Obstacle tool is a digitalized version using the mind maps and the paper version results for tablets. It has a version that	
				n = 16 older adults in their homes		
				Aged over 65 years old	can be used by health professionals or informal caregivers too. The digital version includes (1)	The Obstacle tool was developed and judged to be very useful by occupational therapists. It was highlighted that the Obstacle is adapted for use for persons with dementia and a mini-obstacle tool is under construction and will be digitalized to be available for clients and their informal caregivers. The next step is to make the tool accessible to everybody.
				2nd phase:	the possibility of structuring the screening by adding the rooms in the order of preference for occupational therapists, (2) registration of problems/scores, (3) the addition of photos to the screening, (4) a better overview than in the paper version, (5) the option to store and save data and make a back-up, and (6) connection with the application H-OPP (a digital coach for occupational therapists).	
				n= 31 older adults		
				Aged over 65 years old		
				3rd phase:		
				n= 5 older adults, 5 informal caregivers, 5 professional caregivers		
				Setting = community-dwelling		
				The number of females was not specified		

4.1. A Description of the Type of Participants in Study

This theme was about the types of participants included in the studies (i.e., older adult(s) or occupational therapist(s)), the age of the participants, the percentage of females in the sample, the environment in which the study was conducted (i.e., community-dwelling or an institution), and other interesting details mentioned.

The reviews included in this study did not provide some of the participants' characteristics (see [Table 3](#) details of the participants). In a review by Chase and colleagues^[24], a total of thirty-three studies were included, all randomized controlled trials, while in the review by Stewart and colleagues study^[25], ten studies were included and these were different types of studies.

The original articles [22][23][24][25][26][27][28][29][30][31][32] included 614 older adults, 15 occupational therapists, and 5 informal caregivers. The samples size of the studies included 1 ($n = 1$) [31] to 445 ($n = 1$) older adults [24], 1 ($n = 7$) [22][23][24][29][30][31][32] to 10 ($n = 1$) types of occupational therapists, and 1 ($n = 9$) to 5 ($n = 1$) informal caregivers. There was a large age range age of older adults from 50 years old (study aimed to a maximum of 98 years old) ($n = 1$) [26]. These people were recruited from the community ($n = 8$) [22][23][24][29], an activity center for older adults ($n = 1$) [25], or people attending a gym group on a university campus ($n = 1$) [30]. Regarding the percentage of females, all studies included more females than males, with a percentage of females of between 60 ($n = 2$) [26][32] and 68% ($n = 2$) [22][24]. One hundred percent of the occupational therapists were female [27].

All participants volunteered to participate in the different studies, but in the study by Hamm and colleagues, the occupational therapists received a ten pound voucher [27].

4.2. Type of Intervention and Approach to Falls

This theme included the type of intervention if the study was about a specific risk factor related to falls (extrinsic to the individual, or intrinsic, related to the person, and the aging process) and the type of approach to falls (detection, prevention, or treatment).

Four types of intervention were identified: “home modifications” ($n = 5$) [21][23][24][25], “exercise” ($n = 2$) [25][31], and “educational” ($n = 2$) [22][30]. We considered single-component interventions and focused on fall detection, prevention, and treatment. Interventions that focused on home modification addressed extrinsic factors such as bathroom modifications (i.e., bath-tower shower), appropriate chair height, and indications to include space to move, among others. Assistive technology interventions addressed extrinsic factors that can affect the safety of the person in the home. This included the use of telecare, emergency alarms, and fall detectors (i.e., pendant alarms). Exercise was used as a strategy to address intrinsic factors related to physical condition and the use of educational interventions in these studies was based on extrinsic factors, which consisted of the types of modification the person has to do to be safe at home.

4.3. Type of Technology Used

This theme was about the types of technology addressed in the studies. The technologies used in the studies were classified as software developments, telehealth, multimedia materials, and commercial and technological devices. Software developments ($n = 5$) included augmented reality applications [32], 3D measurement aid prototype applications [29][28], a digital version of the Obstacle Tool [24], and a Falls Sensei 3D game. The use of telehealth was integrated into telecare to reduce the fear of falling [25], and the system was integrated into a watch to provide a factor sensor system to monitor temperature with an analog display, an emergency button, and an accelerometer [23]. The multimedia materials included different videos to identify environmental fall factors [22]. The commercial and technological devices included smart home technology to operate lights, appliances, doors, and windows [21], and these included the use of Kinect [31] and Nintendo Wii [26].

Table 3 shows the relationships between Section 3.2.2 (Type of intervention and approach to falls) and Section 3.2.3 (Type of technology used, and the compatibility of the technology used with the technological devices: Tablet, iPad, computer, laptop, Xbox, Nintendo Wii, and mobile phone).

Table 3. Relationships between Section 3.2.2. and Section 3.2.3.

Types of Technology	Technology Used	Compatibility	Intervention	Falls Risk
Software developments	Augmented reality application	Setting = Not specified	Home modifications	based 2D measurement guide equivalent.
	3D measurement aid prototype application	Tablet, mobile phone, or laptop		Extrinsic factors
	Digital version of Obstacle Tool	Tablet		
	Falls Sensei 3D game	Computer		
Telehealth	Telehealth system	Not applied	Assistive technology	Extrinsic factors
	Telecare			
	Smart home technology			

Types of Technology	Technology Used	This mixed-	Compatibility	Intervention	Falls Risk
Multimedia materials	Videos	method study used Falls Sensei 3D to	n= 15 Computer, laptop, mobile phone or Tablet	Educational Falls Sensei 3D	Extrinsic factors This study offers a promising
Commercial and technological devices	Kinect with Xbox and Nintendo Wii	evaluate the overall game	Kinect with Xbox and Nintendo Wii	game is a first- person 3D exploration game	extrinsic factors using
5. Discussion		usability from an older adult perspective and	Age = between 50 and 80 years	with four levels that correspond to four key living areas within the home.	challenging games to address extrinsic
	Money, A.G. This study presents the first scoping review of occupational therapy interventions, to address older adults' falls using technology. The objectives defined focused on exploring the literature about the topic to determine the type of studies conducted, where these studies have usually been published, and the approaches and strategies used to reduce fall risk by occupational therapists using technology.	to explore older adults' perceptions of using Falls Sensei, the factors affecting the adoption of this application, and the extent to which the modification of digital prevention- related behavior can occur as a consequence of playing the falls prevention game	old participants old Setting = adults attending a 50s gym group on a university campus Female = 60%	bedroom, lounge, and stairs. The appearance was the same raised awareness of home hazard detection, but further research is needed to draw comparisons with established interventions.	factors in fall risk reduction. Data analysis triangulation suggests that the game raised awareness of home hazard detection, but further research is needed to draw comparisons with established interventions.
	Atwal, A.; Boyce, E.; Gaber, S.; Wineault, S.	Decis. Mak.		GameObject, which contains 3D Models and associated approaches are at each game level	
	U.K.				
	Alexandrou, K. 2019 [67]				

4. Thematic Categories

Regarding the type of intervention used to address falls in older adults through occupational therapy, only single-component interventions were used [17], even though different authors have suggested that multifactorial programs help to prevent and reduce falls because of their complexity [10]. None of the studies included an interprofessional team, despite

its importance. The effectiveness of the multifactorial programs is also due to the use of an interprofessional team for fall prevention and treatment^[14].

In terms of multifactorial programs^[10], these involved exercise, as was done in two studies included in this review^{[26][31]}; individualized and comprehensive fall risk assessment about the environment of an older adult, as done in a few studies mentioned in this review^{[22][27][28][29][30][31][32]}; and education on fall prevention, as done in two studies^{[22][31]}. Furthermore, any intervention includes occupational therapy home visits^[45], which can be an essential aspect to include, especially in the cases of home modification^{[22][27][28][29][30][31][32]} and assistive technology^{[21][22][23][24][25]}. The studies that integrated aging in place focused on assistive technology and home modifications because these factors are widely acknowledged as being the primary and preferred interventions during ageing^{[46][47]}. Understanding how older people's needs contribute to improving their quality of life, which is affected after a fall, is necessary ^{[48][49][50][11][51][52][53]}.

A previous systematic review explored the cost-effectiveness of several occupational therapy interventions for older people, concluding that they are useful and cost-effective compared with standard care or other therapies^[54]. In this way, socioeconomic impact is one of the consequences of falls^[8]. However, the results were not focused on aspects related to socioeconomic impact.

Our results reinforce the idea that home modifications, assistive technology, and educational interventions can address extrinsic factors, particularly environmental factors, and exercise can address intrinsic factors. This is in accordance with previous studies about the use of occupational therapy interventions to address fall risk^{[56][57][58][59]}.

Although, as mentioned above, some examples of assistive technology used to prevent falls in older adults include video-monitoring, health monitoring, electronic sensors, and fall detectors^[55] these were not included in the studies mentioned in this review. Moreover, regarding fall interventions globally, there is more focus on exercise options and an extensive range of technologies from virtual reality ^[60] to wearables^[61] that were not included in these studies. Primarily, virtual reality interventions are used in occupational therapy, for example, in children ^[60].

As a result of this review, it is suggested that researchers in this field perform more studies that include the latest technology in the field of falls, so that more studies with a higher level of evidence exist. Interprofessional and multifactorial interventions should be integrated.

Limitations

The present scoping review has few limitations since all those publications related to the topic have been included; regarding the language were included articles in English, Spanish, and Portuguese; and regarding the type of study, any of them were included. The first limitation maybe not including other languages or other databases in the search process. However, we included databases of occupational therapy and socio-health care. Regarding the searches and the inclusion and exclusion of studies, it was carried out by one of the authors, which may be a limitation. In spite that the researcher used a structured process, some data may have been omitted or excluded. As future research, it would be important to integrate more researchers into this process.

References

1. Burton, E.; Lewin, G.; O'Connell, H.; Hill, K. Falls prevention in community care: 10 years on. *Clin. Interv. Aging* 2018, 13, 261–269, doi:10.2147/CIA.S153687.
2. Lach, H.W.; Noimontree, W. Fall prevention among community-dwelling older adults: Current guidelines and older adult responses. *J. Gerontol. Nurs.* 2018, 44, 21–29, doi:10.3928/00989134-20180808-06.
3. Sherrington, C.; Michaleff, Z.A.; Fairhall, N.; Paul, S.S.; Tiedemann, A.; Whitney, J.; Cumming, R.G.; Herbert, R.D.; Close, J.C.T.; Lord, S.R. Exercise to prevent falls in older adults: An updated systematic review and meta-analysis. *Br. J. Sports Med.* 2017, 51, 1749–1757, doi:10.1136/bjsports-2016-096547.
4. Cumming, R.G.; Thomas, M.; Ot, G.D. Home visits by an occupational therapist for assessment and modification of environmental hazards: a randomized trial of falls prevention. *J. Am. Geriatr. Soc.* 1999, 44, 1397–1402, doi: 10.1111/j.1532-5415.1999.tb01556.x.
5. LaStayo, P.C.; Ewy, G.A.; Pierotti, D.D.; Johns, R.K.; Lindstedt, S. The positive effects of negative work: Increased muscle strength and decreased fall risk in a frail elderly population. *J. Gerontol.* 2003, 58, 419–424, doi:10.1093/gerona/58.5.M419.

6. Ejupi, A.; Lord, S.R.; Delbaere, K. New methods for fall risk prediction. *Curr. Opin. Clin. Nutr. Metab. Care* 2014, 17, 407–411, doi:10.1097/MCO.0000000000000081.
7. Layton, N.; Clarke, A.; Pennock, J. "Doing with not doing for": A paradigm shift in home care services and what it means for occupational therapy. *Aust. Occup. Ther. J.* 2014, 61, 11–13, doi:10.1111/1440-1630.12184.
8. Balzer, K.; Bremer, M.; Schramm, S.; Lühmann, D.; Raspe, H. Falls prevention for the elderly. *GMS Health Technol. Assess.* 2012, 8, Doc01, doi:10.3205/hta000099.
9. Cameron, I.D.; Dyer, S.M.; Panagoda, C.E.; Murray, G.R.; Hill, K.D.; Cumming, R.G.; Kerse, N. Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database Syst. Rev.* 2018, doi:10.1002/14651858.CD005465.pub4.
10. Morello, R.T.; Soh, S.-E.; Behm, K.; Egan, A.; Ayton, D.; Hill, K.; Flicker, L.; Etherton-Beer, C.D.; Arendts, G.; Waldron, N.; et al. Multifactorial falls prevention programmes for older adults presenting to the emergency department with a fall: systematic review and meta-analysis. *Inj. Prev.* 2019, 25, 557–564, doi:10.1136/injuryprev-2019-043214.
11. World Health Organization What are the main risk factors for falls amongst older people and what are the most effective interventions to prevent these falls?; 2004. Available online: <http://www.euro.who.int/document/E82552.pdf> (accessed on 1 November 2020).
12. Nieto-Riveiro, L.; Groba, B.; Miranda, M.C.; Concheiro, P.; Pazos, A.; Pousada, T.; Pereira, J. Technologies for participatory medicine and health promotion in the elderly population. *Med. Baltim.* 2018, 97, e10791, doi:10.1097/MD.00000000000010791.
13. Close, J.; Ellis, M.; Hooper, R.; Glucksman, E.; Jackson, S.; Swift, C. Prevention of falls in the elderly trial (PROFET): A randomised controlled trial. *Lancet* 1999, 353, 93–97, doi:10.1016/S0140-6736(98)06119-4.
14. Tanaka, M. Multidisciplinary team approach for elderly patients. *Geriatr. Gerontol. Int.* 2003, 3, 69–72, doi:10.1046/j.1444-1586.2003.00074.x.
15. Weinstein, M.; Booth, J. Preventing Falls in Older Adults: A Multifactorial Approach. *Home Health Care Manag. Pract.* 2006, 19, 45–50, doi:10.1177/1084822306292232.
16. Bleijlevens, M.H.C.; Hendriks, M.R.C.; van Haastregt, J.C.M.; van Rossum, E.; Kempen, G.I.J.M.; Diederiks, J.P.M.; Crebolder, H.F.J.M.; van Eijk, J.T.M. Process factors explaining the ineffectiveness of a multidisciplinary fall prevention programme: A process evaluation. *BMC Public Health* 2008, 8, 332, doi:10.1186/1471-2458-8-332.
17. Elliott, S.; Leland, N.E. Occupational therapy fall prevention interventions for community-dwelling older adults: A systematic review. *Am. J. Occup. Ther.* 2018, 72, 7204190040p1–7204190040p11, doi:10.5014/ajot.2018.030494.
18. Tricco, A.C.; Lillie, E.; Zarin, W.; O'Brien, K.K.; Colquhoun, H.; Levac, D.; Moher, D.; Peters, M.D.J.; Horsley, T.; Weeks, L.; et al. PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Ann. Intern. Med.* 2018, 169, 467, doi:10.7326/M18-0850.
19. Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* 2009, 6, e1000097, doi:10.1371/journal.pmed.1000097.
20. Biblioteca Virtual de Salud España. 1998. Available online: <https://bvsalud.isciii.es/> (accessed on 1 November 2020).
21. Chase, C.A.; Mann, K.; Wasek, S.; Arbesman, M. Systematic review of the effect of home modification and fall prevention programs on falls and the performance of community-dwelling older adults. *Am. J. Occup. Ther.* 2012, 66, 284–91, doi:10.5014/ajot.2012.005017.
22. Horowitz, B.P.; Nochajski, S.M.; Schweitzer, J.A. Occupational therapy community practice and home assessments: Use of the home safety self-assessment tool (HSSAT) to support aging in place. *Occup. Ther. Heal. care* 2013, 27, 216–27, doi:10.3109/07380577.2013.807450.
23. Charness, N. Utilizing technology to improve older adult health. *Occup. Ther. Heal. Care* 2014, 28, 21–30, doi:10.3109/07380577.2013.865859.
24. Arthanat, S.; Wilcox, J.; Macuch, M. Profiles and predictors of smart home technology adoption by older adults. *OTJR Occup. Particip. Heal.* 2019, 39, 247–256, doi:10.1177/1539449218813906.
25. Stewart, L.S.P.; McKinstry, B. Fear of falling and the use of telecare by older people. *Br. J. Occup. Ther.* 2012, 75, 304–312, doi:10.4276/030802212X13418284515758.
26. Glännfjord, F.; Hemmingsson, H.; Larsson Ranada, Å. Elderly people's perceptions of using Wii sports bowling—A qualitative study. *Scand. J. Occup. Ther.* 2017, 24, 329–338, doi:10.1080/11038128.2016.1267259.
27. Hamm, J.; Money, A.; Atwal, A. Fall prevention self-assessments via mobile 3D visualization technologies: Community dwelling older adults' perceptions of opportunities and challenges. *JMIR Hum. Factors* 2017, 4, e15, doi:10.2196/humanfactors.7161.

28. Lemmens, R.; Gielen, C.; Spooren, A. Obstacle: A tool to assess the home environment designed for all. *Stud. Health Technol. Inform.* 2017, 242, 168–174.
29. Hamm, J.; Money, A.G.; Atwal, A. Enabling older adults to carry out paperless falls-risk self-assessments using guidetomeasure-3D: A mixed methods study. 2019, 92, 103135, doi:10.1016/j.jbi.2019.103135.
30. Money, A.G.; Atwal, A.; Boyce, E.; Gaber, S.; Windeatt, S.; Alexandrou, K. Falls sensei: A serious 3D exploration game to enable the detection of extrinsic home fall hazards for older adults. *BMC Med. Inform. Decis. Mak.* 2019, 19, 85, doi:10.1186/s12911-019-0808-x.
31. Khaled, A.B.H.; Khalfallah, A.; Bouhlel, M.S. Fall prevention exergame using occupational therapy based on kinect. In *Smart Innovation, Systems and Technologies*; Springer Science and Business Media Deutschland GmbH: Berlin, Germany, 2020; Volume 146, pp. 479–493.
32. Bianco, M.; Pedell, S.; Renda, G. Augmented reality and home modifications. In *Proceedings of the 28th Australian Conference on Computer-Human Interaction—OzCHI 2016*, Launceston, Australia, 29 November–2 December 2016; ACM Press: New York, NY, USA, 2016; pp. 499–507, doi:10.1145/3010915.3010929.
33. Marston, H.R.; Woodbury, A.; Gschwind, Y.J.; Kroll, M.; Fink, D.; Eichberg, S.; Kreiner, K.; Ejupi, A.; Annegarn, J.; de Rosario, H.; et al. The design of a purpose-built exergame for fall prediction and prevention for older people. *Eur. Rev. Aging Phys. Act.* 2015, 12, doi:10.1186/s11556-015-0157-4.
34. Drobics, M.; Lord, S.R.; Gschwind, Y.J.; Delbaere, K.; Aal, K.; Wieching, R.; de Rosario, H.; Marston, H.R.; Woodbury, A.; Kroll, M.; et al. ICT-based system to predict and prevent falls (iStoppFalls): Results from an international multicenter randomized controlled trial. *Eur. Rev. Aging Phys. Act.* 2015, 12, 10, doi:10.1186/s11556-015-0155-6.
35. Özsungur, F. A research on the effects of successful aging on the acceptance and use of technology of the elderly. *Assist. Technol.* 2019, 18, 1–14, doi:10.1080/10400435.2019.1691085.
36. Boulton, E.; Hawley-Hague, H.; Vereijken, B.; Clifford, A.; Guldemon, N.; Pfeiffer, K.; Hall, A.; Chesani, F.; Mellone, S.; Bourke, A.; et al. Developing the FARSEEING Taxonomy of Technologies: Classification and description of technology use (including ICT) in falls prevention studies. *J. Biomed. Inform.* 2016, 61, 132–140, doi:10.1016/j.jbi.2016.03.017.
37. Boulton, E.; Hawley-Hague, H.; French, D.P.; Mellone, S.; Zacchi, A.; Clemson, L.; Vereijken, B.; Todd, C. Implementing behaviour change theory and techniques to increase physical activity and prevent functional decline among adults aged 61–70: The PreventIT project. *Prog. Cardiovasc. Dis.* 2019, 62, 147–156, doi:10.1016/j.pcad.2019.01.003.
38. World Health Organization Assistive technology Available online: https://www.who.int/health-topics/assistive-technology#tab=tab_1 (accessed on 3 November 2020).
39. Di Monaco, M.; Vallero, F.; De Toma, E.; De Lauso, L.; Tappero, R.; Cavanna, A. A single home visit by an occupational therapist reduces the risk of falling after hip fracture in elderly women: A quasi-randomized controlled trial. *J. Rehabil. Med.* 2008, 40, 446–450, doi:10.2340/16501977-0206.
40. Silva, J.S. e.; Gonçalves, R.; Branco, F.; Pereira, A.; Au-Yong-Oliveira, M.; Martins, J. Accessible software development: a conceptual model proposal. *Univers. Access Inf. Soc.* 2019, 18, 703–716, doi:10.1007/s10209-019-00688-5.
41. Design for All Foundations Design for All Available online: <http://designforall.org/index.php> (accessed on 5 November 2020).
42. Oxford Institute of Population Ageing Bridging the digital divide amongst older adults Available online: <https://www.ageing.ox.ac.uk/blog/digital-divide> (accessed on 5 November 2020).
43. World Health Organization Ageing: Healthy ageing and functional ability Available online: <https://www.who.int/westernpacific/news/q-a-detail/ageing-healthy-ageing-and-functional-ability> (accessed on 5 November 2020).
44. Oxford Martin School Life Expectancy - Our World in Data Available online: <https://ourworldindata.org/life-expectancy> (accessed on 5 November 2020).
45. Brandis, S.J.; Tuite, A.T. Falls prevention: partnering occupational therapy and general practitioners. *Aust. Health Rev.* 2001, 24, 37–42, doi:10.1071/ah010037.
46. Wiles, J.L.; Leibing, A.; Guberman, N.; Reeve, J.; Allen, R.E.S. The Meaning of “Aging in Place” to Older People. *Gerontologist* 2012, 52, 357–366, doi:10.1093/geront/gnr098.
47. National Institute on Aging Aging in Place: Growing Older at Home | National Institute on Aging Available online: <https://www.nia.nih.gov/health/aging-place-growing-older-home#planning> (accessed on 5 November 2020).
48. Lachman, M.E.; Howland, J.; Tennstedt, S.; Jette, A.; Assmann, S.; Peterson, E.W. Fear of falling and activity restriction: The Survey of Activities and Fear of Falling in the Elderly (SAFE). *Journals Gerontol. - Ser. B Psychol. Sci.*

49. Cumming, R.G.; Salkeld, G.; Thomas, M.; Szonyi, G. Prospective study of the impact of fear of falling on activities of daily living, SF-36 scores, and nursing home admission. *Journals Gerontol. - Ser. A Biol. Sci. Med. Sci.* 2000, 55, M299–M305, doi:10.1093/gerona/55.5.M299.
50. Yardley, L.; Smith, H. A prospective study of the relationship between feared consequences of falling and avoidance of activity in community-living older people. *Gerontologist* 2002, 42, 17–23, doi:10.1093/geront/42.1.17.
51. Kerse, N.; Flicker, L.; Pfaff, J.J.; Draper, B.; Lautenschlager, N.T.; Sim, M.; Snowdon, J.; Almeida, O.P. Falls, depression and antidepressants in later life: A large primary care appraisal. *PLoS One* 2008, 3, e2423, doi:10.1371/journal.pone.0002423.
52. Bailey, C.; Foran, T.G.; Nli Scanail, C.; Dromey, B. Older adults, falls and technologies for independent living: A life space approach. *Ageing Soc.* 2011, 31, 829–848, doi:10.1017/S0144686X10001170.
53. Brodie, M.A.; Coppens, M.J.; Ejupi, A.; Gschwind, Y.J.; Annegarn, J.; Schoene, D.; Wieching, R.; Lord, S.R.; Delbaere, K. Comparison between clinical gait and daily-life gait assessments of fall risk in older people. *Geriatr. Gerontol. Int.* 2017, 17, 2274–2282, doi:10.1111/ggi.12979.
54. Nagayama, H.; Tomori, K.; Ohno, K.; Takahashi, K.; Yamauchi, K. Cost-effectiveness of Occupational Therapy in Older People: Systematic Review of Randomized Controlled Trials. *Occup. Ther. Int.* 2016, 23, 103–120, doi:10.1002/oti.1408.
55. Miskelly, F.G. Assistive technology in elderly care. *Age Ageing* 2001, 30, 455–458, doi:10.1093/ageing/30.6.455.
56. Walker, J.E.; Howland, J. Falls and fear of falling among elderly persons living in the community: occupational therapy interventions. *Am. J. Occup. Ther.* 1991, 45, 119–122, doi:10.5014/ajot.45.2.119.
57. Somerville, E.; Smallfield, S.; Stark, S.; Seibert, C.; Arbesman, M.; Lieberman, D. Occupational therapy home modification assessment and intervention. *Am. J. Occup. Ther.* 2016, 70, 7005395010p1-7005395010p3, doi:10.5014/ajot.2016.705002.
58. Lord, S.R.; Menz, H.B.; Sherrington, C. Home environment risk factors for falls in older people and the efficacy of home modifications. *Age and Ageing*, 2006; 35, 55–59.
59. Cumming, R.G.; Thomas, M.; Szonyi, G.; Frampton, G.; Salkeld, G.; Clemson, L. Adherence to occupational therapist recommendations for home modifications for falls prevention. *Am. J. Occup. Ther.* 2001, 55, 641–648, doi:10.5014/ajot.55.6.641.
60. Mirelman, A.; Rochester, L.; Maidan, I.; Del Din, S.; Alcock, L.; Nieuwhof, F.; Rikkert, M.O.; Bloem, B.R.; Pelosin, E.; Avanzino, L.; et al. Addition of a non-immersive virtual reality component to treadmill training to reduce fall risk in older adults (V-TIME): a randomised controlled trial. *Lancet* 2016, 388, 1170–1182, doi:10.1016/S0140-6736(16)31325-3.
61. Vaziri, D.D.; Aal, K.; Gschwind, Y.J.; Delbaere, K.; Weibert, A.; Annegarn, J.; de Rosario, H.; Wieching, R.; Randall, D.; Wulf, V. Analysis of effects and usage indicators for a ICT-based fall prevention system in community dwelling older adults. *Int. J. Hum. Comput. Stud.* 2017, 106, 10–25, doi:10.1016/j.ijhcs.2017.05.004.