

Water Footprint of Food Production

Subjects: Environmental Sciences

Contributor: Mesfin M Mekonnen

Blue water footprint - the volume of fresh surface and groundwater that has been evaporated or incorporated into a product.

Green water footprint - the rainwater that is stored in the root zone of the soil and evapotranspired or incorporated into the product.

Unsustainable blue water footprint - when it exceeds the available renewable blue water, thereby violating the environmental flow standard and depleting groundwater

Water footprint - an indicator of the direct and indirect water use to produce the goods and services we use.

Keywords: sustainable diet, food waste, food production

1. Introduction

Agriculture is by far the largest user of water. Agricultural production needs to increase by almost 50% by 2050 compared to 2012 to meet the rising demand for food, fiber, and biofuels. This is probably going to require more water. Most of the increase in agricultural production is expected to occur in Sub-Saharan Africa and South Asia, where the agricultural output will need to more than double by 2050^[1]. The expected increase in the rest of the world is around 30%. Agricultural production has increased by 260% between 1961 and 2018^[2]. During the same period, the harvested crop area increased by 47%, suggesting that 113% of the increase in production is linked to an increase in crop yield. The increase in crop yield between 1961 and 1990 was 72%, but between 1991 and 2018, the increase was 43%, indicating that yields are now rising at a slower rate than in previous decades^[2]. The increase in crop yields was largely due to increased irrigation, improved crop varieties, agrochemical inputs, and improved soil and water management. However, the increase in crop productivity is not expected to continue indefinitely. In most parts of the world, yields for major crops have begun to stagnate^{[3][4]}. Climate change, soil degradation, and salinization of irrigated areas will potentially limit future increases in production. Ray et al.^[5] have shown that with the current rate of yield increase, it is not possible to meet the expected food demand by 2050. They have argued that some level of cropland expansion is needed to meet the food production deficits but at a higher environmental cost to biodiversity.

The amount of food available for human consumption is affected by the allocation of crops to other nonfood uses such as animal feed, bioenergy, and industrial uses. Globally, only 67% of the crop produced (by mass) or 55% of the calories produced is available for direct human consumption^[6]. The remaining crop was allocated to animal feed (24% by mass) and other industrial use, including bioenergy (9% by weight). Animal production is less efficient than crop production in converting feed to human edible food^{[7][8][9][10]}. As a result, only 12% of the 36% of the global calories used for animal feed will ultimately contribute to human diets^[6].

In 2011, the global water footprint (WF) of agricultural production was 8362 km³/year (80% green, 11% blue, and 9% grey)^[11]. World water demand is expected to increase by 20%–30% between 2010 and 2050^[12]. Demand for land and water resources has increased significantly, and these resources are expected to be scarcer in the future. Efficient water management in agriculture is needed to meet the growing demand for food and reduce poverty and hunger in a sustainable manner. The question is how the world will feed the global population without further impacting the freshwater and ecosystems. Several researchers have advocated for sustainable intensification ^{[10][13][14][15][16]}, dietary changes, and reduction of food waste and loss^{[17][18][19]} to feed the world. A number of studies have shown the value of virtual water trade in global water saving, reducing water scarcity, and it will help to reduce the risk of water scarcity^{[20][21][22][23]}. This paper provides a brief review of the WF of food production, the water demand for different food products and diets, and the WF of food loss and waste.

2. From Quantification to Sustainability Assessment

Over the last few years, the assessment of WFs of national consumption and production has shown significant improvement in terms of product coverage, spatial and temporal detail, and sustainability assessment^[24]. Hoekstra and Hung^[25] were the first to estimate the WF of national consumption of 38 crops for a large number of countries. The second global assessment made a number of improvements in terms of product coverage by including all crops and livestock products and other refinements^{[26][27]}. The third global study made further refinements by assessing the national water footprints of production and consumption at a high spatial and temporal resolution^{[11][28]}. In addition to the global studies on the WF of crop production mentioned earlier ^{[29][30][31][32][33][34][35]}, other national^{[36][37][38][39][40]}, regional or basin^{[41][42]}, and global studies^{[28][43]} have also traced and mapped the WF of consumption per country, but none of these studies assessed the sustainability of the blue WF of consumption at the place of production. The blue WF is unsustainable if it is above the available renewable blue water and violates the environmental flow requirements.

Van Oel et al.^[44] carried out the first assessment of the sustainability of the WF in relation to Dutch consumption. In a case study for France, Ercin et al.^[45] assessed the sustainability of the WF of consumption of France, identifying priority basins and products. Hoekstra and Mekonnen^[46] assessed the sustainability and water-use efficiency of the UK's WF of consumption. There are more recent works that have assessed the unsustainable blue WF of consumption for a single country or EU as a whole^{[47][48]}. A few studies have assessed the sustainability of the WF of crop production and virtual water flows at a global level^{[49][50][51]}. Other studies focused on sustainability of groundwater use^{[52][53][54][55][56]}.

In a more detailed global study, Mekonnen and Hoekstra^[57] estimated that 513 km³/year, or 57% of the blue WF, related to crop production, was unsustainable. Approximately 75% of the global unsustainable blue WF is related to the production of only six crops (Figure 1). These are wheat, rice, cotton, sugar cane, fodder, and maize. Five countries account for about 70% of the unsustainable blue WF (Figure 1), India, China, the US, Pakistan, and Iran. Of the total unsustainable blue WF, 90% was for food and fodder crops, while only 10% was for fiber crops, rubber, and tobacco. Figure 2 shows the sustainable and unsustainable blue WF for global cereal production. The unsustainable portion of the blue WF is large in the Indus and Ganges river basins in India and Pakistan, in the north-eastern part of China, and in the US. High Plains aquifer. The study also showed that some 25% of global blue water can be saved by reducing the WF of each crop to the benchmark level.




Figure 1. The unsustainable blue WF related to crop production: (a) Contribution of different crops toward the global blue unsustainable blue WF of crop production; (b) location of the unsustainable blue WF of crop production. Data source from Mekonnen and Hoekstra^[57].




Figure 2. The sustainable (green) and unsustainable (yellow to dark red) parts of the blue WF of global cereals production. Data source from Mekonnen and Hoekstra^[57].

References

1. FAO. The Future of Food and Agriculture-Trends and Challenges; Food and Agriculture Organization: Rome, Italy, 2017.
2. FAO. FAOSTAT Online Database; FAO: Rome, Italy, 2020.
3. Grassini, P.; Eskridge, K.M.; Cassman, K.G. Distinguishing between yield advances and yield plateaus in historical crop production trends. *Nat. Commun.* 2013, 4.
4. Ray, D.K.; Ramankutty, N.; Mueller, N.D.; West, P.C.; Foley, J.A. Recent patterns of crop yield growth and stagnation. *Nat. Commun.* 2012, 3, 1293. Available online: http://www.nature.com/ncomms/journal/v3/n12/supinfo/ncomms2296_S1.html (accessed on 13 April 2015).
5. Ray, D.K.; Mueller, N.D.; West, P.C.; Foley, J.A. Yield trends are insufficient to double global crop production by 2050. *PLoS ONE* 2013, 8, e66428.
6. Cassidy, E.S.; West, P.C.; Gerber, J.S.; Foley, J.A. Redefining agricultural yields: From tonnes to people nourished per hectare. *Environ. Res. Lett.* 2013, 8, 034015.
7. Mekonnen, M.M.; Neale, C.M.U.; Ray, C.; Erickson, G.E.; Hoekstra, A.Y. Water productivity in meat and milk production in the USA from 1960 to 2016. *Environ. Int.* 2019, 132, 105084.
8. Wirsenius, S. Efficiencies and biomass appropriation of food commodities on global and regional levels. *Agric. Syst.* 2003, 77, 219–255.
9. Bouwman, A.F.; Van der Hoek, K.W.; Eickhout, B.; Soenario, I. Exploring changes in world ruminant production systems. *Agric. Syst.* 2005, 84, 121–153.
10. Tilman, D.; Balzer, C.; Hill, J.; Befort, B.L. Global food demand and the sustainable intensification of agriculture. *Proc. Natl. Acad. Sci. USA* 2011, 108, 20260–20264.
11. Hoekstra, A.Y.; Mekonnen, M.M. The water footprint of humanity. *Proc. Natl. Acad. Sci. USA* 2012, 109, 3232–3237.
12. Burek, P.; Satoh, Y.; Fischer, G.; Kahil, M.T.; Scherzer, A.; Tramberend, S.; Nava, L.F.; Wada, Y.; Eisner, S.; Flörke, M.; et al. Water Futures and Solution-Fast Track Initiative (Final Report); IIASA: Laxenburg, Austria, 2016.
13. Cassman, K.G.; Grassini, P. A global perspective on sustainable intensification research. *Nat. Sustain.* 2020, 3, 262–268.
14. Drechsel, P.; Heffer, P.; Magen, H.; Mikkelsen, R.; Wichelns, D. Managing Water and Fertilizer for Sustainable Agricultural Intensification; International Fertilizer Industry Association (IFA): Paris, France; International Water Management Institute (IWMI): Colombo, Sri Lanka; International Plant Nutrition Institute (IPNI): Peachtree Corners, GA, USA; International Potash Institute (IPI): Horgen, Switzerland, 2015.
15. Garnett, T.; Appleby, M.C.; Balmford, A.; Bateman, I.J.; Benton, T.G.; Bloomer, P.; Burlingame, B.; Dawkins, M.; Dolan, L.; Fraser, D.; et al. Sustainable intensification in agriculture: Premises and policies. *Science* 2013, 341, 33–34.
16. Godfray, H.C.J.; Beddington, J.R.; Crute, I.R.; Haddad, L.; Lawrence, D.; Muir, J.F.; Pretty, J.; Robinson, S.; Thomas, S.M.; Toulmin, C. Food security: The challenge of feeding 9 billion people. *Science* 2010, 327, 812–818.
17. Foley, J.A.; Ramankutty, N.; Brauman, K.A.; Cassidy, E.S.; Gerber, J.S.; Johnston, M.; Mueller, N.D.; O'Connell, C.; Ray, D.K.; West, P.C.; et al. Solutions for a cultivated planet. *Nature* 2011, 478, 337–342.
18. Jalava, M.; Guillaume, J.H.A.; Kummu, M.; Porkka, M.; Siebert, S.; Varis, O. Diet change and food loss reduction: What is their combined impact on global water use and scarcity? *Earth's Future* 2016, 4, 62–78.
19. Kummu, M.; de Moel, H.; Porkka, M.; Siebert, S.; Varis, O.; Ward, P.J. Lost food, wasted resources: Global food supply chain losses and their impacts on freshwater, cropland, and fertiliser use. *Sci. Total Environ.* 2012, 438, 477–489.
20. Liu, W.; Yang, H.; Liu, Y.; Kummu, M.; Hoekstra, A.Y.; Liu, J.; Schulin, R. Water resources conservation and nitrogen pollution reduction under global food trade and agricultural intensification. *Sci. Total Environ.* 2018, 633, 1591–1601.
21. Liu, W.; Antonelli, M.; Kummu, M.; Zhao, X.; Wu, P.; Liu, J.; Zhuo, L.; Yang, H. Savings and losses of global water resources in food-related virtual water trade. *WIREs Water* 2019, 6, e1320.
22. Mekonnen, M.M.; Hoekstra, A.Y. A global and high-resolution assessment of the green, blue and grey water footprint of wheat. *Hydrol. Earth Syst. Sci.* 2010, 14, 1259–1276.
23. Chapagain, A.K.; Hoekstra, A.Y.; Savenije, H.H.G. Water saving through international trade of agricultural products. *HESS* 2006, 10, 455–468.
24. Hoekstra, A.Y. Water footprint assessment: Evolvement of a new research field. *Water Resour. Manage.* 2017, 31, 3061–3081.

25. Hoekstra, A.Y.; Hung, P.Q. *Virtual Water Trade: A Quantification of Virtual Water Flows between Nations in Relation to International Crop Trade*, 11th ed.; IHE: Delft, The Netherlands, 2002.
26. Hoekstra, A.Y.; Chapagain, A.K. *Globalization of Water: Sharing the Planet's Freshwater Resources*; Blackwell: Oxford, UK, 2008.
27. Hoekstra, A.Y.; Chapagain, A.K. Water footprints of nations: Water use by people as a function of their consumption pattern. *Water Resour. Manage.* 2007, 21, 35–48.
28. Mekonnen, M.M.; Hoekstra, A.Y. *National Water Footprint Accounts: The Green, Blue and Grey Water Footprint of Production and Consumption*; UNESCO-IHE: Delft, The Netherlands, 2011.
29. Siebert, S.; Döll, P. Quantifying blue and green virtual water contents in global crop production as well as potential production losses without irrigation. *J. Hydrol.* 2010, 384, 198–217.
30. Liu, J.; Yang, H. Spatially explicit assessment of global consumptive water uses in cropland: Green and blue water. *J. Hydrol.* 2010, 384, 187–197.
31. Liu, J.; Zehnder, A.J.B.; Yang, H. Global consumptive water use for crop production: The importance of green water and virtual water. *Water Resour. Res.* 2009, 45.
32. Hanasaki, N.; Inuzuka, T.; Kanae, S.; Oki, T. An estimation of global virtual water flow and sources of water withdrawal for major crops and livestock products using a global hydrological model. *J. Hydrol.* 2010, 384, 232–244.
33. Fader, M.; Gerten, D.; Thammer, M.; Heinke, J.; Lotze-Campen, H.; Lucht, W.; Cramer, W. Internal and external green-blue agricultural water footprints of nations, and related water and land savings through trade. *Hydrol. Earth Syst. Sci.* 2011, 15, 1641–1660.
34. Mekonnen, M.M.; Hoekstra, A.Y. The green, blue and grey water footprint of crops and derived crop products. *HESS* 2011, 15, 1577–1600.
35. Rost, S.; Gerten, D.; Heyder, U. Human alterations of the terrestrial water cycle through land management. *Adv. Geosci.* 2008, 18, 43–50.
36. Mekonnen, M.M.; Hoekstra, A.Y. Water conservation through trade: The case of Kenya. *Water Int.* 2014, 39, 1–18.
37. Zhuo, L.; Mekonnen, M.M.; Hoekstra, A.Y. Consumptive water footprint and virtual water trade scenarios for China — With a focus on crop production, consumption and trade. *Environ. Int.* 2016, 94, 211–223.
38. Bultink, F.; Hoekstra, A.Y.; Booij, M.J. The water footprint of Indonesian provinces related to the consumption of crop products. *HESS* 2010, 14, 119–128.
39. Marston, L.; Ao, Y.; Konar, M.; Mekonnen, M.M.; Hoekstra, A.Y. High-resolution water footprints of production of the united states. *Water Resour. Res.* 2018, 54, 2288–2316.
40. Schyns, J.; Hamaideh, A.; Hoekstra, A.; Mekonnen, M.; Schyns, M. Mitigating the risk of extreme water scarcity and dependency: The case of Jordan. *Water* 2015, 7, 5705–5730.
41. Mekonnen, M.; Pahlow, M.; Aldaya, M.; Zarate, E.; Hoekstra, A. Sustainability, efficiency and equitability of water consumption and pollution in Latin America and the Caribbean. *Sustainability* 2015, 7, 2086–2112.
42. Zeitoun, M.; Allan, J.A.; Mhieledeen, Y. Virtual water 'flows' of the Nile basin, 1998–2004: A first approximation and implications for water security. *Glob. Environ. Change* 2010, 20, 229–242.
43. Chen, Z.-M.; Chen, G.Q. Virtual water accounting for the globalized world economy: National water footprint and international virtual water trade. *Ecol. Indicators* 2013, 28, 142–149.
44. Van Oel, P.R.; Mekonnen, M.M.; Hoekstra, A.Y. The external water footprint of The Netherlands: Geographically-explicit quantification and impact assessment. *Ecolog. Econ.* 2009, 69, 82–92.
45. Erkin, A.E.; Mekonnen, M.M.; Hoekstra, A.Y. Sustainability of national consumption from a water resources perspective: The case study for France. *Ecolog. Econ.* 2013, 88, 133–147.
46. Hoekstra, A.Y.; Mekonnen, M.M. Imported water risk: The case of the UK. *Environ. Res. Lett.* 2016, 11, 055002.
47. Dolganova, I.; Mikosch, N.; Berger, M.; Núñez, M.; Müller-Frank, A.; Finkbeiner, M. The water footprint of European agricultural imports: Hotspots in the context of water scarcity. *Resources* 2019, 8, 141.
48. Finogenova, N.; Dolganova, I.; Berger, M.; Núñez, M.; Blizniukova, D.; Müller-Frank, A.; Finkbeiner, M. Water footprint of German agricultural imports: Local impacts due to global trade flows in a fifteen-year perspective. *Sci. Total Environ.* 2019, 662, 521–529.
49. Wang, R.; Zimmerman, J. Hybrid analysis of blue water consumption and water scarcity implications at the global, national, and basin levels in an increasingly globalized world. *Environ. Sci. Technol.* 2016, 50, 5143–5153.

50. Jägermeyr, J.; Pastor, A.; Biemans, H.; Gerten, D. Reconciling irrigated food production with environmental flows for sustainable development goals implementation. *Nat. Commun.* 2017, 8, 15900.
51. Rosa, L.; Chiarelli, D.D.; Tu, C.; Rulli, M.C.; D'Odorico, P. Global unsustainable virtual water flows in agricultural trade. *Environ. Res. Lett.* 2019, 14, 114001.
52. Dalin, C.; Taniguchi, M.; Green, T.R. Unsustainable groundwater use for global food production and related international trade. *Glob. Sustain.* 2019, 2, e12.
53. Dalin, C.; Wada, Y.; Kastner, T.; Puma, M.J. Groundwater depletion embedded in international food trade. *Nature* 2017, 543, 700.
54. Wada, Y.; van Beek, L.P.H.; Bierkens, M.F.P. Nonsustainable groundwater sustaining irrigation: A global assessment. *Water Resour. Res.* 2012, 48, W00L06.
55. Scanlon, B.R.; Faunt, C.C.; Longuevergne, L.; Reedy, R.C.; Alley, W.M.; McGuire, V.L.; McMahon, P.B. Groundwater depletion and sustainability of irrigation in the USA high plains and central valley. *Proc. Natl. Acad. Sci. USA* 2012, 109, 9320–9325.
56. Marston, L.; Konar, M.; Cai, X.; Troy, T.J. Virtual groundwater transfers from overexploited aquifers in the United States. *Proc. Natl. Acad. Sci. USA* 2015, 112, 8561–8566.
57. Mekonnen, M.M.; Hoekstra, A.Y. Sustainability of the blue water footprint of crops. *Arizona department Water Resour.* 2020, 143, 103679.

Retrieved from <https://encyclopedia.pub/entry/history/show/7033>