

# CTCs in Gastric Cancer

Subjects: **Oncology**

Contributor: Pham Gia Anh

With over 1 million incidence cases and more than 780,000 deaths in 2018, gastric cancer (GC) was ranked as the 5th most common cancer and the 3rd leading cause of cancer deaths worldwide. Though several biomarkers, including carcinoembryonic antigen (CEA), cancer antigen 19-9 (CA19-9), and cancer antigen 72-4 (CA72-4), have been identified, their diagnostic accuracies were modest. Circulating tumor cells (CTCs), cells derived from tumors and present in body fluids, have recently emerged as promising biomarkers, diagnostically and prognostically, of cancers, including GC. In this review, we present the landscape of CTCs from migration, to the presence in circulation, biologic properties, and morphologic heterogeneities. We evaluated clinical implications of CTCs in GC patients, including diagnosis, prognosis, and therapeutic management, as well as their application in immunotherapy.

circulating tumor cells

gastric cancer

diagnostic

prognostic

treatment

## 1. Introduction

According to data from the International Agency for Research on Cancer, over 1 million new cases of gastric cancer (GC) were found in the past year, and GC was determined to be the third highest cause of cancer-related mortalities worldwide [1][2]. The overall five-year survival rate of GC after curative surgery is approximately 30%. There is a great variation of GC survival rate from developed to developing countries [3], as a patient's prognosis is highly dependent on the stage of the tumor found during surgery. In areas where endoscopic screenings are regularly conducted, such as Japan, a significantly high five-year survival rate is achieved due to the increased possibility for early tumor resection [4]. This survival rate can also drop to less than 5% if the patient is diagnosed at later stages (IIIB to IV) (<https://www.cancer.net/>). Reliable screening and early diagnosis, therefore, play a critical role in the management of GC and patient survival. Despite recent advancement in GC diagnosis and treatment, prognostic outcomes for patients remain quite poor [5]. The asymptomatic characteristics at early stages, high recurrent risk, and rapid metastasis after operation are the main problems that hinder effective diagnosis and treatment of GC.

To date, several biomarkers have been successfully identified and widely used for GC diagnosis and prognosis, including carcinoembryonic antigen (CEA), cancer antigen 19-9 (CA19-9), and cancer antigen 72-4 (CA72-4) [5]. However, those biomarkers display less than 40% positivity rates, and their sensitivities and specificities are modest [5][6]. In this regard, circulating tumor cells (CTCs) emerge as promising markers to monitor treatment responses and to provide progression information in a real-time manner [7][8][9][10].

CTCs are cells that dissociate from the primary tumor and travel through the vasculature. Prior studies have shown that in many cases, the spread and migration of CTCs are heavily responsible for the establishment of distant metastases [11]. CTCs are present in body fluids well before metastasis and they could potentially be used as a predictive indicator of future metastatic formations [11]. Therefore, the use of CTCs for cancer screenings is really promising, as the routine profiling and detection of CTCs from the blood could lead towards the development of a far more efficient, noninvasive system for detection of tumors. Additionally, CTCs exhibit certain properties and unique molecular traits of the primary tumor that could provide valuable insights into possible mechanisms of cancer treatment [12][13][14].

## 2. Landscape of CTCs

### 2.1. CTC Migration

The process of CTC migration is critical towards understanding their behavior in circulation. In the vasculature, CTCs are exposed to attacks from the immune system, high shear stress, and oxidative stress, which pose severely harsh conditions for CTC survival [15][16]. The vast majority of CTCs fail to survive in this foreign environment. However, few CTCs are able to survive and contribute towards the formation of distant metastases, raising the questions of how they are able to do this and what molecular mechanisms are involved.

The traditional model of CTC migration relies heavily upon the epithelial to mesenchymal transition (EMT) process. A major defining characteristic of CTCs in the epithelial state is the expression of e-cadherin, a surface protein heavily involved in the formation of tight junctions and responsible for cellular adhesion [17][18][19]. Consequently, CTCs are inter-locked in a network of intracellular adhesions and lack of migration capabilities. In their mesenchymal state, however, the cells exhibit far greater invasive and migratory properties and are less prone to apoptosis. The increased expression of transforming growth factor  $\beta$  (TGF- $\beta$ ) and several transcriptional factors play critical roles in the induction of EMT and lead to the downregulation of e-cadherin, along with other epithelial markers like alpha and beta catenin [20][21][22][23][24]. The overexpression of extracellular matrix (ECM) proteins such as vimentin and fibronectin results in the induction of certain mesenchymal properties [25][26]. Additional EMT-induced modifications that are likely to help the CTCs adapt in the harsh microenvironments, such as the upregulation of cluster of differentiation 47 (CD47) for increased defense against immune attacks or the downregulation of chaperone calreticulin to dodge immune-surveillance [27]. Further, TGF- $\beta$  has been shown to induce increased coagulation, resulting in the formation of a platelet-rich microclot around CTCs which could physically protect them from the high shear stress in the vasculature as well as provide shielding from NK-mediated attack [28]. EMT has also been shown to activate survival pathways such as Akt and P13k signaling to provide resistance to apoptosis [28].

It is noteworthy that after EMT, CTCs are able to invade the blood vessels and lymph capillaries and migrate towards various organs [29][30]. Once the CTCs are in the vasculature, they can either establish metastases in distant organs or return to the primary tumor, a process known as self-seeding [15]. Self-seeding contributes towards metastatic development, because due to the leaky neovasculature of tumors, "self-seeding" CTCs in

circulation would not require significant adaptations to infiltrate their host tumor [31]. Consequently, the host tumor would be enriched with aggressive CTCs that would lead to an increase in the overall metastatic potential of the tumor. It was found by Kim et al. in a study of MDA231 mammary tumors in mice that self-seeded tumors had a significant increase in the rate of tumor growth, along with increased angiogenesis and stromal recruitment [31].

At their target location, CTCs can accomplish infiltration by undergoing a mesenchymal-to-epithelial cell transition (MET), the reversion of EMT, allowing them to regain their cellular adhesion [32][33]. Yet, even after observing the various modifications that CTCs have undergone for migration, our understanding on how individual CTCs are able to survive in the vasculature and to establish metastases is yet to be elucidated. CTC clusters were generally ignored as there has been a speculation that their large size would enable them to rupture blood vessel walls [16][34]. However, CTC clusters have been observed in the peripheral blood, and it is known that CTC clusters are easier to resist to high shear stress, immune system attacks and anchorage-dependent apoptosis than individual cells [34][35]. Therefore, the migration of CTCs in clusters has gained attention as they could have a far greater impact on colonization and metastatic potential than individual CTCs.

The inner regions of CTC clusters include networks of densely connected cells connected by cell junctions that express various adhesion molecules, while the cells on the outer regions of CTC clusters exhibit far greater migratory properties and are receptive to extracellular inputs directing its migration [29]. Certain received inputs lead to the activation of selected pathways, including mitogen-activated protein kinase (MAPK), extracellular-signal-regulated kinase (ERK), etc., that result in leader cell-type selection in certain CTC subsets and inhibition of leader cell-type selection in the adjacent areas [29]. Unlike individual CTCs, CTC clusters do not completely differentiate from the epithelial to mesenchymal type. Instead, there exists a spectrum of a hybrid epithelial and mesenchymal phenotype. It is also suggested that EMT occurred in a morphogenetic manner, as certain cells in the outer region of CTC cluster tend to lose the majority of their epithelial profile and begin expressing mesenchymal markers, while follower cells are able to retain most of their epithelial phenotype [15][36][37].

It is therefore obvious that CTCs have evolved an extensive and nuanced process for migration. Many of the mechanisms behind the migration of CTCs depend on their highly plastic nature, which is observed in their adaptation to different phenotypes and characteristics according to their microenvironment. Yet, there are still many gaps in our understanding of CTCs migration and thus further research on this area is warranted.

## 2.2. CTCs in Circulation

Once CTCs enter the bloodstream, they have the potential to colonize to the distant organs and establish metastases or return to their primary tumor site(s). However, even with the extensive modifications they have undergone, most CTCs, whether they are circulating individually or in clusters, are unable to survive in the hostile environment of the vasculature before they can extravasate into surrounding tissues [13].

Different studies have demonstrated that the few survivable CTCs which have high metastatic potential, undergo various mechanisms to prolong survival while in circulation. For instance, Shliakhtunou et al. [38] found significant overexpression of the survivin gene in all 36 samples of primary breast carcinoma with malignant tumors compared

with benign ones. Moderate and late-stage tumors, in particular, express survivin at a significantly high level. Survivin suppresses apoptosis by inhibiting caspase 6 and 8 activation and also provides a defense against NK cell cytotoxicity; therefore, its overexpression would provide a survival advantage towards CTCs [39]. Cytokines are also extensively involved with CTCs in the circulation. In a study by Huh et al., human melanoma cells were injected into mice intravenously [40]. Upon administration of human neutrophils, cancer cell retention in the lungs and metastatic growth significantly increased in comparison to non-neutrophil treated mice. Further, the secretion of cytokine IL-8 by melanoma cells attracts neutrophils and promotes B-integrin expression, which facilitates anchorage of the cells to the vascular endothelium [40]. Additionally, Divella et al. found that CTC clusters in patients with metastatic colorectal cancer were significantly associated with elevated levels of cytokines CXCL1 and TGF-beta in circulation [41]. Other studies have also demonstrated this correlation between elevated production of cytokines and CTCs [42][43][44]. Taken together, these findings suggest that CTCs exhibit a cytokine-mediated interaction which aids in CTC survival, transendothelial migration and metastatic development.

It has also been observed that platelet interaction plays an important role in CTC survival. EMT-mediated platelet coating around the CTC surface provides a shield from natural killer (NK)-mediated tumor attack as well from the shear and dynamic stress in the bloodstream [45][46]. Furthermore, platelet depletion enhances the efficacy of NK cytolysis of CTCs [47]. Accordingly, Placke et al. [48] utilized immunoelectron microscopy, immunofluorescent staining, and other techniques to reveal that platelet coating of CTCs results in the transfer of platelet-derived major histocompatibility complex (MHC Class 1) to the cell surface of CTCs, allowing them to mimic the phenotype of a host cell and therefore escape from immunosurveillance. These results suggested that therapies targeting platelet production may have critical effects in reducing CTC survival and preventing metastasis formation. Additionally, conditions such as thrombocytosis associated with excessive platelet formation may be used as prognostic indicators for tumor malignancy and migration, as increased platelet accumulation stemming from these conditions may help facilitate CTC survival and hence promote metastatic formation [49].

Overall, in circulation, there are several mechanisms in place to protect CTCs from the harsh environment of the vasculature. Further understanding these mechanisms could provide valuable insight into possible therapies that can inhibit the progression of certain tumors.

### 2.3. Morphologic Heterogeneity of CTCs

A critical property of CTCs are their heterogeneities. CTCs derived from the same tumor exhibit a wide variety of morphological traits, which include size, ratio of nucleus to cytoplasm, structural irregularities, and apoptotic phenotype. CTC heterogeneity is largely responsible for its metastatic capabilities, as greater diversity in CTC morphology allows clonal selection of CTCs at secondary tumor site [50]. Generally, conventional CTCs that are positive for cytokeratin and EpCAM and negative for CD45 have a larger size and nucleus-to-cytoplasm ratio than leukocytes [51]. They have a distinct cytokeratin signal in the cytoplasm around the nucleus that is larger than the surrounding white blood cell nuclei [13]. CTCs that have a more mesenchymal phenotype and a reduced EpCAM expression and cytokeratin signal exhibit a large and irregularly shaped nucleus [13]. Other CTCs may have a nucleus with no size difference than those of white blood cell (WBC), or they may be apoptotic, showing a

disrupted nucleus and/or possibly cytokeratin stain while also expressing M30, a neoepitope heavily involved in carrying out apoptosis [13][52]. They may have significant implications in assessing metastatic potential of tumors in patients. Kallergi et al., using 56 CTC positive patients, demonstrated that apoptotic CTCs were found under significantly lower amounts in patients that had metastatic disease [53]. Additionally, they found that the percentage of apoptotic CTCs was lower in patients with advanced stages of disease compared to patients who had early stages.

CTCs derived from the same tumor exhibit varying morphological and immunophenotypical characteristics. Given the few examples presented above of CTC heterogeneity, this diversity may have implications in the use of CTCs to analyze the effectiveness of certain therapeutic strategies for disease treatment [52].

### 3. Clinical Implication of CTCs in GC

Although CTCs were discovered 150 years ago, the studies on clinical utilities of CTCs in GC has peaked recently, particular during the past five years. The systematic review in [Table 1](#) shows that CTCs have significant values in the diagnosis, prognosis, and treatment management of GC at various stages.

**Table 1.** Systematic Review of the Clinical Significances of Circulating Tumor Cells (CTCs) in Gastric Cancer (GC) in Studies during 2015–2019 Period.

Author, Country, Year	Cases	Isolation and Identification Tool	Markers	OS and PFS of CTC(+) vs. CTC(-)	Positive Cutoff	Clinical Implications	Reference
Lee (South Korea), 2015	100 metastatic GC patients	anti-EpCAM antibody coated magnetic particles CTC-Profiler (Veridex)	EpCAM, CK8/18/19, CD45	OS: 120 days vs. 220 days; $p = 0.03$ PFS: 59 days vs. 141 days; $p = 0.004$	$\geq 5$ CTCs/7.5 mL	CTCs are associated with poor response to chemotherapy in metastatic gastric cancer patients. CTC positivity was an independent adverse factor for PFS and OS.	[54]
Okabe (Japan), 2016	136 advanced GC patients	semi-automated immunomagnetic separation system CellSearch	EpCAM, CK8/18/19CD45, DAPI	OS: HR 2.20 [95%CI: 1.120–4.03]; $p = 0.009$ PFS: HR 2.03 [95%CI:	$\geq 1$ CTCs/7.5 mL	Detection of CTCs was an independent predictor of a shorter PFS in advanced gastric cancer. Patients who	[55]

Author, Country, Year	Cases	Isolation and Identification Tool	Markers	OS and PFS of CTC(+) vs. CTC(-)	Positive Cutoff	Clinical Implications	Reference
Zhou 2016	1110 GC patients in meta-analysis	-	-	1.13–3.66]; $p = 0.016$	-	require intensive treatment: CTCs could be a valuable biomarker. The combined status of CTC and CY would be useful in selecting patients for radical surgery.	
Mishima(Japan), 2017	101 GC patients 15 advanced GC patients whose primary tumors were HER2-, but CTCs were HER2+	both 3 D-IF-FISH method & CellSearch System3D-IF-FISH only	EpCAM, CK8/18/19, CD45, HER2	OS and PFS of 15 advanced GC patients with CTC-HER2+: 6.1 months (95% CI: 2.1–10.0) and 14.4 months	$\geq 1$ CTCs/7.5 mL	New, non-invasive strategy to select patients who are likely to benefit from trastuzumab-based therapies, despite their primary biopsy	[56]

Author, Country, Year	Cases	Isolation and Identification Tool	Markers	OS and PFS of CTC(+) vs. CTC(-)	Positive Cutoff	Clinical Implications	Reference
				(11.0–17.8), respectively		being HER2-negative.	
Liu (China), 2017	59 GC patients of stage II-IV	CELLection™ Epithelial Enrich kit	EpCAM, CK8/18/19, DAPI	OS: HR = 3.59, 95% CI: 1.655-7.817, $p = 0.001$ PFS: = 2.81, 95% CI: 1.313-5.999, $p = 0.008$	$\geq 2$ CTCs/5 mL	The baseline CTC count of >2 cells/5 mL and an increase of the CTC count after the first cycle of chemotherapy was an independent prognostic marker of poor PFS and OS → patients with a low baseline CTC count or decrease of the CTC count after the first cycle of chemotherapy may benefit significantly from palliative chemotherapy	[58]
Zheng (China), 2017	81 GC patients	ISET-immunofluorescence	CK8/18/19, vimentin	CTM positivity was an independent factor for determining the PFS ( $p = 0.016$ ) and OS ( $p = 0.003$ ) of stage IV patients. CTM correlated with shorter PFS and OS than single CTCs ( $p < 0.05$ )	$\geq 1$ CTCs/5 mL For CTM: $\geq 3$ CTCs	In stage IV patients, CTM positivity was correlated with serum CA125 level. CTM were an independent predictor of shorter PFS and OS in stage IV patients. → CTM detection may be a useful tool to predict prognosis in stage IV patients.	[59]

Author, Country, Year	Cases	Isolation and Identification Tool	Markers	OS and PFS of CTC(+) vs. CTC(-)	Positive Cutoff	Clinical Implications	Reference
Kang (South Korea), 2017	116 patients with gastric cancer patients & 31 healthy volunteers	"FAST disc" centrifugal microfluidic system	EpCAM, CK8/18/19, CD45 DAPI	≥ 2 CTCs/7.5 mL Sensitivity: 85.3 Specificity: 90.3	-	Although the clinical feasibility of CTCs for gastric cancer staging was not proved, these results suggest a potential role of CTCs as an early diagnostic biomarker of gastric cancer.	[60]
Yue (China), 2018	35 patients with different advanced gastrointestinal tumors	Pep MNPs isolated system	CK19, CD45 DAPI, PD-L1	PFS based on baseline PD-L1 <sup>high</sup> CTC count: 4.27 vs. 2.07 months HR = 3.342; 95%CI 1.488–7.505; <i>p</i> = 0.002 PFS based on post-therapeutic PD-L1 <sup>high</sup> CTC count: 3.4 vs. 2.1 months; HR= 0.412; 95%CI 0.177–0.962; <i>p</i> = 0.031)	≥ 2 PD-L1 <sup>high</sup> CTCs/4 mL	The abundance of PD-L1 <sup>high</sup> CTCs at baseline might serve as a predictor to screen patients for PD-1/PD-L1 blockade therapies. Measuring the dynamic changes of CTC could indicate the therapeutic response at early time.	[61]
Yang (China), 2018	40 GC patients	wedge-shaped microfluidic chip (CTC-ΔChip) & three-color immunocytochemistry method	(CK, CD45, Nucleus marker	-	-	CTC-ΔChip exhibited the feasibility of detecting CTCs from different types of solid tumor, and it identified 7.30 ± 7.29 CTCs from 2 mL peripheral	[62]

Author, Country, Year	Cases	Isolation and Identification Tool	Markers	OS and PFS of CTC(+) vs. CTC(-)	Positive Cutoff	Clinical Implications	Reference
						blood with a positive rate of 75% (30/40) in GC patients. Novel CTC-ΔChip shows high performance for detecting CTCs from less volume of blood samples of cancer patients and important clinical significance in GC.	
Li (China), 2018	115 advanced GC patients, including 56 tumor HER2 <sup>+</sup> subjects who received first-line HER2-targeted therapy plus chemotherapy and 59 tumor HER2 <sup>-</sup> subjects who received chemotherapy alone	IF-FISH Cytelligen system	DAPI, HER2, CEP8, and CD45	-	-	CTC HER2 <sup>+</sup> was found in 91.0% of tumor HER2 <sup>+</sup> and 76.2% tumor HER2 <sup>-</sup> patients and was correlated with development of resistance to trastuzumab for the tumor HER2 <sup>+</sup> patients and chemotherapy alone for the tumor HER2 <sup>-</sup> patients. Determining of CTC HER2 showed advantages in real-time monitoring of therapeutic resistance.	[63]
Cheng (China), 2019	32 advanced GC patients	CanPatrol CTC enrichment technique Multiplex RNA in situ hybridization assay	EpCAM, CK8/18/19, CD45 DAPI, PD-L1,	-	≥ 2 PD-L1 <sup>+</sup> CTCs/5 mL	CTCs count was well correlated with clinicopathology	[64]

Similar to other types of cancer, early detection is one of the most important challenges to GC since 80% of patients are asymptomatic during the early stages [68]. Recently, the role of CTCs in GC diagnosis has markedly gained attention. A meta-analysis by Tang et al. [69] recommended that CTC detection alone could not be used for GC screening because of poor sensitivity. However, with a high positive likelihood ratio and especially the specificity of 0.99 and the area under the curve (AUC) of 0.97 in the receiver operating characteristic (ROC) curve, CTCs detection could be useful for the confirmation of GC. Intriguingly, also according to Tang et al. [69], from 1997 until 2012, there was an annually increased trend of sensitivity, suggesting that the advancement in detection technology might help improving the diagnostic accuracy of CTCs. Indeed, using a centrifugal microfluidic system with a new fluid-assisted separation technique, Kang et al. [60] showed that the sensitivity and specificity of CTCs in

Author, Country, Year	Cases	Isolation and Identification Tool	Markers	OS and PFS of CTC(+) vs. CTC(-)	Positive Cutoff	Clinical Implications	Reference	seful for
			Vimentin and Twist			parameters. Enumeration of epithelial CTC subset and its relative abundance in the total CTC pool are highly correlated with clinical efficacy. Monitoring CTC subtypes exhibits higher sensitivity of evaluating the disease status, compared to the traditional methods.		nimation of oectively, osis and ear cells R-421, a asis and ne of the [74] Similarly, miR-106a involving andidate
Lu (China), 2019	42 GC patients of stage III-IV	ISET-ICC method followed by IHC	EpCAM, CK8/18/19, CD45, Vimentin and Twist, E-cadherin	-PFS between	≥ 1 CTM (2)	The threshold number of CTCs is significantly associated with different clinical stages and was positively correlated with the value in U/mL of CA724. CTCs technology based on ISET method has a high detection rate. CTCs are promising predictor for the evaluation and prediction [55] of treatment responses in stage III-IV gastric cancer.	[65]	asis and nt to the owed a including e was a eval-CI: potential C counts y shorter arker for
Abdallah (Brazil), 2019	At diagnosis (55 samples)	ISET and immunocytochemistry	HER2 and plakoglobin,			The analysis of CTM	[66]	

Author, Country, Year	Cases	Isolation and Identification Tool	Markers	OS and PFS of CTC(+) vs. CTC(-)	Positive Cutoff	Clinical Implications	Reference
	before neoadjuvant treatment) After surgery and before adjuvant therapy (33 samples)	& microscopy	CD45	CTM-positive patients vs. CTM-negative patients (18.7 months vs. 21.6 months; $p = 0.258$ ) -PFS between plakoglobin-positive CTM patients vs. plakoglobin-positive CTM patients: 15.9 months vs. 21.3 months; $p = 0.114$	CTCs/4 mL	plakoglobin expression is a promising tool in the understanding the biology and prognosis of GC.	
Gao, 2019	3814 GC patients in meta-analysis	-	-	HR = 1.84, 95%CI 1.50–2.26, $p < 0.001$	-	CTC positivity was associated with poorer OS. <a href="#">[67]</a>	

**Figure 1.** The predictive ability of CTCs in gastric cancer treatment. High baseline or acquired CTC count was associated with decreased OS, PFS, and DC. The HER2 and PD-L1 positivity or high amplification could predict positive response to trastuzumab and checkpoint therapy, respectively. Meanwhile, the CD44<sup>+</sup> CTCs were associated with poor prognosis and metastasis.

Additionally, in a study by Li et al. [\[80\]](#), the post-therapy CTC level was used for monitoring therapeutic response in GC patients and predicting their prognosis. Following 6 weeks of chemotherapy, an unfavorable CTC count ( $\geq 3$  CTCs per 7.5 mL) significantly correlated with the objective response rate ( $p = 0.02$ ) and the disease control rate ( $p = 0.01$ ) and furthermore could independently predict a shorter PFS and OS. In contrast, a continuously favorable CTC count or an early shift into a favorable CTC count following therapy could predict an improved prognosis. Similarly, in Liu's report [\[58\]](#), only those patients with a low baseline CTC level or a decreased CTC level after the first cycle of chemotherapy remarkably benefited from the treatment. Those with a baseline CTC level of  $> 2$  cells/5 mL blood, which was determined as an independent poor prognostic marker for PFS and OS, were spared from the adverse reactions of chemotherapy. The role of CTC count in the prognosis and prediction of chemotherapeutic response was also supported in metastatic GC patients [\[54\]](#). In chemotherapy, not only can CTC count be used to predict recurrence risk after operative therapy, there is also a significant difference in the recurrence rates of those patients with different preoperative and postoperative CTC levels. A cut-off value of 1 CTC/7.5 mL was chosen for CTC measurement. Advanced GC patients with preoperative CTCs  $\geq 1$  and increased postoperatively had the poorest prognosis [\[81\]](#).

It should be noted that, not only the number of CTCs but their genotypes and phenotypes can affect the prognostic significance in GC ([Figure 1](#)). Szczepanik et al. [\[82\]](#) demonstrated that among the cytokeratin positive CTCs, only

those with CD44+ phenotype were correlated with poor prognosis and metastasis. CD44 is a transmembrane glycoprotein that was one of the first molecules identified on GC stem cells and is now known to be a marker for cancer stem cells (CSCs) [83]. The superior malignancy of the CD44<sup>(+)</sup>/CD45<sup>(-)</sup> gastric CTCs over the human GC cells was supported in a comparison by Yuan et al. [84]. Moreover, in an in vitro screening by Yuan et al. [84], such CTCs were found to be relatively sensitive to 5 fluorouracil, cisplatin, and paclitaxel, but relatively resistant to radiation, oxaliplatin, cetuximab, and trastuzumab. Along with the fact that CD44<sup>(+)</sup>/CD45<sup>(-)</sup> CTCs show some degree of chemotherapy resistance, they have also been shown to exhibit other characteristics of cancer stem cells (CSC). For example, Grillet et al. generated ex vivo CTC cell lines and found that they were able to differentiate into distinct heterogenous lineages, indicating the presence of multipotent cell population resembling CSCs. Additionally, they also found significant expression of CD44 and aldehyde dehydrogenase, a CSC marker, in CTC cell lines. This indicates that CD44<sup>(+)</sup>/CD45<sup>(-)</sup> CTCs functionally resemble or potentially are CSCs. Therefore, CTCs can mimic the genetic features of tumors at a given time and represent the current state of disease [85]. CTCs cultured in vitro can be used for drug sensitivity screening, a promising approach toward precision medicines. The idea of using CTC culture like an antibiogram has also been raised with other kinds of cancers, such as advanced metastatic colorectal cancer or breast cancer [86][87].

In precision medicine, patient stratification is essential to select those who respond to a treatment regimen. The response indicators are usually molecular biomarkers in biopsy samples from primary tumors. For instance, HER2 upregulation in primary tumors has been used as a biomarker for personalization of trastuzumab, a HER2 monoclonal antibody in advanced GC [88][89]. Intriguingly, Mishima et al. [57] recently demonstrated that HER2 amplification in CTCs was found even in advanced GC patients with HER2 negativity in tumor biopsy. Those patients received clinical benefits from trastuzumab which were comparable to those from the trastuzumab in combination with chemotherapy of the ToGA clinical trial [89]. Similarly, as reported by Abdallah et al. [66], three out of five nonmetastatic gastric adenocarcinoma patients with disease progression had HER2-negative primary tumors but HER2-positive CTCs. A significant CTC count drop in follow-up was seen for CTC-HER2-positive cases and CTC-plakoglobin-positive cases compared with CTC-HER2-negative cases and CTC-plakoglobin-negative cases.

Genetic signatures of CTCs could reveal the metastatic capability and the malignancy of the cancer cells [14][90]. Mimori et al., using microarray analysis and RT-PCR on 810 GC patients, identified membrane type 1 matrix metalloproteinase (MT1-MMP) as a marker in CTCs for metastatic growth [91]. Further, Li et al. profiled CTCs in 31 patients with advanced GC [92] found that those with greater percentages of aneuploid CTCs had significantly poorer progression-free survival and overall survival. Similarly, Li et al. in karyotyping CTCs from patients with advanced GC found that those with chromosome 8 aneuploidy CTCs displayed decreased sensitivity and increased resistance to paclitaxel- and cisplatin-based therapy [93]. Such findings open a perspective of using CTCs as an alternative biopsy which is less invasive and more effective than the conventional tumor biopsy for GC prognosis and therapeutic management.

### 3.3. The Role of CTCs in GC Immunotherapy

With several recent breakthroughs, immunotherapy is an important facet in the improvement of GC management. Until now, the two major approaches of immunotherapy that have been applied in GC clinical trials are adoptive cell immunotherapy and immune checkpoint inhibitor use [94]. The immune checkpoint blockade is progressing more rapidly with a promising response rate from pembrolizumab, a programmed cell death-1 (PD-1) inhibitor, in advanced GC patients [95]. Now, PD-L1 is the most common biomarker for PD-1/PD-L1 checkpoint blockade therapies. The FDA has approved four immunohistochemistry (IHC)-based assays for predicting therapeutic responses to PD-1/PD-L1 blockers. Nonetheless, the predictive value of PD-L1 IHC is still controversial, since in various types of cancer, including GC, some patients with PD-L1 negative tumors also respond to anti-PD-1 therapy [95][96]. Recently, Cheng et al. [64] found that in GC patients with negative PD-L1 in tumor tissue, the CTC Imaging flow cytometry signal was as varied as that of IHC staining, suggesting that expression of CTC PD-L1 is useful in the immunophenotypic differential diagnosis of tumors and thus can be a potential candidate for anti-PD-1/PD-L1 immune checkpoint therapy [97]. CTC level after checkpoint blockade therapy may be effective to predict poor prognosis in advanced GC [98]. When CTC PD-L1 has been recognized as a biomarker for therapeutic efficacy of immune checkpoint inhibition in non-small cell lung cancer (NSCLC), it is unclear whether it can be a predictive marker for GC immune checkpoint therapy [99][100]. Along this line, Yue et al. [61] demonstrated that PD-L1 level in CTCs can be a potential predictor for PD-1/PD-L1 blockade therapies in patients with advanced gastrointestinal tumors. The disease control (DC) rate in the patients with high CTC PD-L1 level was much higher than those with lower level. Additionally, several PD-L1 high CTCs at the baseline had predictive values for PFS. Moreover, the dynamic changes of PD-L1 positive CTCs can reflect real time response status. Indeed, in Yue's study [61], the count changes of total CTCs, PD-L1 positive CTC, and PD-L1 high CTC significantly correlated with the clinical outcomes (Figure 1). This is an advantage of CTC PD-L1 over tumor PD-L1, which can only provide baseline information for treatment response prediction. Similarly, Cheng et al., using the CanPatrol CTC enrichment technique to validate the blood samples from 32 GC patients, also demonstrated that total CTC pool and CTC-PD-L1 are highly correlated with clinical outcome of checkpoint blockade therapy. In addition, high circulating PD-L1 expression in advanced GC patients is predictive of better five-year OS [101]. Altogether, these evidences support CTC PD-L1 expression as a prognostic factor for checkpoint blockade therapy efficacy. Moreover, recent advances in -OMICs technologies to support single-cell molecular profiling of CTCs and tumor cells using next-generation sequencing may also provide the pool of tools to validate predictive biomarkers including CTC PD-1 and PD-L1.

## References

1. Lyons, K.; Le, L.C.; Pham, Y.T.H.; Borr/on, C.; Park, J.Y.; Tran, C.T.D.; Tran, T.V.; Tran, H.T.T.; Vu, K.T.; Do, C.D.; et al. Gastric cancer: Epidemiology, biology, and prevention: A mini review. *Eur. J. Cancer Prev.* 2019, 28.
2. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *Ca A Cancer J. Clin.* 2018, 68, 394–424.

3. Redaniel, M.T.; Laudico, A.; Mirasol-Lumague, M.R.; Gondos, A.; Pulte, D.; Mapua, C.; Brenner, H. Cancer survival discrepancies in developed and developing countries: Comparisons between the Philippines and the United States. *Br. J. Cancer* 2009, 100, 858.
4. Correa, P. Gastric cancer: Overview. *Gastroenterol Clin. North. Am.* 2013, 42, 211–217.
5. Ting-Ting, L.; Hao, L.; Jiang, Y.; Guang-Yao, S.; Li-Ying, Z.; Guo-Xin, L.; Ting-Ting, L.; Hao, L.; Jiang, Y.; Li-Ying, Z.; et al. Prognostic and predictive blood biomarkers in gastric cancer and the potential application of circulating tumor cells. *World J. Gastroenterol.* 2018, 24, 2236–2246.
6. Li, Y.; Yang, Y.; Lu, M.; Shen, L. Predictive value of serum CEA, CA19-9 and CA72.4 in early diagnosis of recurrence after radical resection of gastric cancer. *Hepatogastroenterology* 2011, 58, 2166–2170.
7. Jin, Z.; Jiang, W.; Wang, L. Biomarkers for gastric cancer: Progression in early diagnosis and prognosis (Review). *Oncol. Lett.* 2015, 9, 1502–1508.
8. Nakamura, K.; Iwatsuki, M.; Kurashige, J.; Ishimoto, T.; Baba, Y.; Miyamoto, Y.; Yoshida, N.; Watanabe, M.; Baba, H. Circulating tumor cells in gastric cancer. *J. Cancer Metastasis Treat.* 2018, 4, 32.
9. Zhou, J.; Ma, X.; Bi, F.; Liu, M. Clinical significance of circulating tumor cells in gastric cancer patients. *Oncotarget* 2017, 8, 25713–25720.
10. Palmirotta, R.; Lovero, D.; Cafforio, P.; Felici, C.; Mannavola, F.; Pellè, E.; Quaresmini, D.; Tucci, M.; Silvestris, F. Liquid biopsy of cancer: A multimodal diagnostic tool in clinical oncology. *Ther. Adv. Med. Oncol.* 2018, 10, 1758835918794630.
11. Yap, T.A.; Lorente, D.; Omlin, A.; Olmos, D.; de Bono, J.S. Circulating tumor cells: A multifunctional biomarker. *Clin. Cancer Res.* 2014, 20, 2553–2568.
12. Krebs, M.G.; Metcalf, R.L.; Carter, L.; Brady, G.; Blackhall, F.H.; Dive, C. Molecular analysis of circulating tumour cells-biology and biomarkers. *Nat. Rev. Clin. Oncol.* 2014, 11, 129–144.
13. Thiele, J.A.; Bethel, K.; Kralickova, M.; Kuhn, P. Circulating Tumor Cells: Fluid Surrogates of Solid Tumors. *Annu. Rev. Pathol.* 2017, 12, 419–447.
14. Paterlini-Brechot, P. Organ-specific markers in circulating tumor cell screening: An early indicator of metastasis-capable malignancy. *Future Oncol.* 2011, 7, 849–871.
15. Giuliano, M.; Shaikh, A.; Lo, H.C.; Arpino, G.; de Placido, S.; Zhang, X.H.; Cristofanilli, M.; Schiff, R.; Trivedi, M.V. Perspective on Circulating Tumor Cell Clusters: Why It Takes a Village to Metastasize. *Cancer Res.* 2018, 78, 845–852.
16. Au, S.H.; Storey, B.D.; Moore, J.C.; Tang, Q.; Chen, Y.L.; Javaid, S.; Sarioglu, A.F.; Sullivan, R.; Madden, M.W.; O'Keefe, R.; et al. Clusters of circulating tumor cells traverse capillary-sized vessels. *Proc. Natl. Acad. Sci. USA* 2016, 113, 4947–4952.

17. Barriere, G.; Fici, P.; Gallerani, G.; Fabbri, F.; Zoli, W.; Rigaud, M. Circulating tumor cells and epithelial, mesenchymal and stemness markers: Characterization of cell subpopulations. *Ann. Transl. Med.* 2014, 2, 109.
18. Maitre, J.L.; Heisenberg, C.P. Three functions of cadherins in cell adhesion. *Curr. Biol.* 2013, 23, R626–R633.
19. Nakamura, T.; Kato, Y.; Fuji, H.; Horiuchi, T.; Chiba, Y.; Tanaka, K. E-cadherin-dependent intercellular adhesion enhances chemoresistance. *Int. J. Mol. Med.* 2003, 12, 693–700.
20. Alonso-Alconada, L.; Muinelo-Romay, L.; Madissoo, K.; Diaz-Lopez, A.; Krakstad, C.; Trovik, J.; Wik, E.; Hapangama, D.; Coenegrachts, L.; Cano, A.; et al. Molecular profiling of circulating tumor cells links plasticity to the metastatic process in endometrial cancer. *Mol. Cancer* 2014, 13, 223.
21. Nagrath, S.; Sequist, L.V.; Maheswaran, S.; Bell, D.W.; Irimia, D.; Ulkus, L.; Smith, M.R.; Kwak, E.L.; Digumarthy, S.; Muzikansky, A.; et al. Isolation of rare circulating tumour cells in cancer patients by microchip technology. *Nature* 2007, 450, 1235–1239.
22. Galletti, G.; Sung, M.S.; Vahdat, L.T.; Shah, M.A.; Santana, S.M.; Altavilla, G.; Kirby, B.J.; Giannakakou, P. Isolation of breast cancer and gastric cancer circulating tumor cells by use of an anti HER2-based microfluidic device. *Lab. Chip* 2014, 14, 147–156.
23. Micalizzi, D.S.; Farabaugh, S.M.; Ford, H.L. Epithelial-mesenchymal transition in cancer: Parallels between normal development and tumor progression. *J. Mammary Gland. Biol. Neoplasia* 2010, 15, 117–134.
24. Petrova, Y.I.; Schecterson, L.; Gumbiner, B.M. Roles for E-cadherin cell surface regulation in cancer. *Mol. Biol. Cell* 2016, 27, 3233–3244.
25. Kalluri, R.; Weinberg, R.A. The basics of epithelial-mesenchymal transition. *J. Clin. Investig.* 2009, 119, 1420–1428.
26. Armstrong, A.J.; Marengo, M.S.; Oltean, S.; Kemeny, G.; Bitting, R.L.; Turnbull, J.D.; Herold, C.I.; Marcom, P.K.; George, D.J.; Garcia-Blanco, M.A. Circulating tumor cells from patients with advanced prostate and breast cancer display both epithelial and mesenchymal markers. *Mol. Cancer Res.* 2011, 9, 997–1007.
27. Williams, A.; Rawal, S.; Ao, Z.; Lu, B.; Torres-Munoz, J.; Rini, B.; Pelley, R.; Budd, G.T.; Borden, E.; Zheng, S.; et al. Abstract 2372: Capture and molecular characterization of CTC in metastatic breast, prostate, colorectal, and renal cancer. *Cancer Res.* 2012, 72, 2372.
28. Francart, M.E.; Lambert, J.; Vanwynsberghe, A.M.; Thompson, E.W.; Bourcy, M.; Polette, M.; Gilles, C. Epithelial-mesenchymal plasticity and circulating tumor cells: Travel companions to metastases. *Dev. Dyn.* 2018, 247, 432–450.

29. Yang, Y.; Zheng, H.; Zhan, Y.; Fan, S. An emerging tumor invasion mechanism about the collective cell migration. *Am. J. Transl. Res.* 2019, 11, 5301–5312.

30. Rejniak, K.A. Circulating Tumor Cells: When a Solid Tumor Meets a Fluid Microenvironment. *Adv. Exp. Med. Biol.* 2016, 936, 93–106.

31. Kim, M.Y.; Oskarsson, T.; Acharyya, S.; Nguyen, D.X.; Zhang, X.H.; Norton, L.; Massague, J. Tumor self-seeding by circulating cancer cells. *Cell* 2009, 139, 1315–1326.

32. Jie, X.X.; Zhang, X.Y.; Xu, C.J. Epithelial-to-mesenchymal transition, circulating tumor cells and cancer metastasis: Mechanisms and clinical applications. *Oncotarget* 2017, 8, 81558–81571.

33. Zhao, X.H.; Wang, Z.R.; Chen, C.L.; Di, L.; Bi, Z.F.; Li, Z.H.; Liu, Y.M. Molecular detection of epithelial-mesenchymal transition markers in circulating tumor cells from pancreatic cancer patients: Potential role in clinical practice. *World J. Gastroenterol.* 2019, 25, 138–150.

34. Hong, B.; Zu, Y. Detecting circulating tumor cells: Current challenges and new trends. *Theranostics* 2013, 3, 377–394.

35. Hou, J.M.; Krebs, M.; Ward, T.; Sloane, R.; Priest, L.; Hughes, A.; Clack, G.; Ranson, M.; Blackhall, F.; Dive, C. Circulating tumor cells as a window on metastasis biology in lung cancer. *Am. J. Pathol.* 2011, 178, 989–996.

36. Jia, D.; Li, X.; Bocci, F.; Tripathi, S.; Deng, Y.; Jolly, M.K.; Onuchic, J.N.; Levine, H. Quantifying Cancer Epithelial-Mesenchymal Plasticity and its Association with Stemness and Immune Response. *J. Clin. Med.* 2019, 8, 725.

37. Yu, M.; Bardia, A.; Wittner, B.S.; Stott, S.L.; Smas, M.E.; Ting, D.T.; Isakoff, S.J.; Ciciliano, J.C.; Wells, M.N.; Shah, A.M.; et al. Circulating breast tumor cells exhibit dynamic changes in epithelial and mesenchymal composition. *Science* 2013, 339, 580–584.

38. Shliakhtunou, Y. Survivin gene expression in the primary tumor and circulating tumor cells—A new biomarker of tumor progression of breast cancer. *Ann. Oncol.* 2016, 27.

39. Végran, F.; Boidot, R. Survivin-3B promotes chemoresistance and immune escape by inhibiting caspase-8 and -6 in cancer cells. *Oncolimmunology* 2013, 2, e26328.

40. Huh, S.J.; Liang, S.; Sharma, A.; Dong, C.; Robertson, G.P. Transiently entrapped circulating tumor cells interact with neutrophils to facilitate lung metastasis development. *Cancer Res.* 2010, 70, 6071–6082.

41. Divella, R.; Daniele, A.; Abbate, I.; Bellizzi, A.; Savino, E.; Simone, G.; Giannone, G.; Giuliani, F.; Fazio, V.; Gadaleta-Caldarola, G.; et al. The presence of clustered circulating tumor cells (CTCs) and circulating cytokines define an aggressive phenotype in metastatic colorectal cancer. *Cancer Causes Control.* 2014, 25, 1531–1541.

42. Killock, D. CTCs ‘piggyback’ off neutrophils. *Nat. Rev. Clin. Oncol.* 2019, 16, 208.

43. Xie, Z.; Gao, X.; Cheng, K.; Yu, L. Correlation between the presence of circulating tumor cells and the pathologic type and staging of non-small cell lung cancer during the early postoperative period. *Oncol. Lett.* 2017, 14, 5825–5830.

44. Vilsmaier, T.; Rack, B.; Konig, A.; Friese, K.; Janni, W.; Jeschke, U.; Weissenbacher, T.; Group, S.S. Influence of Circulating Tumour Cells on Production of IL-1alpha, IL-1beta and IL-12 in Sera of Patients with Primary Diagnosis of Breast Cancer Before Treatment. *Anticancer Res.* 2016, 36, 5227–5236.

45. Wang, W.-C.; Zhang, X.-F.; Peng, J.; Li, X.-F.; Wang, A.-L.; Bie, Y.-Q.; Shi, L.-H.; Lin, M.-B.; -F, X. Zhang, Survival Mechanisms and Influence Factors of Circulating Tumor Cells. *Biomed Res. Int.* 2018, 2018, 9.

46. Mathewson, N.D.; Jenq, R.; Mathew, A.V.; Koenigsknecht, M.; Hanash, A.; Toubai, T.; Oravecz-Wilson, K.; Wu, S.R.; Sun, Y.; Rossi, C.; et al. Gut microbiome-derived metabolites modulate intestinal epithelial cell damage and mitigate graft-versus-host disease. *Nat. Immunol.* 2016, 17, 505–513.

47. Nieswandt, B.; Hafner, M.; Echtenacher, B.; Mannel, D.N. Lysis of tumor cells by natural killer cells in mice is impeded by platelets. *Cancer Res.* 1999, 59, 1295–1300.

48. Placke, T.; Orgel, M.; Schaller, M.; Jung, G.; Rammensee, H.G.; Kopp, H.G.; Salih, H.R. Platelet-derived MHC class I confers a pseudonormal phenotype to cancer cells that subverts the antitumor reactivity of natural killer immune cells. *Cancer Res.* 2012, 72, 440–448.

49. Huong, P.T.; Nguyen, L.T.; Nguyen, X.-B.; Lee, S.K.; Bach, D.-H. The Role of Platelets in the Tumor-Microenvironment and the Drug Resistance of Cancer Cells. *Cancers* 2019, 11, 240.

50. Cleris, L.; Daidone, M.G.; Fina, E.; Cappelletti, V. The Detection and Morphological Analysis of Circulating Tumor and Host Cells in Breast Cancer Xenograft Models. *Cells* 2019, 8, 683.

51. Park, S.; Ang, R.R.; Duffy, S.P.; Bazov, J.; Chi, K.N.; Black, P.C.; Ma, H. Morphological differences between circulating tumor cells from prostate cancer patients and cultured prostate cancer cells. *PLoS ONE* 2014, 9, e85264.

52. Bulfoni, M.; Turetta, M.; del Ben, F.; di Loreto, C.; Beltrami, A.P.; Cesselli, D. Dissecting the Heterogeneity of Circulating Tumor Cells in Metastatic Breast Cancer: Going Far Beyond the Needle in the Haystack. *Int. J. Mol. Sci.* 2016, 17, 1775.

53. Kallergi, G.; Konstantinidis, G.; Markomanolaki, H.; Papadaki, M.A.; Mavroudis, D.; Stournaras, C.; Georgoulias, V.; Agelaki, S. Apoptotic circulating tumor cells in early and metastatic breast cancer patients. *Mol. Cancer* 2013, 12, 1886–1895.

54. Lee, S.J.; Lee, J.; Kim, S.T.; Park, S.H.; Park, J.O.; Park, Y.S.; Lim, H.Y.; Kang, W.K. Circulating tumor cells are predictive of poor response to chemotherapy in metastatic gastric cancer. *Int. J. Biol. Markers* 2015, 30, e382–e386.

55. Okabe, H.; Tsunoda, S.; Hosogi, H.; Hisamori, S.; Tanaka, E.; Tanaka, S.; Sakai, Y. Circulating Tumor Cells as an Independent Predictor of Survival in Advanced Gastric Cancer. *Ann. Surg. Oncol.* 2015, 22, 3954–3961.

56. Zou, K.; Yang, S.; Zheng, L.; Wang, S.; Xiong, B. Prognostic Role of the Circulating Tumor Cells Detected by Cytological Methods in Gastric Cancer: A Meta-Analysis. *Biomed Res. Int.* 2016, 2016, 2765464.

57. Mishima, Y.; Matsusaka, S.; Chin, K.; Mikuniya, M.; Minowa, S.; Takayama, T.; Shibata, H.; Kuniyoshi, R.; Ogura, M.; Terui, Y. Detection of HER2 amplification in circulating tumor cells of HER2-negative gastric cancer patients. *Target. Oncol.* 2017, 12, 341–351.

58. Liu, Y.; Ling, Y.; Qi, Q.; Lan, F.; Zhu, M.; Zhang, Y.; Bao, Y.; Zhang, C. Prognostic value of circulating tumor cells in advanced gastric cancer patients receiving chemotherapy. *Mol. Clin. Oncol.* 2017, 6, 235–242.

59. Zheng, X.; Fan, L.; Zhou, P.; Ma, H.; Huang, S.; Yu, D.; Zhao, L.; Yang, S.; Liu, J.; Huang, A.; et al. Detection of Circulating Tumor Cells and Circulating Tumor Microemboli in Gastric Cancer. *Transl. Oncol.* 2017, 10, 431–441.

60. Kang, H.M.; Kim, G.H. Circulating tumor cells detected by lab-on-a-disc: Role in early diagnosis of gastric cancer. *PLoS ONE* 2017, 12, e0180251.

61. Yue, C.; Jiang, Y.; Li, P.; Wang, Y.; Xue, J.; Li, N.; Li, D.; Wang, R.; Dang, Y.; Hu, Z.; et al. Dynamic change of PD-L1 expression on circulating tumor cells in advanced solid tumor patients undergoing PD-1 blockade therapy. *Oncolimmunology* 2018, 7, e1438111.

62. Yang, C.; Zhang, N.; Wang, S.; Shi, D.; Zhang, C.; Liu, K.; Xiong, B. Wedge-shaped microfluidic chip for circulating tumor cells isolation and its clinical significance in gastric cancer. *J. Transl. Med.* 2018, 16, 139.

63. Li, Y.; Zhang, X.; Liu, D.; Gong, J.; Wang, D.D.; Li, S.; Peng, Z.; Li, Y.; Wang, X.; Lin, P.P.; et al. Evolutionary Expression of HER2 Conferred by Chromosome Aneuploidy on Circulating Gastric Cancer Cells Contributes to Developing Targeted and Chemotherapeutic Resistance. *Clin. Cancer Res.* 2018, 24, 5261–5271.

64. Cheng, B.; Tong, G.; Wu, X.; Cai, W.; Li, Z.; Tong, Z.; He, L.; Yu, S.; Wang, S. Enumeration and Characterization Of Circulating Tumor Cells And Its Application In Advanced Gastric Cancer. *Oncotargets Ther.* 2019, 12, 7887–7896.

65. Lu, R.; Chen, Q.; Liu, X.; Shen, S.; Pan, Z.; Shi, C. Detection of circulating stage III–IV gastric cancer tumor cells based on isolation by size of epithelial tumor: Using the circulating tumor cell biopsy technology. *Transl. Cancer Res.* 2019, 8, 1342–1350.

66. Abdallah, E.A.; Braun, A.C.; Flores, B.; Senda, L.; Urvanegia, A.C.; Calsavara, V.; de Jesus, V.H.F.; Almeida, M.F.A.; Begnami, M.D.; Coimbra, F.J.F.; et al. The Potential Clinical Implications

of Circulating Tumor Cells and Circulating Tumor Microemboli in Gastric Cancer. *Oncologist* 2019, 24, e854–e863.

67. Gao, Y.; Xi, H.; Wei, B.; Cui, J.; Zhang, K.; Li, H.; Cai, A.; Shen, W.; Li, J.; Rosell, R.; et al. Association Between Liquid Biopsy and Prognosis of Gastric Cancer Patients: A Systematic Review and Meta-Analysis. *Front. Oncol.* 2019, 9, 1222.

68. Venerito, M.; Vasapolli, R.; Rokkas, T.; Malfertheiner, P. Gastric cancer: Epidemiology, prevention, and therapy. *Helicobacter* 2018, 23, e12518.

69. Tang, L.; Zhao, S.; Liu, W.; Parchim, N.F.; Huang, J.; Tang, Y.; Gan, P.; Zhong, M. Diagnostic accuracy of circulating tumor cells detection in gastric cancer: Systematic review and meta-analysis. *Bmc Cancer* 2013, 13, 314.

70. Vaiopoulos, A.G.; Kostakis, I.D.; Gkioka, E.; Athanasoula, K.; Pikoulis, E.; Papalambros, A.; Christopoulos, P.; Gogas, H.; Kouraklis, G.; Koutsilieris, M. Detection of circulating tumor cells in colorectal and gastric cancer using a multiplex PCR assay. *Anticancer Res.* 2014, 34, 3083–3092.

71. Hao, N.-B.; He, Y.-F.; Li, X.-Q.; Wang, K.; Wang, R.-L. The role of miRNA and lncRNA in gastric cancer. *Oncotarget* 2017, 8, 81572–81582.

72. Sierzega, M.; Kaczor, M.; Kolodziejczyk, P.; Kulig, J.; Sanak, M.; Richter, P. Evaluation of serum microRNA biomarkers for gastric cancer based on blood and tissue pools profiling: The importance of miR-21 and miR-331. *Br. J. Cancer* 2017, 117, 266.

73. Ge, X.; Liu, X.; Lin, F.; Li, P.; Liu, K.; Geng, R.; Dai, C.; Lin, Y.; Tang, W.; Wu, Z.; et al. MicroRNA-421 regulated by HIF-1alpha promotes metastasis, inhibits apoptosis, and induces cisplatin resistance by targeting E-cadherin and caspase-3 in gastric cancer. *Oncotarget* 2016, 7, 24466–24482.

74. Zhou, H.; Xiao, B.; Zhou, F.; Deng, H.; Zhang, X.; Lou, Y.; Gong, Z.; Du, C.; Guo, J. MiR-421 is a functional marker of circulating tumor cells in gastric cancer patients. *Biomarkers* 2012, 17, 104–110.

75. Cao, Y.; Fan, Y.; Zhuang, W.; Sun, H.; Xuan, B.; Chen, Y.; Xu, W.; Sheng, H. miR-543 functions as a new marker of circulating tumor cells in gastric cancer patients. *Int. J. Clin. Exp. Pathol.* 2017, 10, 473–478.

76. Zhu, M.; Zhang, N.; He, S.; Lui, Y.; Lu, G.; Zhao, L. MicroRNA-106a targets TIMP2 to regulate invasion and metastasis of gastric cancer. *FEBS Lett.* 2014, 588, 600–607.

77. Hu, G.; Lv, Q.; Yan, J.; Chen, L.; Du, J.; Zhao, K.; Xu, W. MicroRNA-17 as a promising diagnostic biomarker of gastric cancer: An investigation combining TCGA, GEO, meta-analysis, and bioinformatics. *FEBS Open Bio.* 2018, 8, 1508–1523.

78. Zhou, H.; Guo, J.M.; Lou, Y.R.; Zhang, X.J.; Zhong, F.D.; Jiang, Z.; Cheng, J.; Xiao, B.X. Detection of circulating tumor cells in peripheral blood from patients with gastric cancer using microRNA as a marker. *J. Mol. Med. (Berl. Ger.)* 2010, 88, 709–717.

79. Huang, X.; Gao, P.; Sun, J.; Chen, X.; Song, Y.; Zhao, J.; Xu, H.; Wang, Z. Clinicopathological and prognostic significance of circulating tumor cells in patients with gastric cancer: A meta-analysis. *Int. J. Cancer* 2015, 136, 21–33.

80. Li, Y.; Gong, J.; Zhang, Q.; Lu, Z.; Gao, J.; Li, Y.; Cao, Y.; Shen, L. Dynamic monitoring of circulating tumour cells to evaluate therapeutic efficacy in advanced gastric cancer. *Br. J. Cancer* 2016, 114, 138–145.

81. Ishiguro, Y. Prognostic significance of circulating tumor cells in patients with gastric cancer: Epithelial mesenchymal transition and perioperative kinetics. *J. Clin. Oncol.* 2019, 37, 59.

82. Szczechaniak, A.; Sierzega, M.; Drabik, G.; Pituch-Noworolska, A.; Kołodziejczyk, P.; Zembala, M. CD44+ cytokeratin-positive tumor cells in blood and bone marrow are associated with poor prognosis of patients with gastric cancer. *Gastric Cancer* 2019, 22, 264–272.

83. Takaishi, S.; Okumura, T.; Tu, S.; Wang, S.S.; Shibata, W.; Vigneshwaran, R.; Gordon, S.A.; Shimada, Y.; Wang, T.C. Identification of gastric cancer stem cells using the cell surface marker CD44. *Stem Cells* 2009, 27, 1006–1020.

84. Yuan, D.; Chen, L.; Li, M.; Xia, H.; Zhang, Y.; Chen, T.; Xia, R.; Tang, Q.; Gao, F.; Mo, X.; et al. Isolation and characterization of circulating tumor cells from human gastric cancer patients. *J. Cancer Res. Clin. Oncol.* 2015, 141, 647–660.

85. Brown, H.K.; Tellez-Gabriel, M.; Cartron, P.F.; Vallette, F.M.; Heymann, M.F.; Heymann, D. Characterization of circulating tumor cells as a reflection of the tumor heterogeneity: Myth or reality? *Drug Discov. Today* 2019, 24, 763–772.

86. Grillet, F.; Bayet, E.; Villeronce, O.; Zappia, L.; Lagerqvist, E.L.; Lunke, S.; Charafe-Jauffret, E.; Pham, K.; Molck, C.; Rolland, N.; et al. Circulating tumour cells from patients with colorectal cancer have cancer stem cell hallmarks in ex vivo culture. *Gut* 2017, 66, 1802–1810.

87. Yu, M.; Bardia, A.; Aceto, N.; Bersani, F.; Madden, M.W.; Donaldson, M.C.; Desai, R.; Zhu, H.; Comaills, V.; Zheng, Z.; et al. Cancer therapy. Ex vivo culture of circulating breast tumor cells for individualized testing of drug susceptibility. *Science (N. Y.)* 2014, 345, 216–220.

88. Fu, X.; Zhang, Y.; Yang, J.; Qi, Y.; Ming, Y.; Sun, M.; Shang, Y.; Yang, Y.; Zhu, X.; Gao, Q. Efficacy and safety of trastuzumab as maintenance or palliative therapy in advanced HER2-positive gastric cancer. *Oncotargets Ther.* 2018, 11, 6091–6100.

89. Bang, Y.J.; van Cutsem, E.; Feyereislova, A.; Chung, H.C.; Shen, L.; Sawaki, A.; Lordick, F.; Ohtsu, A.; Omuro, Y.; Satoh, T.; et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal

junction cancer (ToGA): A phase 3, open-label, randomised controlled trial. *Lancet (Lond. Engl.)* 2010, 376, 687–697.

90. Bernards, R.; Weinberg, R.A. A progression puzzle. *Nature* 2002, 418, 823.

91. Mimori, K.; Fukagawa, T.; Kosaka, Y.; Ishikawa, K.; Iwatsuki, M.; Yokobori, T.; Hirasaki, S.; Takatsuno, Y.; Sakashita, H.; Ishii, H.; et al. A large-scale study of MT1-MMP as a marker for isolated tumor cells in peripheral blood and bone marrow in gastric cancer cases. *Ann. Surg. Oncol.* 2008, 15, 2934–2942.

92. Li, Y.; Zhang, X.; Gong, J.; Zhang, Q.; Gao, J.; Cao, Y.; Wang, D.D.; Lin, P.P.; Shen, L. Aneuploidy of chromosome 8 in circulating tumor cells correlates with prognosis in patients with advanced gastric cancer. *Chin. J. Cancer Res.* 2016, 28, 579–588.

93. Hong, S.H.; Cha, J.M.; Lee, J.I.; Joo, K.R.; Shin, H.P.; Park, J.J.; Jeun, J.W.; Lim, J.U. Association of hyper-LDL cholesterolemia with increased risk of colorectal adenoma. *Hepatogastroenterology* 2014, 61, 1588–1594.

94. Dolcetti, R.; de Re, V. Immunotherapy for Gastric Cancer: Time for a Personalized Approach? *IJMS* 2018, 19, 1602.

95. Fuchs, C.S.; Doi, T.; Jang, R.W.; Muro, K.; Satoh, T.; Machado, M.; Sun, W.; Jalal, S.I.; Shah, M.A.; Metges, J.P.; et al. Safety and Efficacy of Pembrolizumab Monotherapy in Patients With Previously Treated Advanced Gastric and Gastroesophageal Junction Cancer: Phase 2 Clinical KEYNOTE-059 Trial. *JAMA Oncol.* 2018, 4, e180013.

96. Gandini, S.; Massi, D.; Mandala, M. PD-L1 expression in cancer patients receiving anti PD-1/PD-L1 antibodies: A systematic review and meta-analysis. *Crit. Rev. Oncol. Hematol.* 2016, 100, 88–98.

97. Kloten, V.; Lampignano, R.; Krahn, T.; Schlange, T. Circulating Tumor Cell PD-L1 Expression as Biomarker for Therapeutic Efficacy of Immune Checkpoint Inhibition in NSCLC. *Cells* 2019, 8, 9.

98. Schehr, J.L.; Schultz, Z.D.; Warrick, J.W.; Guckenberger, D.J.; Pezzi, H.M.; Sperger, E.; Heninger, A.; Saeed, T.; Leal, K.; Matto, A.M.; et al. Lang, High Specificity in Circulating Tumor Cell Identification Is Required for Accurate Evaluation of Programmed Death-Ligand 1. *PLoS ONE* 2016, 11, e0159397.

99. Wang, Y.; Kim, T.H.; Fouladdel, S.; Zhang, Z.; Soni, P.; Qin, A.; Zhao, L.; Azizi, E.; Lawrence, T.S.; Ramnath, N.; et al. PD-L1 Expression in Circulating Tumor Cells Increases during Radio(chemo)therapy and Indicates Poor Prognosis in Non-small Cell Lung Cancer. *Sci. Rep.* 2019, 9, 566.

100. Kallergi, G.; Vetsika, E.K.; Aggouraki, D.; Lagoudaki, E.; Koutsopoulos, A.; Koinis, F.; Katsarlinos, P.; Trypaki, M.; Messaritakis, I.; Stournaras, C.; et al. Evaluation of PD-L1/PD-1 on circulating

tumor cells in patients with advanced non-small cell lung cancer. *Adv. Med. Oncol.* 2018, **10**, 1758834017750121.

101. Zheng, Z.; Bu, Z.; Liu, X.; Zhang, L.; Li, Z.; Wu, A.; Wu, X.; Cheng, X.; Xing, X.; Du, H.; et al. Level of circulating PD-L1 expression in patients with advanced gastric cancer and its clinical implications. *Chin. J. Cancer Res.* 2014, **26**, 104–111.

Retrieved from <https://www.encyclopedia.pub/entry/history/show/24936>