

1q21.1 CNV in Neuropsychiatric Disorders

Subjects: **Neurosciences**

Contributor: Joy Yoon

The 1q21.1 CNVs, rare and large chromosomal microduplications and microdeletions, are detected in many patients with NDs. Phenotypes of duplication and deletion appear at the two ends of the spectrum. Microdeletions are predominant in individuals with schizophrenia (SCZ) and microcephaly, whereas microduplications are predominant in individuals with autism spectrum disorder (ASD) and macrocephaly.

copy number variation

microdeletion

microduplication

schizophrenia

autism spectrum disorder

microcephaly

macrocephaly

neurodegeneration

synaptic plasticity

1. Introduction

Rare CNVs, such as chromosomal deletions and duplications, have raised much scientific interest in etiological studies of NDs. It has been suggested that genetics play a major role in NDs, with ~52.4% and ~80% of inheritability in ASD and SCZ, respectively. A genetic study has shown that rare and large CNVs are likely to be causative, as they can lead to numerous gene imbalances ^[1]. Case-control studies have demonstrated that rare CNVs occur at higher frequency in cases than in controls, suggesting that patients bear a high CNV burden ^{[2][3]}. Moreover, 17.1% of those who presented abnormal clinical presentations carried pathogenic CNVs ^[4]. Approximately 40% of carriers had de novo mutations, and the majority of the de novo mutations (91%) were pathogenic ^[4]. These patterns show up in most ND studies, including ASD, SCZ, intellectual disability (ID) and attention deficit hyperactivity disorder (ADHD) ^{[5][6][7]}. These findings shed light on the contribution of CNVs to the risks of different NDs.

In general, CNVs are pleiotropic and have variable expressivity, in that different patients carrying CNVs at the same chromosomal regions can show the symptoms of different psychiatric disorders; for example, many ASD-associated CNVs are also found in SCZ patients ^{[3][4][8][9]}. Despite having the same CNV carriers, phenotypes and severity range diversely, and show incomplete penetrance ^[10]. This suggests that there must be other factors involved, such as other genetic components (the two-hit model) ^[11] or environmental factors ^[12]. Hence the complexity of CNVs has been underscored in the etiology of ND.

A recent GWAS has identified risk loci prevalent in NDs, which are rare CNVs seen in cases but not in controls ^[2]. At least eight distinct CNVs, 1q21.1, 2p16.3, 3q29, 7q11.23, 15q13.2, 16p11.2, 22q11.2 and NRXN1, have been consistently reported as risk factors for many NDs ^{[6][7][8][13][14][15]}. Deletions are less frequent but more pathogenic

than duplications. Therefore, an increased odds ratio (OR) was found for deletions (i.e., ORs of 1q21.1 = 11.82 (del) and = 6 (dup)) [15]. The abnormal clinical presentations are postulated to be a result of carrying those pathogenic CNVs. Many genetic studies have attempted to identify the relationships between genetic rearrangements in the regions and clinical phenotypes. As little is known about their effect size, penetrance and genetic predisposition towards a certain phenotype, it is too early to use those rare CNVs for diagnoses of any NDs.

Among the aforementioned associated CNVs, this paper aimed to focus on the 1q21.1 CNV that is found with high incidence in ASD, SCZ, ADHD, ID and epilepsy [16]. Due to its structural complexity and inconsistent clinical phenotypes, this genetic locus has been understudied. A significant and popular finding in 1q21.1 is its mirror effect on neurodevelopment: microdeletions are widely found in the cases of SCZ, and microduplications are widely found in the cases of ASD [17].

2. Chromosomal Mapping and Genetic Pathway of 1q21.1

2.1. Chromosomal Structure

The 1q21.1 CNV is found within a 144 to 148 Mb region [18] (Figure 1a). In contrast to small CNVs, which are less detrimental, larger CNVs (>500 kb in size) can alter the expression levels of multiple genes [19]. It is a complex locus to study in that it not only spans 20–40 putative genes, but the region is also susceptible to genomic rearrangements due to the numbers of low copy repeats (LCRs). The more LCRs in the region, the more prone it is to frequent non-allelic homologous recombination (NAHR) during meiosis [20]. Clustered with LCRs, breakpoints (BPs) divide the locus into four possible segmental blocks and complicate the mapping and prediction of phenotypic expressivity [18]. Many of the LCRs and BPs are located adjacent to the crossing over points, making it difficult to estimate the phenotypes or genomic sequences in any given persons [21]. Through this mechanism, the CNVs, emerging in chromosomal duplications or deletions, can alter some of the dosage-sensitive genes and create a broad range of phenotypic variability [22]. Array comparative genomic hybridization and fluorescent in-situ hybridization analyses mapped out the overall structure of the 1q21.1 in great detail. The 1q21.1 region is associated with mental retardation, autism [23], schizophrenia [24] and microcephaly [21]. Duplication of 1q21.1 is strongly associated with autism [21].




Figure 1. (a) Chromosomal structure of 1q21.1, mapped with four BPs (gray) and two distinct regions (red). (b) An enlargement of the region between 144 Mb and 146 Mb. Known genes commonly found in microduplication and microdeletion carriers are marked with blue bars. The reference locations on the chromosome are based on the March 2006 human reference sequence (NCBI build 36.1). The two distinct regions—TAR and Distal—are indicated by red blocks. (c) The two classes of duplications and deletions are shown with green bars. The size of the bars represents the minimally affected region in each class.

Duplications and deletions are classified into two classes: Class I and Class II. Class I duplication/deletion involves only the distal 1q21.1 region between BP3 and BP4 (1.35 Mb in size), whereas Class II duplication/deletion extends from the distal 1q21.1 to the proximal 1q21.1 commonly detected between BP2 and BP4 (~3 Mb) [21] (Figure 1c). Combined data show enrichment in Class I deletions and duplications with a parental origin, but the components of genes and BPs can be varied after generations [25]. Both analyses discovered two distinct regions: proximal and distal 1q21.1, where a genomic gain or loss occurs (Figure 1b) [21][26]. Microdeletions at proximal 1q21.1 are mainly associated with thrombocytopenia-absent radius (TAR) syndrome and this region is often referred to as the TAR region. In particular, a core exon junction complex gene, *RBM8A*, is located in the TAR region and compound mutations in the *RBM8A* gene cause the TAR syndrome [27] that is comorbid with ID [28]. Other brain dysfunctions, including psychosis, agenesis of the corpus callosum and hypoplasia of the cerebellar vermis, are present in TAR patients [28][29][30]. Consistent with human patient studies, knockdown and knockout of *Rbm8a* in a mouse model revealed the critical role of *RBM8A* in neural progenitor cell (NPC) proliferation, neuronal migration and interneuron development, and loss of function in *RBM8A* in NPCs causes microcephaly [31][32][33]. Moreover, *RBM8A* plays a key role in adult neurogenesis and in regulating anxiety-related behavior [34], further supporting the important role of *RBM8A* in psychiatric diseases.

2.2. Genetic Architecture

The recent advanced genomic assay has deciphered the genes encoded in the region and the position on the locus. The core genes commonly affected in the 1q21.1 CNV carriers are PRKAB2, FMO5, CHD1L, BCL9, ACP6, GJA5, GJA8, GPR89B and PDZK1 [25][35][36] (Figure 2; Table 1). However, the genetic study of the risk genes is far from clear as to the phenotypic consequences. Reported clinical phenotypes of the 1q21.1 duplication and deletion are not consistent, and no single gene has been confirmed to cause a pathologic effect in human studies [36].

This complex expressivity can be explained by a *cis*-epistasis genetic model. In contrast to a single gene CNV model, the gene expression is regulated by one or more CNV drivers and multiple modifiers [37]. Gain or loss of a single gene contributes only a small effect to trigger explicit clinical phenotypes [38]. This was confirmed in a number of genotype–phenotype association studies. A correlation analysis between gene expression and the copy number of 1q21.1 indicated that the candidate genes drew a positively correlated trend, in which a duplication CNV model was likely to have increased gene expression and vice versa [25], but the clinic severity may not have been correlated with the level of gene expression [39]. Harvard et al. conducted a family-based study of 1q21.1 microdeletion and microduplication and showed that individuals with the same CNV exhibited different levels of severity despite the identical gene components and almost identical BPs. Entangled chromatids are increased in lymphoblast cells derived from patients carrying both duplication and deletion of 1q21. To narrow down the causal gene, they identified two candidate genes, *CHD1L* and *PRKAB2*. Knockdown of *CHD1L* led to increased micronuclei in response to a topoisomerase II inhibitor, ICRF-193. However, both deletion and duplication carriers show the same cellular phenotype, suggesting that the gene dosage difference may not correlate with severity of symptoms. These findings once again emphasize the characteristic of the variable expressivity and the *cis*-epistasis model of the 1q21.1 CNVs [40][41]. Nevertheless, understanding of a linkage between genetic imbalance and apparent phenotypes is still incomplete.




Figure 2. A genetic map of the associated genes observed in 1q21.1 microduplications and microdeletions. Blue circles are the 10 affected genes; black circles are the 20 related genes. None of the 10 core risk genes interacts directly with another. A total of 49 genetic linkages are drawn with different widths of green lines and were generated by the GeneMANIA program [42]. Gene expression of the candidate genes is positively correlated with the copy number of 1q21.1 but not with phenotypic severity. Even within the same genetic components, clinical presentations are shown to a different extent in cases, which denies the one gene–one phenotype module. The blue circles are the major genes discussed in the paper. Eight top-ranked genes in the correlation study are not directly linked to each other but are indirectly connected via subtype genes.

Table 1. Genetic function and known phenotypes of dosage-sensitive genes associated with 1q21.1.

Function ¹	Molecular/Cellular Phenotypes	References
CHD1	Chromatin remodeling and DNA damage	Impaired decatenation checkpoint [25]

Function ¹	Molecular/Cellular Phenotypes	References	
	response	activation	
PRKAB2	AMPK regulatory subunit; maintaining energy homeostasis	Neurodegeneration; learning and memory impairment	[42]
GJA8	Gap junction protein; Connexin50	Cataracts; cardiac myopathy; increased risk of SCZ	[43][44]
GJA5	Gap junction protein; Connexin40	Cataracts; cardiac abnormalities	[18][45] [46]
PDZK1	Ion transporter protein; regulates second messenger cascades	Increased risk of ASD and psychosis	[36]
GPR89B	Voltage dependent anion channel	Unknown	
BCL9	Wnt signaling pathway	Increased risk of SCZ	[47]
FMO5	Modulator of metabolic aging		[48][49]
¹			
ACP6	Histidine acid phosphatase protein	Unknown	

These clinical manifestations are associated with the genomic segmental regions on 1q21.1. The frequency of the chromosomal abnormalities was highly skewed to distal regions compared with proximal regions. Minimal deletions in BP2-BP3, known as the TAR syndrome region, however, raised a question of whether this region is benign or pathogenic. The overall chromosomal abnormalities in the proximal region were less frequent than in the distal region. However, the relative enrichment of proximal 1q21.1 in microduplication, especially with a low ratio of de novo inheritance compared with the microdeletions, suggests that the proximal BP2–BP3 region is responsible for clinical microduplication aberrations and is mild enough to maintain fecundity [50][51]. Bearing in mind that developmental delay (DD) is a common history in microdeletions and microduplications, the genes within the proximal BP2–BP3 region account for cerebral development in addition to TAR syndrome [51]. On the other hand, even though the head size was a notable phenotype by dosage, head sizes between the proximal microdeletions and microduplications were not found to be discrete, suggesting that the genes in the proximal region are not

sufficient or not responsible for microcephaly/macrocephaly [51]. These findings confirmed the pathogenicity of the proximal 1q21.1 region; this should be re-evaluated on a large scale to be supportive.

3. Molecular and Cellular Mechanisms Associated with 1q21.1 CNVs

3.1. Effect Range of 1q21.1 CNVs

Studies of CNV pathogenesis have shown that deletions have deleterious effects, while duplications exhibit mild phenotypes [4]. Consistent with its pathogenicity, individuals with deletions have low fecundity and therefore undergo negative selection pressure [52][53]. These features of pathogenic CNVs appear in populations with low frequency and high mutation rates [8][52]. In light of this fact, it has become mainstream in genetic studies to distinguish distinct effect sizes in each ND. In line with the comparable burden of each structural variant, duplications exhibit a smaller burden than deletions in the synaptic pathway; functional clusters of duplications are enriched in NMDA receptor signaling, while functional clusters of deletions are enriched in the nervous system or behavioral phenotypes [15].

Examination of the cellular phenotype is a crucial step in the study of pathogenesis. Because many implicated risk genes in 1q21.1 CNVs are responsible for different cellular processes, including cell signaling, sensing and repair, impairment of these gene functions is expected to disrupt the cellular functions specifically involved with brain development and, in turn, to cause diseases [25]. However, a systematic pathological analysis of postmortem brains carrying 1q21.1 CNVs is still lacking. Due to the clinical manifestation reported among patients, animal models that mimic the genetic deficiency of 1q21.1 CNV could be good tools to provide some mechanistic insights and cellular and molecular targets for further therapeutic development [8][54][55].

3.2. Synaptic Signaling Pathway

Genes for cell signaling are enriched in 1q21.1 [56]. Cell signaling in the brain is impeded by abnormal synaptic plasticity. The dopamine hypothesis has been proposed in many ND studies, including ASD [57] and SCZ [58][59]. A 1q21.1 deletion mouse model recapitulated the function of 1q21.1 CNV in cellular phenotypes [55]. The 1q21.1 CNV accounts for the increased sensitivity to psychostimulants (e.g., amphetamine) and increased dopamine cell firing, and hypersensitivity is not mediated by a different number of D1/D2 receptors [55]. Thus, the findings are consistent with previous studies showing that 1q21.1 deletion shows a higher prevalence in SCZ patients than in ASD [58].

Alteration of the potassium channel function can impair in the whole neural network. Disruption of potassium ion homeostasis often becomes an initiator of the cells' pathological cascade. In light of the crucial function of the potassium channel in neurodevelopment, GWAS has revealed a genetic overlap between rare risk CNVs (e.g., 1q21.1) and genes (e.g., *KCNN3*) encoding the potassium pump, transporter and channel [60][61][62]. The longer CAG repeats within the *KCNN3* gene seem to be associated with SCZ patients [61][62]. However, other studies did not confirm this association [63]. Interestingly, a mutant *KCNN3* channel found in a SCZ patient was localized in the

nucleus and inhibited the current mediated by another potassium channel, KCNN2 [64]. Therefore, the SCZ KCNN3 variant can function as a dominant-negative mutant to suppress endogenous small-conductance K currents and interfere with neuronal firing. Consistent with this notion, dysfunction in astrocyte differentiation derived from SCZ patient-derived induced pluripotent cells (iPSCs) was a result of excessive downregulation of potassium transporters in SCZ glia [60].

3.3. Mitochondrial Functions

Mitochondrial diseases are often associated with ASD children [65]; as a result, creatine kinase, ammonia and aspartate aminotransferase have been used biomarkers for mitochondrial dysfunction in ASD [66]; however, the scale of these studies is still small. In animal studies, AMP-activated protein kinase (AMPK) function is modulated by one of the highest correlated genes, *PRKAB2* [25] in a *Drosophila* model of 1q21.1 [42]. A study confirmed that decreased AMPK activity impaired synaptic plasticity, which is critical for working memory and learning, and leads to sleep dysregulation and shortened lifespan [42]. Loss of AMPK activity also has been associated with the neurodegeneration phenotypes in a fly model of mitochondrial dysfunction [67]. Intriguingly, transcriptomic analyses of the three CNV mouse models—hemizygous deletions in corresponding regions of 1q21, 15q13 and 22q11—have identified that neuronal mitochondrial genes are consistently downregulated across three mutant genotypes and are shared with the transcriptomic changes observed in both SCZ and ASD postmortem brains [68]. This study suggests a previously understudied mitochondrial hypothesis underlying neuropsychiatric diseases associated with CNVs [69][70].

3.4. The WNT Signaling Pathway and BCL9

Epidemiological studies have revealed that the prenatal period is vulnerable to ASD [71][72][73][74][75][76] and SCZ [77][78][79][80][81][82][83][84][85]. Among the key signaling pathways regulating fetal brain development, Wnt proteins play indispensable roles in angiogenesis [86][87][88][89][90], neurogenesis [91][92][93][94][95][96][97][98], cell survival [99][100][101][102], synaptogenesis [103][104][105] and neurite outgrowth [106][107]. The canonical pathway is well known to play a major role in neural development [108]. WNT signaling is regulated by several key components of the canonical Wnt pathway, including β -catenin, whose level determines the activity of canonical Wnt signaling. Recently, mutations in β -catenin have been identified as a frequent cause of ID (OMIM #615075), known as CTNNB1 syndrome [109][110][111][112][113][114], with some individuals also being diagnosed with ASD [115][116][117][118][119]. CTNNB1 syndrome patients are characterized by low IQ, microcephaly and facial dysmorphism that cannot be attributed to a known clinical syndrome [109][110][111][112][113][114]. A β -catenin conditional KO mouse specifically in PV interneurons showed that β -cat cKO mice have increased anxiety, impaired social interactions and elevated repetitive behaviors, which mimic some core symptoms of patients with ASD [120]. In addition, several mouse models with KO of Wnt regulators have shown consistent ASD-like behavioral deficits, including APC [121], DVL1 [122] and PTEN [123][124][125][126]. These data provide compelling evidence that an abnormal Wnt pathway is involved in the development of mental illness.

The *BCL9* gene is located within the 1q21 region and encodes a nuclear retention factor for β -catenin, a critical part of the WNT signaling pathway [127][128][129]. BCL9 is essential for activation of the Wnt signaling in adult

myogenic progenitors and regulates muscle regeneration ^[130]. To determine whether common variants in 1q21 can function as a candidate risk of SCZ, a large-scale GWAS comprising 5772 control and 4187 SCZ patients and 1135 patients with bipolar disorder was conducted in the Chinese Han population ^[47]. Interestingly, multiple SNPs within the *BCL9* gene are significantly associated with SCZ. Consistently, other GWAS and integrative analyses suggest that *BCL9* is associated with negative symptoms in SCZ ^{[131][132]} and is one of top risk genes in CNV ^[133]. As disruption of the BCL9– β -catenin interaction inhibits Wnt activation ^[134], which has been proposed as a therapeutic target for cancer ^{[135][136]}, it remains to be tested if increasing BCL9 levels or fine-tuning WNT signaling could reverse the deficits caused by 1q21 CNV. In addition, several components of the Wnt signaling show an association with SCZ ^{[137][138][139][140][141]} and other psychiatric disorders ^{[115][142][143]}. Among the genetic factors associated with schizophrenia, the *DISC1* ^[144] gene is a genetic risk factor for major mental illness ^{[145][146][147][148][149]}. *DISC1* is a key regulator of NPC proliferation and mouse behavior through modulating the canonical Wnt signaling pathway ^[150]. *DISC1* regulates cortical NPC proliferation and neuronal differentiation via inhibition of GSK3 β . Treatment with pharmacological inhibitors of GSK3 β can completely ameliorate the *DISC1* loss-of-function-induced progenitor proliferation defects and behavioral abnormalities, which illustrates the exciting opportunity to develop small-molecule modulators of the Wnt pathway as prototypical drug treatments for psychiatric diseases.

References

1. Girirajan, S.; Brkanac, Z.; Coe, B.P.; Baker, C.; Vives, L.; Vu, T.H.; Shafer, N.; Bernier, R.; Ferrero, G.B.; Silengo, M.; et al. Relative Burden of Large CNVs on a Range of Neurodevelopmental Phenotypes. *PLoS Genet.* 2011, 7, e1002334.
2. Walsh, T.; McClellan, J.M.; McCarthy, S.E.; Addington, A.M.; Pierce, S.B.; Cooper, G.M.; Nord, A.S.; Kusenda, M.; Malhotra, D.; Bhandari, A.; et al. Rare Structural Variants Disrupt Multiple Genes in Neurodevelopmental Pathways in Schizophrenia. *Science* 2008, 320, 539–543.
3. Kaminsky, E.B.; Kaul, V.; Paschall, J.; Church, D.M.; Bunke, B.; Kunig, D.; Moreno-De-Luca, D.; Moreno-De-Luca, A.; Mulle, J.G.; Warren, S.T.; et al. An evidence-based approach to establish the functional and clinical significance of copy number variants in intellectual and developmental disabilities. *Genet. Med.* 2011, 13, 777–784.
4. Girirajan, S.; Rosenfeld, J.A.; Coe, B.P.; Parikh, S.; Friedman, N.; Goldstein, A.; Filipink, R.A.; McConnell, J.S.; Angle, B.; Meschino, W.S.; et al. Phenotypic Heterogeneity of Genomic Disorders and Rare Copy-Number Variants. *N. Engl. J. Med.* 2012, 367, 1321–1331.
5. Guyatt, A.L.; Stergiakouli, E.; Martin, J.; Walters, J.; O'Donovan, M.; Owen, M.; Thapar, A.; Kirov, G.; Rodriguez, S.; Rai, D.; et al. Association of copy number variation across the genome with neuropsychiatric traits in the general population. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 2018, 177, 489–502.

6. Coe, B.P.; Girirajan, S.; Eichler, E.E. The genetic variability and commonality of neurodevelopmental disease. *Am. J. Med. Genet. Part C Semin. Med. Genet.* 2012, 160C, 118–129.
7. Forsingdal, A.; Jørgensen, T.N.; Olsen, L.; Werge, T.; Didriksen, M.; Nielsen, J. Can Animal Models of Copy Number Variants That Predispose to Schizophrenia Elucidate Underlying Biology? *Biol. Psychiatry* 2019, 85, 13–24.
8. Stefansson, H.; Rujescu, D.; Cichon, S.; Pietiläinen, O.P.H.; Ingason, A.; Steinberg, S.; Fossdal, R.; Sigurdsson, E.; Sigmundsson, T.; Buizer-Voskamp, J.E.; et al. Large recurrent microdeletions associated with schizophrenia. *Nature* 2008, 455, 232–236.
9. Stone, J.L.; O'Donovan, M.C.; Gurling, H.; Kirov, G.K.; Blackwood, D.H.R.; Corvin, A.; Craddock, N.J.; Gill, M.; Hultman, C.M.; Lichtenstein, P.; et al. Rare chromosomal deletions and duplications increase risk of schizophrenia. *Nature* 2008, 455, 237–241.
10. Kirov, G.; Rees, E.; Walters, J.T.R.; Escott-Price, V.; Georgieva, L.; Richards, A.L.; Chambert, K.D.; Davies, G.; Legge, S.E.; Moran, J.L.; et al. The Penetrance of Copy Number Variations for Schizophrenia and Developmental Delay. *Biol. Psychiatry* 2014, 75, 378–385.
11. Girirajan, S.; Rosenfeld, J.A.; Cooper, G.M.; Antonacci, F.; Siswara, P.; Itsara, A.; Vives, L.; Walsh, T.; McCarthy, S.E.; Baker, C.; et al. A recurrent 16p12.1 microdeletion supports a two-hit model for severe developmental delay. *Nat. Genet.* 2010, 42, 203–209.
12. Srirenakumar, V.; Zai, C.C.; Wasim, S.; Barsanti-Innes, B.; Kennedy, J.L.; So, J. Copy number variant syndromes are frequent in schizophrenia: Progressing towards a CNV-schizophrenia model. *Schizophr. Res.* 2019, 209, 171–178.
13. Rees, E.; Walters, J.T.R.; Georgieva, L.; Isles, A.R.; Chambert, K.D.; Richards, A.L.; Mahoney-Davies, G.; Legge, S.E.; Moran, J.L.; McCarroll, S.A.; et al. Analysis of copy number variations at 15 schizophrenia-associated loci. *Br. J. Psychiatry* 2014, 204, 108–114.
14. Takumi, T.; Tamada, K. CNV biology in neurodevelopmental disorders. *Curr. Opin. Neurobiol.* 2018, 48, 183–192.
15. Zhuo, C.; Hou, W.; Lin, C.; Hu, L.; Li, J. Potential Value of Genomic Copy Number Variations in Schizophrenia. *Front. Mol. Neurosci.* 2017, 10.
16. Marshall, C.R.; Howrigan, D.P.; Merico, D.; Thiruvahindrapuram, B.; Wu, W.; Greer, D.S.; Antaki, D.; Shetty, A.; Holmans, P.A.; Pinto, D.; et al. Contribution of copy number variants to schizophrenia from a genome-wide study of 41,321 subjects. *Nat. Genet.* 2017, 49, 27–35.
17. Sanders, S.J.; Ercan-Sencicek, A.G.; Hus, V.; Luo, R.; Murtha, M.T.; Moreno-De-Luca, D.; Chu, S.H.; Moreau, M.P.; Gupta, A.R.; Thomson, S.A.; et al. Multiple recurrent de novo CNVs, including duplications of the 7q11. 23 Williams syndrome region, are strongly associated with autism. *Neuron* 2011, 70, 863–885.

18. Mefford, H.C.; Sharp, A.J.; Baker, C.; Itsara, A.; Jiang, Z.; Buysse, K.; Huang, S.; Maloney, V.K.; Crolla, J.A.; Baralle, D.; et al. Recurrent Rearrangements of Chromosome 1q21.1 and Variable Pediatric Phenotypes. *N. Engl. J. Med.* 2008, 359, 1685–1699.
19. Sullivan, P.F. Schizophrenia and the dynamic genome. *Genome Med.* 2017, 9, 22.
20. Malhotra, D.; Sebat, J. CNVs: Harbingers of a Rare Variant Revolution in Psychiatric Genetics. *Cell* 2012, 148, 1223–1241.
21. Brunetti-Pierri, N.; Berg, J.S.; Scaglia, F.; Belmont, J.; Bacino, C.A.; Sahoo, T.; Lalani, S.R.; Graham, B.; Lee, B.; Shinawi, M.; et al. Recurrent reciprocal 1q21.1 deletions and duplications associated with microcephaly or macrocephaly and developmental and behavioral abnormalities. *Nat. Genet.* 2008, 40, 1466–1471.
22. Owen, M.J. Implications of Genetic Findings for Understanding Schizophrenia. *Schizophr. Bull.* 2012, 38, 904–907.
23. Haldeman-Englert, C.; Jewett, T. 1q21.1 Microdeletion; University of Washington: Seattle, WA, USA, 1993–2021.
24. Ripke, S.; O'Dushlaine, C.; Chambert, K.; Moran, J.L.; Kahler, A.K.; Akterin, S.; Bergen, S.E.; Collins, A.L.; Crowley, J.J.; Fromer, M.; et al. Genome-wide association analysis identifies 13 new risk loci for schizophrenia. *Nat. Genet.* 2013, 45, 1150–1159.
25. Harvard, C.; Strong, E.; Mercier, E.; Colnaghi, R.; Alcantara, D.; Chow, E.; Martell, S.; Tyson, C.; Hrynychak, M.; McGillivray, B.; et al. Understanding the impact of 1q21.1 copy number variant. *Orphanet. J. Rare Dis.* 2011, 6, 54.
26. Torres, F.; Barbosa, M.; Maciel, P. Recurrent copy number variations as risk factors for neurodevelopmental disorders: Critical overview and analysis of clinical implications. *J. Med. Genet.* 2016, 53, 73–90.
27. Albers, C.A.; Paul, D.S.; Schulze, H.; Freson, K.; Stephens, J.C.; Smethurst, P.A.; Jolley, J.D.; Cvejic, A.; Kostadima, M.; Bertone, P.; et al. Compound inheritance of a low-frequency regulatory SNP and a rare null mutation in exon-junction complex subunit RBM8A causes TAR syndrome. *Nat. Genet.* 2012, 44, 435–439.
28. Skorka, A.; Bielicka-Cymermann, J.; Gieruszczak-Bialek, D.; Korniszewski, L. Thrombocytopenia-absent radius (tar) syndrome: A case with agenesis of corpus callosum, hypoplasia of cerebellar vermis and horseshoe kidney. *Genet. Couns.* 2005, 16, 377–382.
29. Weiss Sachdev, S.; Sunde, R.A. Selenium regulation of transcript abundance and translational efficiency of glutathione peroxidase-1 and -4 in rat liver. *Biochem. J.* 2001, 357, 851–858.
30. Ceylan, A.C.; Sahin, I.; Erdem, H.B.; Kayhan, G.; Simsek-Kiper, P.O.; Utine, G.E.; Percin, F.; Boduroglu, K.; Alikasifoglu, M. An eight-case 1q21 region series: Novel aberrations and clinical

- variability with new features. *J. Intellect. Disabil. Res.* 2019, 63, 548–557.
31. Mao, H.; Pilaz, L.-J.; McMahon, J.J.; Golzio, C.; Wu, D.; Shi, L.; Katsanis, N.; Silver, D.L. Rbm8a Haploinsufficiency Disrupts Embryonic Cortical Development Resulting in Microcephaly. *J. Neurosci.* 2015, 35, 7003–7018.
 32. Zou, D.; McSweeney, C.; Sebastian, A.; Reynolds, D.J.; Dong, F.; Zhou, Y.; Deng, D.; Wang, Y.; Liu, L.; Zhu, J.; et al. A critical role of RBM8a in proliferation and differentiation of embryonic neural progenitors. *Neural. Dev.* 2015, 10, 18.
 33. McSweeney, C.; Dong, F.; Chen, M.; Vitale, J.; Xu, L.; Crowley, N.; Luscher, B.; Zou, D.; Mao, Y. Full function of exon junction complex factor, Rbm8a, is critical for interneuron development. *Transl. Psychiatry* 2020, 10, 379.
 34. Alachkar, A.; Jiang, D.; Harrison, M.; Zhou, Y.; Chen, G.; Mao, Y. An EJC factor RBM8a Regulates Anxiety Behaviors. *Curr. Mol. Med.* 2013, 13, 887–899.
 35. Bernier, R.; Steinman, K.J.; Reilly, B.; Wallace, A.S.; Sherr, E.H.; Pojman, N.; Mefford, H.C.; Gerdts, J.; Earl, R.; Hanson, E.; et al. Clinical phenotype of the recurrent 1q21.1 copy-number variant. *Genet. Med.* 2016, 18, 341–349.
 36. Busè, M.; Cuttaia, H.C.; Palazzo, D.; Mazara, M.V.; Lauricella, S.A.; Malacarne, M.; Pierluigi, M.; Cavani, S.; Piccione, M. Expanding the phenotype of reciprocal 1q21.1 deletions and duplications: A case series. *Ital. J. Pediatrics* 2017, 43, 61.
 37. Golzio, C.; Katsanis, N. Genetic architecture of reciprocal CNVs. *Curr. Opin. Genet. Dev.* 2013, 23, 240–248.
 38. Hall, J.; Trent, S.; Thomas, K.L.; O'Donovan, M.C.; Owen, M.J. Genetic Risk for Schizophrenia: Convergence on Synaptic Pathways Involved in Plasticity. *Biol. Psychiatry* 2015, 77, 52–58.
 39. Deshpande, A.; Weiss, L.A. Recurrent reciprocal copy number variants: Roles and rules in neurodevelopmental disorders. *Dev. Neurobiol.* 2018, 78, 519–530.
 40. Crespi, B.J.; Crofts, H.J. Association testing of copy number variants in schizophrenia and autism spectrum disorders. *J. Neurodev. Disord.* 2012, 4, 15.
 41. Warde-Farley, D.; Donaldson, S.L.; Comes, O.; Zuberi, K.; Badrawi, R.; Chao, P.; Franz, M.; Grouios, C.; Kazi, F.; Lopes, C.T.; et al. The GeneMANIA prediction server: Biological network integration for gene prioritization and predicting gene function. *Nucleic Acids Res.* 2010, 38, W214–W220.
 42. Nagy, S.; Maurer, G.W.; Hentze, J.L.; Rose, M.; Werge, T.M.; Rewitz, K. AMPK signaling linked to the schizophrenia-associated 1q21.1 deletion is required for neuronal and sleep maintenance. *PLoS Genet.* 2018, 14, e1007623.

43. Rong, P.; Wang, X.; Niesman, I.; Wu, Y.; Benedetti, L.E.; Dunia, I.; Levy, E.; Gong, X. Disruption of *ja8* ($\alpha 8$ connexin) in mice leads to microphthalmia associated with retardation of lens growth and lens fiber maturation. *Development* 2002, 129, 167–174.
44. Ni, X.; Valente, J.; Azevedo, M.H.; Pato, M.T.; Pato, C.N.; Kennedy, J.L. Connexin 50 gene on human chromosome 1q21 is associated with schizophrenia in matched case–control and family-based studies. *J. Med Genet.* 2007, 44, 532–536.
45. Verhagen, J.M.A.; de Leeuw, N.; Papatsonis, D.N.M.; Grijseels, E.W.M.; de Krijger, R.R.; Wessels, M.W. Phenotypic Variability Associated with a Large Recurrent 1q21.1 Microduplication in a Three-Generation Family. *Mol. Syndromol.* 2015, 6, 71–76.
46. Simon, A.M.; Goodenough, D.A.; Paul, D.L. Mice lacking connexin40 have cardiac conduction abnormalities characteristic of atrioventricular block and bundle branch block. *Curr. Biol.* 1998, 8, 295–298.
47. Li, J.; Zhou, G.; Ji, W.; Feng, G.; Zhao, Q.; Liu, J.; Li, T.; Li, Y.; Chen, P.; Zeng, Z.; et al. Common Variants in the *BCL9* Gene Conferring Risk of Schizophrenia. *Arch. Gen. Psychiatry* 2011, 68, 232–240.
48. Varshavi, D.; Scott, F.H.; Varshavi, D.; Veeravalli, S.; Phillips, I.R.; Veselkov, K.; Strittmatter, N.; Takats, Z.; Shephard, E.A.; Everett, J.R. Metabolic Biomarkers of Ageing in C57BL/6J Wild-Type and Flavin-Containing Monooxygenase 5 (FMO5)-Knockout Mice. *Front. Mol. Biosci.* 2018, 5, 28.
49. Gagliardi, S.; Abel, K.; Bianchi, M.; Milani, P.; Bernuzzi, S.; Corato, M.; Ceroni, M.; Cashman, J.R.; Cereda, C. Regulation of FMO and PON Detoxication Systems in ALS Human Tissues. *Neurotox. Res.* 2013, 23, 370–377.
50. Pang, H.; Yu, X.; Kim, Y.M.; Wang, X.; Jenkins, J.K.; Yin, J.; Li, S.; Gu, H. Disorders Associated With Diverse, Recurrent Deletions and Duplications at 1q21.1. *Front. Genet.* 2020, 11.
51. Rosenfeld, J.A.; Traylor, R.N.; Schaefer, G.B.; McPherson, E.W.; Ballif, B.C.; Klopocki, E.; Mundlos, S.; Shaffer, L.G.; Aylsworth, A.S. Proximal microdeletions and microduplications of 1q21.1 contribute to variable abnormal phenotypes. *Eur. J. Hum. Genet.* 2012, 20, 754–761.
52. Rees, E.; Moskvina, V.; Owen, M.J.; O'Donovan, M.C.; Kirov, G. De Novo Rates and Selection of Schizophrenia-Associated Copy Number Variants. *Biol. Psychiatry* 2011, 70, 1109–1114.
53. Kirov, G.; Pocklington, A.J.; Holmans, P.; Ivanov, D.; Ikeda, M.; Ruderfer, D.; Moran, J.; Chambert, K.; Toncheva, D.; Georgieva, L.; et al. De novo CNV analysis implicates specific abnormalities of postsynaptic signalling complexes in the pathogenesis of schizophrenia. *Mol. Psychiatry* 2012, 17, 142–153.
54. Reinwald, J.R.; Sartorius, A.; Weber-Fahr, W.; Sack, M.; Becker, R.; Didriksen, M.; Stensbøl, T.B.; Schwarz, A.J.; Meyer-Lindenberg, A.; Gass, N. Separable neural mechanisms for the pleiotropic association of copy number variants with neuropsychiatric traits. *Transl. Psychiatry* 2020, 10, 93.

55. Nielsen, J.; Fejgin, K.; Sotty, F.; Nielsen, V.; Mørk, A.; Christoffersen, C.T.; Yavich, L.; Lauridsen, J.B.; Clausen, D.; Larsen, P.H.; et al. A mouse model of the schizophrenia-associated 1q21.1 microdeletion syndrome exhibits altered mesolimbic dopamine transmission. *Transl. Psychiatry* 2017, 7, 1261.
56. Sanders, S.J.; He, X.; Willsey, A.J.; Ercan-Sencicek, A.G.; Samocha, K.E.; Cicek, A.E.; Murtha, M.T.; Bal, V.H.; Bishop, S.L.; Dong, S.; et al. Insights into Autism Spectrum Disorder Genomic Architecture and Biology from 71 Risk Loci. *Neuron* 2015, 87, 1215–1233.
57. Marotta, R.; Risoleo, M.C.; Messina, G.; Parisi, L.; Carotenuto, M.; Vetri, L.; Roccella, M. The Neurochemistry of Autism. *Brain Sci.* 2020, 10, 163.
58. Howes, O.D.; Kambeitz, J.; Kim, E.; Stahl, D.; Slifstein, M.; Abi-Dargham, A.; Kapur, S. The Nature of Dopamine Dysfunction in Schizophrenia and What This Means for Treatment: Meta-analysis of Imaging Studies. *Arch. Gen. Psychiatry* 2012, 69, 776–786.
59. Howes, O.D.; Shotbolt, P.; Bloomfield, M.; Daalman, K.; Demjaha, A.; Diederer, K.M.J.; Ibrahim, K.; Kim, E.; McGuire, P.; Kahn, R.S.; et al. Dopaminergic Function in the Psychosis Spectrum: An [18F]-DOPA Imaging Study in Healthy Individuals with Auditory Hallucinations. *Schizophr. Bull.* 2013, 39, 807–814.
60. Liu, Z.; Osipovitch, M.; Benraiss, A.; Huynh, N.P.T.; Foti, R.; Bates, J.; Chandler-Militello, D.; Findling, R.L.; Tesar, P.J.; Nedergaard, M.; et al. Dysregulated Glial Differentiation in Schizophrenia May Be Relieved by Suppression of SMAD4- and REST-Dependent Signaling. *Cell Rep.* 2019, 27, 3832–3843.e3836.
61. Dror, V.; Shamir, E.; Ghanshani, S.; Kimhi, R.; Swartz, M.; Barak, Y.; Weizman, R.; Avivi, L.; Litmanovitch, T.; Fantino, E.; et al. hKCa3/KCNN3 potassium channel gene: Association of longer CAG repeats with schizophrenia in Israeli Ashkenazi Jews, expression in human tissues and localization to chromosome 1q21. *Mol. Psychiatry* 1999, 4, 254–260.
62. Austin, C.P.; Holder, D.J.; Ma, L.; Mixson, L.A.; Caskey, C.T. Mapping of hKCa3 to chromosome 1q21 and investigation of linkage of CAG repeat polymorphism to schizophrenia. *Mol. Psychiatry* 1999, 4, 261–266.
63. Meissner, B.; Purmann, S.; Schürmann, M.; Zühlke, C.; Lencer, R.; Arolt, V.; Müller-Myhsok, B.; Morris-Rosendahl, D.J.; Schwinger, E. hSKCa3: A candidate gene for schizophrenia? *Psychiatr. Genet.* 1999, 9, 91–96.
64. Miller, M.J.; Rauer, H.; Tomita, H.; Rauer, H.; Gargus, J.J.; Gutman, G.A.; Cahalan, M.D.; Chandy, K.G. Nuclear Localization and Dominant-negative Suppression by a Mutant SKCa3 N-terminal Channel Fragment Identified in a Patient with Schizophrenia. *J. Biol. Chem.* 2001, 276, 27753–27756.

65. Frye, R.E.; Vassall, S.; Kaur, G.; Lewis, C.; Karim, M.; Rossignol, D. Emerging biomarkers in autism spectrum disorder: A systematic review. *Ann. Transl. Med.* 2019, 7, 792.
66. Rossignol, D.A.; Frye, R.E. Mitochondrial dysfunction in autism spectrum disorders: A systematic review and meta-analysis. *Mol. Psychiatry* 2012, 17, 290–314.
67. Ng, C.-H.; Guan, M.S.H.; Koh, C.; Ouyang, X.; Yu, F.; Tan, E.-K.; O'Neill, S.P.; Zhang, X.; Chung, J.; Lim, K.-L. AMP Kinase Activation Mitigates Dopaminergic Dysfunction and Mitochondrial Abnormalities in *Drosophila* Models of Parkinson's Disease. *J. Neurosci.* 2012, 32, 14311–14317.
68. Gordon, A.; Forsingdal, A.; Klewe, I.V.; Nielsen, J.; Didriksen, M.; Werge, T.; Geschwind, D.H. Transcriptomic networks implicate neuronal energetic abnormalities in three mouse models harboring autism and schizophrenia-associated mutations. *Mol. Psychiatry* 2019.
69. Prabakaran, S.; Swatton, J.E.; Ryan, M.M.; Huffaker, S.J.; Huang, J.-J.; Griffin, J.L.; Wayland, M.; Freeman, T.; Dudbridge, F.; Lilley, K.S.; et al. Mitochondrial dysfunction in schizophrenia: Evidence for compromised brain metabolism and oxidative stress. *Mol. Psychiatry* 2004, 9, 684–697.
70. Roberts, R.C. Mitochondrial dysfunction in schizophrenia: With a focus on postmortem studies. *Mitochondrion* 2021, 56, 91–101.
71. Sunyer, J.; Dadvand, P. Pre-natal brain development as a target for urban air pollution. *Basic Clin. Pharmacol. Toxicol.* 2019, 125, 81–88.
72. Windham, G.C.; Pearl, M.; Poon, V.; Berger, K.; Soriano, J.W.; Eyles, D.; Lyall, K.; Kharrazi, M.; Croen, L.A. Maternal Vitamin D Levels During Pregnancy in Association With Autism Spectrum Disorders (ASD) or Intellectual Disability (ID) in Offspring; Exploring Non-linear Patterns and Demographic Sub-groups. *Autism. Res.* 2020, 13, 2216–2229.
73. Jones, H.F.; Ho, A.C.C.; Sharma, S.; Mohammad, S.S.; Kothur, K.; Patel, S.; Brilot, F.; Guastella, A.J.; Dale, R.C.; Group, I.-N.S. Maternal thyroid autoimmunity associated with acute-onset neuropsychiatric disorders and global regression in offspring. *Dev. Med. Child Neurol.* 2019, 61, 984–988.
74. Van den Bergh, B.R.H.; van den Heuvel, M.I.; Lahti, M.; Braeken, M.; de Rooij, S.R.; Entringer, S.; Hoyer, D.; Roseboom, T.; Räikkönen, K.; King, S.; et al. Prenatal developmental origins of behavior and mental health: The influence of maternal stress in pregnancy. *Neurosci. Biobehav. Rev.* 2020, 117, 26–64.
75. Webb, S.J.; Garrison, M.M.; Bernier, R.; McClintic, A.M.; King, B.H.; Mourad, P.D. Severity of ASD symptoms and their correlation with the presence of copy number variations and exposure to first trimester ultrasound. *Autism Res.* 2017, 10, 472–484.
76. Shelton, J.F.; Geraghty, E.M.; Tancredi, D.J.; Delwiche, L.D.; Schmidt, R.J.; Ritz, B.; Hansen, R.L.; Hertz-Picciotto, I. Neurodevelopmental Disorders and Prenatal Residential Proximity to

- Agricultural Pesticides: The CHARGE Study. *Environ. Health Perspect.* 2014, 122, 1103–1109.
77. Pedersen, M.G.; Stevens, H.; Pedersen, C.B.; Norgaard-Pedersen, B.; Mortensen, P.B. Toxoplasma infection and later development of schizophrenia in mothers. *Am. J. Psychiatry* 2011, 168, 814–821.
 78. Dickerson, F.; Kirkpatrick, B.; Boronow, J.; Stallings, C.; Origoni, A.; Yolken, R. Deficit schizophrenia: Association with serum antibodies to cytomegalovirus. *Schizophr. Bull.* 2006, 32, 396–400.
 79. O'Callaghan, E.; Sham, P.; Takei, N.; Glover, G.; Murray, R.M. Schizophrenia after prenatal exposure to 1957 A2 influenza epidemic. *Lancet* 1991, 337, 1248–1250.
 80. Suvisaari, J.; Haukka, J.; Tanskanen, A.; Hovi, T.; Lonnqvist, J. Association between prenatal exposure to poliovirus infection and adult schizophrenia. *Am. J. Psychiatry* 1999, 156, 1100–1102.
 81. Susser, E.S.; Lin, S.P. Schizophrenia after prenatal exposure to the Dutch Hunger Winter of 1944–1945. *Arch. Gen. Psychiatry* 1992, 49, 983–988.
 82. St Clair, D.; Xu, M.; Wang, P.; Yu, Y.; Fang, Y.; Zhang, F.; Zheng, X.; Gu, N.; Feng, G.; Sham, P.; et al. Rates of adult schizophrenia following prenatal exposure to the Chinese famine of 1959–1961. *JAMA* 2005, 294, 557–562.
 83. Shen, Q.; Li, Z.Q.; Sun, Y.; Wang, T.; Wan, C.L.; Li, X.W.; Zhao, X.Z.; Feng, G.Y.; Li, S.; St Clair, D.; et al. The role of pro-inflammatory factors in mediating the effects on the fetus of prenatal undernutrition: Implications for schizophrenia. *Schizophr. Res.* 2008, 99, 48–55.
 84. O'Donnell, K.; O'Connor, T.G.; Glover, V. Prenatal stress and neurodevelopment of the child: Focus on the HPA axis and role of the placenta. *Dev. Neurosci.* 2009, 31, 285–292.
 85. Jablensky, A.V.; Morgan, V.; Zubrick, S.R.; Bower, C.; Yellachich, L.A. Pregnancy, delivery, and neonatal complications in a population cohort of women with schizophrenia and major affective disorders. *Am. J. Psychiatry* 2005, 162, 79–91.
 86. Gradinaru, V.; Zhang, F.; Ramakrishnan, C.; Mattis, J.; Prakash, R.; Diester, I.; Goshen, I.; Thompson, K.R.; Deisseroth, K. Molecular and cellular approaches for diversifying and extending optogenetics. *Cell* 2010, 141, 154–165.
 87. Junge, H.J.; Yang, S.; Burton, J.B.; Paes, K.; Shu, X.; French, D.M.; Costa, M.; Rice, D.S.; Ye, W. TSPAN12 regulates retinal vascular development by promoting Norrin- but not Wnt-induced FZD4/beta-catenin signaling. *Cell* 2009, 139, 299–311.
 88. Ye, X.; Wang, Y.; Cahill, H.; Yu, M.; Badea, T.C.; Smallwood, P.M.; Peachey, N.S.; Nathans, J. Norrin, frizzled-4, and Lrp5 signaling in endothelial cells controls a genetic program for retinal vascularization. *Cell* 2009, 139, 285–298.

89. Phng, L.K.; Potente, M.; Leslie, J.D.; Babbage, J.; Nyqvist, D.; Lobov, I.; Ondr, J.K.; Rao, S.; Lang, R.A.; Thurston, G.; et al. Nrarp coordinates endothelial Notch and Wnt signaling to control vessel density in angiogenesis. *Dev. Cell* 2009, 16, 70–82.
90. Daneman, R.; Agalliu, D.; Zhou, L.; Kuhnert, F.; Kuo, C.J.; Barres, B.A. Wnt/beta-catenin signaling is required for CNS, but not non-CNS, angiogenesis. *Proc. Natl. Acad. Sci. USA* 2009, 106, 641–646.
91. Kalani, M.Y.; Cheshier, S.H.; Cord, B.J.; Bababeygy, S.R.; Vogel, H.; Weissman, I.L.; Palmer, T.D.; Nusse, R. Wnt-mediated self-renewal of neural stem/progenitor cells. *Proc. Natl. Acad. Sci. USA* 2008, 105, 16970–16975.
92. Viti, J.; Gulacsi, A.; Lillien, L. Wnt regulation of progenitor maturation in the cortex depends on Shh or fibroblast growth factor 2. *J. Neurosci.* 2003, 23, 5919–5927.
93. Lyu, J.; Joo, C.K. Wnt signaling enhances FGF2-triggered lens fiber cell differentiation. *Development* 2004, 131, 1813–1824.
94. Kuwabara, T.; Hsieh, J.; Muotri, A.; Yeo, G.; Warashina, M.; Lie, D.C.; Moore, L.; Nakashima, K.; Asashima, M.; Gage, F.H. Wnt-mediated activation of NeuroD1 and retro-elements during adult neurogenesis. *Nat. Neurosci.* 2009, 12, 1097–1105.
95. Munji, R.N.; Choe, Y.; Li, G.; Siegenthaler, J.A.; Pleasure, S.J. Wnt signaling regulates neuronal differentiation of cortical intermediate progenitors. *J. Neurosci.* 2011, 31, 1676–1687.
96. Wang, Q.; Charych, E.I.; Pulito, V.L.; Lee, J.B.; Graziane, N.M.; Crozier, R.A.; Revilla-Sanchez, R.; Kelly, M.P.; Dunlop, A.J.; Murdoch, H.; et al. The psychiatric disease risk factors DISC1 and TNIK interact to regulate synapse composition and function. *Mol. Psychiatry* 2011, 16, 1006–1023.
97. De Rienzo, G.; Bishop, J.A.; Mao, Y.; Pan, L.; Ma, T.P.; Moens, C.B.; Tsai, L.H.; Sive, H. Disc1 regulates both beta-catenin-mediated and noncanonical Wnt signaling during vertebrate embryogenesis. *FASEB J.* 2011, 25, 4184–4197.
98. Lie, D.C.; Colamarino, S.A.; Song, H.J.; Desire, L.; Mira, H.; Consiglio, A.; Lein, E.S.; Jessberger, S.; Lansford, H.; Dearie, A.R.; et al. Wnt signalling regulates adult hippocampal neurogenesis. *Nature* 2005, 437, 1370–1375.
99. L'Episcopo, F.; Serapide, M.F.; Tirolo, C.; Testa, N.; Caniglia, S.; Morale, M.C.; Pluchino, S.; Marchetti, B. A Wnt1 regulated Frizzled-1/beta-Catenin signaling pathway as a candidate regulatory circuit controlling mesencephalic dopaminergic neuron-astrocyte crosstalk: Therapeutical relevance for neuron survival and neuroprotection. *Mol. Neurodegener.* 2011, 6, 49.
100. Faulkner, R.L.; Jang, M.H.; Liu, X.B.; Duan, X.; Sailor, K.A.; Kim, J.Y.; Ge, S.; Jones, E.G.; Ming, G.L.; Song, H.; et al. Development of hippocampal mossy fiber synaptic outputs by new neurons in the adult brain. *Proc. Natl. Acad. Sci. USA* 2008, 105, 14157–14162.

101. Alvarez, A.R.; Godoy, J.A.; Mullendorff, K.; Olivares, G.H.; Bronfman, M.; Inestrosa, N.C. Wnt-3a overcomes beta-amyloid toxicity in rat hippocampal neurons. *Exp. Cell Res.* 2004, 297, 186–196.
102. Chong, Z.Z.; Maiese, K. Targeting WNT, protein kinase B, and mitochondrial membrane integrity to foster cellular survival in the nervous system. *Histol. Histopathol.* 2004, 19, 495–504.
103. Sahores, M.; Gibb, A.; Salinas, P.C. Frizzled-5, a receptor for the synaptic organizer Wnt7a, regulates activity-mediated synaptogenesis. *Development* 2010, 137, 2215–2225.
104. Klassen, M.P.; Shen, K. Wnt signaling positions neuromuscular connectivity by inhibiting synapse formation in *C. elegans*. *Cell* 2007, 130, 704–716.
105. Hall, A.C.; Lucas, F.R.; Salinas, P.C. Axonal remodeling and synaptic differentiation in the cerebellum is regulated by WNT-7a signaling. *Cell* 2000, 100, 525–535.
106. Krylova, O.; Messenger, M.J.; Salinas, P.C. Dishevelled-1 regulates microtubule stability: A new function mediated by glycogen synthase kinase-3beta. *J. Cell Biol.* 2000, 151, 83–94.
107. Lyuksyutova, A.I.; Lu, C.C.; Milanesio, N.; King, L.A.; Guo, N.; Wang, Y.; Nathans, J.; Tessier-Lavigne, M.; Zou, Y. Anterior-posterior guidance of commissural axons by Wnt-frizzled signaling. *Science* 2003, 302, 1984–1988.
108. Zechner, D.; Fujita, Y.; Hülsken, J.; Müller, T.; Walther, I.; Taketo, M.M.; Crenshaw, I.I.I.E.B.; Birchmeier, W.; Birchmeier, C. β -Catenin signals regulate cell growth and the balance between progenitor cell expansion and differentiation in the nervous system. *Dev. Biol.* 2003, 258, 406–418.
109. De Ligt, J.; Willemsen, M.H.; van Bon, B.W.; Kleefstra, T.; Yntema, H.G.; Kroes, T.; Vulto-van Silfhout, A.T.; Koolen, D.A.; de Vries, P.; Gilissen, C.; et al. Diagnostic exome sequencing in persons with severe intellectual disability. *N. Engl. J. Med.* 2012, 367, 1921–1929.
110. Kuechler, A.; Willemsen, M.H.; Albrecht, B.; Bacino, C.A.; Bartholomew, D.W.; van Bokhoven, H.; van den Boogaard, M.J.; Bramswig, N.; Buttner, C.; Cremer, K.; et al. De novo mutations in beta-catenin (CTNNB1) appear to be a frequent cause of intellectual disability: Expanding the mutational and clinical spectrum. *Hum. Genet.* 2015, 134, 97–109.
111. Winczewska-Wiktor, A.; Badura-Stronka, M.; Monies-Nowicka, A.; Nowicki, M.M.; Steinborn, B.; Latos-Bielenska, A.; Monies, D. A de novo CTNNB1 nonsense mutation associated with syndromic atypical hyperekplexia, microcephaly and intellectual disability: A case report. *BMC Neurol.* 2016, 16, 35.
112. Kharbanda, M.; Pilz, D.T.; Tomkins, S.; Chandler, K.; Saggar, A.; Fryer, A.; McKay, V.; Louro, P.; Smith, J.C.; Burn, J.; et al. Clinical features associated with CTNNB1 de novo loss of function mutations in ten individuals. *Eur. J. Med Genet.* 2017, 60, 130–135.

113. Dubruc, E.; Putoux, A.; Labalme, A.; Rougeot, C.; Sanlaville, D.; Edery, P. A new intellectual disability syndrome caused by CTNNB1 haploinsufficiency. *Am. J. Med. Genet. Part A* 2014, 164A, 1571–1575.
114. Tucci, V.; Kleefstra, T.; Hardy, A.; Heise, I.; Maggi, S.; Willemsen, M.H.; Hilton, H.; Esapa, C.; Simon, M.; Buenavista, M.T.; et al. Dominant beta-catenin mutations cause intellectual disability with recognizable syndromic features. *J. Clin. Investig.* 2014, 124, 1468–1482.
115. O’Roak, B.J.; Vives, L.; Girirajan, S.; Karakoc, E.; Krumm, N.; Coe, B.P.; Levy, R.; Ko, A.; Lee, C.; Smith, J.D.; et al. Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations. *Nature* 2012, 485, 246–250.
116. Stessman, H.A.; Xiong, B.; Coe, B.P.; Wang, T.; Hoekzema, K.; Fencikova, M.; Kvarnung, M.; Gerds, J.; Trinh, S.; Cosemans, N.; et al. Targeted sequencing identifies 91 neurodevelopmental-disorder risk genes with autism and developmental-disability biases. *Nat. Genet.* 2017, 49, 515–526.
117. O’Roak, B.J.; Vives, L.; Fu, W.; Egertson, J.D.; Stanaway, I.B.; Phelps, I.G.; Carvill, G.; Kumar, A.; Lee, C.; Ankenman, K.; et al. Multiplex Targeted Sequencing Identifies Recurrently Mutated Genes in Autism Spectrum Disorders. *Science* 2012.
118. Iakoucheva, L.M.; Muotri, A.R.; Sebat, J. Getting to the Cores of Autism. *Cell* 2019, 178, 1287–1298.
119. Krupp, D.R.; Barnard, R.A.; Duffourd, Y.; Evans, S.A.; Mulqueen, R.M.; Bernier, R.; Rivière, J.-B.; Fombonne, E.; O’Roak, B.J. Exonic Mosaic Mutations Contribute Risk for Autism Spectrum Disorder. *Am. J. Hum. Genet.* 2017, 101, 369–390.
120. Dong, F.; Jiang, J.; McSweeney, C.; Zou, D.; Liu, L.; Mao, Y. Deletion of CTNNB1 in inhibitory circuitry contributes to autism-associated behavioral defects. *Hum. Mol. Genet.* 2016, 25, 2738–2751.
121. Mohn, J.L.; Alexander, J.; Pirone, A.; Palka, C.D.; Lee, S.Y.; Mebane, L.; Haydon, P.G.; Jacob, M.H. Adenomatous polyposis coli protein deletion leads to cognitive and autism-like disabilities. *Mol. Psychiatry* 2014, 19, 1133–1142.
122. Lijam, N.; Paylor, R.; McDonald, M.P.; Crawley, J.N.; Deng, C.-X.; Herrup, K.; Stevens, K.E.; Maccaferri, G.; McBain, C.J.; Sussman, D.J.; et al. Social Interaction and Sensorimotor Gating Abnormalities in Mice Lacking Dvl1. *Cell* 1997, 90, 895–905.
123. Shin, S.; Santi, A.; Huang, S. Conditional Pten knockout in parvalbumin- or somatostatin-positive neurons sufficiently leads to autism-related behavioral phenotypes. *Mol. Brain* 2021, 14, 24.
124. Busch, R.M.; Srivastava, S.; Hogue, O.; Frazier, T.W.; Klaas, P.; Hardan, A.; Martinez-Agosto, J.A.; Sahin, M.; Eng, C. Neurobehavioral phenotype of autism spectrum disorder associated with germline heterozygous mutations in PTEN. *Transl. Psychiatry* 2019, 9, 253.

125. Chen, C.-J.; Sgritta, M.; Mays, J.; Zhou, H.; Lucero, R.; Park, J.; Wang, I.C.; Park, J.H.; Kaiparettu, B.A.; Stoica, L.; et al. Therapeutic inhibition of mTORC2 rescues the behavioral and neurophysiological abnormalities associated with Pten-deficiency. *Nat. Med.* 2019, 25, 1684–1690.
126. Lugo, J.N.; Smith, G.D.; Arbuckle, E.P.; White, J.; Holley, A.J.; Floruta, C.M.; Ahmed, N.; Gomez, M.C.; Okonkwo, O. Deletion of PTEN produces autism-like behavioral deficits and alterations in synaptic proteins. *Front. Mol. Neurosci.* 2014, 7.
127. Kramps, T.; Peter, O.; Brunner, E.; Nellen, D.; Froesch, B.; Chatterjee, S.; Murone, M.; Züllig, S.; Basler, K. Wnt/Wingless Signaling Requires BCL9/Legless-Mediated Recruitment of Pygopus to the Nuclear β -Catenin-TCF Complex. *Cell* 2002, 109, 47–60.
128. Brembeck, F.H.; Schwarz-Romond, T.; Bakkers, J.; Wilhelm, S.; Hammerschmidt, M.; Birchmeier, W. Essential role of BCL9-2 in the switch between β -catenin's adhesive and transcriptional functions. *Genes Dev.* 2004, 18, 2225–2230.
129. Mieszczanek, J.; de la Roche, M.; Bienz, M. A role of Pygopus as an anti-repressor in facilitating Wnt-dependent transcription. *Proc. Natl. Acad. Sci. USA* 2008, 105, 19324–19329.
130. Brack, A.S.; Murphy-Seiler, F.; Hanifi, J.; Deka, J.; Eyckerman, S.; Keller, C.; Aguet, M.; Rando, T.A. BCL9 is an essential component of canonical Wnt signaling that mediates the differentiation of myogenic progenitors during muscle regeneration. *Dev. Biol.* 2009, 335, 93–105.
131. Xu, C.; Aragam, N.; Li, X.; Villa, E.C.; Wang, L.; Briones, D.; Petty, L.; Posada, Y.; Arana, T.B.; Cruz, G.; et al. BCL9 and C9orf5 Are Associated with Negative Symptoms in Schizophrenia: Meta-Analysis of Two Genome-Wide Association Studies. *PLoS ONE* 2013, 8, e51674.
132. Kimura, H.; Tanaka, S.; Kushima, I.; Koide, T.; Banno, M.; Kikuchi, T.; Nakamura, Y.; Shiino, T.; Yoshimi, A.; Oya-Ito, T.; et al. Association study of BCL9 gene polymorphism rs583583 with schizophrenia and negative symptoms in Japanese population. *Sci. Rep.* 2015, 5, 15705.
133. Luo, X.; Huang, L.; Han, L.; Luo, Z.; Hu, F.; Tieu, R.; Gan, L. Systematic Prioritization and Integrative Analysis of Copy Number Variations in Schizophrenia Reveal Key Schizophrenia Susceptibility Genes. *Schizophr. Bull.* 2014, 40, 1285–1299.
134. Takada, K.; Zhu, D.; Bird, G.H.; Sukhdeo, K.; Zhao, J.-J.; Mani, M.; Lemieux, M.; Carrasco, D.E.; Ryan, J.; Horst, D.; et al. Targeted Disruption of the BCL9/ β -Catenin Complex Inhibits Oncogenic Wnt Signaling. *Sci. Transl. Med.* 2012, 4, 148ra117.
135. Moor, A.E.; Anderle, P.; Cantù, C.; Rodriguez, P.; Wiedemann, N.; Baruthio, F.; Deka, J.; André, S.; Valenta, T.; Moor, M.B.; et al. BCL9/9L- β -catenin Signaling is Associated With Poor Outcome in Colorectal Cancer. *EBioMedicine* 2015, 2, 1932–1943.
136. Beaulieu, J.-F. Tuning WNT- β -catenin signaling via BCL9 proteins for targeting colorectal cancer cells. *EBioMedicine* 2015, 2, 1846–1847.

137. Miyaoka, T.; Seno, H.; Ishino, H. Increased expression of Wnt-1 in schizophrenic brains. *Schizophr. Res.* 1999, 38, 1–6.
138. Proitsi, P.; Li, T.; Hamilton, G.; Di Forti, M.; Collier, D.; Killick, R.; Chen, R.; Sham, P.; Murray, R.; Powell, J.; et al. Positional pathway screen of wnt signaling genes in schizophrenia: Association with DKK4. *Biol. Psychiatry* 2008, 63, 13–16.
139. Yang, J.; Si, T.; Ling, Y.; Ruan, Y.; Han, Y.; Wang, X.; Zhang, H.; Kong, Q.; Li, X.; Liu, C.; et al. Association study of the human FZD3 locus with schizophrenia. *Biol. Psychiatry* 2003, 54, 1298–1301.
140. Ide, M.; Muratake, T.; Yamada, K.; Iwayama-Shigeno, Y.; Iwamoto, K.; Takao, H.; Toyota, T.; Kaneko, N.; Minabe, Y.; Nakamura, K.; et al. Genetic and expression analyses of FZD3 in schizophrenia. *Biol. Psychiatry* 2004, 56, 462–465.
141. Zhang, Y.; Yu, X.; Yuan, Y.; Ling, Y.; Ruan, Y.; Si, T.; Lu, T.; Wu, S.; Gong, X.; Zhu, Z.; et al. Positive association of the human frizzled 3 (FZD3) gene haplotype with schizophrenia in Chinese Han population. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 2004, 129, 16–19.
142. Lachman, H.M.; Pedrosa, E.; Petruolo, O.A.; Cockerham, M.; Papolos, A.; Novak, T.; Papolos, D.F.; Stopkova, P. Increase in GSK3beta gene copy number variation in bipolar disorder. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 2007, 144, 259–265.
143. Martin, P.M.; Yang, X.; Robin, N.; Lam, E.; Rabinowitz, J.S.; Erdman, C.A.; Quinn, J.; Weiss, L.A.; Hamilton, S.P.; Kwok, P.Y.; et al. A rare WNT1 missense variant overrepresented in ASD leads to increased Wnt signal pathway activation. *Transl. Psychiatry* 2013, 3, e301.
144. Millar, J.K.; Wilson-Annan, J.C.; Anderson, S.; Christie, S.; Taylor, M.S.; Semple, C.A.; Devon, R.S.; Clair, D.M.; Muir, W.J.; Blackwood, D.H.; et al. Disruption of two novel genes by a translocation co-segregating with schizophrenia. *Hum. Mol. Genet.* 2000, 9, 1415–1423.
145. Kim, H.J.; Park, H.J.; Jung, K.H.; Ban, J.Y.; Ra, J.; Kim, J.W.; Park, J.K.; Choe, B.K.; Yim, S.V.; Kwon, Y.K.; et al. Association study of polymorphisms between DISC1 and schizophrenia in a Korean population. *Neurosci. Lett.* 2007, 26, 81–96.
146. Tomppa, L.; Hennah, W.; Lahermo, P.; Loukola, A.; Tuulio-Henriksson, A.; Suvisaari, J.; Partonen, T.; Ekelund, J.; Lonnqvist, J.; Peltonen, L. Association between genes of Disrupted in schizophrenia 1 (DISC1) interactors and schizophrenia supports the role of the DISC1 pathway in the etiology of major mental illnesses. *Biol. Psychiatry* 2009, 65, 1055–1062.
147. Schosser, A.; Gaysina, D.; Cohen-Woods, S.; Chow, P.C.; Martucci, L.; Craddock, N.; Farmer, A.; Korszun, A.; Gunasinghe, C.; Gray, J.; et al. Association of DISC1 and TSNAX genes and affective disorders in the depression case-control (DeCC) and bipolar affective case-control (BACCS) studies. *Mol. Psychiatry* 2009.

148. Hennah, W.; Thomson, P.; McQuillin, A.; Bass, N.; Loukola, A.; Anjorin, A.; Blackwood, D.; Curtis, D.; Deary, I.J.; Harris, S.E.; et al. DISC1 association, heterogeneity and interplay in schizophrenia and bipolar disorder. *Mol. Psychiatry* 2009, 14, 865–873.
149. Kilpinen, H.; Ylisaukko-Oja, T.; Hennah, W.; Palo, O.M.; Varilo, T.; Vanhala, R.; Nieminen-von Wendt, T.; von Wendt, L.; Paunio, T.; Peltonen, L. Association of DISC1 with autism and Asperger syndrome. *Mol. Psychiatry* 2007, 13, 187–196.
150. Mao, Y.; Ge, X.; Frank, C.L.; Madison, J.M.; Koehler, A.N.; Doud, M.K.; Tassa, C.; Berry, E.M.; Soda, T.; Singh, K.K.; et al. Disrupted in schizophrenia 1 regulates neuronal progenitor proliferation via modulation of GSK3beta/beta-catenin signaling. *Cell* 2009, 136, 1017–1031.

Retrieved from <https://encyclopedia.pub/entry/history/show/26661>