

Megalencephaly-Capillary Malformation Syndrome

Subjects: Genetics & Heredity

Contributor: Rita Xu

Megalencephaly-capillary malformation syndrome (MCAP) is a disorder characterized by overgrowth of several tissues in the body. Its primary features are a large brain (megalencephaly) and abnormalities of small blood vessels in the skin called capillaries (capillary malformations).

Keywords: genetic conditions

1. Introduction

In individuals with MCAP, megalencephaly leads to an unusually large head size (macrocephaly), which is typically evident at birth. After birth, the brain and head continue to grow at a fast rate for the first few years of life; then, the growth slows to a normal rate, although the head remains larger than average. Additional brain abnormalities are common in people with MCAP; these can include excess fluid within the brain (hydrocephalus) and abnormalities in the brain's structure, such as those known as Chiari malformation and polymicrogyria. Abnormal brain development leads to intellectual disability in most affected individuals and can also cause seizures or weak muscle tone (hypotonia). In particular, polymicrogyria is associated with speech delays and difficulty chewing and swallowing.

The capillary malformations characteristic of MCAP are composed of enlarged capillaries that increase blood flow near the surface of the skin. These malformations usually look like pink or red spots on the skin. In most affected individuals, capillary malformations occur on the face, particularly the nose, the upper lip, and the area between the nose and upper lip (the philtrum). In other people with MCAP, the malformations appear as patches spread over the body or as a reddish net-like pattern on the skin (cutis marmorata).

In some people with MCAP, excessive growth affects not only the brain but other individual parts of the body, which is known as segmental overgrowth. This can lead to one arm or leg that is bigger or longer than the other or a few oversized fingers or toes. Some affected individuals have fusion of the skin between two or more fingers or toes (cutaneous syndactyly).

Additional features of MCAP can include flexible joints and skin that stretches easily. Some affected individuals are said to have doughy skin because the tissue under the skin is unusually thick and soft.

The gene involved in MCAP is also associated with several types of cancer. Only a small number of individuals with MCAP have developed tumors (in particular, a childhood form of kidney cancer known as Wilms tumor and noncancerous tumors in the nervous system known as meningiomas).

2. Frequency

The prevalence of MCAP is unknown. At least 150 affected individuals have been reported in the medical literature. Because the condition is often thought to be misdiagnosed or underdiagnosed, it may be more common than reported.

3. Causes

MCAP is caused by mutations in the *PIK3CA* gene, which provides instructions for making the p110 alpha (p110 α) protein. This protein is one piece (subunit) of an enzyme called phosphatidylinositol 3-kinase (PI3K), which plays a role in chemical signaling within cells. PI3K signaling is important for many cell activities, including cell growth and division (proliferation), movement (migration) of cells, and cell survival. These functions make PI3K important for the development of tissues throughout the body, including the brain and blood vessels.

PIK3CA gene mutations involved in MCAP alter the p110 α protein. The altered subunit makes PI3K abnormally active, which allows cells to grow and divide continuously. Increased cell proliferation leads to the overgrowth of the brain, blood vessels, and other organs and tissues characteristic of MCAP.

MCAP is one of several overgrowth syndromes, including Klippel-Trenaunay syndrome, that are caused by mutations in the *PIK3CA* gene. Together, these conditions are known as the *PIK3CA*-related overgrowth spectrum (PROS).

3.1. The Gene Associated with Megalencephaly-Capillary Malformation Syndrome

PIK3CA

4. Inheritance

MCAP is not inherited from a parent and does not run in families. In people with MCAP, a *PIK3CA* gene mutation arises randomly in one cell during the early stages of development before birth. As cells continue to divide, some cells will have the mutation and other cells will not. This mixture of cells with and without a genetic mutation is known as mosaicism.

5. Other Names for This Condition

- M-CM
- macrocephaly cutis marmorata telangiectatica congenita
- macrocephaly-capillary malformation syndrome
- MCAP
- MCMTC
- megalencephaly cutis marmorata telangiectatica congenita
- megalencephaly-capillary malformation-polymicrogyria syndrome

References

1. Luks VL, Kamitaki N, Vivero MP, Uller W, Rab R, Bovée JV, Rialon KL, GuevaraCJ, Alomari AI, Greene AK, Fishman S J, Kozakewich HP, Maclellan RA, Mulliken JB, Rahbar R, Spencer SA, Trenor CC 3rd, Upton J, Zurakowski D, Perkins JA, Kirsh A, Bennett JT, Dobyns WB, Kurek KC, Warman ML, McCarroll SA, Murillo R. Lymphatic and other vascular malformative/overgrowth disorders are caused by somatic mutations in *PIK3CA*. *J Pediatr*. 2015 Apr;166(4):1048-54.e1-5. doi:10.1016/j.jpeds.2014.12.069.
2. Martínez-Glez V, Romanelli V, Mori MA, Gracia R, Segovia M, González-Meneses A, López-Gutierrez JC, Gean E, Martorell L, Lapunzina P. Macrocephaly-capillary malformation: Analysis of 13 patients and review of the diagnostic criteria. *Am J Med Genet A*. 2010 Dec;152A(12):3101-6. doi: 10.1002/ajmg.a.33514.
3. Mirzaa GM, Conway RL, Gripp KW, Lerman-Sagie T, Siegel DH, deVries LS, Lev D, Kramer N, Hopkins E, Graham JM Jr, Dobyns WB. Megalencephaly-capillary malformation (MCAP) and megalencephaly-polydactyly-polymicrogyria-hydrocephalus (MPPH) syndromes: two closely related disorders of brain overgrowth and abnormal brain and body morphogenesis. *Am J Med Genet A*. 2012 Feb;158A(2):269-91. doi:10.1002/ajmg.a.34402.
4. Mirzaa GM, Rivière JB, Dobyns WB. Megalencephaly syndromes and activating mutations in the PI3K-AKT pathway: MPPH and MCAP. *Am J Med Genet C Semin Med Genet*. 2013 May;163C(2):122-30. doi: 10.1002/ajmg.c.31361.
5. Rivière JB, Mirzaa GM, O'Roak BJ, Beddaoui M, Alcantara D, Conway RL, St-Onge J, Schwartzentruber JA, Gripp KW, Nikkel SM, Worthylake T, Sullivan CT, Ward TR, Butler HE, Kramer NA, Albrecht B, Armour CM, Armstrong L, Caluseriu O, Cytrynbaum C, Drolet BA, Innes AM, Lauzon JL, Lin AE, Mancini GM, Meschino WS, Reggin JD, Saggar AK, Lerman-Sagie T, Uyanik G, Weksberg R, Zirn B, Beaulieu CL; Finding of Rare Disease Genes (FORGE) Canada Consortium, Majewski J, Bulman DE, O'Driscoll M, Shendure J, Graham JM Jr, Boycott KM, Dobyns WB. De novo germline and postzygotic mutations in *AKT3*, *PIK3R2* and *PIK3CA* cause a spectrum of related megalencephaly syndromes. *Nat Genet*. 2012 Jun 24;44(8):934-40. doi:10.1038/ng.2331.