

Outer Membrane Vesicles

Subjects: Biochemistry & Molecular Biology

Contributor: Gianluigi Franci

OMVs are rounded nanostructures released during their growth by Gram-negative bacteria. Biologically active toxins and virulence factors are often entrapped within these vesicles that behave as molecular carriers. Recently, OMVs have been reported to contain DNA molecules, but little is known about the vesicle packaging, release, and transfer mechanisms.

Keywords: outer membrane vesicles ; horizontal gene transfer ; gram-negative bacteria ; DNA

1. Introduction

Many successful bacterial infections are consequences of bacterial virulence mechanisms associated with antimicrobial escape [1]. The transduction involves bacteriophages that transfer DNA to bacterial cells through infection. Plasmid transfer systems are strictly dependent on the characteristics of the genetic material to be transferred: conjugative plasmids contain genes necessary for pilus formation, while mobilizable plasmids carry mobility genes and transfer origin, lacking pilus coding genes [2][3]. Recent studies have revealed a novel mechanism for gene transfer via OMVs.

2. OMVs: An Overview from Structure to Function

OMVs (**Figure 2**) are spherical bi-layered membrane nanostructures (50–500 nm), secreted by Gram-negative bacteria through bulging and ‘pinching off’ of the outer membrane [4]. Purification techniques have revealed the specific proteins and lipid composition of these vesicles [5]. The lumen of the vesicles contains periplasmic proteins, cytosolic components, and nucleic acids [6]. The composition of OMVs is modulated by growth conditions and is also highly influenced from the interactions between host cells and bacteria [7][8].

Several studies have demonstrated the presence within OMVs of outer membrane proteins, periplasmic proteins, and different virulence factors, engaged in the adhesion and invasion of cell hosts [9]. The first group mainly includes outer membrane proteins such as porins, components of transport systems, adhesins, phospholipases, and proteases. In *Neisseria meningitidis* OMVs, porin A, factor H binding protein, and opacity-associated protein C represent the largest protein fraction [10]. OMVs from *Treponema denticola* contain active proteases that cause damage to host cells [11].

Lipids play an essential role in the structure of OMVs and consist of phospholipids and lipopolysaccharides. While phospholipids constitute the inner sheet of the outer membrane, LPS is exclusively located on the outside surface of the outer membrane. In *P. aeruginosa*, the OMVs phosphatidylglycerol and stearic acid are abundantly detected, proving greater rigidity of the vesicles [12]. *Helicobacter pylori* OMVs have cardiolipin as the main lipid component [9]. The type of LPS band present on OMVs depends on the site of vesicle budding [4].

OMVs carry DNA and RNA on their surface or in the vesicular lumen. A clear difference can be observed by OMV treatment with DNase and RNase: luminal DNA and RNA are preserved even after the enzymatic treatment [13][14]. Furuse et al. have recently identified tRNA fragments in the OMVs of *Chlamydia* and *Legionella* strains, involved in direct subversion of host gene translation and mRNA stability [15]. Although novel information on this topic is piling up at a steady pace, nucleic acid incorporation mechanisms remain to be elucidated.

In the first model, Burdett et al. suggested that the absence or transfer of covalent links between the outer membrane and peptidoglycan layer promotes vesiculation [16]. The second model showed that the accumulation of peptidoglycan fragments or poorly folded proteins in periplasmic space exert pressure on the outer membrane, determining the curvature of the membrane and final budding [17]. Sequestration of positively charged compounds (Mg²⁺ and Ca²⁺ salt) by PQS causes anionic repulsion of lipopolysaccharides, increasing vesiculations [18]. These studies indicate that bacteria developed the OMV production mechanism as a part of stress response to ensure bacterial survival [19].

Several biological functions are attributed to the OMVs. These vesicles represent a long-distance delivery system of biomolecules, such as nucleic acids, enzymes, toxins, and virulence factors, protecting them from extracellular degradation and dilution.

Vesiculation allows intraspecies and interspecies communications and contributes to interaction with the host [20]. In addition, OMVs are involved in the acquisition of nutrients, stress responses, and the formation of a microenvironment necessary for the survival of pathogens [8][21][22]. The packaging of proteases, phosphatase, and glycosidases in OMVs plays an important role in the degradation of complex molecules, promoting nutrient availability [23]. Evans et al. have shown that alkaline phosphatase in *Myxococcus xanthus* OMVs causes the release of phosphate, contributing to the development of multicellular communities [24].

OMVs participate in the formation of the biofilm matrix by the release of exopolysaccharides, thus increasing cell co-aggregation [25][26]. *P. aeruginosa* is able to form biofilms and cause surgical site infections, orthopedic peri-implant bone infections, and lung infection in cystic fibrosis patients. They found that the addition of OMVs promoted biofilm formation in a dose-dependent manner (except for one strain, which turned out to be non-producer). The bacterium–host interactions trigger the release of OMVs, which carry toxins and adhesion and virulence factors [27]. The Shiga toxin in *Escherichia coli* OMVs efficiently inhibits eukaryotic protein synthesis compared to the soluble form [28].

In addition to the reported functions, OMVs have recently been recognized as gene transfer vectors [29]. Several studies have detected plasmids, chromosomal DNA fragments, bacteriophage DNA and RNA fragments in OMVs [30][31][32][33]. Therefore, in the following section, recent evidence on the role of OMVs as carriers for horizontal gene transfer will be reported.

3. Horizontal Gene Transfer Mediated by OMVs

Gene transfer can occur via proven processes of transformation, conjugation, and transduction, as well as through recently identified OMV-mediated mechanisms [34]. Few studies have evaluated the gene transfer potential of OMVs.

were the first to identify OMVs as gene transfer vectors. Polymerase chain reaction (PCR) data revealed the presence of the *eae*, *uidA*, *stx1*, and *stx2* virulence genes in the luminal space. HGT was proved through PCR amplification of virulence genes in transformed *Escherichia coli* JM109 [35]. These first findings laid the foundations for other investigations, deepening the role of OMVs in gene transfer mechanisms.

proved that genetic exchanges through OMVs can also occur between bacteria of different species. Transformation experiments were performed, using *Escherichia coli* JM109 and *Salmonella enterica* serovar Enteritidis as recipient cells, and target genes were determined by colony PCR amplification. The acquisition of virulence genes in the recipient cells resulted in an increase in pathogenicity; the latter was assessed by Vero cell assay. The transformed recipient strains induced a cytotoxicity six times higher than the unprocessed strains, indicating the expression of virulence factors only in transformed strains.

Besides *E. coli*, other Gram-negative species exploit OMVs as HGT vectors. proved that *P. gingivalis* OMVs mediated the transfer of virulence genes between members of the same species. Genes encoding the major subunit of long fimbriae (*fimA*) and superoxide dismutase (*sod*) were detected in the vesicular lumen by PCR analysis, suggesting possible preferential DNA packaging. OMV–HGT experiments were conducted using a mutant *P. gingivalis* 49,417, obtained by introducing a 2.1 Kb segment of the *ermF-ermAM* gene into the *fimA* gene, which conferred resistance to erythromycin.

The involvement of OMVs in the spread of resistance genes was only revealed 10 years later. identified for the first time OMVs as vectors of antibiotic resistance gene transfer. Chatterjee et al. also supported the transfer of antibiotic resistance genes through OMVs. The recipient cells were positive for the *bla*NDM-1 gene and exhibited a broad profile of resistance to β -lactam antibiotics, recording higher MIC values, compared to untransformed strains.

DNA packaging into vesicular lumens protected it from adverse environmental conditions, demonstrating an additional bacterial survival advantage associated with OMV–HGT. After DNase treatment of OMVs, plasmid integrity in the vesicles was evaluated through agarose gel electrophoresis and HindIII plasmid digestion. Vesicular plasmid integrity was not compromised, which indicates that OMVs provide protection for the DNA within them. In the thermal environments and in the presence of DNase, OMVs increased the frequency of transformation, compared to free DNA.

Despite different studies that have shown a high gene transfer potential of OMVs, Renelli et al. suggested that vesicles carry the genetic material but do not promote an effective transformation. PCR amplification of plasmid (pAK1900) and chromosomal (*oprL*) sequences indicated the presence of only plasmid DNA within vesicles. The transformation

experiments were performed using *P. aeruginosa* PAO1 and *Escherichia coli* DH5a as recipient strains. The author speculated that p-OMVs transferred the plasmid into the periplasm of recipient cells, which does not by-pass the plasma membrane for efficient transformation [36].

Although previous studies have highlighted the role of OMVs in HGT, the mechanisms underlying the transfer and the factors influencing the process are not yet clear [31][37]. Few studies have been conducted to understand these aspects.

pLC291 and pUC19 are high-copy number plasmids, while pZS2501 is a low-copy plasmid. Furthermore, they investigated whether the OMVs released by different recipient strains were endowed with different characteristics. Purified vesicles from different recipient strains contained the same protein and plasmid amounts and had a similar size. The vesicles containing pLC291, isolated from *Aeromonas veronii*, *Enterobacter cloacae*, and *Escherichia coli*, were exploited to induce the transformation of five different recipient strains, i.e., *Aeromonas veronii*, *Enterobacter cloacae*, *Escherichia coli*, *Chromobacterium violaceum*, and *P. aeruginosa*.

In a subsequent investigation, Tran et al. evaluated more closely the effect of plasmid features, such as plasmid copy number, size, and origin of replication on OMV-mediated gene transfer. In addition, this study assessed the impact of plasmid size on vesicle loading. Moreover, qPCR results showed that the plasmid size inversely affected the number of plasmid copies in the vesicle. To assess the impact of plasmid origin on OMV production, they constructed three plasmids based on plasmid pLC291, with the same size (3.5 kb) but different origins:

The vesicles purified by treated bacteria were characterized on the basis of protein and DNA contents. Treatment of bacteria with antibiotics and nutrient deprivation caused a significant increase in vesicular diameters. The zeta potential of the OMV produced in the presence of gentamicin showed more negative values compared to the other treatments. These results proved that stress factors can influence the vesicle release, DNA content, and vesicle size [38].

References

1. Beceiro, A.; Tomas, M.; Bou, G. Antimicrobial resistance and virulence: A successful or deleterious association in the bacterial world? *Clin. Microbiol. Rev.* 2013, 26, 185–230.
2. Cabezon, E.; Ripoll-Rozada, J.; Pena, A.; de la Cruz, F.; Arechaga, I. Towards an integrated model of bacterial conjugation. *FEMS Microbiol. Rev.* 2015, 39, 81–95.
3. Fang, Z.; Zhou, H. Identification of the conjugative and mobilizable plasmid fragments in the plasmidome using sequence signatures. *Microb. Genom.* 2020, 6.
4. Kulp, A.; Kuehn, M.J. Biological functions and biogenesis of secreted bacterial outer membrane vesicles. *Annu. Rev. Microbiol.* 2010, 64, 163–184.
5. Klimentova, J.; Stulik, J. Methods of isolation and purification of outer membrane vesicles from gram-negative bacteria. *Microbiol. Res.* 2015, 170, 1–9.
6. Qing, G.; Gong, N.; Chen, X.; Chen, J.; Zhang, H.; Wang, Y.; Wang, R.; Zhang, S.; Zhang, Z.; Zhao, X.; et al. Natural and engineered bacterial outer membrane vesicles. *Biophys. Rep.* 2019, 5, 184–198.
7. Orench-Rivera, N.; Kuehn, M.J. Environmentally controlled bacterial vesicle-mediated export. *Cell Microbiol.* 2016, 18, 1525–1536.
8. Cecil, J.D.; Sirisaengtaksin, N.; O'Brien-Simpson, N.M.; Krachler, A.M. Outer membrane vesicle-host cell interactions. *Microbiol. Spectr.* 2019, 7.
9. Yu, Y.J.; Wang, X.H.; Fan, G.C. Versatile effects of bacterium-released membrane vesicles on mammalian cells and infectious/inflammatory diseases. *Acta Pharm. Sin.* 2018, 39, 514–533.
10. Gerritzen, M.J.H.; Stangowez, L.; van de Waterbeemd, B.; Martens, D.E.; Wijffels, R.H.; Stork, M. Continuous production of *Neisseria meningitidis* outer membrane vesicles. *Appl. Microbiol. Biotechnol.* 2019, 103, 9401–9410.
11. Chi, B.; Qi, M.; Kuramitsu, H.K. Role of dentilisin in *Treponema denticola* epithelial cell layer penetration. *Res. Microbiol.* 2003, 154, 637–643.
12. Baumgarten, T.; Sperling, S.; Seifert, J.; von Bergen, M.; Steiniger, F.; Wick, L.Y.; Heipieper, H.J. Membrane vesicle formation as a multiple-stress response mechanism enhances *Pseudomonas putida* DOT-T1E cell surface hydrophobicity and biofilm formation. *Appl. Environ. Microbiol.* 2012, 78, 6217–6224.
13. Jan, A.T. Outer Membrane Vesicles (OMVs) of gram-negative bacteria: A perspective update. *Front. Microbiol.* 2017, 8, 1053.

14. Fulsundar, S.; Domingues, S.; Nielsen, K.M. Vesicle-Mediated Gene Transfer in *Acinetobacter baumannii*. *Methods Mol. Biol.* 2019, 1946, 87–94.
15. Furuse, Y.; Finethy, R.; Saka, H.A.; Xet-Mull, A.M.; Sisk, D.M.; Smith, K.L.; Lee, S.; Coers, J.; Valdivia, R.H.; Tobin, D.M.; et al. Search for microRNAs expressed by intracellular bacterial pathogens in infected mammalian cells. *PLoS ONE* 2014, 9, e106434.
16. Burdett, I.D.; Murray, R.G. Electron microscope study of septum formation in *Escherichia coli* strains B and B-r during synchronous growth. *J. Bacteriol.* 1974, 119, 1039–1056.
17. Schwechheimer, C.; Kuehn, M.J. Outer-membrane vesicles from Gram-negative bacteria: Biogenesis and functions. *Nat. Rev. Microbiol.* 2015, 13, 605–619.
18. Li, A.; Schertzer, J.W.; Yong, X. Molecular dynamics modeling of *Pseudomonas aeruginosa* outer membranes. *Phys. Chem. Chem. Phys.* 2018, 20, 23635–23648.
19. Macdonald, I.A.; Kuehn, M.J. Stress-induced outer membrane vesicle production by *Pseudomonas aeruginosa*. *J. Bacteriol.* 2013, 195, 2971–2981.
20. Berleman, J.; Auer, M. The role of bacterial outer membrane vesicles for intra- and interspecies delivery. *Environ. Microbiol.* 2013, 15, 347–354.
21. Yanez-Mo, M.; Siljander, P.R.; Andreu, Z.; Zavec, A.B.; Borrás, F.E.; Buzas, E.I.; Buzas, K.; Casal, E.; Cappello, F.; Carvalho, J.; et al. Biological properties of extracellular vesicles and their physiological functions. *J. Extracell. Vesicles* 2015, 4, 27066.
22. Pathirana, R.D.; Kaparakis-Liaskos, M. Bacterial membrane vesicles: Biogenesis, immune regulation and pathogenesis. *Cell Microbiol.* 2016, 18, 1518–1524.
23. Valguarnera, E.; Scott, N.E.; Azimzadeh, P.; Feldman, M.F. Surface exposure and packing of lipoproteins into outer membrane vesicles are coupled processes in bacterioides. *mSphere* 2018, 3.
24. Berleman, J.E.; Allen, S.; Danielewicz, M.A.; Remis, J.P.; Gorur, A.; Cunha, J.; Hadi, M.Z.; Zusman, D.R.; Northen, T.R.; Witkowska, H.E.; et al. The lethal cargo of *Myxococcus xanthus* outer membrane vesicles. *Front. Microbiol.* 2014, 5, 474.
25. Cooke, A.C.; Nello, A.V.; Ernst, R.K.; Schertzer, J.W. Analysis of *Pseudomonas aeruginosa* biofilm membrane vesicles supports multiple mechanisms of biogenesis. *PLoS ONE* 2019, 14, e0212275.
26. Roier, S.; Zingl, F.G.; Cakar, F.; Schild, S. Bacterial outer membrane vesicle biogenesis: A new mechanism and its implications. *Microb. Cell* 2016, 3, 257–259.
27. Bielaszewska, M.; Ruter, C.; Bauwens, A.; Greune, L.; Jarosch, K.A.; Steil, D.; Zhang, W.; He, X.; Lloubes, R.; Fruth, A.; et al. Host cell interactions of outer membrane vesicle-associated virulence factors of enterohemorrhagic *Escherichia coli* O157: Intracellular delivery, trafficking and mechanisms of cell injury. *PLoS Pathog.* 2017, 13, e1006159.
28. Bauwens, A.; Kunsmann, L.; Marejkova, M.; Zhang, W.; Karch, H.; Bielaszewska, M.; Mellmann, A. Intrahost milieu modulates production of outer membrane vesicles, vesicle-associated Shiga toxin 2a and cytotoxicity in *Escherichia coli* O157:H7 and O104:H4. *Environ. Microbiol. Rep.* 2017, 9, 626–634.
29. Domingues, S.; Nielsen, K.M. Membrane vesicles and horizontal gene transfer in prokaryotes. *Curr. Opin. Microbiol.* 2017, 38, 16–21.
30. Perez-Cruz, C.; Carrion, O.; Delgado, L.; Martinez, G.; Lopez-Iglesias, C.; Mercade, E. New type of outer membrane vesicle produced by the Gram-negative bacterium *Shewanella vesiculosa* M7T: Implications for DNA content. *Appl. Environ. Microbiol.* 2013, 79, 1874–1881.
31. Velimirov, B.; Ranftler, C. Unexpected aspects in the dynamics of horizontal gene transfer of prokaryotes: The impact of outer membrane vesicles. *Wien. Med. Wochenschr.* 2018, 168, 307–313.
32. Gaudin, M.; Krupovic, M.; Marguet, E.; Gauliard, E.; Cvirkaitė-Krupovic, V.; Le Cam, E.; Oberto, J.; Forterre, P. Extracellular membrane vesicles harbouring viral genomes. *Environ. Microbiol.* 2014, 16, 1167–1175.
33. Medvedeva, E.S.; Baranova, N.B.; Mouzykantov, A.A.; Grigorieva, T.Y.; Davydova, M.N.; Trushin, M.V.; Chernova, O.A.; Chernov, V.M. Adaptation of mycoplasmas to antimicrobial agents: *Acholeplasma laidlawii* extracellular vesicles mediate the export of ciprofloxacin and a mutant gene related to the antibiotic target. *Sci. World J.* 2014, 2014, 150615.
34. Johnston, C.; Martin, B.; Fichant, G.; Polard, P.; Claverys, J.P. Bacterial transformation: Distribution, shared mechanisms and divergent control. *Nat. Rev. Microbiol.* 2014, 12, 181–196.
35. Kolling, G.L.; Matthews, K.R. Export of virulence genes and Shiga toxin by membrane vesicles of *Escherichia coli* O157:H7. *Appl. Environ. Microbiol.* 1999, 65, 1843–1848.

36. Renelli, M.; Matias, V.; Lo, R.Y.; Beveridge, T.J. DNA-containing membrane vesicles of *Pseudomonas aeruginosa* PAO1 and their genetic transformation potential. *Microbiology* 2004, 150, 2161–2169.
37. Gill, S.; Katchpole, R.; Forterre, P. Extracellular membrane vesicles in the three domains of life and beyond. *FMES Microbiol. Rev.* 2019, 43, 273–303.
38. Fulsundar, S.; Harms, K.; Flaten, G.E.; Johnsen, P.J.; Chopade, B.A.; Nielsen, K.M. Gene transfer potential of outer membrane vesicles of *Acinetobacter baylyi* and effects of stress on vesiculation. *Appl. Env. Microbiol.* 2014, 80, 3469–3483.

Retrieved from <https://encyclopedia.pub/entry/history/show/25741>