

Artemisia Genus as Biopesticides

Subjects: Biochemistry & Molecular Biology

Contributor: Bianca Ivanescu

The *Artemisia* L. genus includes over 500 species with worldwide distribution and diverse chemical composition. Many secondary metabolites of this genus are known for their antimicrobial, insecticidal, parasiticidal, and phytotoxic properties, which recommend them as possible biological control agents against plant pests. Given the negative impact of synthetic pesticides on human health and on the environment, *Artemisia*-derived biopesticides and their nanoformulations emerge as promising ecofriendly alternatives to pest management.

Keywords: antifungal ; antibacterial ; insecticidal ; nematocidal ; phytotoxic

1. Introduction

The *Artemisia* L. genus contains over 500 species, herbaceous plants and shrubs, widespread in the northern hemisphere, in Asia, Europe, and North America. *Artemisia* species are found in various ecosystems, ranging from arid regions to wetland at sea level as well as in the mountains. The largest number of species are located in the steppes of Asia ^[1]. Common names of *Artemisia* species are wormwood, mugwort, and sagebrush. Due to their biological and chemical diversity, *Artemisia* species have numerous applications in the treatment of plant and human diseases, in cosmetic and pharmaceutical industry. In addition, various *Artemisia* species are used all over the world as foods, spices, condiments, and beverages ^[2]. Many important medicinal plants belong to this genus and exert a range of therapeutic actions: antibacterial, antifungal, antiviral, antiprotozoal, anthelmintic, anti-inflammatory, anti-ulcer, appetite stimulating, hepatoprotective, antispasmodic, bronchodilator, hypolipidemic, antihypertensive, analgesic, neuroprotective, neurotrophic, anti-depressant, antioxidant, cytotoxic, antitumor, estrogenic, anti-allergic, immunomodulatory, insecticidal, repellent, and anticonvulsant ^{[3][4][5][6][7][8]}.

Most *Artemisia* species are aromatic plants that produce volatile oil in the secretory hairs on the aerial organs but also through the secretory ducts in the parenchyma tissues. Essential oils could be used as biocontrol agents based on the antibacterial, antifungal, repellent, insecticidal, nematocidal, and phytotoxic effect of volatile compounds. Moreover, the complex mixture of substances with different mechanisms of action, often having synergistic activity, can be effective in preventing the emergence of resistant strains of phytopathogens ^{[9][10][11][12]}.

The global use of synthetic pesticides has many disadvantages, such as high cost, danger to non-target organisms, accumulation of pesticide residues in the environment, the emergence of resistant phytopathogenic strains, and negative impact on human health ^[12]. In contrast, biological pesticides can achieve pest management in an environmentally friendly way and could become safer alternatives for the treatment of crop diseases. Many agents are considered biopesticides, such as viruses, microbes, fungi, entomophagous invertebrates, parasitoids, predators, and substances produced by living organisms such as bacteria, fungi, plants, algae, animals, etc. Throughout this review, we will use the word "biopesticides" for plant-derived substances or extracts. During evolution, plants developed different mechanisms to defend themselves from predators and diseases by producing substances with bactericidal, fungicidal, insecticidal, nematocidal, or repellent activity. At present, these phytochemicals are explored as biocontrol agents for crops integrated pest management. Plant compounds are cheaper, safer for farmers, less toxic to non-target organisms, and rapidly degraded in the environment ^[13].

In this context, numerous researchers have identified new potential biopesticides in plants of the *Artemisia* genus. Since most species are fragrant, the vast majority of investigations have focused on the biological actions of volatile oils and compounds. Essential oils contain a variety of volatile molecules such as mono- and sesquiterpenes as well as phenolic-derived aromatic and aliphatic components ^[14]. The percentage of individual compounds in the essential oil is variable and depends on genetic factors (species, chemotype), plant origin, plant organ, period of harvest or developmental stage, environmental factors (climate, altitude, sun exposure), and cultivation conditions. Qualitative and quantitative differences in the composition of the essential oil can also be caused by drying methods, extraction procedure and time, quantification methods, and conditions of analysis ^[11]. All these elements could change the chemical composition of an essential oil,

leading to changes in activity; thus, standardization is necessary to guarantee the effect, and also for regulatory and marketing purposes. Moreover, plants with desirable pesticide action may give low yields of essential oil, hence the need for new and more efficient extraction methods, which will increase the quantity and quality of extracted oil while reducing the time and cost of extraction [14].

2. *Artemisia* Compounds and Extracts with Pesticide Activity

2.1. Antifungal and Anti-Oomycete Activity

Pathogenic fungi produce almost 30% of crop diseases, threatening the health and food security of a growing human population dependent on substantial agricultural production [15]. Phytopathogenic fungi affect plants during their cultivation or after harvest, causing significant losses in crop plants. In addition, certain fungi (*Aspergillus* spp., *Fusarium* spp., *Alternaria* spp. etc.) produce mycotoxins that endanger the health of consumers through hepatotoxic, nephrotoxic, and carcinogenic effects or even cause death [16]. In an effort to find an ecological solution to this problem, numerous studies have assessed the antifungal effect of *Artemisia* species, focusing especially on volatile oil and compounds. Different methods of evaluation were used in vitro, in planta, or in field conditions, and the results were expressed in various ways: half maximal inhibitory concentration—IC₅₀, minimal inhibitory concentration—MIC, minimum fungicidal concentration—MFC, median effective concentration—ED₅₀, inhibition zone, and percent of inhibition (Table 1).

Table 1. Antifungal activity of *Artemisia* extracts and compounds against phytopathogenic fungi.

<i>Artemisia</i> Species	Extract * or Compound Tested	Fungi	Inhibitory Dose	Type of Study	Reference
<i>A. abrotanum</i> fresh aerial parts	essential oil (eucalyptol)	<i>Sclerotinia sclerotiorum</i>	MIC = 1200 µL/L	in vitro	[17]
<i>A. absinthium</i> aerial parts	essential oil (cis-epoxyocimene, (-)- cis-chrysanthenol, chrysanthenyl acetate, linalool and β- caryophyllene)	<i>Botrytis cinerea</i>	ED ₅₀ = 0.01–0.07 mg/mL	in vitro	[18]
		<i>Fusarium moniliforme</i>	ED ₅₀ = 0.24–0.43 mg/mL		
		<i>F. oxysporum</i>	ED ₅₀ = 0.29–0.40 mg/mL		
		<i>F. solani</i>	ED ₅₀ = 0.24–0.50 mg/mL		
<i>A. absinthium</i> leaves	aqueous extract (1:1)	<i>Alternaria alternata</i>	79.75% inhibition	in vitro	[19]
		<i>Mucor piriformis</i>	73.04% inhibition		
		<i>Penicillium expansum</i>	75.42% inhibition		
<i>A. annua</i> fresh aerial parts	essential oil (artemisia ketone)	<i>Sclerotinia sclerotiorum</i>	MIC = 2400 µL/L	in vitro	[17]
<i>A. annua</i> aerial parts	essential oil (artemisia ketone, α-selinene and γ-terpineol)	<i>Alternaria solani</i>	EC ₅₀ = 21.78 mg/mL	in vitro agar diffusion	[20]
			EC ₅₀ = 14.18 mg/mL	in vitro spore germination	

Artemisia Species	Extract * or Compound Tested	Fungi	Inhibitory Dose	Type of Study	Reference
<i>A. annua</i> leaves	methanol extract (ultrasound-assisted)	<i>Fusarium oxysporum</i>	36.94% inhibition	in vitro	[21]
	essential oil (camphor, germacrene D, β -caryophyllene, camphene)	<i>F. oxysporum</i>	MIC = 0.22 mg/mL		
		<i>F. solani</i>	MIC = 0.37 mg/mL		
	L-camphor	<i>F. oxysporum</i>	MIC = 0.11 mg/mL		
		<i>F. solani</i>	MIC = 0.31 mg/mL		
	DL-camphor	<i>F. oxysporum</i>	MIC = 0.14 mg/mL		
		<i>F. solani</i>	MIC = 0.16 mg/mL		
	β -caryophyllene	<i>F. oxysporum</i>	MIC = 0.13 mg/mL		
		<i>F. solani</i>	MIC = 0.23 mg/mL		
	camphene	<i>F. oxysporum</i>	MIC = 0.16 mg/mL		
		<i>F. solani</i>	MIC = 0.22 mg/mL		
	petroleum ether extract	<i>F. oxysporum</i> , <i>F. solani</i>	27.78% and 25% infection incidence, at 0.25 mg/g and 0.5 mg/g in the culture media, respectively	in vivo on <i>Panax notoginseng</i>	
<i>A. annua</i> whole plant	ethanol extract	<i>Aspergillus flavus</i>	14 mm inhibition zone at 200 μ g/mL	in vitro	[22]
		<i>A. niger</i>	14.5 mm inhibition zone at 200 μ g/mL		
<i>A. annua</i>	artemisinin	<i>Aspergillus fumigatus</i>	IC ₅₀ = 125 μ g/mL IC ₉₀ = 250 μ g/mL	in vitro	[23]
<i>A. arborescens</i>	essential oil (chamazulene, camphor)	<i>Rhizoctonia solani</i>	47.2% inhibition at 12.5 μ L/20 mL medium 100% inhibition at 50 μ L/20 mL medium	in vitro	[24]
<i>A. argyi</i> leaves	essential oil (caryophyllene oxide, neointermedeol, borneol, α -thujone, β -caryophyllene)	<i>Aspergillus niger</i>	MIC = 6.25 μ L/mL	in vitro	[25]
<i>A. argyi</i> inflorescence	essential oil (spathulenol, juniper camphor, caryophyllene oxide, terpineol, 1,8-cineole, borneol, camphor, chamazulene)	<i>Alternaria alternata</i>	84.7% inhibition at 1000 mg/L	in vitro	[26]
		<i>Botrytis cinerea</i>	93.3% inhibition at 1000 mg/L		
<i>A. austriaca</i> fresh aerial parts	essential oil (camphor)	<i>Sclerotinia sclerotiorum</i>	MIC = 2400 μ L/L	in vitro	[17]
<i>A. caerulescens</i> ssp. <i>densiflora</i>	essential oil (terpinen-4-ol, p-cymene, γ -terpinene, 1,8-cyeneole, α -terpineol)	<i>Alternaria</i> spp.	20 mm inhibition zone at 1:2 dilution	in vitro	[27]
		<i>Aspergillus</i> spp.	12 mm inhibition zone at 1:1 dilution		
		<i>Fusarium</i> spp.	16 mm inhibition zone at 1:8 dilution		
<i>A. campestris</i> aerial parts	methanol extracts (1:10)	<i>Aspergillus niger</i>	32.5–33.1 mm inhibition zone at 20 μ g/mL	in vitro	[28]

Artemisia Species	Extract * or Compound Tested	Fungi	Inhibitory Dose	Type of Study	Reference
<i>A. campestris</i> aerial parts	essential oil (α -pinene, β -pinene, β -myrcene, germacrene D)	<i>Aspergillus flavus</i>	MIC = 2.5 μ L/mL MFC = 2.5 μ L/mL	in vitro	[29]
		<i>Aspergillus niger</i>	MIC = 10 μ L/mL MFC >20 μ L/mL		
		<i>Aspergillus ochraceus</i>	MIC = 2.5 μ L/mL MFC = 5 μ L/mL		
		<i>Aspergillus parasiticus</i>	MIC = 2.5 μ L/mL MFC = 5 μ L/mL		
		<i>Fusarium culmorum</i>	MIC = 2.5 μ L/mL MFC = 5 μ L/mL		
		<i>Fusarium graminearum</i>	MIC = 1.25 μ L/mL MFC = 1.25 μ L/mL		
		<i>Fusarium moniliforme</i>	MIC = 2.5 μ L/mL MFC = 2.5 μ L/mL		
		<i>Penicillium citrinum</i>	MIC = 5 μ L/mL MFC > 20 μ L/mL		
		<i>Penicillium expansum</i>	MIC = 2.5 μ L/mL MFC = 2.5 μ L/mL		
		<i>Penicillium viridicatum</i>	MIC = 10 μ L/mL MFC > 20 μ L/mL		
<i>A. chamaemelifolia</i> aerial parts	essential oil (carvacrol, thymol, p-cymene α -cadinol)	<i>Aspergillus oryzae</i>	MIC = 312.5 μ g/mL MFC = 312.5 μ g/mL	in vitro	[30]
		<i>Aspergillus niger</i>	MIC = 2500 μ g/mL MFC = 2500 μ g/mL		
		<i>Byssoschlamys spectabilis</i>	MIC = 625 μ g/mL MFC = 625 μ g/mL		
		<i>Paecilomyces variotii</i>	MIC = 625 μ g/mL MFC = 625 μ g/mL		
		<i>Penicillium chrysogenum</i>	MIC = 625 μ g/mL MFC = 625 μ g/mL		
		<i>Trichoderma harizanum</i>	MIC = 312.5 μ g/mL MFC = 312.5 μ g/mL		
<i>A. dracunculus</i> fresh aerial parts	essential oil (sabinene)	<i>Sclerotinia sclerotiorum</i>	MIC = 2400 μ L/L	in vitro	[17]
<i>A. dracunculus</i> var. <i>pilosa</i> fresh aerial parts	essential oil (borneol)		MIC = 2400 μ L/L		
<i>A. herba-alba</i> aerial parts	essential oil (davanone, camphor, thujone)	<i>Fusarium moniliforme</i>	MIC = 0.5%	in vitro direct contact	[31]
		<i>Fusarium oxysporum</i>	MIC = 0.5%		
		<i>Fusarium solani</i>	MIC = 0.75%		
		<i>Stemphylium solani</i>	MIC = 0.75%		
<i>A. herba-alba</i> leaves	essential oil (β -thujone, α -thujone camphor)	<i>Penicillium aurantiogriseum</i>	100% inhibition at 0.89%	in vitro	[32]
		<i>P. viridicatum</i>	100% inhibition at 1.33%		

Artemisia Species	Extract * or Compound Tested	Fungi	Inhibitory Dose	Type of Study	Reference
<i>A. herba-alba</i> fresh leaves	essential oil	<i>Mucor rouxii</i>	100% inhibition at 1000 µg/mL	in vitro	[33]
		<i>Penicillium citrinum</i>	100% inhibition at 1000 µg/mL		
	carvone	<i>Mucor rouxii</i>	IC ₅₀ = 7 µg/mL		
		<i>Penicillium citrinum</i>	IC ₅₀ = 5 µg/mL		
	piperitone	<i>Mucor rouxii</i>	IC ₅₀ = 1.5 µg/mL		
		<i>Penicillium citrinum</i>	IC ₅₀ = 2 µg/mL		
<i>A. herba-alba</i> aerial parts	chloroform-methanol extract	<i>Fusarium solani</i>	MIC = 62.5 µg/disc	in vitro	[34]
	11-epiartapshin		MIC = 50 µg/disc		
<i>A. incisa</i> aerial parts	santolinyol-3-acetate	<i>Aspergillus flavus</i>	MIC = 300 µg/mL	in vitro	[35]
	santolinyol		MIC = 300 µg/mL		
	<i>trans</i> -ethyl cinnamate		MIC = 500 µg/mL		
	isofraxidin		MIC = 400 µg/mL		
	eupatorin		MIC = 1000 µg/mL		
	scopoletin		inactive		
	esculetin		inactive		
<i>A. judaica</i> aerial parts	essential oil (piperitone, 3-bornanone)	<i>Aspergillus niger</i>	MIC = 1.25 µg/disc	in vitro	[36]
		<i>Fusarium solani</i>	MIC = 2.5 µg/disc		
<i>A. khorasanica</i> aerial parts	essential oil (davanone, p-cymene, Z-citral, β-ascaridol, thymol)	<i>Fusarium moniliforme</i>	MIC = 2000 µL/L	in vitro	[37]
		<i>Fusarium solani</i>	MIC = 1500 µL/L		
		<i>Rhizoctonia solani</i>	MIC = 1000 µL/L		
		<i>Tiarosporella phaseolina</i>	MIC = 2000 µL/L		
<i>A. lavandulaefolia</i> aerial parts	essential oil (eucalyptol, (-)-terpinen-4-ol, α-terpineol)	<i>Alternaria solani</i>	EC ₅₀ = 10.45 mg/mL	in vitro agar diffusion	[20]
			EC ₅₀ = 6.64 mg/mL	in vitro spore germination	
<i>A. lerchiana</i> fresh aerial parts	essential oil (eucalyptol)	<i>Sclerotinia sclerotiorum</i>	MIC = 2400 µL/L	in vitro	[17]
<i>A. maritima</i> aerial parts	essential oil (1,8-cineole, chrysanthenone, germacrene D, borneol)	<i>Aspergillus flavus</i>	35.4% inhibition at 10 µL/plate	in vitro	[38]
		<i>A. niger</i>	60.6% inhibition at 10 µL/plate		
		<i>A. ochraceus</i>	56.1% inhibition at 10 µL/plate		
		<i>A. parasiticus</i>	32.45% inhibition at 10 µL/plate		
		<i>A. terreus</i>	58.3% inhibition at 10 µL/plate		
		<i>Fusarium moniliforme</i>	33.9% inhibition at 10 µL/plate		
		<i>Penicillium chrysogenum</i>	28.6% inhibition at 10 µL/plate		

Artemisia Species	Extract * or Compound Tested	Fungi	Inhibitory Dose	Type of Study	Reference
<i>A. nilagirica</i> shoot	essential oil (camphor, β -caryophyllene, α -thujone, sabinene)	<i>Aspergillus flavus</i> , <i>A. niger</i> , <i>A. ochraceus</i>	MIC = 0.29 μ L/mL MFC = 0.58 μ L/mL	in vitro	[39]
			100% mycotoxin inhibition at 0.16 μ L/mL		
		<i>Aspergillus terreus</i> , <i>Cladosporium cladosporioides</i> , <i>Fusarium moniliforme</i> , <i>Fusarium oxysporum</i> , <i>Mucor mucedo</i> , <i>Penicillium expansum</i> , <i>P. funiculosum</i> , <i>Rhizopus stolonifer</i>	100% inhibition at 0.29–0.58 μ L/mL	in vitro	
<i>A. nilagirica</i> aerial parts	essential oil (1,5-heptadiene-4-one, 3,3,6-trimethyl, artemisia alcohol, α -ionone, benzene, methyl (1-methylethyl))	<i>Aspergillus flavus</i> toxigenic strain	MIC = 1.4 μ L/mL MFC = 4.0 μ L/mL	in vitro	[40]
		<i>Alternaria alternata</i> , <i>Aspergillus flavus</i> , <i>A. minutus</i> , <i>A. niger</i> , <i>A. sydowii</i> , <i>A. terreus</i> , <i>Cheatomium spirale</i> , <i>Curvularia lunata</i> , <i>Mucor</i> spp., <i>Mycelia sterilia</i> <i>Penicillium italicum</i> , <i>P. purpurogenum</i> , <i>Rhizopus stolonifer</i> ,	70–100% inhibition at 1.4 μ L/mL	in vitro	
			71% protection from fungal contamination at 1.4 μ L/mL in air	in situ on <i>Eleusine coracana</i> seeds, 12 months storage	
<i>A. nilagirica</i> aerial parts	essential oil (α -thujone, β -thujone, germacrene D, 4-terpineol, β -caryophyllene, camphene, borneol)	<i>Macrophomina phaseolina</i>	ED ₅₀ = 93.23 mg/L	in vitro	[41]
		<i>Rhizoctonia solani</i>	ED ₅₀ = 85.75 mg/L		
		<i>Sclerotium rolfsii</i>	ED ₅₀ = 87.63 mg/L		
<i>A. nilagirica</i> leaves	essential oil (α -thujone, borneol, β -thujone, 1,8-cineole)	<i>Phytophthora capsici</i>	100% inhibition at 100 ppm	in vitro	[42]
<i>A. pallens</i> leaves	methanol extract 1:10	<i>Sclerospora graminicola</i>	Inhibition of zoosporangium formation	in vitro	[43]
<i>A. parviflora</i> twigs	methanol extract 1:1	<i>Sclerospora graminicola</i>	Inhibition of zoosporangium formation		
<i>A. pontica</i> fresh aerial parts	essential oil (eucalyptol)	<i>Sclerotinia sclerotiorum</i>	MIC = 2400 μ L/L	in vitro	[17]
<i>A. proceriformis</i> fresh leaves	essential oil (α -thujone)	<i>Aspergillus carbonarius</i>	MIC = 10.6 mg/mL	in vitro	[44]
		<i>Aspergillus niger</i>	MIC = 21.2 mg/mL		
		<i>Fusarium graminearum</i>	MIC = 10.6 mg/mL		
		<i>F. verticillioides</i>	MIC = 10.6 mg/mL		
		<i>Septoria glycines</i>	MIC = 2.7 mg/mL		
		<i>Septoria tritici</i>	MIC = 2.7 mg/mL		
<i>A. santonica</i> fresh aerial parts	essential oil (α -thujone)	<i>Sclerotinia sclerotiorum</i>	MIC = 2400 μ L/L	in vitro	[17]
<i>A. scoparia</i> aerial parts	essential oil (acenaphthene, curcumene, (+) caryophyllene oxide, spathulenol, methyl eugenol, β -caryophyllene)	<i>Alternaria solani</i>	EC ₅₀ = 12.2 mg/mL	in vitro agar diffusion	[20]
			EC ₅₀ = 3.8 mg/mL	in vitro spore germination	

Artemisia Species	Extract * or Compound Tested	Fungi	Inhibitory Dose	Type of Study	Reference
<i>A. sieberi</i> aerial parts	1R, 8S-dihydroxy-11R,13-dihydrobalchanin	<i>Fusarium solani</i>	6 mm inhibition zone at 200 µg/10 µL	in vitro	[45]
	11-epiartapshin		7 mm inhibition zone at 200 µg/10 µL		
	3'-hydroxygenkwanin		8 mm inhibition zone at 200 µg/10 µL		
<i>A. sieberi</i> aerial parts	essential oil (camphor, 1,8-cineole, camphene, chrysanthenone)	<i>Botrytis cinerea</i>	100% inhibition at 1000 µl/L	in vitro	[46]
<i>A. stricta</i> f. <i>stricta</i> aerial parts	essential oil (capillene, spathulenol, β-caryophyllene)	<i>Aspergillus flavus</i> , <i>Aspergillus niger</i> , <i>Sporothrix schenckii</i>	MIC = 0.625 mg/mL	in vitro	[47]
<i>A. terrae-albae</i> leaves	camphor, 1,8-cineole, camphene, β-thujone	<i>Aspergillus carbonarius</i>	MIC > 1.20 mg/mL	in vitro	[48]
		<i>Aspergillus niger</i>	MIC > 1.20 mg/mL		
		<i>Fusarium graminearum</i>	MIC = 0.60–1.20 mg/mL		
		<i>Fusarium verticillioides</i>	MIC = 0.60 mg/mL		
<i>A. turanica</i> aerial parts	essential oil (1,8-cineol, cis-verbenyl acetate, camphor)	<i>Aspergillus niger</i>	68.6% inhibition at 1 µL/mL	in vitro	[49]
<i>A. vulgaris</i> whole plant	crude methanol extract (1:10)	<i>Botrytis cinerea</i>	60% inhibition at 2 mg/mL	in vivo on <i>Cucumis sativus</i>	[50]
		<i>Blumeria graminis</i> f. sp. <i>hordei</i>	25% inhibition at 2 mg/mL	in vivo on <i>Hordeum sativum</i>	
		<i>Magnaporthe grisea</i>	16% inhibition at 2 mg/mL	in vivo on <i>Oryza sativa</i>	
		<i>Phytophthora infestans</i>	32% inhibition at 2 mg/mL	in vivo on <i>Lycopersicon esculentum</i>	
		<i>Puccinia recondita</i>	52% inhibition at 2 mg/mL	in vivo on <i>Triticum aestivum</i>	
		<i>Thanatephorus cucumeris</i>	9.3% inhibition at 2 mg/mL	in vivo on <i>Oryza sativa</i>	
<i>A. vulgaris</i> leaves	methanol extract 1:1	<i>Sclerospora graminicola</i>	Inhibition of zoosporangium formation	in vitro	[43]
<i>A. vulgaris</i> fresh aerial parts	essential oil (germacrene D)	<i>Sclerotinia sclerotiorum</i>	MIC = 2400 µL/L	in vitro	[17]

* To highlight the active compounds, the major constituents of the volatile oils were noted in parentheses.

The in vitro antifungal activity was frequently determined by the agar diffusion test, which involves placing the tested plant extract in wells or paper discs on the agar plate previously inoculated with the pathogen [24][25]. Since essential oils diffuse less in the culture medium, it was preferred to include them in agar after prior solubilization, followed by inoculation of the pathogen [20][31][42]. Moreover, for volatile compounds, the fumigation method was used [20]. In vivo antifungal evaluations involved treating the plants with the tested compounds/extracts by spraying them followed by inoculation with the fungal pathogen or by including the compounds in the soil and then planting the inoculated seedling in the treated soil. The disease severity was assessed after a period of infection [21][50]. In situ antifungal efficacy against postharvest pathogens was determined by fumigation in the case of stored foods [39][40].

The extraction method influences the antifungal activity of the volatile oil, as can be seen from the investigation carried out by Julio et al. [18]: *A. absinthium* oil obtained by steam pressure extraction was more effective in inhibiting mycelium growth than that obtained by hydrodistillation, which was due to a different ratio of the major volatile compounds. Similarly, *A. argyi* essential oil obtained by simultaneous distillation–extraction had a higher antifungal activity compared to oils prepared by subcritical extraction or hydrodistillation. Although regardless of the extraction method, the oils had the same five major compounds, in the oil obtained by simultaneous distillation–extraction, the sesquiterpene compounds predominated [25]. Conversely, in the case of *A. chamaemelifolia* essential oil, the method of extraction—microwave-assisted hydrodistillation and classical hydrodistillation—had no influence on the inhibitory effect against the tested fungi. Both oils contained the same major compounds in comparable ratio [30].

The type of extract, the part of the plant used, and the time of harvest also influence the antifungal activity, as underlined in a study carried out with methanol, ethanol, and hexane extracts of *Artemisia annua* against *Aspergillus niger* and *A. flavus*. Whole plant extract was the most efficient in inhibiting the growth of the two fungi, regardless of the type of extract, compared to root, leaf, or stem extracts. Regarding the extraction solvent, ethanol extract had the highest inhibitory effect, followed by methanol and hexane, on both fungal species. Although the harvesting period of the plant had little influence on the antifungal activity, most of the extracts made with the plant collected during anthesis were more active [22].

From analyzing literature data, it appears that sesquiterpenes components of the oil have significant antifungal activity. Oxygenated sesquiterpenes were the major components of *A. khorasanica* volatile oil active against four soil-borne phytopathogenic fungi [37]. *Artemisia scoparia* essential oil, rich in sesquiterpenes, was more efficient in inhibiting mycelial growth and spore germination of *Alternaria solani* compared to *A. lavandulaefolia* and, especially, *A. annua* oils, where monoterpenes were the major compounds. Furthermore, the mode of volatile oil administration influences the outcome: *A. lavandulaefolia* oil was more effective when applied by fumigation than when mixed in the agar medium [20].

Alongside the sesquiterpenes, it seems that thujones present in high amounts in the volatile oil are associated with intense antifungal activity [32][42]. To prove this point, Shafi et al. [42] used a mixture of thujones (α -thujone, β -thujone, and fenchone) at the same concentration instead of *A. nilagirica* oil to achieve the same result against *Phytophthora capsici*—100% inhibition. Borneol was also tested in the aforementioned study and showed no antifungal activity. On the other hand, the antifungal property of *A. terrae-albae* essential oil against *Fusarium* spp. was associated with the presence of camphor, 1,8-cineole, camphene, α - and β -thujone, borneol, and the high content of oxygenated monoterpenes [48]. Other oxygenated monoterpenes, piperitone and carvone, were correlated with the antifungal activity on *Penicillium citrinum* and *Mucor rouxii*; the two ketones are major components of *A. herba-alba* volatile oil [33].

Some volatile compounds (L-camphor; DL-camphor, β -caryophyllene, and camphene) from *A. annua* oil were as efficient as synthetic antifungal products such as flutriafol and hymexazol against *Fusarium oxysporum* and *F. solani*, in vitro [21]. Different compounds isolated from the methanol extract of *A. incisa* were tested against *Aspergillus flavus* with various results: two monoterpenes and one phenolic acid derivative were more active compared to flavones and coumarins, the latter being less active [35].

Moreover, the synergistic action of essential oils and chemical fungicides was evaluated. Thus, *A. annua* essential oil combined with flutriafol exhibits additive inhibitory effect against *Fusarium solani*, while with hymexazol, it manifests synergistic activity on *F. solani* and additive action on *F. oxysporum* [21].

Most *Artemisia* extracts were tested on *Fusarium*, *Alternaria*, *Aspergillus*, and *Penicillium* species. Fungi have different susceptibility to varied antifungal compounds: for example, *Fusarium solani* was moderately sensitive to the action of isolated substances from *A. sieberi* (two sesquiterpene lactones and one methoxylated flavone), while *Alternaria alternata* and *Aspergillus niger* were resistant [45]. In an analogous manner, *Aspergillus niger* was sensitive to the methanol extract of *A. campestris* and resistant to *A. vulgaris* extract, despite similar quantities of flavonoids and phenolic compounds. Quercetin was reported in higher amounts in *A. campestris* extract and seems to be correlated with antifungal activity [28].

Few studies assessed the antifungal activity in vivo. Ma et al. [21] showed that the petroleum ether extract of *A. annua*, imitating the composition of the essential oil, decreased the incidence of infected *Panax notoginseng* plants when added in the culture mixture. *A. vulgaris* crude methanol extract exhibited weak to moderate antifungal activity against *Magnaporthe grisea*, *Thanatephorus cucumeris*, *Botrytis cinerea*, *Phytophthora infestans*, *Puccinia recondite*, and *Blumeria graminis* when tested on plants grown in greenhouse conditions [50].

Stored foods can be degraded by fungi such as *Alternaria* spp., *Penicillium* spp., and *Mucor* spp., which reduce their quality and make them unsuitable or even toxic for consumption. The use of chemical products for the control of postharvest pathogens endangers the environment, human health, and can induce resistance to fungicides. Such being

the case, some investigations tried to estimate the reduction of postharvest fungal spoilage after treatment with *Artemisia* extracts. Fumigation of table grapes with *A. nilagirica* essential oil (200–300 μ L) decreased the weight loss, berry shrinkage, and berry browning, increasing the shelf life for up to 10 days [39]. In addition, *A. nilagirica* volatile oil at a concentration of 1.4 μ L/mL in airtight containers provided 71% protection from fungal contamination after 12 months of storage to millet grains [40].

In addition to the direct inhibition of postharvest phytopathogenic fungi, some studies also evaluated the mycotoxins suppression ability of plant extracts. For instance, *Artemisia herba-alba* keto-rich essential oil completely inhibited the toxin production (penicillic acid, terrestric acid, brevianamide A, aurantiamine, xanthomegnin) for *P. aurantiogriseum* at 0.44% and for *P. viridicatum* at 0.22% [32]. Similarly, *Artemisia nilagirica* essential oil inhibited the production of aflatoxin B₁ by *Aspergillus flavus* toxigenic strain at 1 μ L/mL. A common seed contaminant, aflatoxin B₁ is a powerful human carcinogen and a serious health risk; it also contributes to food deterioration by lipid peroxidation [40]. In another experiment, *A. nilagirica* volatile oil (0.16 μ L/mL) completely inhibited the production of aflatoxin B₁ by *Aspergillus flavus* and ochratoxin A by *A. niger* and *A. ochraceus* [39].

The phytochemicals mechanism of action against fungi involves the inhibition of enzymes that control energy or structural compounds production, degeneration of fungal cell wall with loss of cytoplasm, and plasma membrane dysfunction. Due to their lipophilic nature, components of essential oils can penetrate cell walls, increase cellular membranes permeability and disturb the fungal cells metabolism, causing their death [11]. Monoterpenes delay sclerotic differentiation and promote the generation of lipid peroxides, which can lead to cell death, while phenols present in the essential oil bond to the active sites of fungal enzymes through their hydroxyl group [51].

In addition, spore germination and germ tube growth are negatively influenced by terpenes from the essential oil. *A. annua* volatile oil arrested mycelia growth and conidia germination of *Fusarium oxysporum* and *Fusarium solani* [21]. Electron microscope observations proved that *A. argyi* essential oil affected the cell morphology and the structure of cell walls in *Aspergillus niger* [25]. An earlier study showed that *Artemisia herba-alba* essential oil inhibited mycelium growth, spore germination, and sporulation of *Zygorrhynchus* spp., *Aspergillus niger* and *Penicillium italicum* [52].

The antifungal mode of action of *A. nilagirica* essential oil was investigated by Kumar et al. The fungal cells treated with 1.4 μ L/mL volatile oil exhibited important deformity and shrinkage, detachment of plasma membrane from the cell wall, and development of lomasomes. At the same dose, *A. nilagirica* essential oil completely inhibited ergosterol synthesis in the cell membrane of *Aspergillus flavus* and provoked the leakage of Ca²⁺, K⁺, and Mg²⁺ ions from the cell [40].

It is worth mentioning that in addition to the secondary antifungal metabolites produced by plants, certain endophytic organisms present in *Artemisia* species are able to inhibit the development of phytopathogenic fungi. Thus, in the root, stem, and leaves of *A. argyi*, researchers identified endophytes (*Bacillus subtilis*, *B. cereus*, *Paenibacillus polymyxa*) that produce substances capable of inhibiting the growth of the mycelium of *Fusarium oxysporum*, *Magnaporthe grisea*, and *Alternaria alternata* [53].

2.2. Antibacterial Activity

Only a small number of studies investigated the effect of *Artemisia* spp. extracts on phytopathogenic bacteria. For instance, different *A. nilagirica* leaves extracts were tested in vitro against four phytopathogenic bacteria, *Erwinia* spp., *Clavibacter michiganense*, *Pseudomonas syringae*, and *Xanthomonas campestris*, which cause diseases in potato, tomato, leafy greens, carrot, onion, and green pepper. The hexane extract was the most efficient in inhibiting all tested bacteria with MIC of 32 μ g/mL. The ethanol, methanol, diethyl ether, and chloroform extracts were moderately active against the four bacteria, while the petroleum ether extract was the least effective [54]. Methanol, ethanol, and chloroform extracts from leaves of *Artemisia parviflora* (1:6 w/v) were almost ineffective against *Xanthomonas vesicatoria* and *Ralstonia solanacearum*, with inhibition zones of 1 and 2 mm [55].

The essential oil of *Artemisia turanica* exhibited inhibitory activity at 2% (v/v) concentration against tumor galls induced by *Agrobacterium tumefaciens* on potato discs, but it did not demonstrate antibacterial activity in vitro against *A. tumefaciens* at the same dose [49]. In addition, the methanol extracts of roots, leaves, and flowers of *Artemisia fragrans* inhibited tumor growth in different percentages at 10, 100, and 1000 ppm. Leaves and flowers extract had the highest inhibition at all concentration (20, 38, 46%) compared to root extract (15, 24, 34%). No extract had any significant effect on the viability of *A. tumefaciens* when tested by agar diffusion assay [56].

Dadasoglu et al. [57] evaluated the antibacterial activities of essential oils, hexane, chloroform, acetone, and methanol extracts from the aerial parts of *A. santonicum*, *A. spicigera*, and *A. absinthium* against 25 plant pathogenic bacterial

strains. *A. spicigera* essential oil was only active (MIC = 500 µL/mL) against *Erwinia amylovora*, *Pseudomonas syringae* pv. *syringae*, and *Xanthomonas axonopodis* pv. *vesicatoria*. The volatile oil of *A. absinthium* exhibited moderate activity (MIC = 250–500 µL/mL) against most of the phytopathogenic bacteria. *A. santonicum* essential oil was the most effective with MIC values 125–250 µL/mL on 22 out of 25 bacteria tested, with the exception of *Pseudomonas aeruginosa*, *P. cichorii*, and *Clavibacter michiganensis* subsp. *michiganensis*. None of the *Artemisia* solvent extracts manifested antibacterial activity on the tested strains. The main constituents of *A. absinthium* oil were chamazulene, nuciferol butanoate, nuciferol propionate, and caryophyllene oxide, while *A. santonicum* and *A. spicigera* oil shared similar major components: camphor, 1,8-cineole, cubenol, borneol, terpinen-4-ol, and α-terpineol.

In the previously mentioned study, some constituents isolated from the essential oils were evaluated individually for their antibacterial activity. Caryophyllene oxide, camphor, borneol, and 1,8-cineole did not show activity against the phytopathogenic bacteria. Terpinen-4-ol inhibited the growth of all tested bacteria with MIC values ranging from 60 to 110 µL/mL and linalool blocked the development of 22 bacterial strains with MIC values in the 50–110 µL/mL domain. α-Terpineol was active (MIC = 60–70 µL/mL) only on *Pseudomonas cichorii*, *P. huttiensis*, *P. syringae* pv. *syringae*, and *Xanthomonas axonopodis* pv. *vesicatoria* [57].

The essential oil extracted from fresh leaves of *Artemisia proceriformis* manifested weak antimicrobial activity against four bacteria: *Erwinia carotovora* (MIC = 21.2 mg/mL), *Pseudomonas corrugate* (MIC = 21.2 mg/mL), *Pseudomonas syringae* (MIC = 5.31 mg/mL), and *Xanthomonas vesicatoria* (MIC > 42.5 mg/mL). The major component was α-thujone, in proportion of 66.9% [44].

Terpenes and phenolic compounds found in the essential oils are responsible for the intense antimicrobial activity. Terpenes have the ability to increase membrane permeability by infiltrating the phospholipidic bilayer; the damage to the bacterial membrane causes the loss of cytoplasmic components, which leads to cell death. Plant extracts are studied not only as inhibitors of bacterial growth, but also for the prevention of biofilm formation. Such is the case of *A. herba-alba*, *A. absinthium*, and *A. campestris* essential oils that can reduce biofilm formation by up to 70% [58].

2.3. Insecticidal Activity

Insects are the more diverse group of animals on Earth, and only 0.5% are considered pests. Nonetheless, herbivorous insects destroy every year one-fifth of the world's crop production. Synthetic chemicals used to control insect pests are toxic to humans, animals, and the environment through accumulation. In addition, the development of insecticide resistance and the migration of harmful insects require the search for an alternative for plant protection. Considering these facts, botanical insecticides represent a viable substitute with low toxicity toward humans and the environment [59].

Plant-derived substances or plant extracts usually have a lower acute toxicity toward insects compared to synthetic insecticides. Nevertheless, their subacute toxicity was frequently noted and is important because it can limit insect spreading (diminished fertility, fecundity, vitality, or shorter lifespan) and decrease crop loss due to repellent, suppressant, or deterrent activity. These effects are generally called “antifeedant” and are manifested in insects by lower weight and body size, decreased fertility, and altered behavior [60].

Artemisia compounds can influence insects by direct contact or fumigation, can repel insects or keep them from feeding, or can hinder their reproduction. Volatile compounds can induce toxicity to insects via inhalation or direct contact by forming an impermeable film on the cuticle leading to suffocation. Some volatile components can penetrate through the cuticle, affecting cellular membrane function and oxidative phosphorylation [61]. Phytochemicals such as cinnamyl alcohol, eugenol, and trans-anethole can activate octopamine receptors, interfering with the normal activity of octopamine, a neurotransmitter, neuromodulator, and neurohormone in an invertebrate system [62]. Furthermore, volatile compounds can interfere with the γ-aminobutyric acid (GABA) receptor in insects [14]. Other studies reported the inhibition of acetylcholinesterase by 1,8-cineole, (-)-citronellal, limonene, α-pinene, pulegone, and 4-terpineol [63] or inhibition of adenosinetriphosphatase by essential oils [64]. In addition, plant substances may cause the suppression of cytochrome P450 in insects (the enzymes responsible for phase I metabolism of xenobiotics) and may alter various biochemical processes, which shift the balance of the endocrine system [14].

The activity of *Artemisia* compounds and extracts depends on the solvent used, the susceptibility of pest species to the active substance, the development stage of the insect, whether it is male or female, and the method of application. Table 2 lists the more recent studies on insecticidal activity of *Artemisia* genus. Essential oils and volatile compounds can be applied via fumigation, which is a procedure used frequently in the pest management of stored products. This method has obvious advantages such as the possibility to spread the substance evenly, even in unreachable places, and the ability to maintain an effective level of insecticides within a closed space [60]. Some of the shortcomings of natural insecticides are

poor water solubility and rapid degradation in the environment, leading to low persistence and poor efficiency. To solve these problems, plant insecticides may be formulated as micro- and nanocapsules, nanoparticles, or nanoemulsions. These nanoformulations can increase the solubility, persistence, and stability of bioinsecticides, enhancing their activity and, at the same time, limiting their negative impact on the environment [65].

Table 2. Insecticidal activity of *Artemisia* compounds and extracts.

<i>Artemisia</i> spp.	Extract or Compound Tested	Target Species	Reference
<i>A. absinthium</i>	essential oil	<i>Leptinotarsa decemlineata</i> <i>Myzus persicae</i> <i>Rhopalosiphum padi</i> <i>Spodoptera littoralis</i>	[18]
	essential oil	<i>Trialeurodes vaporariorum</i> <i>Tuta absoluta</i>	[66]
	essential oil	<i>Tetranychus cinnabarinus</i>	[67]
	essential oil	<i>Diaphania hyalinata</i>	[68]
	methanol extract	<i>Sitophilus oryzae</i>	[69]
	essential oil	<i>Orysaephilus surinamensis</i> <i>Tribolium castaneum</i>	[70]
	powdered plant	<i>Oryzaephilus surinamensis</i>	[71]
	water extract ethanol extract	<i>Hyphantria cunea</i>	[72]
	supercritical extracts	<i>Spodoptera littoralis</i>	[73]
	essential oil	<i>Myzus persicae</i>	[74]
	essential oil carvacrol (-)- α -bisabolol chamazulene	<i>Diaphorina citri</i>	[75]
<i>A. annua</i>	methanol extract essential oil	<i>Helicoverpa armigera</i>	[76]
	methanol extract artemisinic acid artemisinin scopoletin arteannuin-B deoxy-artemisinin artemetin casticin chrysosplenetin	<i>Helicoverpa armigera</i>	[77]
	essential oil	<i>Glyphodes pyloalis</i>	[78]
	methanol extract	<i>Pieris rapae</i>	[79]
	methanol extract	<i>Hyphantria cunea</i>	[80]
	methanol extract	<i>Glyphodes pyloalis</i>	[81]
	essential oil	<i>Diaphania hyalinata</i>	[68]
<i>A. arborescens</i>	essential oil	<i>Rhyssopertha dominica</i>	[24]
<i>A. argyi</i>	ethanol extract	<i>Brevicoryne brassicae</i>	[82]
	essential oil	<i>Diaphania hyalinata</i>	[68]
	water extract ethanol extract	<i>Hyphantria cunea</i>	[72]
	essential oil	<i>Plodia interpunctella</i>	[83]

Artemisia spp.	Extract or Compound Tested	Target Species	Reference
<i>A. frigida</i>	essential oil	<i>Liposcelis bostrychophila</i> <i>Sitophilus zeamais</i>	[84]
	essential oil terpinen-4-ol verbenone camphene α -terpineol α -terpinyl acetate	<i>Lasioderma serricorne</i> <i>Liposcelis bostrychophila</i> <i>Tribolium castaneum</i>	[85]
<i>A. herba-alba</i>	essential oil	<i>Orysaephilus surinamensis</i> <i>Tribolium castaneum</i>	[70]
<i>A. judaica</i>	essential oil	<i>Sitophilus orizae</i>	[64]
<i>A. lavandulaefolia</i>	essential oil 1,8-cineole chamazulene β -caryophyllene	<i>Lasioderma serricorne</i>	[86]
<i>A. monosperma</i>	essential oil	<i>Sitophilus orizae</i>	[64]
	essential oil	<i>Aphis nerii</i>	[87]
<i>A. nilagirica</i>	cow urine extract	<i>Scirpophaga incertulas</i>	[88]
<i>A. spicigera</i>	essential oil	<i>Dendroctonus micans</i>	[89]
<i>A. vulgaris</i>	essential oil	<i>Callosobruchus maculatus</i> <i>Rhyzopertha dominica</i> <i>Tribolium castaneum</i>	[90]
	essential oil	<i>Diaphania hyalinata</i>	[68]
	water extract ethanol extract	<i>Hyphantria cunea</i>	[72]

2.4. Nematicidal Activity

Plant parasitic nematodes cause severe yield losses in different crops, especially in tropics and subtropics. Frequent nematodes that affect plants include *Meloidogyne* (root-knot nematodes), *Pratylenchus* (lesion nematodes), *Xiphinema* (dagger nematodes), *Aphelenchoides* (foliar nematodes), *Globodera* (potato cyst nematodes), and *Heterodera* (soybean cyst nematodes). *Meloidogyne* species induce histological damages to roots, with the appearance of visible galls. Some phytoparasitic nematodes act as vectors for plant viruses, such as *Xiphinema* species [91].

Various *Artemisia* species were evaluated for nematicidal activity, some with promising results. For instance, *A. judaica* essential oil (1 μ L/L) caused 85% mortality on *Meloidogyne javanica* second-stage juveniles and inhibited the hatching of eggs. The main component of the essential oil was artemisia ketone. In the same study, *A. arborescens* and *A. dracuncululus* essential oils were poorly active on the root-knot nematode [92]. In vitro toxicity of *Artemisia annua* essential oil was evaluated against second-stage juveniles of *Meloidogyne incognita* and pre-adults of *Rotylenchulus reniformis* (reniform nematode). Concentrations of 500 and 250 ppm induced 100% mortality in both nematode species [93]. Moreover, there are reports of nematicidal activity exhibited by the alcoholic and aqueous extracts of *Artemisia annua* against *Meloidogyne incognita* and *Pratylenchus loosi* (tea root lesion nematode) [91].

Artemisia herba-alba essential oil produced 94.4% mortality on *Meloidogyne incognita* second-stage juveniles at 15 μ g/mL and 100% mortality on *Xiphinema index* females at 2 μ g/mL, after 24 h exposure. However, mixed-age infective specimens of *Pratylenchus vulnus* were more resistant to the activity of *A. herba-alba* essential oil with mortality values ranging from 56.8% to 67% after 24 to 96 h of exposure. The major components of the essential oil were cis- and trans-thujone, camphor, 1,8-cineole, trans-chrysantenyl acetate, and camphene. In an additional test, the three nematode species were exposed to various compounds of the essential oils of four plants, including *A. herba-alba*. Borneol and α -pinene manifested poor to moderate activity, while limonene lack activity on the three nematode species. Camphor exhibited a moderate nematicidal effect, whilst thymol and thujone (mixture of cis-thujone, 70% and trans-thuione) displayed strong activity against *M. incognita*, and less so on *P. vulnus* and *X. index*. The fact that the activity of the components of the volatile oil is weaker than that of the whole oil suggests a possible synergistic action of the mixture. In addition, soil treatments with 100 or 200 μ g/kg *A. herba-alba* essential oil, by fumigation or application of water solution,

significantly inhibited nematode density on tomato roots and in soil and also increased the plant biomass. Fumigation was proven to be more effective than drenching treatment [94].

A. absinthium essential oil (β -thujone 51% and linalyl acetate 24%) had over 99% mortality rate at 0.25 and 0.5% concentrations (v/v) against *Meloidogyne javanica* juveniles in an in vitro test. Furthermore, in vivo experiments were conducted in order to assess the ability of the essential oil to inhibit root-knot nematode development after being absorbed by the tomato plants. It was observed that spraying the oil on tomato leaves actually increased the number of galls and eggs in treated plants, and applying the essential oil into the soil at 0.25% and 0.5% concentrations did not lower the number of galls or nematode eggs in tomato plants. The authors believe that the nematicidal compounds could have been volatilized or degraded by microorganisms in the soil or by the plant, or possibly, the root exudates were modified by the absorbed essential oil, making the tomato plants more appealing to the nematodes [95]. In another study, commercially available *A. absinthium* volatile oil had only a slight effect on *Meloidogyne javanica* in vitro (the median lethal dose LC₅₀ of 937 μ g/mL at 48 h and 734 μ g/mL at 72 h). The major components of the oil were borneol acetate, β -terpineol, 1,8-cineol, linalool, sabinene, and o-cymene [96].

The nematicidal activity of *Artemisia absinthium* hydrolate, a by-product of essential oil extraction, was evaluated on the root-knot nematode, *Meloidogyne javanica*. The hydrolate caused high mortality of second-stage juvenile and suppression of egg hatching, proving the ability of the *A. absinthium* hydrolate to penetrate the gelatinous matrix of eggs. In vivo tests showed a strong inhibition of juveniles' penetration in the tomato roots. Soil treatment with *A. absinthium* hydrolate (60% and 20% concentrations) significantly reduced the reproductive capacity of root-knot nematode and the infection frequency. The main component of the hydrolate, responsible for the nematicidal activity, was identified as (5Z)-2,6-dimethylocta-5,7-dien-2,3-diol [97].

Kalaiselvi et al. [98] showed that essential oils of *A. nilagirica* plants collected from high and low altitude have different composition and different nematicidal activity against *Meloidogyne incognita* (LC_{50/48h} of 5.75 and 10.23 μ g/mL, respectively). α -thujone, α -myrcene, and linalyl isovalerate were the main components of high-altitude *A. nilagirica* volatile oil, while the low-altitude plants produced an oil composed mostly of camphor, caryophyllene oxide, eucalyptol, humulene epoxide II, α -humulene, and β -caryophyllene. Experiments carried out in vivo by soil irrigation with the essential oil revealed that both volatile oils significantly reduced the infection of tomato plant (number of nematode juveniles and eggs) and enhanced plant growth (fresh weight of aerial parts and roots) at 20 μ g/mL. Again, the effect was greater for the oil originated from high-altitude *A. nilagirica*. Moreover, the ethanol extract of flowering parts of *A. nilagirica* (1 mg/mL) exhibited nematicidal activity against *Meloidogyne incognita*, as reported by an earlier study [99].

Various hypotheses have been advanced as explanations for the nematicidal effects of essential oils: disruption of cell membrane permeability and obstruction of its functions, irreversible modifications of proteins structures from the nematode surface induced by aldehydes, inhibition of acetylcholinesterase with build-up of neurotransmitter in the central nervous system of the nematode followed by convulsion, paralysis, and death [11]. Research on *A. nilagirica* essential oil ascribe the nematicidal action to an increased generation of intracellular reactive oxygen species, activation of signaling pathway of apoptosis, and DNA damage prompting cell death [98].

In addition to the essential oils and their volatile compounds, few other substances from *Artemisia* genus have been tested for their activity against plant nematodes. Thirteen chemical compounds (apigenin, bonanzin, nepetin, dihydroluteolin, scopoletin, isoscapoletin, benzoic acid, β -sitosterol, γ -sitosterol, betulinic acid, friedelin, linoleic acid, and a long chain ketone) isolated from *Artemisia elegantissima* and *Artemisia incisa* were tested in vitro and in vivo for nematicidal activity against *M. incognita*. All phytochemicals significantly inhibited egg hatching and induced high mortality of second-stage juveniles at the tested concentrations (0.1, 0.2, and 0.3 mg/mL). Isoscapoletin was even more effective than the positive control carbofuran. In addition, application of the compounds as a root drench (0.1 mg/mL) on potted tomato plants caused a marked reduction of galls, galling index, and egg masses on plant roots, numbers of juveniles in the rhizosphere soil, and also improved tomato plant growth parameters (shoot and root length and weight). Isoscapoletin and apigenin were the most active compounds [100].

2.5. Herbicidal Activity

One of the most influential groups of plant secondary metabolites is the allelochemicals. They are released into the environment in order to affect the germination, growth, behavior, survival, and reproduction of competing plants, which is a process better known as allelopathy. They are produced mainly in the plant's roots, seeds, flowers, and leaves, and their synthesis depends on the changes of the climate conditions as well as exposure to biotic or abiotic stress. Allelochemicals activity can be harmful or beneficial for the growth and survival of target species [101]. The destructive effect of allelochemicals is crucial for defending plants against herbivores and providing an advantage in the competition for

resources [102]. In agroecosystems, allelopathy can influence weed management, and plant allelochemicals could be employed as bioherbicides in order to reduce the negative impact of chemical herbicides on the environment [103].

The allelopathic properties of *Artemisia* species are well known [104][105][106][107][108][109][110], so it was expected that numerous studies would investigate their herbicide potential on various weeds. Most researchers focused on the volatile oils, and only a few dealt with aqueous or alcoholic extracts (Table 3). The phytotoxic effect of essential oils is owed to multiple mechanisms of action: inhibition of cell division, decrease of mitochondrial respiration, reduction of photosynthetic pigments and photosynthesis, generation of radical oxygen species in excess and oxidative impairment, destruction of waxy cuticular layer, inhibition of enzymes activity, water uptake, and alteration of gibberellic acid content [102][111][112]. Most of these actions are correlated with the presence of oxygenated monoterpenes. For example, 1,8-cineole and camphor inhibit DNA synthesis, cell proliferation, and elongation [113].

Table 3. Phytotoxic activity of *Artemisia* compounds and extracts.

<i>Artemisia</i> Species	Extract * or Compound Tested	Weed/Target Plant	Observed Effect	Reference
<i>A. absinthium</i> aerial parts	essential oil (cis-epoxyocimene, (-)-cis-chrysanthenol, chrysanthenyl acetate, linalool and β -caryophyllene)	<i>Lolium perene</i>	Suppression of root and leaf growth No effect on seed germination	[18]
		<i>Lactuca sativa</i>	Suppression of root and leaf growth No effect on seed germination	
<i>A. absinthium</i> fresh aerial parts	essential oil (β -thujone, chamazulene)	<i>Sinapis arvensis</i>	Complete inhibition of seed germination and seedling growth at 2 μ L/mL	[114]
<i>A. absinthium</i> leaves	aqueous extract 1:10 w/v	<i>Parthenium hysterophorus</i>	Inhibition of seed germination, shoot and root growth, reduction of chlorophyll and carotenoid content, at 25, 50, 75, and 100% Enhanced malondialdehyde levels, phenolic content and increased activity of antioxidative enzymes, at 25, 50, 75, and 100%	[105]
<i>A. absinthium</i> shoot and root	aqueous extract	<i>Chenopodium album</i>	Decreases growth criteria (root and shoot length and fresh weight, number of leaves) at 1–100 mg/mL No effect on seed germination Increased peroxidase and superoxide dismutase activity in root	[115]
<i>A. afra</i> leaves	aqueous extract	<i>Triticum aestivum</i>	No effect on seed germination	[116]
		<i>Brassica napus</i>	Complete inhibition of seed germination	
		<i>Medicago sativa</i>	Increased germination rate	
		resistant and non-resistant <i>Lolium</i> spp.	Significant inhibition of seed germination	

Artemisia Species	Extract * or Compound Tested	Weed/Target Plant	Observed Effect	Reference
<i>A. annua</i> flower heads	essential oil (1,8-cineole, trans-sabinyl acetate, artemisia ketone, camphor α -pinene)	<i>Amaranthus retroflexus</i>	In vitro, complete inhibition of seed germination, at 10 and 100 μ g/L In vivo, plant death, at the cotyledon stage (100 mg/L) and true leaf stage (1000 mg/L)	[117]
		<i>Setaria viridis</i>	In vitro, complete inhibition of seed germination, at 100 μ g/L In vivo, plant death, at the cotyledon stage (100 mg/L) and true leaf stage (1000 mg/L)	
<i>A. annua</i> aerial parts	artemisinin arteannuin B artemisinic acid	<i>Secale cereale</i> , <i>Hordeum vulgare</i> , <i>Artemisia annua</i> , <i>Portulaca oleracea</i> , <i>Amaranthus blitun</i> , <i>Lactuca sativa</i> , <i>Raphanus sativus</i>	Inhibition of seed germination Inhibition of shoot and root growth	[118]
<i>A. annua</i>	artemisinin	<i>Lactuca sativa</i>	Inhibition of root and shoot elongation, reduced cell division and cell viability in root tips, at 10 μ M Reduced chlorophyll a and b levels Increased malondialdehyde and proline levels, at 1 μ M	[119]
<i>A. annua</i>	artemisinin	<i>Arabidopsis thaliana</i>	Reduction of fresh biomass, chlorophyll a, b, and leaf mineral contents at 40–160 μ M Reduction of photosynthetic efficiency, yield, and electron transport rate, calcium and nitrogen levels at 80 and 160 μ M Elevated lipid peroxidation (malondialdehyde contents) at 80 and 160 μ M	[120]
<i>A. arborescens</i> shoot	sesamin ashantin	<i>Agrostis stolonifera</i> , <i>Lactuca sativa</i>	Growth inhibition at 1 mg/mL	[107]
	sesamin	<i>Lemna paucicostata</i>	Growth inhibition IC ₅₀ = 401 μ M	
	ashantin	<i>Lemna paucicostata</i>	Growth inhibition IC ₅₀ = 224 μ M	
<i>A. arborescens</i> leaf litter	crude methanol extract	<i>Lactuca sativa</i> , <i>Raphanus sativus</i> , <i>Amaranthus retroflexus</i> , <i>Cynodon dactylon</i>	Inhibition of seed germination ED ₅₀ = 1.61–3.05 mg/mL Inhibition of root growth ED ₅₀ = 1.22–3.14 mg/mL	[121]
	hexane, chloroform, and ethyl acetate fractions		Inhibition of seed germination ED ₅₀ = 1.19–6.25 mg/mL Inhibition of root growth ED ₅₀ = 0.92–3.98 mg/mL	
<i>A. arborescens</i> aerial part	crude methanol and aqueous extracts	<i>Lactuca sativa</i>	Inhibition of seed germination and root growth ED ₅₀ = 0.5–2.8 mg/mL	[122]
	ethyl acetate, n-hexane, chloroform, n-butanol fractions		Inhibition of seed germination and root growth ED ₅₀ = 0.4–5.4 mg/mL	

Artemisia Species	Extract * or Compound Tested	Weed/Target Plant	Observed Effect	Reference
<i>A. argyi</i> leaves	water extract (caffeic acid, schaftoside, 4-caffeoylquinic acid, 5-caffeoylquinic acid, 3,5-dicaffeoylquinic acid and 3-caffeoylquinic acid)	<i>Brassica pekinensis</i> , <i>Lactuca sativa</i> , <i>Oryza sativa</i>	Inhibition of germination, root and stem growth, and biomass (at 50, 100, and 150 ng/mL)	[108]
		<i>Brassica pekinensis</i> , <i>Lactuca sativa</i> , <i>Oryza sativa</i> , <i>Portulaca oleracea</i> , <i>Oxalis corniculata</i> , <i>Setaria viridis</i>	Inhibition of germination and growth in pot experiment (<i>A. argyi</i> powder mixed into sand soil at the ratio 100:0, 100:2, 100:4, and 100:8)	
<i>A. campestris</i> leaves	essential oil (β -pinene, 1, 8-cineole, p-cymene, myrcene)	<i>Daucus carota</i> , <i>Cicer arietinum</i> , <i>Phaseolus vulgaris</i> , <i>Triticum sativum</i>	Reduces seed germination at 1000–2000 ppm Enhances seed germination at 100 ppm Delays the germination of <i>D. carota</i> seeds	[123]
<i>A. dracunculus</i> aerial parts	essential oil	<i>Medicago minima</i> , <i>Rumex crispus</i> , <i>Taraxacum officinale</i>	No effect on seed germination at 0.3–1.2 mg/L	[124]
<i>A. dracunculus</i>	leachate	<i>Lactuca sativa</i>	Radicle growth inhibition	[125]
<i>A. fragrans</i> aerial parts	essential oil (α -thujone, camphor, 1,8-cineole, β -thujone)	<i>Convolvulus arvensis</i>	Important reduction in the shoot, root, and plant length, shoot and root fresh weight, shoot and root dry weight Inhibited seed germination Significant decrease of photosynthetic pigments and antioxidant enzymes Increased production of H ₂ O ₂ and malondialdehyde content, and membrane leakage	[126]
<i>A. fragrans</i> roots, leaves, and flowers	methanol extracts	<i>Raphanus raphanistrum</i>	Inhibition of root growth at 1000 ppm Inhibition of seed germination at 7500 ppm	[56]
<i>A. frigida</i>	volatile organic compounds (1,8-cineole, camphene, (E)-3-hexen-1-ol acetate, α -terpineol, β -terpineol)	<i>Melilotus suaveolens</i> , <i>Sorghum sudanense</i> , <i>Elymus dahuricus</i> , <i>Agropyron cristatum</i>	Significantly decreases the seed germination and seedling growth	[127]
<i>A. judaica</i> aerial parts	essential oil (piperitone, 3-bornanone)	<i>Lactuca sativa</i>	Reduced seed germination, shoot and root growth at 250–1000 μ L/L	[36]
<i>A. lavandulaefolia</i> leaves	aqueous extract	<i>Lactuca sativa</i> , <i>Artemisia princeps</i> , <i>Achyranthes japonica</i> , <i>Oenothera odorata</i> , <i>Plantago asiatica</i> , <i>Aster yomena</i> , <i>Elsholzia ciliata</i> , <i>Raphanus sativus</i>	Inhibition of root growth Inhibition of seed germination	[128]
	essential oil (1,8-cineole, α -terpineol, α -terpinene, camphor, azulene, 2-buten-1-ol)			
<i>A. monosperma</i> aerial parts	aqueous extract	<i>Phaseolus vulgaris</i>	Stimulation of seed germination at 1% and 2% concentration Inhibition of seed germination at 3% and 4% concentration Inhibition of amylase and protease activity	[129]

Artemisia Species	Extract * or Compound Tested	Weed/Target Plant	Observed Effect	Reference
<i>A. monosperma</i> aerial parts	aqueous extract	<i>Medicago polymorpha</i>	Reduction of germination percentage, plumule and radicle growth, and seedling dry weight	[130]
	crude plant powder mixed with clay loam soil		Inhibitory effects on leaf area index, total photosynthetic pigments, total available carbohydrates and total protein, in pot culture bioassay	
<i>A. scoparia</i> fresh leaves	essential oil (β -myrcene, (+)-limonene, (Z)- β -ocimene, γ -terpinene)	<i>Avena fatua</i> , <i>Cyperus rotundus</i> , <i>Phalaris minor</i>	Important reduction in germination, seedling growth, and dry matter at 0.07–0.7 mg/mL	[131]
<i>A. scoparia</i> fresh leaves	essential oil (p-cymene, β -myrcene, (+)-limonene)	<i>Achyranthes aspera</i> , <i>Cassia occidentalis</i> , <i>Parthenium hysterophorus</i> , <i>Echinochloa crus-galli</i> , <i>Ageratum conyzoides</i>	Inhibition of seed germination, root and shoot growth at 10, 25, and 50 μ g oil/g sand Chlorosis, necrosis and complete wilting of plants 1 to 7-days after spraying with oil (2%, 4%, and 6%, v/v) Significant decline in chlorophyll content and cellular respiration, electrolyte leakage	[132]
<i>A. sieversiana</i> fresh aerial parts	essential oil (α -thujone, eucalyptol)	<i>Amaranthus retroflexus</i> , <i>Medicago sativa</i> , <i>Poa annua</i> , <i>Pennisetum alopecuroides</i>	Inhibition of root and shoot growth IC ₅₀ = 1.89–4.69 mg/mL	[133]
	α -thujone		IC ₅₀ = 1.55–6.21 mg/mL	
	eucalyptol		IC ₅₀ = 1.42–17.81 mg/mL	
	α -thujone and eucalyptol mixture		IC ₅₀ = 0.23–1.05 mg/mL	
<i>A. terrae-albae</i> aerial parts	essential oil (α -thujone, β -thujone, eucalyptol, camphor)	<i>Amaranthus retroflexus</i>	Reduces root and shoot growth at 1.5 μ g/mL Completely inhibits seed germination at 3 μ g/mL	[134]
		<i>Poa annua</i>	Reduces root and shoot growth at 1.5 μ g/mL Completely inhibits seed germination at 5 μ g/mL	
<i>A. verlotiorum</i> flower heads	essential oil (chrysanthenone, 1,8-cineole, β -pinene, camphor, 2,6-dimethyl phenol, β -caryophyllene)	<i>Amaranthus retroflexus</i>	In vitro, complete inhibition of seed germination, at 10 and 100 μ g/L In vivo, plant death, at the cotyledon stage (100 mg/L) and true leaf stage (1000 mg/L)	[117]
		<i>Setaria viridis</i>	In vitro, inhibition of seed germination, at 10 and 100 μ g/L In vivo, plant death, at the cotyledon stage (1000 mg/L) and true leaf stage (1000 mg/L)	

Artemisia Species	Extract * or Compound Tested	Weed/Target Plant	Observed Effect	Reference
<i>A. vulgaris</i> aerial parts	aqueous extract	<i>Amaranthus retroflexus</i>	Inhibition of seed germination, radicle, and hypocotyl length at 7.5% to 10% w/v, in Petri dish bioassays Inhibition of seedling emergence and plant growth, in pot culture bioassays	[104]
		<i>Zea mays</i>	Stimulation of radicle and mesocotyl growth at 7.5% to 10% w/v, in Petri dish bioassays Stimulation of plant biomass, in pot culture bioassays	
<i>A. vulgaris</i> leaves and flowers	essential oil	<i>Agrostemma githago</i> , <i>Amaranthus retroflexus</i> , <i>Cardaria draba</i> , <i>Chenopodium album</i> , <i>Echinochloa crus-galli</i> , <i>Reseda lutea</i> , <i>Rumex crispus</i> , <i>Trifolium pratense</i>	Inhibition of root and shoot growth and reduction of germination rate (at 2, 5, 10 and 20 µL/plate)	[135]
<i>A. vulgaris</i> root	aqueous extracts	<i>Triticum aestivum</i> (winter wheat)	Inhibition of shoot and root growth by all concentrations (1:6250 to 1:10)	[136]
<i>A. vulgaris</i> aerial parts		<i>Brassica napus</i> spp. <i>oleifera</i> var. <i>biennis</i> (winter oilseed rape)	Significant inhibition of germination at the 1:10 concentration Significant inhibition of root growth at 1:10 concentration Stimulation of shoot growth	

* To highlight the active compounds, the major constituents of the volatile oils were noted in parentheses.

Artemisia fragrans essential oil inhibited seed germination and growth of *Convolvulus arvensis* at 1–4% concentration in a Petri dish and pot experiment. It significantly reduced the level of photosynthetic pigments (chlorophyll a, chlorophyll b, and carotenoids) and of antioxidant enzymes (catalase, peroxidase, ascorbate peroxidase, superoxide dismutase), as well as enhancing the production of hydrogen peroxide and malondialdehyde. It seems that volatile oil compounds—mostly oxygenated monoterpenes—inhibited the electron transport chain and affected the process of photosynthesis, leading to an increased production of oxygen reactive species. In turn, these intensified the lipid peroxidation of the cell membrane followed by electrolyte leakage [126].

Oxygenated monoterpenes were the major ingredients of *Artemisia sieversiana* essential oil (α -thujone 64.46% and eucalyptol 10.15%) that suppressed seedling growth of *Amaranthus retroflexus*, *Medicago sativa*, *Poa annua*, and *Pennisetum alopecuroides*. The experiment showed that the mixture of the major constituents, in the same ratio as found in the oil, was more phytotoxic compared to each individual compound, indicating a possible synergistic effect of α -thujone and eucalyptol [133].

Although oxygenated monoterpenes were the major constituents of *A. judaica* essential oils obtained by hydro-distillation and microwave-assisted extraction, the oil extracted by hydro-distillation exhibited greater phytotoxicity on *Lactuca sativa* seed germination and plant growth [36], showing that the extraction method impacts the phytotoxic activity of volatile oils.

Major constituents of *A. terrae-albae* essential oil were tested on seed germination, root and shoot growth of *Poa annua* and *Amaranthus retroflexus*. The phytotoxic effect of α -thujone, eucalyptol, camphor, and the mixture of these compounds was inferior to that of the essential oil, which suggests that probably other volatile components are causing the herbicidal activity of the oil [134]. α -Terpinen and β -pinene, compounds of *A. lavandulaefolia* essential oil, exhibited strong phytotoxic activity on seed germination test against eight target plants (Table 3), whereas β -caryophyllene and myrcene only inhibited *Achyranthes japonica* seed germination [128].

Artemisia scoparia essential oil inhibits germination and plant growth through the production of oxidative stress related to membrane disruption, increased lipid peroxidation, and buildup of hydrogen peroxide. It also interferes in cellular respiration and photosynthesis processes [132].

Field experiments in *Triticum aestivum* used pre-emergence application of *Artemisia vulgaris* aqueous extract (20% w/v) together with chlorsulfuron. This treatment permitted lowering the dose of the herbicide up to 80%, while manifesting an inhibitory effect of 70% against *Lolium multiflorum* [137]. Another field trial demonstrated that *A. argyi* water extract markedly suppressed the growth of weeds in *Chrysanthemum morifolium* field with no adverse effect on the growth of *C. morifolium*. The investigations showed that *A. argyi* inhibited weed growth and germination through inhibition of chlorophyll synthesis and photosynthesis [108]. Conversely, field treatment of *Triticum turgidum* L. subsp. *durum* Desf. with *A. absinthium* aqueous extract exerted a stimulating effect on weed presence and reduced wheat growth and yield [106].

The sensitivity of different weed species to a certain herbicide varies greatly. Among eight weeds tested in a study, *Amaranthus retroflexus*, *Echinochloa crus-galli*, and *Reseda lutea* were more susceptible to the action of *A. vulgaris* essential oil, compared to *Rumex crispus*, *Agrostemma githago*, *Trifolium pretense*, *Chenopodium album*, and *Cardaria draba*, which were more resistant [135]. Similarly, *Parthenium hysterophorus* and *Ageratum conyzoides* were more vulnerable to the inhibitory effect of *Artemisia scoparia* volatile oil, in comparison with *Cassia occidentalis*, under laboratory conditions. In another test, *Echinochloa crus-galli* and *Parthenium hysterophorus* were more affected by post-emergence application of the oil [132].

The phytotoxicity of isolated compounds from *Artemisia annua* was evaluated against two monocots and five dicots (Table 3). The suppression of germination and seedling growth varies in the order: artemisinin>arteannuin B>artemisinic acid. *Raphanus sativus* was the most resistant to the action of tested compounds, followed by *Secale cereale*. The weaker activity of arteannuin B and artemisinic acid—molecules without an endoperoxide bridge—implies that the moiety is important for the phytotoxic effect [118]. Artemisinin reduces many physiological and biochemical processes in the target plant and affects mitosis by inhibiting microtubules formation [120][138].

The incorporation of artemisinin into soil inhibited the growth of above-ground lettuce plants, with $EC_{50} = 2.5$ mg/Kg sandy soil, but the germination was not arrested up to 100 mg/Kg soil [139]. Furthermore, adding *A. annua* leaves containing 0.81–0.22% artemisinin in soil led to the inhibition of *Zea mays* growth [140]. Artemisinin is phytotoxic in concentrations comparable to those of commercial herbicides and has a good activity in soil [110].

In vivo tests proved that artemisinin is a potent suppressor of photosynthetic activity through the formation of a highly reactive artemisinin-metabolite that is able to inhibit the photosynthetic electron flow [141]. Other investigations showed that artemisinin enhances the generation of radical oxygen species and lipid peroxidation, which leads to cell death and arrest of mitotic phases in *Lactuca sativa* seedlings [119]. When added to the culture medium of *Arabidopsis thaliana* seedlings, artemisinin (1, 2, 5, 20, 100 μ M) reduced the root gravitropic responses, elongation of primary and lateral roots, root hairs density, and length. Furthermore, artemisinin diminished starch grain and auxin concentrations and affected auxin redistribution in root tips [142].

2.6. Activity on Non-Target Organisms

Since biopesticides and bioherbicides are of natural origin, they are considered to be less harmful to the environment and the health of applicators and consumers. Usually, plant-based formulations are mixtures of compounds, and they do not consist of a single substance, which should prevent resistance in target organisms. In addition, some phytochemicals are rapidly degraded in nature, so there is no risk of their accumulation in the environment, as is the case with chemical pesticides. Consequently, plant-based pesticides and herbicides are regarded as generally safe. Still, these products can affect the non-target organism directly or indirectly by influencing biodiversity and species interactions, so it is imperative to assess their safety [13][143].

Little information is available regarding the ecotoxicity of *Artemisia* compounds and extracts. Pino-Otin et al. [13] evaluated the toxicity of hydrolate and organic extracts from *A. absinthium* on three aquatic ecotoxicity indicator organisms: an invertebrate (*Daphnia magna*), a marine bacterium (*Vibrio fischeri*), and a unicellular freshwater alga (*Chlamydomonas reinhardtii*). The wormwood hydrolate, a by-product of essential oil extraction, is a promising biopesticide with nematicidal effect due to (5Z)-2,6-dimethylocta-5,7-dien-2,3-diol [97]. *A. absinthium* hydrolate caused acute toxicity on non-target organisms: *D. magna* ($LC_{50} = 0.236\%$) > *V. fischeri* ($LC_{50} = 1.85\%$) > *C. reinhardtii* ($LC_{50} = 16.49\%$). Moreover, the wormwood ethanol extract was highly toxic to *D. magna* ($LC_{50} = 0.093$ mg/L). However, the effect of wormwood hydrolate on a river microbial community, composed mainly of Proteobacteria, was negligible, causing only small changes in metabolic diversity and a slight inhibition of bacterial growth. It is possible that natural freshwater microbial populations are more resistant to 2,6-dimethylocta-5,7-diene-2,3-diol action because of the modified bioavailability of compounds in the river water and particular sensitivity of the various microbial species [13].

The same *A. absinthium* hydrolate was tested on non-target soil organisms: natural microbial communities, the earthworm *Eisenia fetida*, and the plant *Allium cepa*. The hydrolate was toxic in low concentrations: it caused substantial inhibition of onion root growth ($LC_{50} = 3.87\%$ v/v), high mortality of the earthworm *E. fetida* ($LC_{50} = 0.07$ mL/g), and decreased bacterial metabolism ($LC_{50} = 25.72\%$ v/v after 1 day of exposure). All these effects were exhibited at inferior concentrations than those needed to contain the target organism. Probably, 2,6-dimethylocta-5,7-diene-2,3-diol is able to penetrate biological membranes and thus affect the survival and metabolic processes of soil organism from different trophic levels [13].

The methanol extracts of *Artemisia fragrans* manifested significant toxicity in the brine shrimp (*Artemia salina*) lethality assay, with $ED_{50} = 19.7$ ppm for the root extract and $ED_{50} = 11.99$ ppm for flowers and leaves extract [56]. In another study, the aqueous extracts from *Artemisia ordosica* leaves were tested on two algae from the biological soil crusts, *Chlorella vulgaris* and *Nostoc* spp. The less concentrated extract (1 g/L) stimulated *C. vulgaris* growth but did not significantly affect *Nostoc* spp., indicating that *C. vulgaris* might utilize the sugars and other carbon sources in the extract to promote self-growth. The highly concentrated extract (5 and 10 g/L) inhibited the growth of both algae [109].

The safety profile of the *Artemisia nilagirica* essential oil was determined in terms of mammalian toxicity on male mice (*Mus musculus*) and millet (*Eleusine coracana*) seeds viability. The essential oil showed low toxicity on mice ($LD_{50} = 7528.10$ μ L/kg) and no effect on millet seed germination. Thus, the oil is suitable as a food preservative for both consumption and sowing purposes [40]. More so, *Artemisia nilagirica* essential oil did not cause any significant changes in the physicochemical and sensory properties of table grapes when applied by fumigation on the fruits [39].

Artemisia absinthium essential oil, a potential biopesticide, was evaluated for toxicity against non-target organisms: the honey bee (*Apis mellifera*) and tomato plant (*Solanum lycopersicum*). Honeybee toxicity ($EC_{50} = 0.26$ mg/cm²) is reached at lower concentrations of *A. absinthium* oil than the ones necessary for controlling the leaf miner *Tuta absoluta* ($EC_{50} = 0.5$ mg/cm²), but not at rates needed to control the whitefly *Trialeurodes vaporariorum* ($EC_{50} = 0.08$ mg/cm²). A similar phenomenon was noted for the phytotoxic effect on tomato; seed germination and root growth were inhibited at oil concentrations needed to control the leaf miner, but not the whitefly [66].

Investigations to date have shown that biopesticides derived from *Artemisia* are most likely to have some toxicity toward non-target organisms, and further studies are needed to assess the risk in natural communities in order to ensure the safe use of biopesticides in agricultural practices.

Choosing the right formulation can reduce toxicity as well as increase the stability and effectiveness of *Artemisia* biopesticides. For instance, terpenoids are lipophilic, volatile, and thermolabile compounds that are easily oxidized or hydrolyzed, so they can be affected during extraction, storage, and transport. Furthermore, after application onto plants, they volatilized quickly and start degrading, leading to short persistence and low efficacy in the field. These drawbacks can be overcome by a suitable formulation through encapsulation or nanoparticles synthesis. A product formulation is a homogeneous and stable mixture of components put together according to a specific procedure with the purpose of increasing the biological activity, stability, persistence, and efficiency, while decreasing the toxicity of the product. The selected formulation depends on the intended use and mode of application, the targeted phytopathogen, and the degradation factors present in the ecosystem [16].

References

1. Abad, M.J.; Bedoya, L.M.; Apaza, L.; Bermejo, P. The *Artemisia* L. Genus: A review of bioactive essential oils. *Molecules* 2012, 17, 2542–2566.
2. Trendafilova, A.; Moujir, L.M.; Sousa, P.M.C.; Seca, A.M.L. Research advances on health effects of edible *Artemisia* Species and some sesquiterpene lactones constituents. *Foods* 2020, 10, 65.
3. Septembre-Malaterre, A.; Lalarizo Rakoto, M.; Marodon, C.; Bedoui, Y.; Nakab, J.; Simon, E.; Hoarau, L.; Savriama, S.; Strasberg, D.; Guiraud, P.; et al. *Artemisia annua*, a traditional plant brought to light. *Int. J. Mol. Sci.* 2020, 21, 4986.
4. Ekiert, H.; Pajor, J.; Klin, P.; Rzepiela, A.; Ślesak, H.; Szopa, A. Significance of *Artemisia vulgaris* L. (Common Mugwort) in the history of medicine and its possible contemporary applications substantiated by phytochemical and pharmacological studies. *Molecules* 2020, 25, 4415.
5. Szopa, A.; Pajor, J.; Klin, P.; Rzepiela, A.; Elansary, H.O.; Al-Mana, F.A.; Mattar, M.A.; Ekiert, H. *Artemisia absinthium* L.—Importance in the history of medicine, the latest advances in phytochemistry and therapeutical, cosmetological and culinary uses. *Plants* 2020, 9, 1063.

6. Pandey, A.K.; Singh, P. The Genus *Artemisia*: A 2012–2017 Literature review on chemical composition, antimicrobial, insecticidal and antioxidant activities of essential oils. *Medicines* 2017, 4, 68.
7. Liu, S.-J.; Liao, Z.-X.; Tang, Z.-S.; Cui, C.-L.; Liu, H.-B.; Liang, Y.-N.; Zhang, Y.; Shi, H.-X.; Liu, Y.-R. Phytochemicals and biological activities of *Artemisia sieversiana*. *Phytochem. Rev.* 2017, 16, 441–460.
8. Dib, I.; Angenot, L.; Mihamou, A.; Ziyat, A.; Tits, M. *Artemisia campestris* L.: Ethnomedicinal, phytochemical and pharmacological review. *J. Herb. Med.* 2017, 7, 1–10.
9. Martinez, A.J. Natural fungicides obtained from plants. In *Fungicides for Plant and Animal Diseases*; InTech: London, UK, 2012; pp. 3–28.
10. Bora, K.S.; Sharma, A. The genus *Artemisia*: A comprehensive review. *Pharm. Biol.* 2011, 49, 101–109.
11. Basaid, K.; Chebli, B.; Mayad, E.H.; Furze, J.N.; Bouharroud, R.; Krier, F.; Barakate, M.; Paulitz, T. Biological activities of essential oils and lipopeptides applied to control plant pests and diseases: A review. *Int. J. Pest Manag.* 2021, 67, 155–177.
12. Saroj, A.; Oriyomi, O.V.; Nayak, A.K.; Haider, S.Z. Phytochemicals of plant-derived essential oils. In *Natural Remedies for Pest, Disease and Weed Control*; Elsevier: Amsterdam, The Netherlands, 2020; pp. 65–79.
13. Pino-Otín, M.R.; Ballester, D.; Navarro, E.; González-Coloma, A.; Val, J.; Mainar, A.M. Ecotoxicity of a novel biopesticide from *Artemisia absinthium* on non-target aquatic organisms. *Chemosphere* 2019, 216, 131–146.
14. Pavla, R.; Benelli, G. Essential oils as ecofriendly biopesticides? Challenges and constraints. *Trends Plant Sci.* 2016, 21, 1000–1007.
15. Jain, A.; Sarsaiya, S.; Wu, Q.; Lu, Y.; Shi, J. A review of plant leaf fungal diseases and its environment speciation. *Bioengineered* 2019, 10, 409–424.
16. Raveau, R.; Fontaine, J.; Lounès-Hadj Sahraoui, A. Essential oils as potential alternative biocontrol products against plant pathogens and weeds: A review. *Foods* 2020, 9, 365.
17. Badea, M.L.; Delian, E. In vitro antifungal activity of the essential oils from *Artemisia* spp. L. on *Sclerotinia sclerotiorum*. *Rom. Biotechnol. Lett.* 2014, 19, 9345–9352.
18. Julio, L.F.; Burillo, J.; Giménez, C.; Cabrera, R.; Díaz, C.E.; Sanz, J.; González-Coloma, A. Chemical and biocidal characterization of two cultivated *Artemisia absinthium* populations with different domestication levels. *Ind. Crops Prod.* 2015, 76, 787–792.
19. Parveen, S.; Wani, A.H.; Ganie, A.A.; Pala, S.A.; Mir, R.A. Antifungal activity of some plant extracts on some pathogenic fungi. *Arch. Phytopathol. Plant Prot.* 2014, 47, 279–284.
20. Huang, X.; Chen, S.; Zhang, Y.; Wang, Y.; Zhang, X.; Bi, Z.; Yuan, H. Chemical composition and antifungal activity of essential oils from three *Artemisia* species against *Alternaria solani*. *J. Essent. Oil Bear. Plants* 2019, 22, 1581–1592.
21. Ma, Y.-N.; Chen, C.-J.; Li, Q.-Q.; Xu, F.-R.; Cheng, Y.-X.; Dong, X. Monitoring antifungal agents of *Artemisia annua* against *Fusarium oxysporum* and *Fusarium solani*, associated with *Panax notoginseng* root-rot disease. *Molecules* 2019, 24, 213.
22. Jhansi Rani, S.; Supraja, P.; Sujitha, A.; Kiranmayee, P.; Usha, R. Evaluation of antibacterial and antifungal activity of *Artemisia annua* during pre and post flowering stages. *Int. J. Curr. Res.* 2015, 7, 21581–21587.
23. Gautam, P.; Upadhyay, S.K.; Hassan, W.; Madan, T.; Sirdeshmukh, R.; Sundaram, C.S.; Gade, W.N.; Basir, S.F.; Singh, Y.; Sarma, P.U. Transcriptomic and proteomic profile of *Aspergillus fumigatus* on exposure to artemisinin. *Mycopathologia* 2011, 172, 331–346.
24. Bouzenna, H.; Krichen, L. *Pelargonium graveolens* L'Her. and *Artemisia arborescens* L. essential oils: Chemical composition, antifungal activity against *Rhizoctonia solani* and insecticidal activity against *Rhyssopertha dominica*. *Nat. Prod. Res.* 2013, 27, 841–846.
25. Guan, X.; Ge, D.; Li, S.; Huang, K.; Liu, J.; Li, F. Chemical composition and antimicrobial activities of *Artemisia argyi* Lévl. et Vant essential oils extracted by simultaneous distillation-extraction, subcritical extraction and hydrodistillation. *Molecules* 2019, 24, 483.
26. Wenqiang, G.; Shufen, L.; Ruixiang, Y.; Yanfeng, H. Comparison of composition and antifungal activity of *Artemisia argyi* Lévl. et Vant inflorescence essential oil extracted by hydrodistillation and supercritical carbon dioxide. *Nat. Prod. Res.* 2006, 20, 992–998.
27. Petretto, G.L.; Chessa, M.; Piana, A.; Masia, M.D.; Foddai, M.; Mangano, G.; Culeddu, N.; Afifi, F.U.; Pintore, G. Chemical and biological study on the essential oil of *Artemisia caerulea* L. ssp. *densiflora* (Viv.). *Nat. Prod. Res.* 2013, 27, 1709–1715.

28. Karabegović, I.; Nikolova, M.; Veličković, D.; Stojičević, S.; Veljković, V.; Lazić, M. Comparison of antioxidant and antimicrobial activities of methanolic extracts of the *Artemisia* sp. recovered by different extraction techniques. *Chin. J. Chem. Eng.* 2011, 19, 504–511.
29. Houicher, A.; Hechachna, H.; Özogul, F. In vitro determination of the antifungal activity of *Artemisia campestris* essential oil from Algeria. *Int. J. Food Prop.* 2016, 19, 1749–1756.
30. Eblaghi, M.; Khajehie, N.; Golmakani, M.-T.; Eskandari, M.H. Investigating the effects of microwave-assisted hydrodistillation on antioxidant and antifungal activities of *Tanacetum polycephalum* and *Artemisia chamaemelifolia* essential oils. *J. Essent. Oil Res.* 2016, 28, 528–539.
31. Goudjil, M.B.; Ladjel, S.; Bencheikh, S.E.; Hammoya, F.; Bensaci, M.B.; Zighmi, S.; Mehani, M. Bioactivity of *Artemisia herba alba* essential oil against plant pathogenic fungi. *Der Pharma Chem.* 2016, 8, 46–52.
32. Khaddor, M.; Lamarti, A.; Tantaoui-Elaraki, A.; Ezziyani, M.; Castillo, M.-E.C.; Badoc, A. Antifungal activity of three essential oils on growth and toxigenesis of *Penicillium aurantiogriseum* and *Penicillium viridicatum*. *J. Essent. Oil Res.* 2006, 18, 586–589.
33. Saleh, M.; Belal, M.; El-Baroty, G. Fungicidal activity of *Artemisia herba alba* Asso (Asteraceae). *J. Environ. Sci. Health Part B Pestic. Food Contam. Agric. Wastes* 2006, 41, 237–244.
34. El-Wassimy, M.T.M.; Ahmed, M.M.; Younes, S.H.H.; Hegazy, M.-E.F. In vitro antiproliferative and antimicrobial activity of 11-epiartapshin compound isolated from *Seriphidium herba-alba*. *J. Pharm. Appl. Chem.* 2018, 4, 147–154.
35. Rashid, M.U.; Alamzeb, M.; Ali, S.; Shah, Z.A.; Naz, I.; Khan, A.A.; Semaan, D.; Khan, M.R. A new irregular monoterpene acetate along with eight known compounds with antifungal potential from the aerial parts of *Artemisia incisa* Pamp (Asteraceae). *Nat. Prod. Res.* 2017, 31, 428–435.
36. Elshamy, A.; Abd-ElGawad, A.; Mohamed, T.; El Gendy, A.E.; Abd El Aty, A.A.; Saleh, I.; Moustafa, M.F.; Hussien, T.A.; Pare, P.W.; Hegazy, M. Extraction development for antimicrobial and phytotoxic essential oils from Asteraceae species: *Achillea fragrantissima*, *Artemisia judaica* and *Tanacetum sinaicum*. *Flavour Fragr. J.* 2021, 36, 352–364.
37. Hadian, J.; Ramak-Masoumi, T.; Farzaneh, M.; Mirjalili, M.-H.; Nejad-Ebrahimi, S.; Ghorbani, M. Chemical compositions of essential oil of *Artemisia khorasanica* Podl. and its antifungal activity on soil-born phytopathogens. *J. Essent. Oil Bear. Plants* 2007, 10, 53–59.
38. Mohan, M.; Pandey, A.K.; Singh, P.; Nautiyal, M.K.; Gupta, S. Evaluation of *Artemisia maritima* L. essential oil for its chemical and biological properties against some foodborne pathogens. *Anal. Chem. Lett.* 2016, 6, 47–54.
39. Sonker, N.; Pandey, A.K.; Singh, P. Efficiency of *Artemisia nilagirica* (Clarke) Pamp. essential oil as a mycotoxin against postharvest mycobiota of table grapes. *J. Sci. Food Agric.* 2015, 95, 1932–1939.
40. Kumar, M.; Dwivedy, A.K.; Sarma, P.; Dkhar, M.S.; Kayang, H.; Raghuwanshi, R.; Dubey, N.K. Chemically characterized *Artemisia nilagirica* (Clarke) Pamp. essential oil as a safe plant-based preservative and shelf-life enhancer of millets against fungal and aflatoxin contamination and lipid peroxidation. *Plant Biosyst. An Int. J. Deal. Asp. Plant Biol.* 2020, 154, 269–276.
41. Sati, S.C.; Sati, N.; Ahluwalia, V.; Walia, S.; Sati, O.P. Chemical composition and antifungal activity of *Artemisia nilagirica* essential oil growing in northern hilly areas of India. *Nat. Prod. Res.* 2013, 27, 45–48.
42. Shafi, P.M.; Nambiar, M.K.G.; Clery, R.A.; Sarma, Y.R.; Veena, S.S. Composition and antifungal activity of the oil of *Artemisia nilagirica* (Clarke) Pamp. *J. Essent. Oil Res.* 2004, 16, 377–379.
43. Deepak, S.A.; Oros, G.; Sathyanarayana, S.G.; Shetty, N.P.; Shetty, H.S.; Sashikanth, S. Antisporulant activity of leaf extracts of Indian plants against *Sclerotinia graminicola* causing downy mildew disease of pearl millet. *Arch. Phytopathol. Plant Prot.* 2005, 38, 31–39.
44. Sampietro, D.A.; Lizarraga, E.F.; Ibatayev, Z.A.; Omarova, A.B.; Suleimen, Y.M.; Catalán, C.A.N. Chemical composition and antimicrobial activity of essential oils from *Acantholippia deserticola*, *Artemisia proceriformis*, *Achillea micrantha* and *Libanotis buchtormensis* against phytopathogenic bacteria and fungi. *Nat. Prod. Res.* 2016, 30, 1950–1955.
45. Mohamed, T.A.; Hegazy, M.-E.F.; Abd El Aty, A.A.; Ghabbour, H.A.; Alsaid, M.S.; Shahat, A.A.; Paré, P.W. Antimicrobial sesquiterpene lactones from *Artemisia sieberi*. *J. Asian Nat. Prod. Res.* 2017, 19, 1093–1101.
46. Ghasemi, G.; Alirezalu, A.; Ishkeh, S.R.; Ghosta, Y. Phytochemical properties of essential oil from *Artemisia sieberi* Besser (Iranian accession) and its antioxidant and antifungal activities. *Nat. Prod. Res.* 2020, 1–5.
47. Manika, N.; Chanotiya, C.S.; Darokar, M.; Singh, S.; Bagchi, G.D. Compositional characters and antimicrobial potential of *Artemisia stricta* Edgew. f. *stricta* Pamp. essential oil. *Rec. Nat. Prod.* 2016, 10, 40–44.
48. Sampietro, D.A.; de los Angeles Gomez, A.; Jimenez, C.M.; Lizarraga, E.F.; Ibatayev, Z.A.; Suleimen, Y.M.; Catalán, C.A. Chemical composition and antifungal activity of essential oils from medicinal plants of Kazakhstan. *Nat. Prod. Res.* 2017,

49. Behravan, J.; Ramezani, M.; Hassanzadeh, M.K.; Eliaspour, N.; Sabeti, Z. Cytotoxic and antimycotic activities of essential oil of *Artemisia turanica* Krasch from Iran. *J. Essent. Oil Bear. Plants* 2006, 9, 196–203.
50. Park, I.-K.; Kim, J.; Lee, Y.-S.; Shin, S.-C. In vivo fungicidal activity of medicinal plant extracts against six phytopathogenic fungi. *Int. J. Pest Manag.* 2008, 54, 63–68.
51. da Cruz Cabral, L.; Fernández Pinto, V.; Patriarca, A. Application of plant derived compounds to control fungal spoilage and mycotoxin production in foods. *Int. J. Food Microbiol.* 2013, 166, 1–14.
52. Tantaoui-Elaraki, A.; Ferhout, H.; Errifi, A. Inhibition of the fungal asexual reproduction stages by three Moroccan essential oils. *J. Essent. Oil Res.* 1993, 5, 535–545.
53. Xu, Y.; Zhao, L.; Chen, P.; Jiang, X.; Wei, G. Isolation, screening and characterization of phytopathogen antagonistic endophytes from wild *Artemisia argyi*. *Acta Ecol. Sin.* 2013, 33, 3697–3705.
54. Ahameethunisa, A.R.; Hopper, W. Antibacterial activity of *Artemisia nilagirica* leaf extracts against clinical and phytopathogenic bacteria. *BMC Complement. Altern. Med.* 2010, 10.
55. Sukanya, S.L.; Sudisha, J.; Hariprasad, P.; Niranjana, S.R.; Prakash, H.S.; Fathima, S.K. Antimicrobial activity of leaf extracts of Indian medicinal plants against clinical and phytopathogenic bacteria. *Afr. J. Biotechnol.* 2009, 8, 6677–6682.
56. Inayatullah, S.; Irum, R.; Ateeq-Ur-Rehman; Chaudhary, M.F.; Mirza, B. Biological evaluation of some selected plant species of Pakistan. *Pharm. Biol.* 2007, 45, 397–403.
57. Dadasoglu, F.; Kotan, R.; Cakir, A.; Cakmakci, R.; Kordali, S.; Ozer, H.; Karagoz, K.; Dikbas, N. Antibacterial activities of essential oils, extracts and some of their major components of *Artemisia* spp. L. against seed-borne plant pathogenic bacteria. *Fresenius Environ. Bull.* 2015, 24, 2715–2724.
58. Mathlouthi, A.; Saadaoui, N.; Pennacchiotti, E.; De Biase, D.; Ben-Attia, M. Essential oils from *Artemisia* species inhibit biofilm formation and the virulence of *Escherichia coli* EPEC 2348/69. *Biofouling* 2021, 37, 174–183.
59. Hikal, W.M.; Baeshen, R.S.; Said-Al Ahl, H.A.H. Botanical insecticide as simple extractives for pest control. *Cogent Biol.* 2017, 3, 1404274.
60. Spochacz, M.; Chowański, S.; Walkowiak-Nowicka, K.; Szymczak, M.; Adamski, Z. Plant-derived substances used against beetles—pests of stored crops and food—and their mode of action: A Review. *Compr. Rev. Food Sci. Food Saf.* 2018, 17, 1339–1366.
61. Regnault-Roger, C.; Philogène, B.J.R. Past and current prospects for the use of botanicals and plant allelochemicals in integrated pest management. *Pharm. Biol.* 2008, 46, 41–52.
62. Rattan, R.S. Mechanism of action of insecticidal secondary metabolites of plant origin. *Crop Prot.* 2010, 29, 913–920.
63. Abdelgaleil, S.A.M.; Badawy, M.E.I.; Mahmoud, N.F.; Marei, A.E.-S.M. Acaricidal activity, biochemical effects and molecular docking of some monoterpenes against two-spotted spider mite (*Tetranychus urticae* Koch). *Pestic. Biochem. Physiol.* 2019, 156, 105–115.
64. Abdelgaleil, S.A.M.; Mohamed, M.I.E.; Shawir, M.S.; Abou-Taleb, H.K. Chemical composition, insecticidal and biochemical effects of essential oils of different plant species from Northern Egypt on the rice weevil, *Sitophilus oryzae* L. *J. Pest Sci.* 2016, 89, 219–229.
65. Kumar, S.; Nehra, M.; Dilbaghi, N.; Marrazza, G.; Hassan, A.A.; Kim, K.H. Nano-based smart pesticide formulations: Emerging opportunities for agriculture. *J. Control. Release* 2019, 294, 131–153.
66. Umpiérrez, M.L.; Paullier, J.; Porrini, M.; Garrido, M.; Santos, E.; Rossini, C. Potential botanical pesticides from Asteraceae essential oils for tomato production: Activity against whiteflies, plants and bees. *Ind. Crops Prod.* 2017, 109, 686–692.
67. Cheng, Z.H.; Duan, H.J.; Zhu, X.R.; Fan, F.F.; Li, R.; Li, S.C.; Ma, X.Y.; Zhang, E.J.; Liu, Y.K.; Wang, J.Y. Effects of patchouli and wormwood oils on the bioassays and behaviors of *Tetranychus cinnabarinus* (Boisduval) (Acari: Tetranychida). *Int. J. Pest Manag.* 2020, 66, 271–278.
68. Seixas, P.T.L.; Demuner, A.J.; Alvarenga, E.S.; Barbosa, L.C.A.; Marques, A.; de Farias, E.S.; Picanço, M.C. Bioactivity of essential oils from *Artemisia* against *Diaphania hyalinata* and its selectivity to beneficial insects. *Sci. Agric.* 2018, 75, 519–525.
69. Dane, Y.; Mouhouche, F.; Canela-Garayoa, R.; Delpino-Rius, A. Phytochemical analysis of methanolic extracts of *Artemisia absinthium* L. 1753 (Asteraceae), *Juniperus phoenicea* L., and *Tetraclinis articulata* (Vahl) Mast, 1892 (Cupressaceae) and evaluation of their biological activity for stored grain protection. *Arab. J. Sci. Eng.* 2016, 41, 2147–2158.
70. Bachrouch, O.; Ferjani, N.; Haouel, S.; Jemâa, J.M.B. Major compounds and insecticidal activities of two Tunisian *Artemisia* essential oils toward two major coleopteran pests. *Ind. Crops Prod.* 2015, 65, 127–133.

71. Kłyś, M.; Przystupińska, A. The mortality of *Oryzaephilus surinamensis* Linnaeus, 1758 (Coleoptera: Silvanidae) induced by powdered plants. *J. Plant Prot. Res.* 2015, 55, 110–116.
72. Brudea, V.; Rîșca, I.; Enea, C.; Tomescu, C. Efficacy of some biopesticides and plant secondary metabolites against fall webworm *Hyphantria Cunea* Drury (F. Arctiidae-Lepidoptera) in the lab conditions. *Cercet. Agron. Mold.* 2012, 45, 73–80.
73. Martín, L.; González-Coloma, A.; Burillo, J.; Palavra, A.M.F.F.; Urieta, J.S.; Mainar, A.M. Microcalorimetric determination of the activity of supercritical extracts of wormwood (*Artemisia absinthium* L.) over *Spodoptera littoralis*. *J. Therm. Anal. Calorim.* 2013, 111, 1837–1844.
74. Czerniewicz, P.; Chrzanowski, G.; Sprawka, I.; Sytykiewicz, H. Aphicidal activity of selected Asteraceae essential oils and their effect on enzyme activities of the green peach aphid, *Myzus persicae* (Sulzer). *Pestic. Biochem. Physiol.* 2018, 145, 84–92.
75. Rizvi, S.A.H.; Ling, S.; Tian, F.; Xie, F.; Zeng, X. Toxicity and enzyme inhibition activities of the essential oil and dominant constituents derived from *Artemisia absinthium* L. against adult Asian citrus psyllid *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae). *Ind. Crops Prod.* 2018, 121, 468–475.
76. Anshul, N.; Kalra, A.; Singh, D. Biological effect of sweet wormwood, *Artemisia annua* methanol extracts and essential oil against *Helicoverpa armigera* Hub. (Lepidoptera : Noctuidae). *J. Entomol. Zool. Stud.* 2014, 2, 304–307.
77. Anshul, N.; Bhakuni, R.S.; Gaur, R.; Singh, D. Isomeric flavonoids of *Artemisia annua* (Asterales: Asteraceae) as insect growth inhibitors against *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Florida Entomol.* 2013, 96, 897–903.
78. Oftadeh, M.; Sendi, J.J.; Ebadollahi, A. Toxicity and deleterious effects of *Artemisia annua* essential oil extracts on mulberry pyralid (*Glyphodes pyloalis*). *Pestic. Biochem. Physiol.* 2020, 170, 104702.
79. Hasheminia, S.M.; Sendi, J.J.; Jahromi, K.T.; Moharramipour, S. The effects of *Artemisia annua* L. and *Achillea millefolium* L. crude leaf extracts on the toxicity, development, feeding efficiency and chemical activities of small cabbage *Pieris rapae* L. (Lepidoptera: Pieridae). *Pestic. Biochem. Physiol.* 2011, 99, 244–249.
80. Zibae, A.; Zibae, I.; Bandani, A.R. *Artemisia annua* L. (Asteraceae) changes some biochemical compounds in the hemolymph of *Hyphantria cunea* Drury (Lepidoptera: Arctiidae). *J. Med. Plants Res.* 2011, 5, 3229–3235.
81. Khosravi, R.; Sendi, J.J.; Ghadamyari, M.; Yezdani, E. Effect of sweet wormwood *Artemisia annua* crude leaf extracts on some biological and physiological characteristics of the lesser mulberry pyralid, *Glyphodes pyloalis*. *J. Insect Sci.* 2011, 11, 1–13.
82. Ahmed, M.; Peiwen, Q.; Gu, Z.; Liu, Y.; Sikandar, A.; Hussain, D.; Javeed, A.; Shafi, J.; Iqbal, M.F.; An, R.; et al. Insecticidal activity and biochemical composition of *Citrullus colocynthis*, *Cannabis indica* and *Artemisia argyi* extracts against cabbage aphid (*Brevicoryne brassicae* L.). *Sci. Rep.* 2020, 10, 1–10.
83. Ebadollahi, A.; Ashouri, S. Toxicity of essential oils isolated from *Achillea millefolium* L., *Artemisia dracunculus* L. and *Heracleum persicum* Desf. against adults of *Plodia interpunctella* (Hübner) (Lepidoptera: Pyralidae) in Islamic Republic of Iran. *Ecol. Balk.* 2011, 3, 41–48.
84. Liu, X.C.; Li, Y.; Wang, T.; Wang, Q.; Liu, Z.L. Chemical composition and insecticidal activity of essential oil of *Artemisia frigida* Willd (Compositae) against two grain storage insects. *Trop. J. Pharm. Res.* 2014, 13, 587–592.
85. Zhang, Z.; Pang, X.; Guo, S.; Cao, J.; Wang, Y.; Chen, Z.; Feng, Y.; Lei, N.; Du, S. Insecticidal activity of *Artemisia frigida* Willd. essential oil and its constituents against three stored product insects. *Rec. Nat. Prod.* 2019, 13, 176–181.
86. Zhou, J.; Zou, K.; Zhang, W.; Guo, S.; Liu, H.; Sun, J.; Li, J.; Huang, D.; Wu, Y.; Du, S.; et al. Efficacy of compounds isolated from the essential oil of *Artemisia lavandulaefolia* in control of the cigarette beetle, *Lasioderma serricorne*. *Molecules* 2018, 23, 343.
87. Hussein, H.S.; Tawfeek, M.E.; Abdelgaleil, S.A.M. Chemical composition, aphicidal and antiacetylcholinesterase activities of essential oils against *Aphis nerii* Boyer de Fonscolombe (Hemiptera: Aphididae). *J. Asia. Pac. Entomol.* 2021.
88. Yumnam, S.; Singh, K.I.; Ray, D.C. Effect of indigenous plant extracts on the incidence of *Scirpophaga incertulas* (Walker) (Lepidoptera: Pyralidae) in Kharif rice ecosystem. *J. Sci. Ind. Res.* 2017, 76, 494–500.
89. Gokturk, T.; Kordali, S.; Calmasur, O.; Tozlu, G. Insecticidal effects of essential plant oils against larvae of great spruce bark beetle, *Dendroctonus micans* (Kugelann) (Coleoptera: Curculionidae: Scolytinae). *Fresenius Environ. Bull.* 2011, 20, 2365–2370.
90. Sharifian, I.; Hashemi, S.M.; Darvishzadeh, A. Fumigant toxicity of essential oil of mugwort (*Artemisia vulgaris* L.) against three major stored product beetles. *Arch. Phytopathol. Plant Prot.* 2013, 46, 445–450.
91. Jalali Sendi, J.; Khosravi, R. Recent developments in controlling insect, acari, nematode, and plant pathogens of agricultural and medical importance by *Artemisia annua* L. (Asteraceae). In *Artemisia Annua—Pharmacology and Biotechnology*

gy; Springer: Berlin/Heidelberg, Germany, 2014; pp. 229–247.

92. Oka, Y.; Nacar, S.; Putievsky, E.; Ravid, U.; Yaniv, Z.; Spiegel, Y. Nematicidal activity of essential oils and their components against the root-knot nematode. *Phytopathology* 2000, 90, 710–715.
93. Shakil, N.A.; Prasad, D.; Saxena, D.B.; Gupta, A.K. Nematicidal activity of essential oils of *Artemisia annua* against root-knot and reniform nematodes. *Ann. Plant Prot. Sci.* 2004, 12, 397–402.
94. Avato, P.; Laquale, S.; Argentieri, M.P.; Lamiri, A.; Radicci, V.; D'Addabbo, T. Nematicidal activity of essential oils from aromatic plants of Morocco. *J. Pest Sci.* 2017, 90, 711–722.
95. Amora, D.X.; de Podestá, G.S.; Grupioni, P.H.F.; das Nasu, É.G.C.; de Figueiredo, L.D.; Ferreira, F.C.; de Freitas, L.G.; Lopes, E.A.; Ferraz, S. Effect of essential oils on the root-knot nematode. *Rev. Agri Environ. Sci.* 2017, 3, 15–23.
96. Kundu, A.; Dutta, A.; Mandal, A.; Negi, L.; Malik, M.; Puramchatwad, R.; Antil, J.; Singh, A.; Rao, U.; Saha, S.; et al. A Comprehensive in vitro and in silico analysis of nematicidal action of essential oils. *Front. Plant Sci.* 2021, 11.
97. Julio, L.F.; González-Coloma, A.; Burillo, J.; Diaz, C.E.; Andrés, M.F. Nematicidal activity of the hydrolate byproduct from the semi industrial vapor pressure extraction of domesticated *Artemisia absinthium* against *Meloidogyne javanica*. *Crop Prot.* 2017, 94, 33–37.
98. Kalaiselvi, D.; Mohankumar, A.; Shanmugam, G.; Thiruppathi, G.; Nivitha, S.; Sundararaj, P. Altitude-related changes in the phytochemical profile of essential oils extracted from *Artemisia nilagirica* and their nematicidal activity against *Meloidogyne incognita*. *Ind. Crops Prod.* 2019, 139, 111472.
99. Suresh, J.; Mahesh, N.M.; Ahuja, J.; Santilna, K.S. Review on *Artemisia nilagirica* (Clarke) Pamp. *J. Biol. Act. Prod. Nat.* 2011, 1, 97–104.
100. Khan, R.; Naz, I.; Hussain, S.; Khan, R.A.A.; Ullah, S.; Rashid, M.U.; Siddique, I. Phytochemical management of root knot nematode (*Meloidogyne incognita*) on kofoid and white chitwood by *Artemisia* spp. in tomato (*Lycopersicon esculentum* L.). *Braz. J. Biol.* 2020, 80, 829–838.
101. Farooq, N.; Abbas, T.; Tanveer, A.; Jabran, K. Allelopathy for weed management. In *Co-Evolution of Secondary Metabolites*. Reference Series in Phytochemistry; Springer: Berlin/Heidelberg, Germany, 2020; pp. 505–519.
102. Chen, H.; Singh, H.; Bhardwaj, N.; Bhardwaj, S.K.; Khatri, M.; Kim, K.-H.; Peng, W. An exploration on the toxicity mechanisms of phytotoxins and their potential utilities. *Crit. Rev. Environ. Sci. Technol.* 2020, 1–41.
103. Muzell Trezzi, M.; Vidal, R.A.; Balbinot Junior, A.A.; von Hertwig Bittencourt, H.; da Silva Souza Filho, A.P. Allelopathy: Driving mechanisms governing its activity in agriculture. *J. Plant Interact.* 2016, 11, 53–60.
104. Pannacci, E.; Masi, M.; Farneselli, M.; Tei, F. Evaluation of mugwort (*Artemisia vulgaris* L.) aqueous extract as a potential bioherbicide to control *Amaranthus retroflexus* L. in maize. *Agriculture* 2020, 10, 642.
105. Kapoor, D.; Rinzim; Tiwari, A.; Sehgal, A.; Landi, M.; Brestic, M.; Sharma, A. Exploiting the allelopathic potential of aqueous leaf extracts of *Artemisia absinthium* and *Psidium guajava* against *Parthenium hysterophorus*, a widespread weed in India. *Plants* 2019, 8, 552.
106. Carrubba, A.; Labruzzo, A.; Comparato, A.; Muccilli, S.; Spina, A. Use of plant water extracts for weed control in durum wheat (*Triticum turgidum* L. Subsp. durum Desf.). *Agronomy* 2020, 10, 364.
107. Labruzzo, A.; Cantrell, C.L.; Carrubba, A.; Ali, A.; Wedge, D.E.; Duke, S.O. Phytotoxic lignans from *Artemisia arborescens*. *Nat. Prod. Commun.* 2018, 13, 1934578X1801300.
108. Li, J.; Chen, L.; Chen, Q.; Miao, Y.; Peng, Z.; Huang, B.; Guo, L.; Liu, D.; Du, H. Allelopathic effect of *Artemisia argyi* on the germination and growth of various weeds. *Sci. Rep.* 2021, 11, 4303.
109. Zhou, X.; Zhang, Y.; An, X.; De Philippis, R.; Ma, X.; Ye, C.; Chen, L. Identification of aqueous extracts from *Artemisia ordosica* and their allelopathic effects on desert soil algae. *Chemoecology* 2019, 29, 61–71.
110. Knudsmark Jessing, K.; Duke, S.O.; Cedergreen, N. Potential ecological roles of artemisinin produced by *Artemisia annua* L. *J. Chem. Ecol.* 2014, 40, 100–117.
111. Radhakrishnan, R.; Alqarawi, A.A.; Abd Allah, E.F. Bioherbicides: Current knowledge on weed control mechanism. *Ecotoxicol. Environ. Saf.* 2018, 158, 131–138.
112. Bordin, E.R.; Frumi Camargo, A.; Stefanski, F.S.; Scapini, T.; Bonatto, C.; Zanivan, J.; Preczeski, K.; Modkovski, T.A.; Reichert Júnior, F.; Mossi, A.J.; et al. Current production of bioherbicides: Mechanisms of action and technical and scientific challenges to improve food and environmental security. *Biocatal. Biotransform.* 2020, 1–14.
113. Verdeguer, M.; Sánchez-Moreiras, A.M.; Araniti, F. Phytotoxic effects and mechanism of action of essential oils and terpenoids. *Plants* 2020, 9, 1571.

114. Fouad, R.; Bousta, D.; Lalami, A.E.O.; Chahdi, F.O.; Amri, I.; Jamoussi, B.; Greche, H. Chemical composition and herbicidal effects of essential oils of *Cymbopogon citratus* (DC) Stapf, *Eucalyptus cladocalyx*, *Origanum vulgare* L. and *Artemisia absinthium* L. cultivated in Morocco. *J. Essent. Oil Bear. Plants* 2015, 18, 112–123.
115. Aryakia, E.; Naghavi, M.R.; Farahmand, Z.; Shahzadeh Fazeli, A.A.H. Evaluating allelopathic effects of some plant species in tissue culture media as an accurate method for selection of tolerant plant and screening of bioherbicides. *J. Agric. Sci. Technol.* 2015, 17, 1011–1023.
116. Ammann, N.; Pieterse, P.J. Effects of *Artemisia afra* leaf extracts on seed germination of selected crop and weed species. *S. Afr. J. Plant Soil* 2005, 22, 263–265.
117. Benvenuti, S.; Cioni, P.L.; Flamini, G.; Pardossi, A. Weeds for weed control: Asteraceae essential oils as natural herbicides. *Weed Res.* 2017, 57, 342–353.
118. Paramanik, R.C.; Chikkaswamy, B.K.; Roy, D.G.; Achinto, P.; Venkatesh, K. Effects of biochemicals of *Artemisia annua* in plants. *J. Phytol. Res.* 2008, 21, 11–18.
119. Yan, Z.-Q.; Wang, D.-D.; Ding, L.; Cui, H.-Y.; Jin, H.; Yang, X.-Y.; Yang, J.-S.; Qin, B. Mechanism of artemisinin phytotoxicity action: Induction of reactive oxygen species and cell death in lettuce seedlings. *Plant Physiol. Biochem.* 2015, 88, 53–59.
120. Hussain, M.I.; Reigosa, M.J. Characterization of xanthophyll pigments, photosynthetic performance, photon energy dissipation, reactive oxygen species generation and carbon isotope discrimination during artemisinin-induced stress in *Arabidopsis thaliana*. *PLoS ONE* 2015, 10, e0114826.
121. Araniti, F.; Gullì, T.; Marrelli, M.; Statti, G.; Gelsomino, A.; Abenavoli, M.R. *Artemisia arborescens* L. leaf litter: Phytotoxic activity and phytochemical characterization. *Acta Physiol. Plant.* 2016, 38, 128.
122. Araniti, F.; Lupini, A.; Sorgonà, A.; Conforti, F.; Marrelli, M.; Statti, G.A.; Menichini, F.; Abenavoli, M.R. Allelopathic potential of *Artemisia arborescens*: Isolation, identification and quantification of phytotoxic compounds through fractionation-guided bioassays. *Nat. Prod. Res.* 2013, 27, 880–887.
123. Dhifi, W.; Jilani, I.B.H.; Bellili, S.; Jazi, S.; El Beyrouthy, M.; Mnif, W. Essential oil chemical characterization and allelopathic potential of *Artemisia campestris* L. growing in Tunisia. *J. Microbiol. Biotechnol. Food Sci.* 2017, 7, 302–305.
124. da Silva Saraiva, C.S. Bioherbicidal Effect of Plant Aqueous Extracts and Essential Oils; Escola Superior Agrária de Coimbra: Coimbra, Portugal, 2019.
125. Sekine, T.; Appiah, K.S.; Azizi, M.; Fujii, Y. Plant growth inhibitory activities and volatile active compounds of 53 spices and herbs. *Plants* 2020, 9, 264.
126. Pouresmaeil, M.; Nojadeh, M.S.; Movafeghi, A.; Maggi, F. Exploring the bio-control efficacy of *Artemisia fragrans* essential oil on the perennial weed *Convolvulus arvensis*: Inhibitory effects on the photosynthetic machinery and induction of oxidative stress. *Ind. Crops Prod.* 2020, 155, 112785.
127. Zhang, R.M.; Zuo, Z.J.; Gao, P.J.; Hou, P.; Wen, G.S.; Gao, Y. Allelopathic effects of VOCs of *Artemisia frigida* Willd. on the regeneration of pasture grasses in Inner Mongolia. *J. Arid Environ.* 2012, 87, 212–218.
128. Kil, B.S.; Han, D.M.; Lee, C.H.; Kim, Y.S.; Yun, K.Y.; Yoo, H.G. Allelopathic effects of *Artemisia lavandulaefolia*. *Korean J. Ecol.* 2000, 23, 149–155.
129. Al-Watban, A.; Salama, H.M.H. Physiological effects of allelopathic activity of *Artemisia monosperma* on common bean (*Phaseolus vulgaris* L.). *Int. Res. J. Plant Sci.* 2012, 3, 158–163.
130. Algandaby, M.M.; El-Darier, S.M. Management of the noxious weed; *Medicago polymorpha* L. via allelopathy of some medicinal plants from Taif region, Saudi Arabia. *Saudi J. Biol. Sci.* 2018, 25, 1339–1347.
131. Singh, H.P.; Kaur, S.; Mittal, S.; Batish, D.R.; Kohli, R.K. Essential oil of *Artemisia scoparia* inhibits plant growth by generating reactive oxygen species and causing oxidative damage. *J. Chem. Ecol.* 2009, 35, 154–162.
132. Kaur, S.; Singh, H.P.; Mittal, S.; Batish, D.R.; Kohli, R.K. Phytotoxic effects of volatile oil from *Artemisia scoparia* against weeds and its possible use as a bioherbicide. *Ind. Crops Prod.* 2010, 32, 54–61.
133. Jiang, C.-Y.; Zhou, S.-X.; Toshmatov, Z.; Mei, Y.; Jin, G.-Z.; Han, C.-X.; Zhang, C.; Shao, H. Chemical composition and phytotoxic activity of the essential oil of *Artemisia sieversiana* growing in Xinjiang, China. *Nat. Prod. Res.* 2020, 1–6.
134. Shao, H.; Hu, Y.; Han, C.; Wei, C.; Zhou, S.; Zhang, C.; Zhang, C. Chemical composition and phytotoxic activity of *Seriphidium terrae-albae* (Krasch.) Poljakov (Compositae) essential oil. *Chem. Biodivers.* 2018, 15, e1800348.
135. Önen, H.; Özer, Z.; Telci, I. Bioherbicidal effects of some plant essential oils on different weed species. *J. Plant Dis. Prot.* 2002, 18, 597–605.
136. Marcinkeviciene, A.; Kriauciuniene, Z.; Velicka, R.; Kosteckas, R.; Fujii, Y. Allelopathic effect of *Artemisia vulgaris* on winter wheat and winter oilseed rape. *Fresenius Environ. Bull.* 2018, 27, 727–732.

137. Pannacci, E.; Pettorossi, D.; Regni, L.; Tei, F. Allelopathic potential of mugwort (*Artemisia vulgaris* L.) to control the Italian ryegrass (*Lolium multiflorum* Lam.) in winter wheat. *Allelopath. J.* 2015, 36, 257–272.
138. Dayan, F.E.; Duke, S.O. Biological activity of allelochemicals. In *Plant-Derived Natural Products*; Springer: New York, NY, USA, 2009; pp. 361–384.
139. Jessing, K.K.; Cedergreen, N.; Jensen, J.; Hansen, H.C.B. Degradation and ecotoxicity of the biomedical drug artemisinin in soil. *Environ. Toxicol. Chem.* 2009, 28, 701–710.
140. Delabays, N.; Slacanin, I.; Bohren, C. Herbicidal potential of artemisinin and allelopathic properties of *Artemisia annua* L: From the laboratory to the field. *J. Plant Dis. Protection* 2008, 317–322.
141. Bharati, A.; Kar, M.; Sabat, S.C. Artemisinin inhibits chloroplast electron transport activity: Mode of action. *PLoS ONE* 2012, 7, e38942.
142. Yan, Z.; Wang, D.; Cui, H.; Sun, Y.; Yang, X.; Jin, H.; Zhao, Y.; Li, X.; Xie, M.; Liu, J.; et al. Effects of artemisinin on root gravitropic response and root system development in *Arabidopsis thaliana*. *Plant Growth Regul.* 2018, 85, 211–220.
143. Shao, H.; Zhang, Y. Non-target effects on soil microbial parameters of the synthetic pesticide carbendazim with the biopesticides cantharidin and norcantharidin. *Sci. Rep.* 2017, 7, 5521.

Retrieved from <https://encyclopedia.pub/entry/history/show/27359>