ROS-Responsive Cell Harvesting Methods

Subjects: Materials Science, Biomaterials Contributor: Jong-Chul Park

Cell sheet engineering has evolved rapidly over the last few years and as a new approach for cell-based therapy. Cell sheet harvest technology is important for producing viable, transplantable cell sheets and applying them to tissue engineering. To date, various approaches have been reported for harvesting cell sheets by inducing property changes of the culture surface such as wettability, pH, electricity, and magnetism. This section provides a comprehensive introduction to reactive oxygen species (ROS) and the progress of new strategies for applying ROS in cell sheet detachment.

Keywords: reactive oxygen species ; cell detachment ; extracellular matrix ; focal adhesion kinase

1. Definition of ROS

Reactive oxygen species (ROS) are natural byproducts of cellular oxidative metabolism and are involved in the regulation of cell survival, cell death, differentiation, cell signaling, and inflammation-related factor production ^{[1][2]}. Biologically important ROS elements include free radicals, such as singlet oxygen (¹O₂), superoxide (O₂•–), hydroxyl (HO•), hydroperoxyl (HO₂•), carbonate (CO₃•–), peroxyl (RO₂•), alkoxyl (RO•), and carbon dioxide radicals (CO₂•–), and nonradicals, such as hydrogen peroxide (H₂O₂), hypobromous acid (HOBr), hypochlorous acid (HOCI), ozone (O₃), organic peroxides (ROOH), peroxynitrite (ONOO–), peroxynitrate (O₂NOO–), peroxynitrous acid (ONOOH), peroxomonocarbonate (HOOCO₂–), nitric oxide (NO), and hypochlorite (OCI–) ^{[3][4][5]}. Originally, only phagocytic cells were known to be responsible for ROS production in host cell defense mechanisms. Recent studies have shown that ROS play a role in cell signaling, including apoptosis, gene expression, and the activation of cell signaling cascades ^[6]. In particular, ROS acts differently in cells depending on the concentration. Low levels of ROS activate cell signaling pathways to initiate biological processes ^[7]. However, high levels of ROS cause cellular and DNA damage, and activation of cell death processes, such as apoptosis, depending on the severity and duration of exposure.

2. Source of ROS Generation

The main sources of intracellular ROS are mitochondria, the endoplasmic reticulum (ER), peroxisomes, microsomes, and nicotinamide adenine dinucleotide phosphate (NADPH) oxidase (NOX) complexes in cell membranes ^{[2][8][9]}. In particular, mitochondria are the main intrinsic source of ROS production via the mitochondrial electron-transport system ^[10]. Extracellular sources also contribute to ROS generation, such as radiation, pollutants, nanoparticles (NPs), and various drugs, and certain types of other chemical compounds also play a role ^{[11][12][13]}. NPs of metals can induce the generation of the radical reactive superoxide by donating an electron to molecular oxygen; the superoxide then triggers a cascade of radical forming reactions ^[14], as shown in Figure 1.




Figure 1. Mechanisms for the generation of intracellular and extracellular ROS (reactive oxygen species). Extracellular ROS are generated from environmental pollutants, drugs, xenobiotic substances, or radiation. Intracellular ROS are known to be generated through multiple mechanisms inside the cell (metabolic by-products of biological systems).

Photodynamic action (PDA) is a method of inducing the generation of ROS. Integral to PDA are a photosensitizer (PS), a light-absorbing molecule, and a light source with a suitable wavelength. With light irradiation, the PS absorbs the light energy and transfers to an excited state. The excited PS then undergoes a photochemical reaction (PR) with a biological environment to generate ROS, which is called PDA ^[15]. There are two main types of PDA: type I reaction involves electron transfer PR to generate radical and radical anion species, while type II reaction directs PR through energy transfer between oxygen and excited PS, to produce singlet oxygen (Figure 2)^{[16][17][18][19]}.




Figure 2. Schematic illustrations of the process of generating extracellular ROS from a photosensitizer under light irradiation. PS: photosensitizer.

3. Cell Harvesting Methods by Extracellular ROS

Studies have reported that ROS act as mediators of cell adhesion ^[20], and an increase in intracellular ROS levels can lead to cell detachment ^[21]. The Möhwald group^[22] reported the use of light to release fibroblasts cultured on gold nanoparticle-based surfaces. Gold nanoparticles (AuNP) have strong light absorption in the green spectral range. Thus, the authors irradiated the surface with a green laser (532 nm) to produce extracellular ROS by a photochemical mechanism (Figure 3A). The ROS damaged cell membranes at the cell–surface interface, detaching the cells from the substrate. The cells did not immediately detach from the surface, but took up to 24 h to completely detach. One of the advantages of this system is that the surface can be recovered, allowing the cells to reattach in the irradiated areas within 72 h. This property is able to spatially pattern cells, and control the area where the green laser is irradiated to produce cocultured cell sheets that can be reattached by seeding different cell types after the surface has recovered (Figure 3B). This method has been reported to be applicable to the individual detachment of cells from the culture surface, but has not shown the results of detaching the cells in sheet form. Recently, Koo et al. ^[23] reported a new ROS-induced strategy for direct transfer of intact cell sheets to target sites without intermediate harvesting processes based on hematoporphyrin-incorporated polyketone (Hp-PK) film. After green light emitting diode (LED) (510 nm) irradiation for 5–10 min, exogenous ROS generated from the Hp-PK film induced cell sheet detachment and transfer simultaneously. Briefly, this process is carried out by placing a cell cultured film on the target site, irradiating light, and then peeling only the film (Figure 3C). They have successfully applied the detachment of various cell types in a simple way to control the irradiation power and time. In addition, this strategy was applied to the in vivo transplantation onto subcutaneous tissue. Cell sheets can adhere to tissues because of the presence of extracellular matrix (ECM). Therefore, They stacked the cell sheets in multiple layers and transplanted them by the ROS-induced method, and confirmed that the stacked cell sheets were well grafted at the transplant site without cell loss. In particular, multi-layered stem cell sheets accelerated wound healing in full-thickness skin defects ^[24]. The currently reported single cells or cell sheet detachment for a long time ^{[25][26][27]}. In addition, the ligands and magnetic NPs, which are used for cell detachment, are released together with the cells. Thus, there is a limit in harvesting only an intact cell sheet. This system addresses these limitations and demonstrates the possibility of efficient cell transplantation into the lesion site for tissue regeneration and reconstruction, the ultimate goal of cell sheet engineering.

The advantage of the methods using extracellular ROS is the ease of spatio-temporal control of cell detachment. ROS have a short half-life (<40 ns) and can only act close to the site of generation (<20 nm), which is significantly less than cell dimensions ^{[28][29]}. Thus, on a molecular scale, ROS can only act chemically at a small distance from the site of their production. Due to this property, the Möhwald group's research ^[22] mentioned above shows that spatial control is possible by ensuring that the cells attached to the surface of AuNP undergo patterned detachment according to the ring-like profile of the laser beam. In addition, the PS-incorporated film used in Koo's study ^[23] suggested that the occurrence of unintentional cell detachment can be avoided by spatially controlling the generation of ROS and controlling the production time with or without light stimulation.




Figure 3. (A) Schematics of cell detachment by laser irradiation on AuNP-based surfaces. (B) Phase contrast microscopic images showing cell detachment areas according to the laser power. The images were taken 24 h after irradiation (Reprinted with permission from ^[22]. Copyright (2019) American Chemical Society). (C) Schematic illustration of the ROS-induced cell sheet detachment and transfer procedure on Hp-PK films (Reprinted from ^[23] with permission on Elsevier and Copyright Clerance Center.)

4. Current Limitation for Clarifying the Mechanism by ROS

Because cell sheet detachment by extracellular ROS occurs gradually, some studies have reported that this change is due to cell signaling initiated by ROS ^{[30][31]}. The downstream effect of ROS production is the generally reversible oxidation of proteins ^{[20][32]}. Redox-sensitive proteins, which include protein tyrosine phosphatases (PTPs) as the active site cysteine, are the target of specific oxidation by various oxidants ^[33]. Focal adhesion kinase (FAK) is a non-receptor protein tyrosine kinase that plays an important role in signal transduction from integrin-enriched focal adhesion (FA) sites that mediate cellular contact with the extracellular matrix. Multiple protein–protein interaction sites of FAK mediate association with adapters and structural proteins ^{[34][35]}. Chiarugi et al. ^[20] provide evidence that ROS take a role in

integrin signaling. In the role for oxidative species in integrin signaling, ROS generated during cell adhesion induce upregulation of FAK. However, increasing of intracellular ROS up to a threshold level delays cell adhesion to ECM proteins, and results in a negative function of PTP on FA development and cytoskeleton organization. Koo et al. ^[36] previously reported that extracellular ROS increase the amount of intracellular ROS. H_2O_2 can diffuse through specific aquaporins (AQP) in the plasma membrane, and superoxide anion (O_2^-) can penetrate the cell membrane through anion channels (CI⁻ channel-3) to initiate intracellular signal transduction ^{[36][37][38][39]}. There are many ways in which extracellular ROS can be transported into cells, but the exact mechanism of ROS that affects cell detachment is unknown. In contrast, Koo's study using FTIR spectroscopy demonstrates that ROS-induced cell detachment is due to secondary structural changes in proteins adsorbed on the Hp-PK films ^[23]. Thus, they have supposed that these conformational changes are caused by extracellular ROS in ECM proteins present in the cell sheet. However, the type of ROS that affects cell detachment has not been identified.

ROS are difficult to distinguish and quantify from each element by specific assays. These other reactive molecules have properties of overlapping or distinguishing from each other. Some scientists have reported that various types of ROS can be distinguished by specific probes and categorized several probes for each ROS (Table 1) ^{[40][41][42][43]}. There are various ROS probes that can be analyzed by flow cytometry or microscopy ^{[44][45]}. However, most of these are not specific to a particular ROS species, are unstable, and can be affected by other factors distinct from the oxidants. Therefore, when using these probes, it is necessary to carefully interpret the data derived by comparing the various methods.

Probe	Specificity	Advantages	Disadvantages
Fluorescent probes	02* , H ₂ O ₂	Cell permeable, intensity	Products complex, low specificity, interfered by ONOO [—]
		quantifiable, product stable	
Chemiluminescent probes	O2 ^{•—} , ∙OH	Cell permeable	Low selectivity and sensitivity, intermediates not stable
Spectrophotometry methods	O2*—, H ₂ O ₂	Sensitive, fast, single product	Low specificity
Chromatography methods	·OH	Fast, sensitive	Products complex
Electrochemical biosensors	02*	Sensitive, fast	Complex to prepare
Electron spin resonance	ROS, RNS	Specific, sensitive	Expensive
Fluorescent proteins	H ₂ O ₂ , redox status changes	Dynamic, real-time, cell friendly	Slow in reaction, restriction in receptor cells, non-sensitive

Table 1. Methods for the detection of ROS. (Reproduced from ^[43] by permission of The Royal Society of Chemistry).

References

- 1. Rhian M Touyz; Molecular and cellular mechanisms in vascular injury in hypertension: role of angiotensin II editorial r eview. *Current Opinion in Nephrology and Hypertension* **2005**, *14*, 125-131, <u>10.1097/00041552-200503000-00007</u>.
- Cornelius F.H. Mueller; Karine Laude; J. Scott McNally; David G. Harrison; Redox Mechanisms in Blood Vessels. Arteri osclerosis, Thrombosis, and Vascular Biology 2005, 25, 274-278, <u>10.1161/01.atv.0000149143.04821.eb</u>.

- 3. Ahmed Abdal Dayem; Mohammed Kawser Hossain; Soo Lee; Kyeongseok Kim; Subbroto Kumar Saha; Gwang-Mo Ya ng; Hye Yeon Choi; Ssang-Goo Cho; Soo Bin Lee; The Role of Reactive Oxygen Species (ROS) in the Biological Activit ies of Metallic Nanoparticles. *International Journal of Molecular Sciences* **2017**, *18*, 120, <u>10.3390/ijms18010120</u>.
- Augusto, O.; Miyamoto, S.; Pantopoulos, K.; Schipper, H.. Principles of Free Radical Biomedicine; Nova Science Publis hers: Brazil, 2011; pp. 19-42.
- 5. Haohao Wu; Jun-Jie Yin; Wayne G. Wamer; Mingyong Zeng; Y. Martin Lo; Reactive oxygen species-related activities of nano-iron metal and nano-iron oxides. *Journal of Food and Drug Analysis* **2014**, *22*, 86-94, <u>10.1016/j.jfda.2014.01.007</u>.
- J. T. Hancock; R. Desikan; S.J. Neill; Role of reactive oxygen species in cell signalling pathways. *Biochemical Society Transactions* 2001, 29, 345-349, <u>10.1042/bst0290345</u>.
- 7. Michael Schieber; Navdeep S. Chandel; ROS function in redox signaling and oxidative stress.. *Current Biology* **2014**, *2* 4, R453-62, <u>10.1016/j.cub.2014.03.034</u>.
- 8. Dunyaporn Trachootham; Jérôme Alexandre; Peng Huang; Targeting cancer cells by ROS-mediated mechanisms: a ra dical therapeutic approach?. *Nature Reviews Drug Discovery* **2009**, *8*, 579-591, <u>10.1038/nrd2803</u>.
- 9. Victor J. Thannickal; Barry L. Fanburg; Reactive oxygen species in cell signaling. *American Journal of Physiology-Lun g Cellular and Molecular Physiology* **2000**, *279*, L1005-L1028, <u>10.1152/ajplung.2000.279.6.11005</u>.
- 10. Toren Finkel; Signal Transduction by Mitochondrial Oxidants. *Journal of Biological Chemistry* **2012**, *287*, 4434-4440, <u>1</u> <u>0.1074/jbc.R111.271999</u>.
- 11. V Vallyathan; X Shi; The role of oxygen free radicals in occupational and environmental lung diseases. *Environmental Health Perspectives* **1997**, *105*, 165-177, <u>10.1289/ehp.97105s1165</u>.
- 12. James C. Bonner; Lung Fibrotic Responses to Particle Exposure. *Toxicologic Pathology* **2007**, *35*, 148-153, <u>10.1080/01</u> <u>926230601060009</u>.
- 13. Lotte Risom; Peter Møller; Steffen Loft; Oxidative stress-induced DNA damage by particulate air pollution. *Mutation Re search/Fundamental and Molecular Mechanisms of Mutagenesis* **2005**, *592*, 119-137, <u>10.1016/j.mrfmmm.2005.06.012</u>.
- Lucía Z. Flores-López; Heriberto Espinoza-Gómez; Ratnasamy Somanathan; Silver nanoparticles: Electron transfer, re active oxygen species, oxidative stress, beneficial and toxicological effects. Mini review. *Journal of Applied Toxicology* 2 019, 39, 16-26, <u>10.1002/jat.3654</u>.
- 15. Maharajan Sivasubramanian; Yao Chen Chuang; Leu-Wei Lo; Evolution of Nanoparticle-Mediated Photodynamic Thera py: From Superficial to Deep-Seated Cancers. *Molecules* **2019**, *24*, 520, <u>10.3390/molecules24030520</u>.
- 16. Martijn Triesscheijn; Paul Baas; Jan H. M. Schellens; Fiona A. Stewart; Photodynamic Therapy in Oncology. *The Oncol* ogist **2006**, *11*, 1034-1044, <u>10.1634/theoncologist.11-9-1034</u>.
- Suneesh C. Karunakaran; P. S. Saneesh Babu; Bollapalli Madhuri; Betsy Marydasan; Albish K. Paul; Asha S. Nair; K. S ridhar Rao; Alagar Srinivasan; Tavarekere K. Chandrashekar; Ch. Mohan Rao; et al. In Vitro Demonstration of Apoptosi s Mediated Photodynamic Activity and NIR Nucleus Imaging through a Novel Porphyrin. ACS Chemical Biology 2013, 8, 127-132, 10.1021/cb3004622.
- Kai Liu; Xiaomin Liu; Qinghui Zeng; YouLin Zhang; Langping Tu; Tao Liu; Xianggui Kong; Yinghui Wang; Feng Cao; Sa skia A. G. Lambrechts; et al. Covalently Assembled NIR Nanoplatform for Simultaneous Fluorescence Imaging and Ph otodynamic Therapy of Cancer Cells. ACS Nano 2012, 6, 4054-4062, <u>10.1021/nn300436b</u>.
- Ana P. Castano; Tatiana N. Demidova; Michael R. Hamblin; Mechanisms in photodynamic therapy: part two-cellular sig naling, cell metabolism and modes of cell death.. *Photodiagnosis and Photodynamic Therapy* 2005, *2*, 1-23, <u>10.1016/S</u> <u>1572-1000(05)00030-X</u>.
- 20. Paola Chiarugi; Giovambattista Pani; Elisa Giannoni; Letizia Taddei; Renata Colavitti; Giovanni Raugei; Mark Symons; Silvia Borrello; Tommaso Galeotti; Giampietro Ramponi; et al. Reactive oxygen species as essential mediators of cell a dhesion. *The Journal of Cell Biology* **2003**, *161*, 933-944, <u>10.1083/jcb.200211118</u>.
- 21. Heesang Song; Min-Ji Cha; Byeong-Wook Song; Il-Kwon Kim; Woochul Chang; Soyeon Lim; Eun Ju Choi; Onju Ham; Se-Yeon Lee; Namsik Chung; et al. Reactive Oxygen Species Inhibit Adhesion of Mesenchymal Stem Cells Implanted i nto Ischemic Myocardium via Interference of Focal Adhesion Complex. *STEM CELLS* 2010, *28*, 555-563, <u>10.1002/stem.302</u>.
- Tatiana A. Kolesnikova; Dorothee Köhler; Andre G. Skirtach; Helmuth Möhwald; Laser-Induced Cell Detachment, Patter ning, and Regrowth on Gold Nanoparticle Functionalized Surfaces. ACS Nano 2012, 6, 9585-9595, <u>10.1021/nn302891</u>
 <u>U</u>.
- 23. Min-Ah Koo; Mi Hee Lee; Byeong-Ju Kwon; Gyeung Mi Seon; Min Sung Kim; Dohyun Kim; Ki Chang Nam; Jong-Chul Park; Exogenous ROS-induced cell sheet transfer based on hematoporphyrin-polyketone film via a one-step process.

Biomaterials 2018, 161, 47-56, 10.1016/j.biomaterials.2018.01.030.

- 24. Min-Ah Koo; Seung Hee Hong; Mi Hee Lee; Byeong-Ju Kwon; Gyeung Mi Seon; Min Sung Kim; Dohyun Kim; Ki Chang Nam; Jong-Chul Park; Effective stacking and transplantation of stem cell sheets using exogenous ROS-producing film f or accelerated wound healing. *Acta Biomaterialia* **2019**, *95*, 418-426, <u>10.1016/j.actbio.2019.01.019</u>.
- 25. Anice C. Lowen; John Steel; Roles of Humidity and Temperature in Shaping Influenza Seasonality. *Journal of Virology* **2014**, *88*, 7692-7695, <u>10.1128/JVI.03544-13</u>.
- 26. Minxiong Li; Jun Ma; Yanbin Gao; Lei Yang; Cell sheet technology: a promising strategy in regenerative medicine. *Cyto therapy* **2019**, *21*, 3-16, <u>10.1016/j.jcyt.2018.10.013</u>.
- 27. Hai Xiao; Tsai-Kun Li; Jin-Ming Yang; Leroy F. Liu; Acidic pH induces topoisomerase II-mediated DNA damage. *Procee dings of the National Academy of Sciences* **2003**, *100*, 5205-5210, <u>10.1073/pnas.0935978100</u>.
- Kai Han; Qi Lei; Shi-Bo Wang; Jing-Jing Hu; Wen-Xiu Qiu; Jing-Yi Zhu; Wei-Na Yin; Xu Luo; Xian-Zheng Zhang; Dual-S tage-Light-Guided Tumor Inhibition by Mitochondria-Targeted Photodynamic Therapy. *Advanced Functional Materials* 2 015, 25, 2961-2971, <u>10.1002/adfm.201500590</u>.
- C.A. Robertson; D. Hawkins Evans; Heidi Abrahamse; Photodynamic therapy (PDT): A short review on cellular mechan isms and cancer research applications for PDT. *Journal of Photochemistry and Photobiology B: Biology* 2009, 96, 1-8, <u>10.1016/j.jphotobiol.2009.04.001</u>.
- 30. Andre E.X. Brown; Dennis E. Discher; Conformational changes and signaling in cell and matrix physics.. *Current Biolog y* **2009**, *19*, R781-9, <u>10.1016/j.cub.2009.06.054</u>.
- 31. Alexis J Torres; Min Wu; David Holowka; Barbara Baird; Nanobiotechnology and cell biology: micro- and nanofabricate d surfaces to investigate receptor-mediated signaling. *Annual Review of Biophysics* **2008**, *37*, 265-288, <u>10.1146/annur ev.biophys.36.040306.132651</u>.
- 32. Toren Finkel; Reactive Oxygen Species and Signal Transduction. *IUBMB Life* **2001**, *52*, 3-6, <u>10.1080/15216540252774</u> 694.
- 33. Dong Xu; Ilsa I Rovira; Toren Finkel; Oxidants painting the cysteine chapel: redox regulation of PTPs.. *Developmental Cell* **2002**, *2*, 251-252, <u>10.1016/s1534-5807(02)00132-6</u>.
- 34. Cord Brakebusch; Daniel Bouvard; Fabio Stanchi; Takao Sakai; Reinhard Fässler; Integrins in invasive growth. *Journal* of *Clinical Investigation* **2002**, *109*, 999-1006, <u>10.1172/JCI15468</u>.
- 35. Christof R. Hauck; Datsun A. Hsia; David D. Schlaepfer; The Focal Adhesion Kinase--A Regulator of Cell Migration and Invasion. *IUBMB Life* **2002**, 53, 115-119, <u>10.1080/15216540211470</u>.
- 36. Min-Ah Koo; Bong-Jin Kim; Mi Hee Lee; Byeong-Ju Kwon; Min Sung Kim; Gyeung Mi Seon; Dohyun Kim; Ki Chang Na m; Kang-Kyun Wang; Yong-Rok Kim; et al. Controlled Delivery of Extracellular ROS Based on Hematoporphyrin-Incorp orated Polyurethane Film for Enhanced Proliferation of Endothelial Cells. ACS Applied Materials & Interfaces 2016, 8, 2 8448-28457, 10.1021/acsami.6b07628.
- 37. Aron B. Fisher; Redox Signaling Across Cell Membranes. *Antioxidants & Redox Signaling* **2009**, *11*, 1349-1356, <u>10.108</u> <u>9/ars.2008.2378</u>.
- 38. Bryan C Dickinson; Christopher J Chang; Chemistry and biology of reactive oxygen species in signaling or stress responses.. *Nature Chemical Biology* **2011**, 7, 504-11, <u>10.1038/nchembio.607</u>.
- 39. Helmut Sies; Role of Metabolic H2O2Generation. *Journal of Biological Chemistry* **2014**, *289*, 8735-8741, <u>10.1074/jbc.r</u> <u>113.544635</u>.
- Christine C. Winterbourn; The challenges of using fluorescent probes to detect and quantify specific reactive oxygen sp ecies in living cells. *Biochimica et Biophysica Acta (BBA) - General Subjects* 2014, 1840, 730-738, <u>10.1016/j.bbagen.2</u> 013.05.004.
- 41. J.F. Woolley; J. Stanicka; T.G. Cotter; Recent advances in reactive oxygen species measurement in biological systems. *Trends in Biochemical Sciences* **2013**, *38*, 556-565, <u>10.1016/j.tibs.2013.08.009</u>.
- Jacek Zielonka; Joy Joseph; Adam Sikora; Balaraman Kalyanaraman; Real-Time Monitoring of Reactive Oxygen and N itrogen Species in a Multiwell Plate Using the Diagnostic Marker Products of Specific Probes. *Methods in Enzymology* 2013, 526, 145-157, <u>10.1016/b978-0-12-405883-5.00009-0</u>.
- 43. Yinfeng Zhang; Menghong Dai; Zonghui Yuan; Methods for the detection of reactive oxygen species. *Analytical Method* s **2018**, *10*, 4625-4638, <u>10.1039/c8ay01339</u>j.
- 44. Christine Winterbourn; Current methods to study reactive oxygen species Pros and cons. *Biochimica et Biophysica Acta (BBA) General Subjects* **2014**, *1840*, 707, <u>10.1016/j.bbagen.2013.09.021</u>.

45. Michael P. Murphy; Arne Holmgren; Nils-Göran Larsson; Barry Halliwell; Christopher J. Chang; Balaraman Kalyanaram an; Sue Goo Rhee; Paul J. Thornalley; Linda Partridge; David Gems; et al. Unraveling the biological roles of reactive o xygen species.. *Cell Metabolism* **2011**, *13*, 361-366, <u>10.1016/j.cmet.2011.03.010</u>.

Retrieved from https://encyclopedia.pub/entry/history/show/35509