

Aquaporins in Mesenchymal Stem Cells

Subjects: [Cell Biology](#) | [Physiology](#)

Contributor: alessandra pelagalli

Aquaporins (AQPs) are a family of membrane water channel proteins that control osmotically-driven water transport across cell membranes. Recent studies have focused on the assessment of fluid flux regulation in relation to the biological processes that maintain mesenchymal stem cell (MSC) physiology. In particular, AQPs seem to regulate MSC proliferation through rapid regulation of the cell volume. Furthermore, several reports have shown that AQPs play a crucial role in modulating MSC attachment to the extracellular matrix, their spread, and migration. This review aims to describe the recent findings on AQPs role in MSCs physiology taking into account their reflex for potential application in regenerative medicine.

aquaporins

water channel

mesenchymal stem cells

physiology

1. Introduction

Aquaporins (AQPs) are a family of transmembrane proteins that form water channels and work as regulators of intra- and inter-cellular water transport [\[1\]](#). To date, thirteen AQPs that are widely distributed in specific cell types of various tissues have been characterised [\[2\]](#). The major roles of AQPs have been investigated in both physiological and pathological conditions, and the results highlight their involvement in the transfer of water, gases, and small solutes (urea and glycerol), to maintain cell homeostasis [\[3\]](#)[\[4\]](#)[\[5\]](#). These proteins regulate many biological processes through their intrinsic activity including cell proliferation, migration, apoptosis, and mitochondrial metabolism.

In addition, several studies have focused on the involvement of AQPs in intriguing aspects of cell biology and have demonstrated that they are involved in a variety of physiological processes and pathophysiological conditions [\[6\]](#)[\[7\]](#)[\[8\]](#).

Here, we review the current understanding about the roles played by AQPs in mesenchymal stem cell (MSC) functions and highlight their involvement in stem cell proliferation, migration, and differentiation.

Specific features of MSCs rely on their self-renewal ability, low immunogenicity, and the ability to migrate, proliferate, and differentiate in different cell types [\[9\]](#)[\[10\]](#). Notably, the biological activities associated with MSCs migration and proliferation are of particular importance because they are involved in tissue regeneration. Following tissue damage, MSCs are able to mobilise from the tissue of origin and migrate through the peripheral circulation to the injured site, where they proliferate and differentiate, thus facilitating the healing process through the activation of various mechanisms [\[11\]](#).

Such processes require the orchestration of multiple signals induced by mechanical (hemodynamic forces applied to the vessel walls through shear stress, vascular cyclic stretching, and extracellular matrix stiffness) and chemical factors (chemokines, and growth factors), that can act simultaneously. MSCs can migrate through 3D tissue and regulate forces that induce cell deformation and act on physical tissue constraints from the mechano-microenvironment [12].

It has also been demonstrated that the tissue source, growth factors, ageing, the microenvironment, and hormones can influence the MSC proliferation rate. In particular, in vitro and in vivo studies have shown that the tissue source and aging affect the properties of MSCs, including their proliferative capacity, lifespan, and ability to differentiate efficiently [13][14][15]. In addition, Zhu et al. observed that a hypoxic microenvironment can increase the proliferation of placenta-derived MSCs via the mitogen-activated protein kinase (MAPK)/extracellular signal-regulated kinase (ERK) pathway [16]. These aspects, along with the various growth factors, cytokines, exosomes, and microvesicles secreted by MSCs [17][18] should be taken into consideration when designing strategies to enable the efficient use of MSCs for repairing dysfunctional organs.

2. AQP

AQPs are a family of thirteen integral-membrane water channel proteins (AQP0 to AQP12) found in humans, animals, and plants. They can be classified into three main functional subfamilies based on their ability to facilitate transport: AQPs, aquaglyceroporins, and a third family that is comprised of AQPs with uncharacterised functions [19] (Table 1).

AQP0, AQP1, AQP2, AQP4, AQP5, AQP6, and AQP8 are considered as the classical water-specific channel proteins. In contrast, AQP3, AQP7, AQP9, and AQP10, which belong to the aquaglyceroporins sub-family, are characterised by their permeability to water, glycerol, urea, and a few small neutral solutes [20]. AQP11 and AQP12 represent the most distantly related paralogs as they show low amino acid sequence identity with the other AQP family members [21]. These AQPs show non-canonical subcellular localisation and functions.

To date, two functional motifs, namely, Asn-Pro-Ala (NPA) [22][23][24][25] and the aromatic-arginine regions have been identified in the AQP sequence. The NPA region, located in the middle of the channel, is involved in proton exclusion and contributes to the localisation of AQPs to the plasma membrane (Figure 1). The aromatic-arginine region acts as a selectivity filter on the extracellular side of the AQP channel and blocks the entry of molecules larger than water.

AQPs are structurally organised as homotetramers, where each monomer (ranging from 26 to 34 kDa) is primarily composed of six transmembrane domains such that both the amino- and carboxy-termini of the protein lie inside the cell. The highly conserved NPA motifs are located in cytoplasmic loop B and in extracellular loop E [26]. These NPA motifs contribute to a monomeric pore structure that facilitates selective, bi-directional, and single-file transport of water in the classical AQPs and water and glycerol in aquaglyceroporins [27].

AQPs are mainly localised to the plasma membrane, and to a lesser extent in cytosolic compartments, where they can be transported to the plasma membrane in response to hormones and kinase activation [28]. This localisation allows AQPs to regulate distinct processes occurring in different cellular compartments. Studies performed in humans and other animal species have shown that AQPs participate in a wide range of physiological functions, including water/salt homeostasis, exocrine fluid secretion, and epidermal hydration. Moreover, AQPs contribute to pathological conditions such as glaucoma, cancer, epilepsy, obesity in which water and small solute transport may be involved. Of note, AQPs seem to act as modulators of adipocyte biology in obesity by facilitating glycerol release from adipose tissue. Mechanisms associated with cell regulation as well as cell migration and signalling have been linked to AQPs in cancer biology and metastasis development.

Molecular evidence has highlighted the importance of AQPs in facilitating multiple cellular processes such as (1) transepithelial fluid flow regulated by osmotic water transport across cell membranes, (2) cell migration and neuroexcitation, (3) cell proliferation mediated by glycerol transport, (4) adipocyte metabolism, and (5) epidermal water retention [29].

In response to environmental stimuli, AQPs regulate the flow of water and small molecules including glycerol, ammonia and urea in a tissue-specific manner to maintain their homeostasis (Table 2).




Figure 1. Schematic representation of the Asn-Pro-Ala (NPA) motif in the aquaporin (AQP) structure.

Table 1. Classification and permeability characteristics of AQPs.

AQP Classification	Isoform	Permeability						
		H ₂ O	Glycerol	NO	H ₂ O ₂	NH ₃ and/or Ammonia	Urea	Uncertain

	AQP0	+	/	/	/	+	/	/
	AQP1	+	/	+	+	+	/	/
	AQP2	+	/	/	/	/	/	/
AQPs	AQP4	+	/	+	/	/	/	/
	AQP5	+	/	/	+	/	/	/
	AQP6	+	+	/	/	+	+	/
	AQP8	+	+	/	+	+	+	/
	AQP3	+	+	/	+	+	+	/
Aquaglyceroporins	AQP7	+	+	/	/	+	+	/
	AQP9	+	+	/	+	+	+	/
	AQP10	+	+	/	/	+	+	/
	AQP11	/	/	/	/	/	/	+
Unorthodox-AQPs	AQP12	/	/	/	/	/	/	+

Bold is refers to each subgroup of AQPs comprising a number of isoforms. The symbol + indicates that the specific AQP isoform is permeable to the specific molecule indicated.

Table 2. Distribution and principal roles of AQPs in the main body tissues.

Tissue Localisation	Aqp Isoform	Main AQP Roles	Main References
Brain	AQP1, AQP3, AQP4, AQP6	<ul style="list-style-type: none"> - regulation of water homeostasis - control of osmotic pressure for efficient axonal conductance 	[30][31]
Eye	AQP0, AQP1, AQP3, AQP4, AQP5,	<ul style="list-style-type: none"> - water balance maintenance in ocular tissues to ensure transparency in cornea and lens, in corneal wound healing - regulation of tear film osmolarity to produce aqueous humor, and to maintain retinal homeostasis 	[32][33]
Reproductive Tract			
Female	AQP1, AQP3, AQP4, AQP5, AQP7, AQP8, AQP9, AQP10, AQP11	<ul style="list-style-type: none"> - secretive (vagina) and absorptive processes (utero) - maternal-fetal fluid exchange 	[34][35]
Male	AQP1, AQP2, AQP3, AQP6, AQP7, AQP9, AQP11, AQP12	<ul style="list-style-type: none"> - fluid regulation for spermatogenesis, spermatozoa maturation and storage 	[36][37]
Heart	AQP1, AQP3, AQP4, AQP6, AQP7, AQP9, AQP11	<ul style="list-style-type: none"> - contribution to the transcellular water flux across the endothelial membranes - involvement in the calcium signaling machinery at level of the cardiac and skeletal muscle 	[38][39]

Kidney	AQP1, AQP2, AQP3, AQP4, AQP5, AQP6, AQP7, AQP11	- maintain normal urine concentration function, tissue development and substance metabolism	[40] [41]
Intestine			
Small	AQP1, AQP3, AQP4, AQP5, AQP8, AQP9, AQP10	- involvement in fluid absorption and secretion	[42]
Large	AQP1, AQP3, AQP4, AQP5, AQP6, AQP7, AQP8, AQP9, AQP11	-regulation of water absorption	[42]
Liver	AQP1, AQP3, AQP7, AQP8, AQP9, AQP11	- canalicular and ductal bile formation	[43] [44] [45]
Lung	AQP1, AQP3, AQP4, AQP5	- regulation via transcellular pathway of water across the lung microvascular endothelium and epithelia	[46]
Salivary glands	AQP1, AQP3, AQP4, AQP5, AQP8	-saliva secretion process	[47] [48]
Skin	AQP1, AQP3, AQP5, AQP9, AQP10	-hydration, wound healing, and skin epidermis homeostasis	[49]
Stomach	AQP1, AQP3, AQP4, AQP5, AQP7, AQP11	- contribution in secretion of gastric fluid	[50]

Bold indicates the tissue containing different Aqp isoforms.

Water and fluid transport across the cell membrane are an important prerequisite for maintaining cell homeostasis and volume, which are factors that regulate several cell functions. Hoffmann and co-workers showed the importance of AQPs in the biological processes associated with changes in cell volume such as migration, inflammation, proliferation and cell death [51].

In particular, it has been observed that several factors (osmolality, modifications in the transmembrane ion gradients that generate an osmotic imbalance) that regulate cell volume through AQPs also influence processes that affect cell proliferation [52][53][54][55][56].

Of note, many findings support the hypothesis that AQP expression correlates with the different phases of the cell cycle [57][58][59][60][61]. The overexpression of AQP1 and AQP3 has been shown to affect the expression of essential checkpoint proteins, such as cyclins, and modify the levels of transcription factors and cytokines [62][63]. Delporte et al. [57] showed the involvement of AQP1 in the cell cycle regulation of an epithelial cell line. High levels of AQP1 mRNA and protein expression were found during the G₀/G₁ phase, whereas a significant decrease in mRNA and protein expression was observed in the S and G₂/M phases. Moreover, AQP2 expression has been shown to accelerate the proliferation and cell cycle progression of the renal cortical collecting-duct cells by decreasing the transit time through the S and G₂/M phases, possibly by increasing the cell volume [60].

Additionally, the altered function of AQPs is often correlated with pathological conditions. Nephrogenic diabetes insipidus and congenital cataracts are genetic diseases caused by loss-of-function mutations in AQPs [64][65]. Pathogenic autoantibodies against astrocyte specific AQP4 contribute to the development of the neuroinflammatory demyelinating disease, neuromyelitis optica [66]. Recent studies have shown an association between AQP polymorphisms and diseases such as cognition-related disorders, Alzheimer's [67], and mesothelioma [68]. While the functional link between AQPs and these diseases needs to be investigated further, these data suggest that AQPs can be used as potential biomarkers for severe and disabling diseases. Moreover, there is a great interest in the development of therapeutic strategies using small-molecule modulators with the ability to target AQP function.

3. Physiological Roles of AQPs in Driving MSC Function

Research performed over the last decade has found that AQPs are expressed in several stem cell types, which suggests their involvement in a variety of physiological processes [69][70]. However, their exact functional role in MSCs remains to be completely clarified, albeit some aspects of AQPs role in regulating MSC behaviour in certain disorders (acute lung injury, hepatocarcinoma) have been studied using animal models of disease. These studies have facilitated a better understanding of the molecular pathways underlying MSC involvement in these diseases [71][72][73].

The presence of AQPs in the apical membranes of different MSCs, their co-localisation with other systems such as the Na⁺/K⁺ ATP pump, and the effect of AQP inhibitors including HgCl₂ and TEA suggest that AQPs may regulate the migration and differentiation of MSCs [69]. The involvement of AQPs in various aspects of stem cell biology is related to their capacity to regulate the flux of fluids between the outer and inner cellular compartments to maintain

homeostasis. Interestingly, the control of stem cell volume is an important prerequisite for the regulation of various stem cell properties, including proliferation, migration and differentiation.

Experiments using human BM-MSCs demonstrated that the attachment, spread, and migration of stem cells are accompanied by water efflux and cell volume reduction, which in turn, correlate with the strength of the attachment [74]. The introduction of the AQP1 gene increases the migration ability of MSCs. These results have been confirmed by in vivo experiments showing that the injection of MSCs expressing AQP1 in a rat tibia fracture model enhances bone healing compared to the injection of non-transfected MSCs. This study suggests that the AQP1-mediated improvement in MSC migration likely occurs through the modulation of β -catenin and FAK expression [75] (Table 3). In addition, our recent findings highlight the crucial role played by AQP1 along with C-X-C chemokine receptor type 4 (CXCR4) in regulating ovine MSC (oMSC) migration through the activation of the serine/threonine protein kinase (Akt) and extracellular signal-regulated kinase (Erk) intracellular signal pathways [76] (Table 3).

Table 3. Involvement of AQPs in different aspects of stem cell biology in health and diseases.

Stem Cell Type	Aqp Isoform	AQP Role	Reference
MSCs	AQP1	-Migration	[75][77]
	AQP1, AQP3, AQP4, AQP7	- Differentiation	[77][78]
progenitor MSC	AQP3	- Differentiation	[77][79]
glioblastoma stem-cell	AQP4, AQP9	- AQP9 involvement in the tumorigenicity process	[80]
NSCs	AQP4	- Proliferation, migration and differentiation	[81][82]
	AQP8	- Mitochondrial volume regulation during NSC differentiation	[83]
	AQP9	- Differentiation	[81]

EPSCs	AQP5	-	Regulation of the balance between proliferation and differentiation	[70]
		-	Proliferation and invasion	
LCSCs	AQP3	-	Stemness, differentiation and apoptosis	[84][85]
		-	Possible role in inducing CSCs death	[86]
Lung CSCs	AQP4	-	Migration	[87]
		-	Migration	[87]

Accumulating evidence has shown that AQP s are involved in the function of the brain and central nervous system (CNS) by either maintaining water homeostasis or regulating cerebrospinal fluid secretion and absorption, as well as fluid transport across neutrophils, cell volume regulation, and central osmo-reception [88]. Among the AQP s, AQP1 [89], AQP4 [90][91][92], and AQP9 [93][94] play a central role in CNS-related diseases including cerebral oedema, brain tumours and epilepsy.

More recently, further studies have demonstrated that alteration in AQP s expression is the main cause of water balance dysregulation in the CNS at both the cellular and subcellular levels, and it is responsible for the pathogenetic profile of different diseases such as focal oedemas, brain tumours, brain ischaemia, and traumatic injuries [95][96][97][98]. The brain water content and swelling of the astrocyte foot have been found to be significantly reduced in a brain oedema mouse model that lacks AQP4 [99].

AQP s seem to play a specific role during neurogenesis as neural stem cells (NSCs) are able to move considerable amounts of water across the cell, thus rapidly changing the cell volume. La Porta et al. [83] (Table 3) showed that AQP8 (localised in mitochondria) regulates water balance in adult NSCs (ANSCs) by mediating the osmotic movement of water between the cytoplasm and the mitochondrial compartment. Interestingly, in vivo studies carried out on rat models have reported similar localisation of AQP8 in other cells of the brain, which suggests a possible role of this mitochondrial isoform in many physiological functions including metabolism and apoptosis, as well as in the pathogenesis of neurological diseases such as Parkinson's [100]. Cavazzin et al. used molecular and phenotypical characterisation to show that AQP4 and AQP9 expression is important in the differentiation of murine ANSCs present in the subventricular zone (SVZ). In particular, immunohistochemistry analysis demonstrated that

ANSC-derived glial cells express low levels of AQP4 and display high levels of glial fibrillary acidic protein (GFAP), when astrocytes express high levels of AQP4 and low levels of GFAP. In contrast, while the occurrence of AQP9 is limited to a few cells, those cells that express AQP4 and AQP9 also co-express GFAP. These differences in the subcellular localisation of AQPs among the various ANSC-derived glial cells have led to the hypothesis that the physiological activity of AQPs is cell-specific (Table 3). Li et al. [101] showed that AQP4 is involved in NSC regulation and co-localises with gamma-aminobutyric acid A receptors (GABAARs) in the subependymal zone (SEZ). This result is intriguing considering that the activation of GABAARs induces hyperpolarisation and osmotic swelling in precursor cells, and thus promotes surface expression of the epidermal growth factor receptor and cell cycle entry. Other studies have investigated the role played by AQP4 in the proliferation, migration, and differentiation of ANSCs in vitro using AQP4-knockout mice. Evaluation of connexin 43 and Cav.1.2 expression during the proliferation or differentiation of ANSCs showed that AQP4 in knockout mice causes significant downregulation of the expression of these proteins, which highlights the involvement of AQP4 in the self-renewal, migration and proliferation of ANSCs. In addition, the authors hypothesised that AQP4 regulates the migration of ANSCs through a multi-step process involving: (1) Ca^{2+} influx; (2) actin depolymerisation; (3) an increase in cytoplasmic osmolality; (4) cell membrane expansion with subsequent protrusion formation, and (5) actin repolymerisation to stabilise the emerging protrusion. Taken together, these findings suggest that AQP4 can regulate the fundamental properties of ANSCs through the Ca^{2+} -related signalling pathway (Table 3).

The characterisation of the neural differentiation process of (Ad)-MSCs demonstrated the involvement of AQPs in this process and suggested their role in helping the cells to achieve rapid regulation of the volume during differentiation. In particular, immunohistochemistry, Western blotting, and RT-PCR-based data showed that differentiated neuronal cells arising from sources other than Ad-MSCs, which express AQP1, also express AQP4, and AQP7 based on the cell type, thus suggesting a correlation between neural differentiation and the AQP expression profile during adult neurogenesis (Table 3). Ma and co-workers [102] underlined the importance of AQP5 in modulation of the BM-MSCs differentiation using a mouse model of bone fracture. Accordingly, they found that a lack of AQP5 significantly increases the levels of adipogenic, osteogenic, and chondrogenic differentiation markers in mutant BM-MSCs. Notably, BM-MSCs derived from knockout mice treated with the apoptotic drug, paclitaxel displayed an improvement in the bone healing process as well as a lower apoptosis rate, thus suggesting a modulatory role of AQP5 in slowing down BM-MSC differentiation.

In a previous study, it was reported that AQP1 and AQP3 expression is modulated during human MSC differentiation into chondrocytes. These studies showed that during MSC differentiation, AQP1 and AQP3 expression increases substantially and is associated with high concentrations of collagen type II, aggrecan, and lubricin, thus demonstrating the relationship between these channel proteins and chondrocyte- ECM adhesion and migration (Table 3).

Recently, Chen et al. [103] showed that AQP1 is expressed in rat tendon stem/progenitor cells (TSPCs) and that this expression is regulated during TSPC senescence through the JAK-STAT pathway. Similarly, Zhou J. et al. found that AQP5 expression in human epidermal stem cells (EPSCs) decreases with skin ageing, suggesting that this channel protein has a critical role in regulating the balance between proliferation and differentiation (Table 3).

AQPs are also regarded as key modulators of cancer stem cells (CSCs). Recent studies have shown that high levels of AQP expression in cancer cells and CSCs correlate with metastasis and resistance to therapies [104]. It has been reported that AQP3 affects the Wnt/GSK-3 β /β-catenin pathway in liver CSCs (LCSCs), and modulates their stemness, differentiation, and apoptosis. Other studies have identified AQP4 as the most abundant AQP in nasopharyngeal and lung CSCs and have shown its involvement in the regulation of large volumetric changes related to an increase in death rate [105]. Using a novel electro-osmotic microfluidic system that controls cell osmolarity gradients, AQP4 expression has been shown to be directly correlated with the speed of cell migration (Table 3). In addition, when the expression of AQP4 is suppressed via siRNA, the CSC migration capability as well as the expression of stemness biomarkers such as Sox-2 or Oct-4 are significantly reduced. Other studies have shown that AQP0–AQP12 are highly expressed in human glioblastoma stem-like cell lines and primary tumours [106][107]. Fosdal et al. have described the involvement of AQP1, AQP4, and AQP9 in CSC function in glioblastoma and observed an upregulation of AQP9 levels and a downregulation of AQP1 and AQP4 levels (Table 3). These data indicate the specific role of AQP9 in facilitating CSC migration towards the surrounding normal tissues.

References

1. Benga, G. Water channel proteins (later called aquaporins) and relatives: Past, present, and future. *IUBMB Life* 2009, **61**, 112–133.
2. Benga, G. On the definition, nomenclature and classification of water channel proteins (aquaporins and relatives). *Mol. Aspects Med.* 2012, **33**, 514–517.
3. Day, R.E.; Kitchen, P.; Owen, D.S.; Bland, C.; Marshall, L.; Conner, A.C.; Bill, R.M.; Conner, M.T. Human aquaporins: Regulators of transcellular water flow. *Biochim. Biophys. Acta* 2014, **1840**, 1492–1506.
4. Kitchen, P.; Day, R.E.; Salman, M.M.; Conner, M.T.; Bill, R.M.; Conner, A.C. Beyond water homeostasis: Diverse functional roles of mammalian aquaporins. *Biochim. Biophys. Acta* 2015, **1850**, 2410–2421.
5. Pelagalli, A.; Squillaciotti, C.; Mirabella, N.; Meli, R. Aquaporins in health and disease: An overview focusing on the gut of different species. *Int. J. Mol. Sci.* 2016, **17**, 1213.
6. Verkman, A. Aquaporins at a glance. *J. Cell. Sci.* 2011, **124**, 2107–2112.
7. Login, F.H.; Jensen, H.H.; Pedersen, G.A.; Koffman, J.S.; Kwon, T.H.; Parsons, M.; Nejsum, L.N. Aquaporins differentially regulate cell-cell adhesion in MDCK cells. *FASEB J.* 2019, **33**, 6980–6994.
8. Elkhider, A.; Wang, B.; Ouyang, X.; Al-Azab, M.; Walana, W.; Sun, X.; Li, H.; Tang, Y.; Wei, J.; Li, X. Aquaporin 5 promotes tumor migration and angiogenesis in non-small cell lung cancer cell line H1299. *Oncol. Lett.* 2020, **19**, 1665–1672.

9. Pittenger, M.F.; Mackay, A.M.; Beck, S.C.; Jaiswal, R.K.; Douglas, R.; Mosca, J.D.; Moorman, M.A.; Simonetti, D.W.; Craig, S.; Marshak, D.R. Multilineage potential of adult human mesenchymal stem cells. *Science* 1999, 284, 143–147.

10. Abdelrazik, H.; Giordano, E.; Barbanti Brodano, G.; Griffoni, C.; De Falco, E.; Pelagalli, A. Substantial overview on Mesenchymal Stem Cell biological and physical properties as an opportunity in translational medicine. *Int. J. Mol. Sci.* 2019, 20, 5386.

11. Fu, X.; Liu, G.; Halim, A.; Ju, Y.; Luo, Q.; Song, G. Mesenchymal stem cell migration and tissue repair. *Cells* 2019, 8, 784.

12. Wolf, K.; Te lindert, M.; Krause, M.; Alexander, S.; Te Riet, J.; Willis, A.L.; Hoffman, R.M.; Figdor, C.G.; Weiss, S.J.; Friedl, P. Physical limits of cell migration: Control by ECM space and nuclear deformation and tuning by proteolysis and traction force. *J. Cell Biol.* 2013, 201, 1069–1084.

13. Donders, R.; Bogie, J.F.J.; Ravanidis, S.; Gervois, P.; Vanheusden, M.; Marée, R.; Schryinemakers, M.; Smeets, H.J.M.; Pinxteren, J.; Gijbels, K.; et al. Human Wharton's jelly-derived stem cells display a distinct immunomodulatory and proregenerative transcriptional signature compared to bone marrow-derived stem cells. *Stem Cells Develop.*, 2018, 27, 65–84.

14. Beeravolu, N.; McKee, C.; Alamri, A.; Mikhael, S.; Brown, C.; Perez-Cruet, M.; Rasul Chaudhry, G. Isolation and characterization of mesenchymal stromal cells from human umbilical cord and fetal placenta. *J. Vis. Exp.* 2017, 122, 55224.

15. Ganguly, P.; El-Jawhari, J.J.; Burska, A.N.; Ponchel, F.; Giannoudis, P.V.; Jones, E.A. The analysis of in vivo aging in human bone marrow mesenchymal stromal cells using colony-forming unit-fibroblast assay and the cd45(low)cd271(+) phenotype. *Stem Cells Int.* 2019, 5197983.

16. Zhu, C.; Yu, J.; Pan, Q.; Yang, J.; Hao, G.; Wang, Y.; Li, L.; Cao, H. Hypoxia-inducible factor-2 promotes the proliferation of human placenta-derived mesenchymal stem cells through the MAPK/ERK signaling pathway. *Sci. Rep.* 2016, 6, 35489.

17. Fontaine, M.J.; Shih, H.; Schäfer, R.; Pittenger, M.F. Unraveling the mesenchymal stromal cells' paracrine immunomodulatory effects. *Transfus. Med. Rev.* 2016, 30, 37–43.

18. Konala, V.B.R.; Mamidi, M.K.; Bhonde, R.; Das, A.K.; Pochampally, R.; Pal, R. The current landscape of the mesenchymal stromal cell secretome: A new paradigm for cell-free regeneration. *Cytotherapy*, 2016, 18, 13–24.

19. Verkman, A.S.; Mitra, A.K. Structure and function of aquaporin water channels. *Am. J. Physiol. Renal Physiol.* 2000, 278, F13–F28.

20. Rojek, A.; Praetorius, J.; Frokiaer, J.; Nielsen, S.; Fenton, R.A. A current view of the mammalian aquaglyceroporins. *Annu. Rev. Physiol.*, 2008, 70, 301–327.

21. Ishibashi, K.; Hara, S.; Kondo, S. Aquaporin water channels in mammals. *Clin. Exp. Nephrol.* 2009, 13, 107–117.

22. Murata, K.; Mitsuoka, K.; Hirai, T.; Walz, T.; Agre, P.; Heymann, J.B.; Engel, A.; Fujiyoshi, Y. Structural determinants of water permeation through aquaporin-1. *Nature* 2000, 407, 599–605.

23. Sui, H.; Han, B.-G.; Lee, J.K.; Walian, P.; Jap, B.K. Structural basis of water-specific transport through the AQP1 water channel. *Nature* 2001, 414, 872–878.

24. Hub, J.S.; de Groot, B.L. Mechanism of selectivity in aquaporins and aquaglyceroporins. *Proc. Natl. Acad. Sci. USA* 2008, 105, 1198–1203.

25. Kreida, S.; Törnroth-Horsefield, S. Structural insights into aquaporin selectivity and regulation. *Curr. Opin. Struct. Biol.* 2015, 33, 126–134.

26. Jung, J.S.; Preston, G.M.; Smith, B.L.; Guggino, W.B.; Agre, P. Molecular structure of the water channel through aquaporin CHIP. The hourglass model. *J. Biol. Chem.* 1994, 269, 14648–14654.

27. Jensen, M.Ø.; Tajkhorshid, E.; Schulten, K. The mechanism of glycerol conduction in aquaglyceroporins. *Structure* 2001, 9, 1083–1093.

28. Conner, A.C.; Bill, R.M.; Conner, M.T. An emerging consensus on aquaporin translocation as a regulatory mechanism. *Mol. Membr. Biol.* 2013, 30, 1–12.

29. Verkman, A.S. Aquaporins in Clinical Medicine. *Annu. Rev. Med.* 2012, 63, 303–316.

30. Dasdelen, D.; Mogulkoc, R.; Baltaci, A.K. Aquaporins and roles in brain health and brain injury. *Mini Rev. Med. Chem.* 2020, 20, 498–512.

31. Badaut, J.; Lasbennes, F.; Magistretti, P.J.; Regli, L. Aquaporins in brain: Distribution, physiology, and pathophysiology. *J. Cereb. Blood Flow Metab.* 2002, 22, 367–378.

32. Schey, K.L.; Wang, Z.; Wenke, J.L.; Qi, Y. Aquaporins in the eye: Expression, function, and roles in ocular disease. *Biochim. Biophys. Acta* 2014, 1840, 1513–1523.

33. Lamagna, B.; Ciaramella, P.; Lamagna, F.; Di Loria, A.; Brunetti, A.; Pelagalli, A. Aquaporin 1 (AQP1) expression in healthy dog tears. *Animals* 2020, 10, 820.

34. Huang, H.F.; He, R.H.; Sun, C.C.; Zhang, Y.; Meng, Q.X.; Ma, Y.Y. Function of aquaporins in female and male reproductive systems. *Hum Reprod. Update* 2006, 12, 785–795.

35. Zhu, C.; Jiang, Z.; Bazer, F.W.; Johnson, G.A.; Burghardt, R.C.; Wu, G. Aquaporins in the female reproductive system of mammals. *Front. Biosci.*, 2015, 20, 838–871.

36. Morató, M.Y.R.; Rodríguez-Gil, J.E.; Bonet, S.; Prieto-Martínez, N. Aquaporins in the male reproductive tract and sperm: Functional implications and cryobiology. *Reprod. Dom. Anim.* 2017, 52, 12–27.

37. Carrageta, D.F.; Bernardino, R.L.; Soveral, G.; Calamita, G.; Alves, M.G.; Oliveira, P.F. Aquaporins and male (in)fertility: Expression and role throughout the male reproductive tract. *Arch. Biochem. Biophys.* 2020, 679, 108222.

38. Rutkovskiy, A.; Valen, G.; Vaage, J. Cardiac aquaporins. *Basic Res. Cardiol.* 2013, 108, 393.

39. Verkerk, A.O.; Lodder, E.M.; Wilders, R. Aquaporin Channels in the Heart—Physiology and Pathophysiology. *Int. J. Mol. Sci.* 2019, 20, 2039.

40. Noda, Y.; Sohara, E.; Ohta, E.; Sasaki, S. Aquaporins in kidney pathophysiology. *Nat. Rev. Nephrol.* 2010, 6, 168–178.

41. Li, Y.; Wang, W.; Jiang, T.; Yang, B. Aquaporins in Urinary System. *Adv. Exp. Med. Biol.* 2017, 969, 131–148.

42. Laforenza, U. Water channel proteins in the gastrointestinal tract. *Mol. Aspects Med.* 2012, 33, 642–650.

43. Huebert, R.C.; Splinter, P.L.; García, F.; Marinelli, R.A.; LaRusso, N.F. Expression and localization of aquaporin water channels in rat hepatocytes. Evidence for a role in canalicular bile secretion. *J. Biol. Chem.* 2002, 277, 22710–22717.

44. Marinelli, R.A.; Gradilone, S.A.; Carreras, F.I.; Calamita, G.; Lehmann, G.L. Liver aquaporins: Significance in canalicular and ductal bile formation. *Ann. Hepatol.* 2004, 3, 130–136.

45. Portincasa, P.; Calamita, G. Water channel proteins in bile formation and flow in health and disease: When immiscible becomes miscible. *Mol. Aspects Med.* 2012, 33, 651–664.

46. Wittekindt, O.H.; Dietl, P. Aquaporins in the lung. *Pflügers Archiv Eur. J. Physiol.* 2019, 471, 519–532.

47. Delporte, C.; Steinfeld, S. Distribution and roles of aquaporins in salivary glands. *Biochim. Biophys. Acta* 2006, 1758, 1061–1070.

48. D'Agostino, C.; Elkashty, O.A.; Chivasso, C.; Perret, J.; Tran, S.D.; Delporte, C. Insight into salivary gland aquaporins. *Cells* 2020, 9, 1547.

49. Boury-Jamot, M.; Daraspe, J.; Bonté, F.; Perrier, E.; Schnebert, S.; Dumas, M.; Verbavatz, J.M. Skin aquaporins: Function in hydration, wound healing, and skin epidermis homeostasis. *Handb. Exp. Pharmacol.* 2009, 190, 205–217.

50. Zhu, C.; Chen, Z.; Jiang, Z. Expression, distribution and role of aquaporin water channels in human and animal stomach and intestines. *Int. J. Mol. Sci.* 2016, 17, 1399.

51. Hoffmann, E.K.; Lambert, I.H.; Pedersen, S.F. Physiology of cell volume regulation in vertebrates. *Physiol. Rev.* 2009, 89, 193–277.

52. Meyer, M.; Maly, K.; Uberall, F.; Hoflacher, J.; Grunicke, H. Stimulation of K⁺ transport systems by Ha-ras. *J. Biol. Chem.* 1991, 266, 8230–8235.

53. Lang, F.; Ritter, M.; Gamper, N.; Huber, S.; Fillon, S.; Tanneur, V.; Lepple-Wienhues, A.; Szabo, I.; Gulbins, E. Cell volume in the regulation of cell proliferation and apoptotic cell death. *Cell Physiol. Biochem.* 2000, 10, 417–428.

54. Lang, F.; Föller, M.; Lang, K.S.; Lang, P.A.; Ritter, M.; Gulbins, E.; Vereninov, A.; Huber, S.M. Ion channels in cell proliferation and apoptotic cell death. *J. Membrane Biol.* 2005, 205, 147–157.

55. Rouzaire-Dubois, B.; Dubois, J.M. K⁺ channel block-induced mammalian neuroblastoma cell swelling: A possible mechanism to influence proliferation. *J. Physiol.* 1998, 510, 93–102.

56. Kunzelmann, K. Ion channels and cancer. *J. Membrane Biol.* 2005, 205, 159–173.

57. Delporte, C.; Chen, Z.J.; Baum, B.J. Aquaporin 1 Expression during the Cell Cycle in A5 Cells. *Biochem. Biophys. Res. Commun.* 1996, 228, 223–228.

58. Yoneda, K.; Yamamoto, N.; Asai, K.; Sobue, K.; Fujita, Y.; Fujita, M.; Mase, M.; Yamada, K.; Nakanishi, M.; Tada, T.; Miura, Y.; Kato, T. Regulation of aquaporin in astrocytes. *Mol. Brain Res.* 2001, 89, 94–102.

59. Rivarola, V.; Flamenco, P.; Melamud, L.; Galizia, L.; Ford, P.; Capurro, C. Adaptation to alkalosis induces cell cycle delay and apoptosis in cortical collecting duct cells: Role of aquaporin-2. *J. Cell Physiol.* 2010, 224, 405–413.

60. Di Giusto, G.; Flamenco, P.; Rivarola, V.; Fernández, J.; Melamud, L.; Ford, P.; Capurro, C. Aquaporin 2-increased renal cell proliferation is associated with cell volume regulation. *J. Cell Biochem.* 2012, 113, 3721–3729.

61. Shimizu, H.; Shiozaki, A.; Ichikawa, D.; Fujiwara, H.; Konishi, H.; Ishii, H.; Komatsu, S.; Kubota, T.; Okamoto, K.; Kishimoto, M.; et al. The expression and role of Aquaporin 5 in esophageal squamous cell carcinoma. *J. Gastroenterol.* 2014, 49, 655–666.

62. Galán-Cobo, A.; Ramírez-Lorca, R.; Serna, A.; Echevarría, M. Overexpression of AQP3 modifies the cell cycle and the proliferation rate of mammalian cells in culture. *PLoS ONE* 2015, 10, e0137692.

63. Galán-Cobo, A.; Ramírez-Lorca, R.; Toledo-Aral, J.J.; Echevarría, M. Aquaporin-1 plays important role in proliferation by affecting cell cycle progression. *J. Cell Physiol.* 2016, 231, 243–256.

64. Bichet, D.G. Genetics and diagnosis of central diabetes insipidus. *Ann. Endocrinol.* 2012, 73, 117–127.

65. Varadaraj, K.; Kumari, S.S.; Patil, R.; Wax, M.B.; Mathias, R.T. Functional characterization of a human aquaporin 0 mutation that leads to a congenital dominant lens cataract. *Exp Eye Res.* 2008, 87, 9–21.

66. Papadopoulos, M.C.; Verkman, A.S. Aquaporin 4 and neuromyelitis optica. *Lancet Neurol.* 2012, 11, 535–544.

67. Burfeind, K.G.; Murchison, C.F.; Westaway, S.K.; Simon, M.J.; Erten-Lyons, D.; Kaye, J.A.; Quinn, J.F.; Iliff, J.J. The effects of noncoding aquaporin-4 single-nucleotide polymorphisms on cognition and functional progression of Alzheimer's disease. *Alzheimers Dement.* 2017, 3, 348–359.

68. Senk, B.; Goricar, K.; Kovac, V.; Dolzan, V.; Franko, A. Genetic polymorphisms in aquaporin 1 as risk factors for malignant mesothelioma and biomarkers of response to cisplatin treatment. *Radiol. Oncol.* 2019, 53, 96–104.

69. Juuti-Uusitalo, K.; Delporte, C.; Grégoire, F.; Perret, J.; Huhtala, H.; Savolainen, V.; Nymark, S.; Hyttinen, J.; Uusitalo, H.; Willermain, F.; et al. Aquaporin expression and function in human pluripotent stem cell–derived retinal pigmented epithelial cells. *Ophthalmol. Vis. Sci.* 2013, 54, 3510–3519.

70. Zhou, J.; Dong, Y.; Liu, J.; Ren, J.; Wu, J.; Zhu, N. AQP5 regulates the proliferation and differentiation of epidermal stem cells in skin aging. *Braz. J. Med. Biol. Res.* 2020, 53, e10009.

71. Gao, P.; Yang, J.; Gao, X.; Xu, D.; Niu, D.; Li, J.; Wen, Q. Salvianolic acid B improves bone marrow-derived mesenchymal stem cell differentiation into alveolar epithelial cells type I via Wnt signaling. *Mol. Med. Rep.* 2015, 12, 1971–1976.

72. Pelagalli, A.; Nardelli, A.; Fontanella, R.; Zannetti, A. Inhibition of AQP1 hampers osteosarcoma and hepatocellular carcinoma progression mediated by bone marrow-derived mesenchymal stem cells. *Int. J. Mol. Sci.* 2016; 17, 1102.

73. Gouda, Z.A.; Ali Khalifa, M.E.; Shalaby, S.M.; Hussein, S. Mechanistic effect of human umbilical cord blood derived mesenchymal stem cells on the submandibular salivary gland in ovariectomized rats. *Biochem. Cell Biol.* 2018, 96, 57–67.

74. Beijer, N.R.M.; Nauryzgaliyeva, Z.M.; Arteaga, E.M.; Pieuchot, L.; Anselme, K.; van de Peppel, J.; Vasilevich, A.S.; Groen, N.; Roumans, N.; Hebel, D.G.A.J.; et al. Dynamic adaptation of mesenchymal stem cell physiology upon exposure to surface micropatterns. *Sci. Rep.* 2019, 9, 9099.

75. Meng, F.; Rui, Y.; Xu, L.; Wan, C.; Jiang, X.; Li, G. Aqp1 enhances migration of bone marrow mesenchymal stem cells through regulation of FAK and β -catenin. *Stem Cells Dev.* 2014, 23, 66–75.

76. Pelagalli, A.; Nardelli, A.; Lucarelli, E.; Zannetti, A.; Brunetti, A. Autocrine signals increase ovine mesenchymal stem cells migration through Aquaporin-1 and CXCR4 overexpression. *J. Cell Physiol.* 2018, 233, 6241–6249.

77. Graziano, A.C.E.; Avola, R.; Pannuzzo, G.; Cardile, A. Aquaporin 1 and 3 modification as a result of chondrogenic differentiation of human mesenchymal stem cell. *J. Cell Physiol.* 2018, 233, 2279–

2291.

78. Avola, R.; Graziano, A.C.E.; Pannuzzo, G.; Cardile, V. Human mesenchymal stem cells from adipose tissue differentiated into neuronal or glial phenotype express different aquaporins. *Mol. Neurobiol.* 2017, 54, 8308–8320.

79. Avril-Delplanque, A.; Casal, I.; Castillon, N.; Hinnrasky, J.; Puchelle, E.; Péault, B. Aquaporin-3 expression in human fetal airway epithelial progenitor cells. *Stem Cells* 2005, 23, 992–1001.

80. Fosdal, G.; Vik-Mo, E.O.; Sandberg, C.; Varghese, M.; Kaarbø, M.; Telmo, E.; Langmoen, I.A.; Murrell, W. Aqp 9 and brain tumour stem cells. *Sci. World J.* 2012, 915176.

81. Cavazzin, C.; Ferrari, D.; Facchetti, F.; Russignan, A.; Vescovi, A.L.; La Porta, C.A.M.; Gritti, A. Unique expression and localization of aquaporin- 4 and aquaporin-9 in murine and human neural stem cells and in their glial progeny. *Glia* 2006, 53, 167–181.

82. Kong, H.; Fan, Y.; Xie, J.; Ding, J.; Sha, L.; Shi, X.; Sun, X.; Hu, G. AQP4 Knockout impairs proliferation, migration and neuronal differentiation of adult neural stem cells. *J. Cell Sci.* 2008, 121, 4029–4036.

83. La Porta, C.A.M.; Gena, P.; Gritti, A.; Fascio, U.; Svelto, M.; Calamita, G. Adult murine CNS stem cells express aquaporin channels. *Biol. Cell* 2006, 98, 89–94.

84. Wang, Y.; Wu, G.; Fu, X.; Xu, S.; Wang, T.; Zhang, Q.; Yang, Y. Aquaporin 3 maintains the stemness of CD133+ hepatocellular carcinoma cells by activating STAT3. *Cell Death Dis.* 2019, 10, 465.

85. Liu, C.; Liu, L.; Zhang, Y.; Jing, H. Molecular mechanism of aquaporin (AQP3) in regulating differentiation and apoptosis of lung cancer stem cells through Wnt/GSK-3 β /β-Catenin pathway. *J BUON* 2020, 25, 828–834.

86. Zheng, X.; Li, C.; Yu, K.; Shi, S.; Chen, H.; Qian, Y.; Mei, Z. Aquaporin-9, mediated by IGF2, suppresses liver cancer stem cell properties via augmenting ros/β-catenin/foxo3a signaling. *Mol. Cancer Res.* 2020, doi:10.1158/1541-7786.MCR-19-1180.

87. Hui, T.H.; Cho, W.C.; Fong, H.W.; Yu, M.; Kwan, K.W.; Ngan, K.C.; Wong, K.H.; Tan, Y.; Yao, S.; Jiang, H.; et al. An electro-osmotic microfluidic system to characterize cancer cell migration under confinement. *J. R. Soc. Interface* 2019, 16, 20190062.

88. Pasantes-Morales, H.; Cruz-Rangel, S. Brain volume regulation: Osmolytes and aquaporin perspectives. *Neuroscience* 2010, 168, 871–884.

89. Nielsen, S.; Smith, B.L.; Christensen, E.I.; Agre, P. Distribution of the aquaporin CHIP in secretory and resorptive epithelia and capillary endothelia. *Proc. Natl. Acad. Sci. USA.* 1993, 90, 7275–7279.

90. Nielsen, S.; Nagelhus, E.A.; Amiry-Moghaddam, M.; Bourque, C.; Agre, P.; Ottersen, O.P. Specialized membrane domains for water transport in glial cells: High-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J. Neurosci.* 1997, 17, 171–180.

91. Nagelhus, E.A.; Veruki, M.L.; Torp, R.; Haug, F.M.; Laake, J.H.; Nielsen, S.; Agre, P.; Ottersen, O.P. Aquaporin-4 water channel protein in the rat retina and optic nerve: Polarized expression in Muller cells and fibrous astrocytes. *J. Neurosci.* 1998, 18, 2506–2519.

92. Frigeri, A.; Nicchia, G.P.; Nico, B.; Quondamatteo, F.; Herken, R.; Roncali, L.; Svelto, M. Aquaporin-4 deficiency in skeletal muscle and brain of dystrophic mdx mice. *FASEB J.* 2001, 15, 90–98.

93. Elkjaer, M.L.; Vajda, Z.; Nejsum, L.N.; Kwon, T.; Jensen, U.B.; Amiry-Moghaddam, M.; Frokiaer, J.; Nielsen, S. Immunolocalization of AQP9 in liver, epididymis, testis, spleen, and brain. *Biochem. Biophys. Res. Commun.* 2000, 276, 1118–1128.

94. Nicchia, G.P.; Frigeri, A.; Nico, B.; Ribatti, D.; Svelto, M. Tissue distribution and membrane localization of aquaporin-9 water channel: Evidence for sex-linked differences in liver. *J. Histochem. Cytochem.* 2001, 49, 1547–1556.

95. Nicchia, G.P.; Frigeri, A.; Liuzzi, G.M.; Svelto, M. Inhibition of aquaporin-4 expression in astrocytes by RNAi determines alteration in cell morphology, growth, and water transport and induces changes in ischemia-related genes. *FASEB J.* 2003, 17, 1508–1510.

96. Binder, D.K.; Papadopoulos, M.C.; Haggie, P.M.; Verkman, A.S. In vivo measurement of brain extracellular space diffusion by cortical surface photobleaching. *J. Neurosci.* 2004, 24, 8049–8056.

97. Fujii, F.; Nodasaka, Y.; Nishimura, G.; Tamura, M. Anoxia induces matrix shrinkage accompanied by an increase in light scattering in isolated brain mitochondria. *Brain Res.* 2004, 999, 29–39.

98. Griesdale, D.E.; Honey, C.R. Aquaporins and brain edema. *Surg. Neurol.* 2004, 61, 418–421.

99. Manley, G.T.; Fujimura, M.; Ma, T.; Noshita, N.; Filiz, F.; Bollen, A.W.; Chan, P.; Verkman, A.S. Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke. *Nat. Med.* 2000, 6, 159–163.

100. Calamita, G.; Ferri, D.; Gena, P.; Liquori, G.E.; Cavalier, A.; Thomas, D.; Svelto, M. The inner mitochondrial membrane has Aquaporin-8 water channels and is highly permeable to water. *J. Biol. Chem.* 2005, 280, 17149–17153.

101. Li, Y.; Schmidt-Edelkraut, U.; Poetz, F.; Oliva, I.; Mandl, C.; Hözl-Wenig, G.; Schönig, K.; Bartsch, D.; Ciccolini, F. γ -aminobutyric A receptor (GABA(A)R) regulates Aquaporin 4 expression in the subependymal zone: Relevance to neural precursors and water exchange. *J. Biol. Chem.* 2015, 290, 4343–4355.

102. Yi, F.; Khan, M.; Gao, H.; Hao, F.; Sun, M.; Zhong, L.; Lu, C.; Feng, X.; Ma, T. Increased differentiation capacity of bone marrow-derived mesenchymal stem cells in aquaporin-5 deficiency. *Stem Cells Dev.* 2012, 21, 2495–2507.

103. Chen, M.; Li, Y.; Xiao, L.; Dai, G.; Lu, P.; Wang, Y.; Rui, Y. AQP1 modulates tendon stem/progenitor cells senescence during tendon aging. *Cell Death Dis.* 2020, 11, 193.

104. De Ieso, M.L.; Yool, A.J. Mechanisms of aquaporin-facilitated cancer invasion and metastasis. *Front. Chem.* 2018, 6,135.

105. Hui, T.H.; Kwan, K.W.; Yip, T.T.C.; Fong, H.W.; Ngan, K.C.; Yu, M.; Yao, S.; Ngan, A.H.W.; Lin, Y. Regulating the membrane transport activity and death of cells via electroosmotic manipulation. *Biophys. J.* 2016, 110, 2769– 2778.

106. Günther, H.S.; Schmidt, N.O.; Phillips, H.S.; Kemming, D.; Kharbanda, S.; Soriano, R.; Modrusan, Z.; Meissner, H.; Westphal, M.; Lamszus, K. Glioblastoma-derived stem cell-enriched cultures form distinct subgroups according to molecular and phenotypic criteria. *Oncogene* 2007, 27, 2897–2909.

107. Schulte, A.; Günther, H.S.; Phillips, H.S.; Kemming, D.; Martens, T.; Kharbanda, S.; Soriano, R.H.; Modrusan, Z.; Zapf, S.; Westphal, M.; et al. A distinct subset of glioma cell lines with stem cell – like properties reflects the transcriptional phenotype of glioblastomas and overexpresses CXCR4 as therapeutic target. *Glia* 2011, 59, 590–602.

Retrieved from <https://encyclopedia.pub/entry/history/show/13827>