

# Craniometaphyseal Dysplasia

Subjects: Genetics & Heredity

Contributor: Nicole Yin

Craniometaphyseal dysplasia is a rare condition characterized by thickening (overgrowth) of bones in the skull (cranium) and abnormalities in a region at the end of long bones known as the metaphysis. The abnormal bone growth continues throughout life. Except in the most severe cases, the lifespan of people with craniometaphyseal dysplasia is normal.

Keywords: genetic conditions

---

## 1. Introduction

Bone overgrowth in the head causes many of the signs and symptoms of craniometaphyseal dysplasia. Affected individuals typically have distinctive facial features such as a wide nasal bridge, a prominent forehead, wide-set eyes (hypertelorism), and a prominent jaw. Excess bone formation in the jaw can delay teething (dentition) or result in absent (non-erupting) teeth. Infants with craniometaphyseal dysplasia may have breathing or feeding problems caused by narrow nasal passages. In severe cases, abnormal bone growth can pinch (compress) the nerves that extend from the brain to various areas of the head and neck (cranial nerves). Compression of the cranial nerves can lead to paralyzed facial muscles (facial nerve palsy), blindness, or deafness.

The x-rays of individuals with craniometaphyseal dysplasia show unusually shaped long bones, particularly long bones in the legs. The ends of these bones are wider and appear less dense than usual in people with this condition.

There are two types of craniometaphyseal dysplasia, which are distinguished by their pattern of inheritance and genetic cause. They are known as the autosomal dominant and autosomal recessive types.

## 2. Frequency

Craniometaphyseal dysplasia is a very rare disorder; its incidence is unknown.

## 3. Causes

Mutations in the *ANKH* gene cause autosomal dominant craniometaphyseal dysplasia. The *ANKH* gene provides instructions for making a protein that plays a role in the development and function of cells that build bones (osteoblasts) and cells that break down bone (osteoclasts). Osteoclasts are involved in bone remodeling, a normal process in which old bone is removed and new bone is created to replace it. In addition, the ANKH protein transports a molecule called pyrophosphate out of cells. The pyrophosphate found outside of cells (extracellular pyrophosphate) helps control bone formation by preventing mineralization, the process by which minerals such as calcium and phosphorus are deposited in developing bones. The ANKH protein may have other, unknown functions.

Mutations in the *ANKH* gene that cause autosomal dominant craniometaphyseal dysplasia impair the maturation (differentiation) of osteoclasts, which likely disrupts bone remodeling. Reduced breakdown of bone tissue can contribute to the bone thickening characteristic of craniometaphyseal dysplasia. *ANKH* gene mutations may also reduce the protein's ability to transport pyrophosphate out of cells. A shortage of extracellular pyrophosphate can increase bone mineralization, which may also contribute to the bone abnormalities.

A mutation in the *GJA1* gene causes some cases of autosomal recessive craniometaphyseal dysplasia. This gene provides instructions for making a protein called connexin 43, which is involved in the development of many tissues in the body, including bone. The protein may be involved in bone remodeling. It is unclear how a mutation in the *GJA1* gene leads to the particular bone abnormalities of craniometaphyseal dysplasia.

The genetic cause of many cases of autosomal recessive craniometaphyseal dysplasia is unknown. It is likely that other, unidentified genes are involved in this form of the disorder.

### 3.1. The Genes Associated with Craniometaphyseal Dysplasia

- ANKH
- GJA1

## 4. Inheritance

When caused by mutations in the *ANKH* gene, craniometaphyseal dysplasia follows an autosomal dominant pattern, which means one altered copy of the *ANKH* gene in each cell is sufficient to cause the disorder. Individuals with autosomal dominant craniometaphyseal dysplasia typically have one parent who also has the condition. Less often, cases result from new mutations in the gene and occur in people with no history of the disorder in their family.

When caused by mutations in the *GJA1* gene, craniometaphyseal dysplasia has an autosomal recessive inheritance pattern, which means both copies of the *GJA1* gene in each cell are altered. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the disorder.

## 5. Other Names for This Condition

- autosomal dominant craniometaphyseal dysplasia
- autosomal recessive craniometaphyseal dysplasia
- CMD
- CMDD
- CMDJ
- CMDR
- craniometaphyseal dysplasia, Jackson type

---

## References

1. Addison WN, Azari F, Sørensen ES, Kaartinen MT, McKee MD. Pyrophosphate inhibits mineralization of osteoblast cultures by binding to mineral, up-regulating osteopontin, and inhibiting alkaline phosphatase activity. *J Biol Chem*. 2007 May 25;282(21):15872-83.
  2. Hu Y, Chen IP, de Almeida S, Tiziani V, Do Amaral CM, Gowrishankar K, Passos-Bueno MR, Reichenberger EJ. A novel autosomal recessive *GJA1* missense mutation linked to Craniometaphyseal dysplasia. *PLoS One*. 2013 Aug 12;8(8):e73576. doi: 10.1371/journal.pone.0073576.
  3. Iughetti P, Alonso LG, Wilcox W, Alonso N, Passos-Bueno MR. Mapping of the autosomal recessive (AR) craniometaphyseal dysplasia locus to chromosome region 6q21-22 and confirmation of genetic heterogeneity for mild AR spondylocostal dysplasia. *Am J Med Genet*. 2000 Dec 18;95(5):482-91.
  4. Nürnberg P, Thiele H, Chandler D, Höhn W, Cunningham ML, Ritter H, Leschik G, Uhlmann K, Mischung C, Harrop K, Goldblatt J, Borochowitz ZU, Kotzot D, Westermann F, Mundlos S, Braun HS, Laing N, Tinschert S. Heterozygous mutations in *ANKH*, the human ortholog of the mouse progressive ankylosis gene, result in craniometaphyseal dysplasia. *Nat Genet*. 2001 May;28(1):37-41.
  5. Reichenberger E, Chen IP. Craniometaphyseal Dysplasia, Autosomal Dominant. 2007 Aug 27 [updated 2020 Jun 11]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. *GeneReviews*® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1461/>
  6. Reichenberger E, Tiziani V, Watanabe S, Park L, Ueki Y, Santanna C, Baur ST, Shiang R, Grange DK, Beighton P, Gardner J, Hamersma H, Sellars S, Ramesar R, Lidral AC, Sommer A, Raposo do Amaral CM, Gorlin RJ, Mulliken JB, Olsen BR. Autosomal dominant craniometaphyseal dysplasia is caused by mutations in the transmembrane protein ANK. *Am J Hum Genet*. 2001 Jun;68(6):1321-6.
  7. Talbot J, Brion R, Lamora A, Mullard M, Morice S, Heymann D, Verrecchia F. Connexin43 intercellular communication drives the early differentiation of human bone marrow stromal cells into osteoblasts. *J Cell Physiol*. 2018 Feb;233(2):946-957. doi: 10.1002/jcp.25938.
  8. Zaka R, Williams CJ. Role of the progressive ankylosis gene in cartilage mineralization. *Curr Opin Rheumatol*. 2006 Mar;18(2):181-6. Review.
-

