

Bacteriophage-Mediated Control of Phytopathogenic Xanthomonads

Subjects: Microbiology | Agronomy | Horticulture

Contributor: Emilio Stefani

Xanthomonads, members of the family Xanthomonadaceae, are economically important plant pathogenic bacteria responsible for infections of over 400 plant species. Bacteriophage-based biopesticides can provide an environmentally friendly, effective solution to control these bacteria. Bacteriophage-based biocontrol has important advantages over chemical pesticides, and treatment with these biopesticides is a minor intervention into the microflora. However, bacteriophages' agricultural application has limitations rooted in these viruses' biological properties as active substances. These disadvantageous features, together with the complicated registration process of bacteriophage-based biopesticides, means that there are few products available on the market.

Keywords: bacteriophages ; bacteriophage therapy ; biological control ; Xanthomonas spp. ; sustainable agriculture ; novel biopesticides

1. Introduction

Plant diseases in pre- and post-harvest frequently account for 20% or more product losses, both in emerging countries as well as in developed areas ^[1]. Although less numerous than fungal diseases, bacterial diseases are often difficult to manage, due to their frequent polycyclic nature and the lack of systemic antibacterial substances ^[1]. Copper compounds and antibiotics are the only antibacterial choices to control phytopathogenic bacteria that are readily available in a large part of the world ^{[2][3]}. Copper presents several risks and unexpected consequences in agricultural systems and for the environment, e.g., phytotoxicity, negative effects on pollinating insects and other beneficial organisms, bioaccumulation in soil and surface water and reduction of microbial biodiversity ^{[4][5][6]}. Antibiotics, such as mainly streptomycin, kasugamycin and tetracyclines, as active substances in agriculture may also pose unacceptable risks when used as pesticides ^[3]. Indeed, although they do not accumulate or cause adverse effects on plants, they may incite the development of resistant traits in bacterial populations, including in the target pathogen(s), and transfer them to bacteria of clinical interest ^[7]. The urgent need to tackle pathogen control in agricultural systems using a more sustainable approach has directed research towards different strategies, among them the development and implementation of microbial biocontrol agents and bacteriophages ^{[8][9]}. In this review, we present the available knowledge on the use of bacteriophages in the management of xanthomonads, the largest group of phytopathogenic bacteria that are often the causal agents of devastating diseases in important crops. This review presents current knowledge on xanthomonads, bacteriophages, host-microbe interaction and ecology interactions. This information, -together with the description of results of relevant laboratory, greenhouse and field trials- supports the understanding of factors influencing the effectivity of bacteriophage-based biopesticides in the fields.

1.1. Xanthomonads

Xanthomonads are Gram negative bacteria belonging to the family of *Xanthomonadaceae*. Within this family *Xanthomonas* emerges as one of the most important genera in phyto bacteriology, for it comprises around forty bacterial species pathogenic to over 400 plant species ^[10]. In turn, several *Xanthomonas* species are further taxonomically classified into different subspecies and pathovars, thus confirming a particular adaptation to plants. Such phytopathological adaptation is due to the expression of virulence factors ^{[11][12]}. Most *Xanthomonas* sp. strains are characterized by their production of xanthomonadin, a yellow pigment that represents the most useful diagnostic feature used for their identification ^[13], although a few pathovars are reported that do not produce such pigment, e.g.: *X. axonopodis* pv. *manihotis*, *X. campestris* pv. *mangiferaeindicae* and *X. campestris* pv. *viticola* ^{[14][15]}. Over the past 25 years, *Xanthomonas* species have undergone thorough changes in nomenclature based on phenotypic and conventional molecular techniques and, more recently, whole-genome sequencing (WGS) ^{[16][17]}. Indeed, evolutionary dynamics renders *Xanthomonas* species as rapidly evolving microbes and they are particularly successful as plant pathogens ^{[14][18]}.

Several devastating plant diseases are caused by xanthomonads, for example *X. oryzae* pv. *oryzae* is the causal agent of bacterial blight, the most serious disease of rice. Together with pv. *oryzicola*, the causal agent of bacterial leaf streak, both pathogens frequently represent a limiting factor constraining rice production in tropical and subtropical regions [19]. Both pathogens exhibit large genetic variation among isolates, thus accounting for a high genetic plasticity [12].

The bacterial canker of citrus, incited by *X. citri* subsp. *citri* affects all commercial varieties of citrus [20]. Two other major crops are affected by xanthomonads: bananas (all types), affected by bacterial wilt caused by *X. vasicola* pv. *musacearum* and cassava, affected by bacterial wilt caused by *X. phaseoli* pv. *manihotis* [21]. International trade and climate change appear fundamental to support dissemination of xanthomonads worldwide and their adaptation and establishment in new areas, as several recent findings confirm [22][23][24].

1.2. Biological Control of Xanthomonads

Biological control of plant pathogenic bacteria may be implemented in several ways, for example (1) using microbial antagonists producing specific substances, such as bacteriocins (antibiosis), (2) using beneficial bacteria to efficiently compete for nutritional resources in planta [25], or (3) applying microbes that produce anti-Quorum Sensing factors [26], or (4) act as hyperparasites [27]. Emerging biocontrol strategies for plant pathogens, and for xanthomonads in particular, increasingly rely on the use of selected microbial biocontrol agents, or microbiome engineering [28][29]. Several microorganisms can efficiently control xanthomonads, both in vitro and in vivo, with some also showing plant growth promoting traits [30]. Specifically, bacterial species belonging to the genera *Pseudomonas* and *Bacillus* are reported to be effective against several *Xanthomonas* spp. A large number of papers describe satisfactory results on the biocontrol of *X. citri* pv. *citri*, *X. campestris* pv. *campestris* and *X. vesicatoria* [28], but most described results were obtained in vitro or in a controlled environment. Conversely, reproducibility of such published results in agricultural systems is not as good as expected, possibly due to the differences in agricultural context and the cropping systems. Nonetheless, a few commercial products based on microbial biocontrol agents that have satisfactory antibacterial activity are readily available on the market. For instance, Serenade® and Serenade® Max (Bayer Crop Science, Leverkusen, Germany) based on a selected strain of *Bacillus subtilis*, are indicated for the biological control of *X. arboricola* pv. *pruni*. Similarly, Double Nickel™ LC (Certis, Columbia, MD, USA) based on a strain of *Bacillus amyloliquefacies*, is indicated for the biological control of the tomato spot disease (*X. perforans*).

1.3. Bacteriophages

Bacteriophages are viruses that specifically infect bacteria and have no direct negative effects on animals or plants. Bacteriophages are widely distributed on the Earth and are measurable components of the natural microflora [31]. In agricultural environments there are multiple sources of bacteriophages, such as healthy and diseased plant organs, soil, surface water, sewage and sludge, particularly from processing plants [32]. Bacteriophages may have different life cycles in natural environments. This includes a lytic life cycle, where a bacteriophage infects its bacterial host cell and rapidly induces its breakdown and a lysogenic cycle, where they are able to integrate their injected DNA into the bacterial genome [33].

Together with research on bacteriophages as prospective biocontrol agents, a number of studies were devoted to elucidating bacterial taxonomy. Bacteriophages have been used as tools to identify and characterize phytopathogenic bacteria [34]. Then, the use of specific bacteriophages appeared to be essential for population studies of phytopathogenic bacteria, in order to unravel key epidemiological factors. This supported the successful use of phages in controlling bacterial diseases [35].

Recent publications on isolation and characterization of bacteriophages against xanthomonads are summarized in [Table 1](#).

Table 1. List of recent publications on bacteriophages against *Xanthomonas* spp.

Host Bacteria, Disease Name and Host Plant	Description of Works Performed	Reference
<i>Xanthomonas fragariae</i> Angular leaf spot in strawberry	Isolation and whole genome sequence analysis of N4-like bacteriophage, named RiverRider, including its host range.	[36]
<i>Xanthomonas citri</i> Asian citrus canker	Isolation and genome sequence analysis of <i>Xanthomonas</i> virus XacN1, a novel jumbo myovirus, showing a wider host range than other <i>X. citri</i> bacteriophages.	[37]

Host Bacteria, Disease Name and Host Plant	Description of Works Performed	Reference
<i>Xanthomonas oryzae</i> pv. <i>oryzae</i> Bacterial leaf blight of rice	Characterization of a novel phage Xoo-sp2, isolated from soil and its potential as a prophylactic agent in biocontrol of the disease.	[38]
	Isolation and complete genome sequence analysis of bacteriophage Xoo-sp13.	[39]
	Isolation and complete genome sequence analysis of a jumbo bacteriophage, Xoo-sp14.	[40]
	Isolation and analysis of the complete genome sequences of 10 OP2-like <i>X. oryzae</i> pv. <i>oryzae</i> bacteriophages	[41]
<i>Xanthomonas campestris</i> pv. <i>campestris</i> Black rot disease of kohlrabi	Evaluation of lytic activity of Xccφ1 bacteriophage in combination with 6-pentyl-α-pyrone (a secondary metabolite produced by <i>Trichoderma atroviride</i> P1) and the mineral hydroxyapatite for the prevention and eradication of bacterial biofilms.	[42]
	Isolation and characterization of specific bacteriophage (Xccφ1) able to control disease, and investigation of <i>X. campestris</i> pv. <i>campestris</i> and Xccφ1, applied singly or combined, on plant metabolome.	[43]
<i>Xanthomonas campestris</i> pv. <i>Campestris</i> Black rot of crucifers	Isolation of phage infecting <i>X. campestris</i> pv. <i>campestris</i> and characterization of the bacteriophage Xcc9SH3.	[44]
<i>Xanthomonas campestris</i> pv. <i>Campestris</i> Black rot of cauliflower	Isolation and morphological, molecular and phylogenetic characterization of <i>X. campestris</i> pv. <i>campestris</i> specific bacteriophage named “Xanthomonas virus XC 2”	[45]
<i>Xanthomonas arboricola</i> pv. <i>Juglandis</i> Walnut blight	Isolation of 24 phages from soil and infected walnut aerial tissues. Two polyvalent bacteriophages, were characterized by their morphological, physiological and genomic analyses.	[46]
	Isolation and complete genome analysis of three bacteriophages, f20-Xaj, f29-Xaj and f30-Xaj, specific to <i>X. arboricola</i> pv. <i>juglandis</i>	[47]
<i>Xanthomonas vesicatoria</i> Bacterial spot of pepper	Isolation and complete genome sequence of a filamentous bacteriophage XaF13 infecting <i>X. vesicatoria</i>	[48]
	Isolation and complete genome sequence of <i>X. vesicatoria</i> bacteriophage ΦXaF18	[49]

2. *Xanthomonas*-Host Plant and Bacteriophage-Host Bacterium Interactions and Their Possible Influence on Bacteriophage-Based Biocontrol Strategies

No species is an island, as each individual organism is constantly in contact with others [50]. Here we discuss bacteriophage—host bacterium interactions and the factors that influence the possible outcomes of bacterial infection of the host plant. The presented data is helpful when identifying the non-satisfactory efficacy of bacteriophage-based pesticides when applied on the field and maybe useful when designing integrated plant management (e.g., with the involvement of other biopesticides). We provide possible solutions and explain why bacteriophage products may have distinct efficacies when applied on different fields. We will also analyze the applicable *Xanthomonas*-plant interactions from the point of view of biocontrol and the relevant bacteriophage-bacterium interactions. Finally, we will investigate the mechanisms of bacteriophage resistance of bacteria.

2.1. *Xanthomonas*-Host Plant Interactions

Bacteriophage-based biocontrol treatments of xanthomonads intend to interfere with a plant-pathogenic *Xanthomonas* spp. system. This subsection contains essential information on this system.

Xanthomonads live part of their life cycle outside the host plant as epiphytes in the lesions of fallen leaves or associated to plant debris in the soil [51]. They are an essential component of the soil microbiome, with 2–7% relative abundance in the bacterial community [52].

The infection cycle of *Xanthomonas* spp. starts with an epiphytic phase followed by entering the host plant through natural openings (stomata, hydathodes) and wounds to start its internal colonization (endophytic phase) [53]. When introduced into the plant surface, xanthomonads use a variety of adhesion strategies to attach to the plant [54][55][56][57][58][59]. Plants

have also evolved various defence mechanisms to protect themselves from pathogens [60]. They respond to pathogen associated molecular patterns (PAMPs) by activating PAMP-triggered immunity (PTI) or effector-triggered immunity (ETI) mediated by pathogen-specific receptors [61]. As a result, a systemic acquired resistance (SAR) status may be established, potentially increasing resistance to subsequent attacks in the entire plant [62][63][64].

A first key element of bacterial survival in the phyllosphere is the biofilm formation, creating a microenvironment that can protect bacteria against environmental stress conditions [58][65]. This is an important virulence factor of phytopathogenic *Xanthomonas* spp. [66][67]. A biofilm, in addition to the cells, is primarily made up of proteins, lipids and extracellular polysaccharides (EPS) [68][69]. The formation of a biofilm may provide resistance to host defence mechanisms and vascular bacteria attachment to xylem vessels, or contribute to bacterial epiphytic survival prior to colonisation of the plant intercellular space [70]. The gum operon, a massive transcriptional unit containing 12 enzyme coding genes (*gumB-gumM*), mediates xanthan gum biosynthesis [71]. A study revealed that biofilm production deficient mutants (particularly *gumB* and *gumD*) showed significantly lower leaf surface survival than wild type *X. citri* pv. *citri* and *X. axonopodis* pv. *manihotis* [72][73][74]. The study of many *Xanthomonas* spp. have shown that gum genes contribute to bacterial in planta growth, epiphytic survival and disease symptom formation [72][75][76][77][78].

The assembly and dispersal of biofilms are partly mediated by the Quorum-sensing (QS) signal molecule, or diffusible signal factor (DSF). DSF positively influences the disruption of biofilms [79].

One survival strategy of bacteria during unfavorable conditions is the formation of persister cells. Persisters are a small fraction (0.001%–0.1%, or up to 1% in biofilms) of cells in a metabolically inactive, dormant state that are resistant against a wide range of antibiotics [80]. *X. campestris* pv. *campestris* and *X. citri* subsp. *citri* can form persister cells under different stress conditions [81][82]. Importantly, bacteriophages can also infect persisters [83].

LPS, as major components of the bacterial outer membrane, protect the cell from harmful environments and are another surface-associated virulence factor in *Xanthomonas* spp. Importantly, LPS not only function as virulence factors but also induce plant defense responses, such as pathogenesis-related gene expression, cell wall thickening and oxidative burst [84][85]. Mutations in LPS gene clusters make bacteria more susceptible to adverse environmental conditions, which may result in a reduction in bacterial virulence, as shown for *X. campestris* pv. *campestris* [86][87][88].

Xanthomonas species have a plethora of potential mechanisms that aid bacterial fitness in diverse environments, including the six different extracellular protein secretion systems (referred to as type I–VI, or T1SS–T6SS) that export proteins via the bacterial multilayer cell envelope and, in some cases, into host target cells. The conserved structural components that characterize these secretion systems, as well as the characteristics of their substrates and the pathway that these substrates take during the export process, distinguish them. T6SS was recently discovered and is involved in at least 25% of all sequenced gram-negative bacterial genomes [89]. The *Hcp* and *VgrG* proteins are essential components of T6SS that mimic the bacteriophage tail and needle complex, respectively [90]. Yang et al. [90] investigated the evolution of the T6SS in the *Xanthomonas* genus and assessed the relevance of the T6SS for virulence and in vitro motility in *X. phaseoli* pv. *manihotis* (*Xpm*), the causal agent of cassava bacterial blight. According to their phylogenetic analyses, the T6SS may have been obtained through a very ancient event of horizontal gene transfer (HGT) and preserved through evolution, implying their significance for host adaptation. They also showed that the T6SS of *Xpm* is functional and immensely contributes to motility and virulence.

Transcription activation-like effectors (TALEs) ensure plasticity in host adaptation for xanthomonads. TALEs have a repetitive domain governing the binding to promoters of host genes [91]. Novel TALEs could be created because this repetitive region is shared among TALEs, and recombination frequently occurs, as it was recently demonstrated in *X. oryzae* pv. *oryzae* [92]. These novel TALE encoding genes could be changed by HGT between bacteria, strengthening their host adaptation abilities [93].

2.2. Bacteriophage-Host Bacterium Interactions

When investigating ecological roles of bacteriophages in a *Xanthomonas* spp. population, it should be highlighted that the relationship between bacteriophages and their hosts could be both antagonistic and mutualistic, and the long-term survival of a bacteriophage population does not always require the lysis of its host. Therefore, bacteriophages are not predators, but either parasites or parasitoids of the host [94].

Bacteriophages can infect bacteria located in biofilms, albeit biofilms can provide a barrier for bacteriophage attacks compared to planktonic bacteria. This barrier is due to the physiological heterogeneity of the bacteria composing the biofilms, the secreted EPS, and the differential display of receptors on the host cell' surface [94]. Bacteriophages can

interact with biofilms of xanthomonads at several points. In a recent study Yoshikawa et al. [37] isolated the *X. citri* jumbo bacteriophage XacN1. They showed that the XacN1 genome encodes potential lytic enzymes such as cell wall hydrolases, C1 family peptidase, M23 family peptidases, lipase and chitinase. According to proteomic analysis, lipase, chitinase, and M23 family peptidases were discovered in the bacteriophage XacN1. They concluded that these enzymes may be necessary to disrupting the biofilm and initiating bacteriophage infection. Bacteriophages have evolved to counteract the biofilm barrier by using depolymerase enzymes on their capsids, and can also induce host lysis, allowing bacteriophages to degrade biofilm [95]. Furthermore, bacteriophage genomes carrying QS genes were detected in *Clostridium difficile* bacteriophage phiCDHM1 and three *Paenibacillus* bacteriophage genomes [96][97][98]. These genes can modify the biofilm disruption and other QS-mediated responses, including the decision on the lysogenic or lytic lifecycle of bacteriophages [97] or even the synthesis of virulence genes, as demonstrated in *X. campestris* [98].

Generally, the diversity of bacterial communities can support their adaptation to environmental circumstances [99]. If a community is more diverse, it is more stable as it can better adapt to the changing environment [100]. Prokaryotic viruses are essential in driving processes in microbial ecosystems [101][102]. In the absence of bacteriophages, one or several strains could become dominant in the niche, and other strains could be extinct, as was demonstrated in in vitro experiments [101][103]. Bacteriophages most likely infect the most abundant host strain, causing a decrease in its abundance ("kill the winner" principle). A consequence of this action will be a fluctuating selection, that increases diversity [103] and strengthens the community's stability or adaptation ability. This may cause that bacteriophage-based pesticides can support the presence of xanthomonads on the fields when not applied carefully. Integrated disease management together with the application of carefully selected bacteriophages timed appropriately could be one solution.

The genome of lysed cells will be available for surrounding bacteria, providing them novel genetic information, which may also include pathogenicity-related genes, as recently shown in the case of the cherry pathogen *Pseudomonas syringae* pv. *morsprunorum* or in *X. albilineans* [104][105]. Lytic bacteriophages increase the mutation rate in their host's genome, even in genes not related to bacteriophage resistance/immunity [101]. This effect can drive both adaptation (short term) or evolution (long term) processes. These from point of biocontrol disadvantageous features of lytic bacteriophages (i.e., providing novel genetic material for surrounding bacteria, increasing the mutation rate in the host's genome) could be managed by an integrated disease management. However, the mentioned drawbacks are less serious, for example, when lysogenic bacteriophages are applied in the fields. Lysogenic bacteriophages can protect bacteria carrying their genomes from superinfection (Superinfection: A second (delayed) bacteriophage infection of an already bacteriophage-infected bacterium) [106]. Horizontal gene transfer is one of the major factors (together with the mutations in avirulence genes) to evade host resistance [107][108][109]. The fact that 5–25% of the genome of *Xanthomonas* spp. originates from recombination events [110] highlights its importance in xanthomonads evolution and adaptation processes. Exchange of virulence factors between *Xanthomonas* spp. via HGT was observed in several cases [12].

The complexity of these HGT actions is demonstrated in the genome of a *X. anoxopodis* strain that contains a truncated bacteriophage genome carrying a gene resembling a plant protein that is induced during citrus blight disease [111].

As bacteriophages are often strain-specific, they can also act on the population level, influencing the population's intraspecific composition. Consequently, lysogens can contribute to the colonization of new niches. When lysis is induced in a small portion of the lysogenic cells, from superinfection-protected bacterial populations, and the bacteria originally located in the niche to be colonized are not protected from the infection, the new population can use their lysogenic bacteriophages as a weapon against the indigenous cells ("kill the relatives" principle) [101]. On the contrary, native bacteria can protect themselves against colonization by sacrificing a part of the population and inducing their prophages' lytic cycle [101]. Lysogenic bacteria can use their prophage weapon effectively, as observed in an in vitro experiment recently, where a lysogenic-lytic switch of bacteriophages to QS autoinducers strongly influenced the viral and bacterial abundance and diversity in soil communities [112].

There are examples of how lytic induction is carried out to optimize the multiplicity of infection (MOI). QS, encoded by either bacteria or bacteriophages, can influence this process [113][114]. Moreover, some bacteriophage genomes contain their own density monitoring equipment (the *arbitrium* system) and encode for small oligopeptides with which the bacteriophage density can be measured, as described in *Bacillus* bacteriophages [114][115]. Lysogeny is preferred when bacteriophages are abundant. Based on the described features of lysogenic and transducing bacteriophages, their field application may contribute to the adaptation and pathogenicity of xanthomonads, i.e., it may lead to unwanted effects. Therefore, the application of well-characterized, strictly lytic bacteriophages is advisable for bacteriophage-based biocontrol.

As bacteriophages and their hosts are not alone in the microflora, bacteriophages will meet their hosts with rare frequency when the living cell number of the host is low. Thus, one important consequence of the "kill the winner" principle is that bacteriophages cannot reduce the living cell number of their hosts to zero in a community [116], a property which differs from most chemical antibacterial compounds.

We mentioned examples in this subsection, how bacteriophages (both lytic and lysogenic ones) can alter the strain and/or species abundancies in communities. The composition of *Xanthomonas* spp. population and/or the microbial community may be distinct in different fields which may be differentially influenced by the described effects of bacteriophages. In addition to the environmental factors, a result of this divergent influence may lead to a distinct outcome of bacteriophage-based biocontrol in fields, at least in several cases [117].

2.3. Bacteriophage Resistance in Bacteria

Bacteriophage-resistance mutations in bacteria usually come with a fitness cost, such as a decrease in virulence, which results in less disease severity. This is because many of the molecules taking part in bacteriophage attachment are also engaged in the virulence mechanism. As a result, mutations that lead to resistance commonly compromise virulence. There are a few examples of how mutations in bacteria surface structures lead to decreased virulence, such as mutation in the *X. campestris* *xanA* gene needed for xanthan and lipopolysaccharide synthesis, which significantly decreases the effectiveness of bacteriophage L7 adsorption [118].

Bacteriophage resistance in bacteria is one of the main concerns regarding the bacteriophage-based biocontrol strategies. A detailed understanding of bacterial resistance to bacteriophages and their interaction with plants play an important role in the design of bacteriophage-based biocontrol strategies of xanthomonads. To survive bacteriophage infections, bacteria have developed a wide range of protection strategies, including spontaneous mutations, restriction modification systems (R–M systems), and adaptive immunity through the CRISPR-Cas system [106]. The key mechanisms driving bacteriophage resistance are spontaneous mutations, which can grant bacteriophage resistance by altering the structure of bacterial surface components that function as bacteriophage receptors [119]. Furthermore, bacteria can acquire resistance through lysogenic bacteriophages that carry sequences in their genetic material which encode bacterial resistance or toxins and incorporated into the bacterial genome [120]. The mechanisms by which bacteriophages counteract the anti-bacteriophage systems of bacteria are poorly understood. Bacteriophages with the ability to acquire new receptor tropism can modify their receptor-binding protein, which means that when a host receptor changes to a mutated form, bacteriophages can recognize the altered receptor structure and thus overcome disturbance in receptors for bacteriophage adsorption [121]. Bacteriophages use various anti-restriction strategies to avoid the wide range of R–M systems. These modification genes encode a small protein that is transmitted to the cell with the viral genome, or it may instantly neutralize the host immune system by intervening with the formation or function of the CRISPR–Cas ribonucleoprotein [122]. Bacteriophages may use bacterial CRISPR–Cas systems to promote their own replication, allowing the phage to complete its lytic cycle [123]. When a bacterium develops resistance to a specific bacteriophage, it retains sensitivity to bacteriophages with various cell surface receptors. Bacteriophage-mediated selection can be used in disease management, for example, by combining various bacteriophages to broaden the host range and suppress resistance evolution [124] and/or reasonably combining bacteriophages and chemical control to establish synergies and decrease the likelihood of resistance evolution [125]. This implies that the application of a bacteriophage cocktail may be beneficial, even if bacteria quickly develop resistance, since resistant strains may be less fit, thus more treatable using another combined method.

3. Bacteriophage-Based Biocontrol of *Xanthomonas* spp.

3.1. Examples for Greenhouse and Field Trials

Shortly after their discovery, bacteriophages were evaluated for control of plant diseases, including those caused by *Xanthomonas* spp. Some of the first studies were conducted by Mallman and Hemstreet (1924) who isolated the "cabbage-rot organism" *X. campestris* pv. *campestris* from rotting cabbage and showed that the filtrate from the decomposed tissue could inhibit pathogen growth in vitro [126].

From the 1960s, a considerable number of studies explored the efficacy of phages for the control of bacterial spot of peach, caused by *X. arboricola* pv. *pruni* [127][128][129][130]. Civerolo and Keil [127] applied bacteriophages 1 h prior to inoculation by the pathogen and reduced bacterial spot severity on peach leaves to 22% compared to 58% for control plants under greenhouse conditions. Civerolo [128] found that preinoculation of peach seedling foliage with crude lysates of the bacteriophage mixtures resulted in 6–8% fewer infected leaves and a 17–31% reduction of disease compared to control plants. Application of premixed bacteriophage–pathogen suspension immediately before inoculation resulted in a

51–54% decrease of bacterial spot symptoms in peach seedlings. Zaccardelli et al., isolated eight bacteriophages active against *X. arboricola* pv. *pruni*, examined their host range and lytic ability, and selected a lytic bacteriophage strain with the broadest host range for disease control [129][130]. By weekly bacteriophage treatment they significantly reduced fruit spot incidence on peaches [130].

Significant achievements have been made in bacteriophage application for control of bacterial spot of tomato caused by *X. campestris* pv. *vesicatoria* in greenhouse and field conditions [131][132][133][134][135][136][137][138]. Flaherty et al. [131] used a mixture of host range mutant bacteriophages and effectively controlled tomato bacterial spot in greenhouse and field conditions. Moreover, bacteriophage application increased total weight of extra-large fruit comparing to nontreated control or plants treated with chemical bactericides. Balogh et al. [133] improved the efficacy of bacteriophage treatments in field and greenhouse experiments by using protective formulations that significantly increased bacteriophage longevity on the plant surface. Bacteriophage mixture formulated either with 0.5% pregelatinized corn flour, Cascrete NH-400 with 0.25% pregelatinized corn flour, or 0.75% powdered skim milk with 0.5% sucrose, provided significant disease control compared to untreated control. However, in greenhouse experiments skim milk gave the best results, while Cascrete performed best in the field [133].

In order to improve bacteriophage efficacy and provide consistent disease control, bacteriophages of *X. campestris* pv. *vesicatoria* have been studied as a part of integrated disease management practices [138]. Obradovic et al., tested various combinations of plant inducers and biological agents for control of tomato bacterial spot [139]. Acibenzolar-S-methyl applied in combination with bacteriophages formulated with skim milk and sucrose, reduced bacterial spot of tomato in a greenhouse [136] as well as in the field [135]. Recently, Abrahamian et al. [140] evaluated 19 different chemical agents, biological control agents, plant defense activators, and novel products for their ability to manage bacterial spot on tomato caused by *X. perforans*. They reported that combination of bacteriophages, cymoxanil, famoxadone and phosphoric acid, significantly improved the disease management compared to the copper-based standard treatment. All these studies led to bacteriophage treatment, integrated with other disease management practices (e.g., late blight), becoming a part of a standard integrated management program for tomato bacterial spot in Florida [138][139].

Gašić et al. [141] studied the efficacy of bacteriophage KΦ1 in the control of pepper bacterial spot caused by *X. euvesicatoria*. They found that double bacteriophage application, before and after challenge inoculation, significantly reduced disease incidence when compared to untreated control. However, integrated application of bacteriophages 2 h before and copper hydroxide 24 h before inoculation was the most efficient treatment. The same bacteriophage strain was used as a part of integrated disease management and combined with other biocontrol agents, copper compounds, antibiotics and plant inducers to control pepper bacterial spot [142]. Bacteriophage combination with copper-hydroxide and acibenzolar-S-methyl was the most effective treatment reducing the disease severity by 96–98% compared to control [142].

Similar studies were performed to develop management strategies for efficient and sustainable control of leaf blight of onion, caused by *X. axonopodis* pv. *allii*. Lang et al. [143] reported that biweekly or weekly applications of bacteriophages reduced disease severity in the field by 26 to 50%: similar to results achieved by weekly applications of copper-mancozeb. Therefore, integrated application of bacteriophage mixtures with acibenzolar-S-methyl could be a promising strategy for managing *Xanthomonas* leaf blight of onion and contribute to reduced use of chemical bactericides [143].

Comprehensive research was done on bacteriophage-mediated control of Asiatic citrus canker caused by *X. axonopodis* pv. *citri*, and citrus bacterial spot *X. axonopodis* pv. *citrumelo* [144][145][146]. Bacteriophage treatment, without skim milk formulation, provided an average 59% reduction in citrus canker severity in greenhouse experiments. In nursery, bacteriophage treatment reduced disease, but was less effective than copper-mancozeb, while bacteriophage integration with copper-mancozeb resulted in equal or less control than copper-mancozeb application alone [145]. Similar results were obtained in the management of citrus bacterial spot, where bacteriophage treatment provided significant disease reduction on moderately sensitive Valencia oranges while it was ineffective on the highly susceptible grapefruit [145]. Ibrahim et al. [146] reported that successful control of Asiatic citrus canker in greenhouse and field can be obtained by combination of bacteriophage mixture formulated with skim milk-sucrose and acibenzolar-S-methyl.

Initial research of bacteriophage infecting *X. oryzae* pv. *oryzae*, the causal agent of bacterial blight of rice, was conducted by Kuo et al., who applied purified bacteriophages 1, 3, and 7 days before inoculation, and obtained 100%, 96% and 86% reductions of bacterial leaf blight, respectively [147]. Recently, Chae et al. [148] significantly reduced the occurrence of bacterial leaf blight to 18.1% compared to 87% in untreated control by treatment with skim milk formulated bacteriophages. Ogunyemi et al. [149] reported the bacteriophage X3 was more effective in disease severity reduction (83.1%) if sprayed before inoculation rather than after (28.9–73.9%) it. However, seed treatment with bacteriophages reduced disease by 95.4%.

Other results on using bacteriophages specific to *Xanthomonadaceae* in plant disease control includes reduction of incidence of bacterial blight of geraniums caused by *X. campestris* pv. *pelargonii* with foliar application of h-mutant bacteriophages [150]. Nagai et al. [151] found that a non-pathogenic *Xanthomonas* sp. strain mixed with bacteriophages effectively controlled black rot of broccoli caused by *X. campestris* pv. *campestris* in field trials. Orynbayev et al. (2020) studied effects of bacteriophage suspensions mixed with different UV-protectants in control of black rot caused by *X. campestris* pv. *campestris* on cabbage seedlings. In two-year greenhouse experiments, bacteriophage DB1 mixed with 0.75% skimmed milk showed an average efficacy of 71.1% in control of the disease, compared to 59.1% efficacy of Kocide 2000 treatment [152].

References

1. Oerke, E.C. Crop losses to pests. *J. Agric. Sci.* 2006, 144, 31–34.
2. Taylor, P.; Reeder, R. Antibiotic use on crops in low and middle-income countries based on recommendations made by agricultural advisors. *CABI Agric. Biosci.* 2020, 1, 1.
3. Sundin, G.W.; Castiblanco, L.F.; Yuan, X.; Zeng, Q.; Yang, C.H. Bacterial disease management: Challenges, experience, innovation and future prospects: Challenges in Bacterial Molecular Plant Pathology. *Mol. Plant. Pathol.* 2016, 17, 1506–1518.
4. Alengebawy, A.; Abdelkhalek, S.T.; Qureshi, S.R.; Wang, M.Q. Heavy Metals and Pesticides Toxicity in Agricultural Soil and Plants: Ecological Risks and Human Health Implications. *Toxics* 2021, 9, 42.
5. Vloek, V.; Pohanka, M. Adsorption of copper in soil and its dependence on physical and chemical properties. *Acta Univ. Agric. Silv. Mendel. Brun.* 2018, 66, 219–224.
6. Wang, L.; Xia, X.; Zhang, W.; Wang, J.; Zhu, L.; Wang, J.; Wei, Z.; Ahmad, Z. Separate and joint eco-toxicological effects of sulfadimidine and copper on soil microbial biomasses and ammonification microorganisms abundances. *Chemosphere* 2019, 228, 556–564.
7. Sundin, G.W.; Wang, N. Antibiotic Resistance in Plant-Pathogenic Bacteria. *Annu. Rev. Phytopathol.* 2018, 56, 161–180.
8. Köhl, J.; Kolnaar, R.; Ravensberg, W.J. Mode of Action of Microbial Biological Control Agents Against Plant Diseases: Relevance Beyond Efficacy. *Front. Plant Sci.* 2019, 10, 845.
9. Svircev, A.; Roach, D.; Castle, A. Framing the Future with Bacteriophages in Agriculture. *Viruses* 2018, 10, 218.
10. Hayward, A. *The Hosts of Xanthomonas*; Springer: Dordrecht, The Netherlands, 1993.
11. Medina, C.A.; Reyes, P.A.; Trujillo, C.A.; Gonzalez, J.L.; Bejarano, D.A.; Montenegro, N.A.; Jacobs, J.M.; Joe, A.; Restrepo, S.; Alfano, J.R.; et al. The role of type III effectors from *Xanthomonas axonopodis* pv. *manihotis* in virulence and suppression of plant immunity. *Mol. Plant Pathol.* 2018, 19, 593–606.
12. Timilsina, S.; Potnis, N.; Newberry, E.A.; Liyanapathirana, P.; Iruegas-Bocardo, F.; White, F.F.; Goss, E.M.; Jones, J.B. *Xanthomonas* diversity, virulence and plant-pathogen interactions. *Nat. Rev. Microbiol.* 2020, 18, 415–427.
13. Poplawsky, A.R.; Kawalek, M.D.; Schaad, N. A xanthomonadin-encoding gene cluster for the identification of pathovars of *Xanthomonas campestris*. *Mol. Plant Microbe Interact.* 1993, 6, 545.
14. Midha, S.; Patil, P.B. Genomic insights into the evolutionary origin of *Xanthomonas axonopodis* pv. *citri* and its ecological relatives. *Appl. Environ. Microbiol.* 2014, 80, 6266–6279.
15. Ferreira, M.A.S.V.; Bonneau, S.; Briand, M.; Cesbron, S.; Portier, P.; Darrasse, A.; Gama, M.A.S.; Barbosa, M.A.G.; Mariano, R.L.R.; Souza, E.B.; et al. *Xanthomonas citri* pv. *viticola* affecting grapevine in Brazil: Emergence of a successful monomorphic pathogen. *Front. Plant Sci.* 2019, 10, 489.
16. Rademaker, J.L.; Louws, F.J.; Schultz, M.H.; Rossbach, U.; Vauterin, L.; Swings, J.; de Bruijn, F.J. A comprehensive species to strain taxonomic framework for *Xanthomonas*. *Phytopathology* 2005, 95, 1098–1111.
17. Timilsina, S.; Kara, S.; Jacques, M.A.; Potnis, N.; Minsavage, G.V.; Vallad, G.E.; Jones, J.B.; Fischer-Le Saux, M. Reclassification of *Xanthomonas gardneri* (ex Šutič 1957) Jones et al. 2006 as a later heterotypic synonym of *Xanthomonas cynarae* Trébaol et al. 2000 and description of *X. cynarae* pv. *cynarae* and *X. cynarae* pv. *gardneri* based on whole genome analyses. *Int. J. Syst. Evol. Microbiol.* 2019, 69, 343–349.
18. Bansal, K.; Kumar, S.; Patil, P.B. Complete Genome Sequence Reveals Evolutionary Dynamics of an Emerging and Variant Pathovar of *Xanthomonas euvesicatoria*. *Genome Biol. Evol.* 2018, 10, 3104–3109.
19. Niño-Liu, D.O.; Ronald, P.C.; Bogdanove, A.J. *Xanthomonas oryzae* pathovars: Model pathogens of a model crop. *Mol. Plant Pathol.* 2006, 7, 303–324.

20. Martins, P.M.M.; de Oliveira Andrade, M.; Benedetti, C.E. *Xanthomonas citri* subsp. *citri*: Host interaction and control strategies. *Trop. Plant Pathol.* 2020, 45, 213–236.
21. Verdier, V.; López, C.; Bernal, A. Cassava bacterial blight, caused by *Xanthomonas axonopodis* pv. *manihotis*. In *Cassava in the Third Millennium: Modern Production, Processing, Use, and Marketing Systems*; Latin American and Caribbean Consortium to support Cassava Research and Development; Centro Internacional de Agricultura Tropical: Cali Colombia; Technical Centre for Agricultural and Rural Cooperation: Wageningen, The Netherlands, 2012.
22. Altin, I.; Casoli, L.; Stefani, E. First report of bacterial spot caused by *Xanthomonas cucurbitae* on pumpkin in Italy. *New Dis. Rep.* 2020, 41, 21.
23. Bultreys, A.; Gheysen, I. First report of *Xanthomonas phaseoli* pv. *phaseoli* and *Xanthomonas citri* pv. *fuscans* causing common bacterial blight of bean in Belgium. *New Dis. Rep.* 2020, 41, 6.
24. Popović, T.; Menković, J.; Prokić, A.; Obradović, A. First Report of *Xanthomonas arboricola* pv. *pruni* Causing Leaf and Fruit Spot on Apricot (*Prunus armeniaca* L.) in Montenegro. *Plant Dis.* 2021.
25. Andrews, J.H. Biological control in the phyllosphere. *Annu. Rev. Phytopathol.* 1992, 30, 603–635.
26. Helman, Y.; Chernin, L. Silencing the mob: Disrupting quorum sensing as a means to fight plant disease. *Mol. Plant Pathol.* 2015, 16, 316–329.
27. McNeely, D.; Chanyi, R.M.; Dooley, J.S.; Moore, J.E.; Koval, S.F. Biocontrol of Burkholderia cepacia complex bacteria and bacterial phytopathogens by *Bdellovibrio bacteriovorus*. *Can. J. Microbiol.* 2017, 63, 350–358.
28. Marin, V.R.; Ferrarezi, J.H.; Vieira, G.; Sass, D.C. Recent advances in the biocontrol of *Xanthomonas* spp. *World J. Microbiol. Biotechnol.* 2019, 35, 72.
29. Martínez-Hidalgo, P.; Maymon, M.; Pule-Meulenberg, F.; Hirsch, A.M. Engineering root microbiomes for healthier crops and soils using beneficial, environmentally safe bacteria. *Can. J. Microbiol.* 2019, 65, 91–104.
30. Vurukonda, S.S.K.P.; Stefani, E. Endophytic colonization by a streptomycete and a pseudomonad mediated plant growth promotion and enhanced antagonistic activities in tomato plants against *Xanthomonas vesicatoria*. *Can. J. Plant Pathol.* 2021, in press.
31. Mushegian, A.R. Are There 10³¹ Virus Particles on Earth, or More, or Fewer? *J. Bacteriol.* 2020, 202.
32. Clokie, M.R.; Millard, A.D.; Letarov, A.V.; Heaphy, S. Phages in nature. *Bacteriophage* 2011, 1, 31–45.
33. Abedon, S.T. *Phages, Ecology, Evolution*; Cambridge University Press: Cambridge, UK, 2008.
34. Billing, E. Further studies on the phage sensitivity and the determination of phytopathogenic *Pseudomonas* spp. *J. Appl. Bacteriol.* 1970, 33, 478–491.
35. Myung, I.S.; Cho, Y.; Lee, Y.H.; Kwon, H.M. Phage typing and lysotype distribution of *Xanthomonas axonopodis* pv. *citri*, the causal agent of citrus bacterial canker in Korea. *Plant Pathol. J.* 2001, 17, 336–341.
36. Miller, M.; Deiulio, A.; Holland, C.; Douthitt, C.; McMahon, J.; Wiersma-Koch, H.; Turechek, W.W.; D'Elia, T. Complete genome sequence of *Xanthomonas* phage RiverRider, a novel N4-like bacteriophage that infects the strawberry pathogen *Xanthomonas fragariae*. *Arch. Virol.* 2020, 165, 1481–1484.
37. Yoshikawa, G.; Askora, A.; Blanc-Mathieu, R.; Kawasaki, T.; Li, Y.; Nakano, M.; Ogata, H.; Yamada, T. *Xanthomonas citri* jumbo phage XacN1 exhibits a wide host range and high complement of tRNA genes. *Sci. Rep.* 2018, 8, 4486.
38. Dong, Z.; Xing, S.; Liu, J.; Tang, X.; Ruan, L.; Sun, M.; Tong, Y.; Peng, D. Isolation and characterization of a novel phage Xoo-sp2 that infects *Xanthomonas oryzae* pv. *oryzae*. *J. Gen. Virol.* 2018, 99, 1453–1462.
39. Nazir, A.; Dong, Z.; Liu, J.; Tahir, R.A.; Rasheed, M.; Qing, H.; Peng, D.; Tong, Y. Genomic analysis of bacteriophage Xoo-sp13 infecting *Xanthomonas oryzae* pv. *oryzae*. *Arch. Virol.* 2021, 166, 1263–1265.
40. Nazir, A.; Dong, Z.; Liu, J.; Zhang, X.; Tahir, R.A.; Ashraf, N.; Qing, H.; Peng, D.; Tong, Y. Sequence Analysis of a Jumbo Bacteriophage, Xoo-sp14, That Infects *Xanthomonas oryzae* pv. *oryzae*. *Microbiol. Resour. Announc.* 2020, 9.
41. Kovacs, T.; Molnar, J.; Varga, I.; Nagy, I.K.; Valappil, S.K.; Papp, S.; Vera Cruz, C.M.; Oliva, R.; Vizi, T.; Schneider, G.; et al. Complete Genome Sequences of 10 *Xanthomonas oryzae* pv. *oryzae* Bacteriophages. *Microbiol. Resour. Announc.* 2019, 8.
42. Papanianni, M.; Ricciardelli, A.; Fulgione, A.; d'Errico, G.; Zoina, A.; Lorito, M.; Woo, S.L.; Vinale, F.; Capparelli, R. Antibiofilm Activity of a *Trichoderma* Metabolite against *Xanthomonas campestris* pv. *campestris*, Alone and in Association with a Phage. *Microorganisms* 2020, 8, 620.
43. Papanianni, M.; Paris, D.; Woo, S.L.; Fulgione, A.; Rigano, M.M.; Parrilli, E.; Tutino, M.L.; Marra, R.; Manganiello, G.; Casillo, A.; et al. Plant Dynamic Metabolic Response to Bacteriophage Treatment After *Xanthomonas campestris* pv. *campestris* Infection. *Front Microbiol.* 2020, 11, 732.

44. Bhojar, M.S.; Singh, U.B.; Sahu, U.; Nagrale, D.T.; Sahu, P.K. Characterization of lytic bacteriophage XCC9SH3 infecting *Xanthomonas campestris* pv. *campestris*. *J. Plant Pathol.* 2017, 99, 233–238.
45. da Silva, F.P.; Xavier, A.D.S.; Bruckner, F.P.; de Rezende, R.R.; Vidigal, P.M.P.; Alfenas-Zerbini, P. Biological and molecular characterization of a bacteriophage infecting *Xanthomonas campestris* pv. *campestris*, isolated from brassica field. *Arch. Virol.* 2019, 164, 1857–1862.
46. Domotor, D.; Frank, T.; Rakhely, G.; Doffkay, Z.; Schneider, G.; Kovacs, T. Comparative analysis of two bacteriophages of *Xanthomonas arboricola* pv. *juglandis*. *Infect. Genet. Evol.* 2016, 43, 371–377.
47. Retamales, J.; Vasquez, I.; Santos, L.; Segovia, C.; Ayala, M.; Alvarado, R.; Nunez, P.; Santander, J. Complete Genome Sequences of Lytic Bacteriophages of *Xanthomonas arboricola* pv. *juglandis*. *Genome Announc.* 2016, 4.
48. Solis-Sanchez, G.A.; Quinones-Aguilar, E.E.; Fraire-Velazquez, S.; Vega-Arreguin, J.; Rincon-Enriquez, G. Complete Genome Sequence of XaF13, a Novel Bacteriophage of *Xanthomonas vesicatoria* from Mexico. *Microbiol. Resour. Annu.* 2020, 9.
49. Rios-Sandoval, M.; Quinones-Aguilar, E.E.; Solis-Sanchez, G.A.; Enriquez-Vara, J.N.; Rincon-Enriquez, G. Complete Genome Sequence of *Xanthomonas vesicatoria* Bacteriophage PhiXaF18, a Contribution to the Biocontrol of Bacterial Spot of Pepper in Mexico. *Microbiol. Resour. Annu.* 2020, 9.
50. Edwards, J.; Johnson, C.; Santos-Medellín, C.; Lurie, E.; Podishetty, N.K.; Bhatnagar, S.; Eisen, J.A.; Sundaresan, V. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc. Natl. Acad. Sci. USA* 2015, 112, E911–E920.
51. Zhao, Y.; Damicone, J.P.; Bender, C.L. Detection, Survival, and Sources of Inoculum for Bacterial Diseases of Leafy Crucifers in Oklahoma. *Plant Dis.* 2002, 86, 883–888.
52. Bulgarelli, D.; Garrido-Oter, R.; Munch, P.C.; Weiman, A.; Droge, J.; Pan, Y.; McHardy, A.C.; Schulze-Lefert, P. Structure and function of the bacterial root microbiota in wild and domesticated barley. *Cell Host Microbe* 2015, 17, 392–403.
53. Taulé, C.; Vaz-Jauri, P.; Battistoni, F. Insights into the early stages of plant–endophytic bacteria interaction. *World J. Microbiol. Biotechnol.* 2021, 37, 13.
54. Petrocelli, S.; Tondo, M.L.; Daurelio, L.D.; Orellano, E.G. Modifications of *Xanthomonas axonopodis* pv. *citri* lipopolysaccharide affect the basal response and the virulence process during citrus canker. *PLoS ONE* 2012, 7, e40051.
55. Pradhan, B.B.; Ranjan, M.; Chatterjee, S. XadM, a novel adhesin of *Xanthomonas oryzae* pv. *oryzae*, exhibits similarity to Rhs family proteins and is required for optimum attachment, biofilm formation, and virulence. *Mol. Plant. Microbe Interact.* 2012, 25, 1157–1170.
56. Dunger, G.; Guzzo, C.R.; Andrade, M.O.; Jones, J.B.; Farah, C.S. *Xanthomonas citri* subsp. *citri* type IV Pilus is required for twitching motility, biofilm development, and adherence. *Mol. Plant. Microbe Interact.* 2014, 27, 1132–1147.
57. Petrocelli, S.; Arana, M.R.; Cabrini, M.N.; Casabuono, A.C.; Moyano, L.; Beltramino, M.; Moreira, L.M.; Couto, A.S.; Orellano, E.G. Deletion of *pilA*, a Minor Pilin-Like Gene, from *Xanthomonas citri* subsp. *citri* Influences Bacterial Physiology and Pathogenesis. *Curr. Microbiol.* 2016, 73, 904–914.
58. An, S.Q.; Potnis, N.; Dow, M.; Vorholter, F.J.; He, Y.Q.; Becker, A.; Teper, D.; Li, Y.; Wang, N.; Bleris, L.; et al. Mechanistic insights into host adaptation, virulence and epidemiology of the phytopathogen *Xanthomonas*. *FEMS Microbiol. Rev.* 2020, 44, 1–32.
59. Bostock, R.M.; Pye, M.F.; Roubtsova, T.V. Predisposition in plant disease: Exploiting the nexus in abiotic and biotic stress perception and response. *Annu. Rev. Phytopathol.* 2014, 52, 517–549.
60. Boller, T.; He, S.Y. Innate immunity in plants: An arms race between pattern recognition receptors in plants and effectors in microbial pathogens. *Science* 2009, 324, 742–744.
61. Návarová, H.; Bernsdorff, F.; Döring, A.C.; Zeier, J. Pipecolic acid, an endogenous mediator of defense amplification and priming, is a critical regulator of inducible plant immunity. *Plant Cell* 2012, 24, 5123–5141.
62. Shah, J.; Chaturvedi, R.; Chowdhury, Z.; Venables, B.; Petros, R.A. Signaling by small metabolites in systemic acquired resistance. *Plant J.* 2014, 79, 645–658.
63. Schwachtje, J.; Fischer, A.; Erban, A.; Kopka, J. Primed primary metabolism in systemic leaves: A functional systems analysis. *Sci. Rep.* 2018, 8, 216.
64. Koczan, J.M.; Lenneman, B.R.; McGrath, M.J.; Sundin, G.W. Cell surface attachment structures contribute to biofilm formation and xylem colonization by *Erwinia amylovora*. *Appl. Environ. Microbiol.* 2011, 77, 7031–7039.
65. Yu, X.; Lund, S.P.; Scott, R.A.; Greenwald, J.W.; Records, A.H.; Nettleton, D.; Lindow, S.E.; Gross, D.C.; Beattie, G.A. Transcriptional responses of *Pseudomonas syringae* to growth in epiphytic versus apoplastic leaf sites. *Proc. Natl. Acad. Sci. USA* 2013, 110, E425–E434.

66. Li, J.; Wang, N. Foliar application of biofilm formation-inhibiting compounds enhances control of citrus canker caused by *Xanthomonas citri* subsp. *citri*. *Phytopathology* 2014, 104, 134–142.
67. Stoodley, P.; Sauer, K.; Davies, D.G.; Costerton, J.W. Biofilms as complex differentiated communities. *Annu. Rev. Microbiol.* 2002, 56, 187–209.
68. Sutherland, I.W. The biofilm matrix--an immobilized but dynamic microbial environment. *Trends Microbiol.* 2001, 9, 222–227.
69. Branda, S.S.; Vik, S.; Friedman, L.; Kolter, R. Biofilms: The matrix revisited. *Trends Microbiol.* 2005, 13, 20–26.
70. Vojnov, A.A.; Slater, H.; Daniels, M.J.; Dow, J.M. Expression of the gum operon directing xanthan biosynthesis in *Xanthomonas campestris* and its regulation in planta. *Mol. Plant Microbe Interact.* 2001, 14, 768–774.
71. Vojnov, A.A.; Zorreguieta, A.; Dow, J.M.; Daniels, M.J.; Dankert, M.A. Evidence for a role for the gumB and gumC gene products in the formation of xanthan from its pentasaccharide repeating unit by *Xanthomonas campestris*. *Microbiology* 1998, 144 Pt 6, 1487–1493.
72. Dunger, G.; Relling, V.M.; Tondo, M.L.; Barreras, M.; Ielpi, L.; Orellano, E.G.; Ottado, J. Xanthan is not essential for pathogenicity in citrus canker but contributes to *Xanthomonas epiphytic* survival. *Arch. Microbiol.* 2007, 188, 127–135.
73. Rigano, L.A.; Siciliano, F.; Enrique, R.; Sendin, L.; Filippone, P.; Torres, P.S.; Questa, J.; Dow, J.M.; Castagnaro, A.P.; Vojnov, A.A.; et al. Biofilm formation, epiphytic fitness, and canker development in *Xanthomonas axonopodis* pv. *citri*. *Mol. Plant Microbe Interact.* 2007, 20, 1222–1230.
74. Fonseca, N.P.; Patane, J.S.L.; Varani, A.M.; Felestrino, E.B.; Caneschi, W.L.; Sanchez, A.B.; Cordeiro, I.F.; Lemes, C. G.C.; Assis, R.A.B.; Garcia, C.C.M.; et al. Analyses of Seven New Genomes of *Xanthomonas citri* pv. *aurantifolii* Strains, Causative Agents of Citrus Canker B and C, Show a Reduced Repertoire of Pathogenicity-Related Genes. *Front. Microbiol.* 2019, 10, 2361.
75. Chou, F.L.; Chou, H.C.; Lin, Y.S.; Yang, B.Y.; Lin, N.T.; Weng, S.F.; Tseng, Y.H. The *Xanthomonas campestris* gumD gene required for synthesis of xanthan gum is involved in normal pigmentation and virulence in causing black rot. *Biochem. Biophys. Res. Commun.* 1997, 233, 265–269.
76. Katzen, F.; Ferreira, D.U.; Oddo, C.G.; Ielmini, M.V.; Becker, A.; Pühler, A.; Ielpi, L. *Xanthomonas campestris* pv. *campestris* gum mutants: Effects on xanthan biosynthesis and plant virulence. *J. Bacteriol.* 1998, 180, 1607–1617.
77. Dharmapuri, S.; Sonti, R.V. A transposon insertion in the gumG homologue of *Xanthomonas oryzae* pv. *oryzae* causes loss of extracellular polysaccharide production and virulence. *FEMS Microbiol. Lett.* 1999, 179, 53–59.
78. Kemp, B.P.; Horne, J.; Bryant, A.; Cooper, R.M. *Xanthomonas axonopodis* pv. *manihotis* gumD gene is essential for EPS production and pathogenicity and enhances epiphytic survival on cassava (*Manihot esculenta*). *Physiol. Mol. Plant Pathol.* 2004, 64, 209–218.
79. Dow, J.M.; Crossman, L.; Findlay, K.; He, Y.Q.; Feng, J.X.; Tang, J.L. Biofilm dispersal in *Xanthomonas campestris* is controlled by cell-cell signaling and is required for full virulence to plants. *Proc. Natl. Acad. Sci. USA* 2003, 100, 10995–11000.
80. Martins, P.M.M.; Merfa, M.V.; Takita, M.A.; De Souza, A.A. Persistence in Phytopathogenic Bacteria: Do We Know Enough? *Front. Microbiol.* 2018, 9, 1099.
81. Ghezzi, J.I.; Steck, T.R. Induction of the viable but non-culturable condition in *Xanthomonas campestris* pv. *campestris* in liquid microcosms and sterile soil. *FEMS Microbiol. Ecol.* 1999, 30, 203–208.
82. Martins, P.M.M.; Wood, T.K.; de Souza, A.A. Persister Cells Form in the Plant Pathogen *Xanthomonas citri* subsp. *citri* under Different Stress Conditions. *Microorganisms* 2021, 9, 384.
83. Harper, D.R.; Parracho, H.M.R.T.; Walker, J.; Sharp, R.; Hughes, G.; Werthen, M.; Lehman, S.; Morales, S. Bacteriophages and Biofilms. *Antibiotics* 2014, 3, 270–284.
84. Dow, M.; Newman, M.A.; von Roepenack, E. The Induction and Modulation of Plant Defense Responses by Bacterial Lipopolysaccharides. *Annu. Rev. Phytopathol.* 2000, 38, 241–261.
85. Meyer, A.; Pühler, A.; Niehaus, K. The lipopolysaccharides of the phytopathogen *Xanthomonas campestris* pv. *campestris* induce an oxidative burst reaction in cell cultures of *Nicotiana tabacum*. *Planta* 2001, 213, 214–222.
86. Kingsley, M.T.; Gabriel, D.W.; Marlow, G.C.; Roberts, P.D. The opsX locus of *Xanthomonas campestris* affects host range and biosynthesis of lipopolysaccharide and extracellular polysaccharide. *J. Bacteriol.* 1993, 175, 5839–5850.
87. Dow, J.M.; Osbourn, A.E.; Wilson, T.J.; Daniels, M.J. A locus determining pathogenicity of *Xanthomonas campestris* is involved in lipopolysaccharide biosynthesis. *Mol. Plant. Microbe Interact.* 1995, 8, 768–777.
88. Newman, M.A.; Dow, J.M.; Daniels, M.J. Bacterial lipopolysaccharides and plant–pathogen interactions. *Eur. J. Plant Pathol.* 2001, 107, 95–102.

89. Gerlach, R.G.; Hensel, M. Protein secretion systems and adhesins: The molecular armory of Gram-negative pathogens. *Int. J. Med. Microbiol.* 2007, 297, 401–415.
90. Yang, X.; Long, M.; Shen, X. Effector–Immunity Pairs Provide the T6SS Nanomachine its Offensive and Defensive Capabilities. *Molecules* 2018, 23, 1009.
91. Boch, J.; Scholze, H.; Schornack, S.; Landgraf, A.; Hahn, S.; Kay, S.; Lahaye, T.; Nickstadt, A.; Bonas, U. Breaking the code of DNA binding specificity of TAL-type III effectors. *Science* 2009, 326, 1509–1512.
92. Lang, J.M.; Perez-Quintero, A.L.; Koebnik, R.; DuCharme, E.; Sarra, S.; Doucoure, H.; Keita, I.; Ziegler, J.; Jacobs, J. M.; Oliva, R.; et al. A Pathovar of *Xanthomonas oryzae* Infecting Wild Grasses Provides Insight Into the Evolution of Pathogenicity in Rice Agroecosystems. *Front. Plant. Sci.* 2019, 10, 507.
93. Ruh, M.; Briand, M.; Bonneau, S.; Jacques, M.A.; Chen, N.W.G. *Xanthomonas* adaptation to common bean is associated with horizontal transfers of genes encoding TAL effectors. *BMC Genom.* 2017, 18, 670.
94. Dennehy, J.J.; Abedon, S.T. Bacteriophage Ecology. In *Bacteriophages*; Harper, D.R., Abedon, S.T., Burrowes, B.H., McConville, M.L., Eds.; Springer Nature: Cham, Switzerland, 2021; pp. 253–294.
95. Born, Y.; Fieseler, L.; Klumpp, J.; Eugster, M.R.; Zurfluh, K.; Duffy, B.; Loessner, M.J. The tail-associated depolymerase of *Erwinia amylovora* phage L1 mediates host cell adsorption and enzymatic capsule removal, which can enhance infection by other phage. *Environ. Microbiol.* 2014, 16, 2168–2180.
96. Hargreaves, K.R.; Kropinski, A.M.; Clokie, M.R. What does the talking?: Quorum sensing signalling genes discovered in a bacteriophage genome. *PLoS ONE* 2014, 9, e85131.
97. Silpe, J.E.; Bassler, B.L. A Host-Produced Quorum-Sensing Autoinducer Controls a Phage Lysis-Lysogeny Decision. *Cell* 2019, 176, 268.e13–280.e13.
98. He, Y.W.; Zhang, L.H. Quorum sensing and virulence regulation in *Xanthomonas campestris*. *FEMS Microbiol. Rev.* 2008, 32, 842–857.
99. Konopka, A.; Lindemann, S.; Fredrickson, J. Dynamics in microbial communities: Unraveling mechanisms to identify principles. *ISME J.* 2015, 9, 1488–1495.
100. Louca, S.; Doebeli, M. Taxonomic variability and functional stability in microbial communities infected by phages. *Environ. Microbiol.* 2017, 19, 3863–3878.
101. Braga, L.P.; Soucy, S.M.; Amgarten, D.E.; da Silva, A.M.; Setubal, J.C. Bacterial Diversification in the Light of the Interactions with Phages: The Genetic Symbionts and Their Role in Ecological Speciation. *Front. Ecol. Evol.* 2018, 6.
102. Molnar, J.; Magyar, B.; Schneider, G.; Laczi, K.; Valappil, S.K.; Kovacs, A.L.; Nagy, I.K.; Rakhely, G.; Kovacs, T. Identification of a novel archaea virus, detected in hydrocarbon polluted Hungarian and Canadian samples. *PLoS ONE* 2020, 15, e0231864.
103. Blount, Z.D.; Borland, C.Z.; Lenski, R.E. Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 2008, 105, 7899–7906.
104. Hulin, M.T.; Armitage, A.D.; Vicente, J.G.; Holub, E.B.; Baxter, L.; Bates, H.J.; Mansfield, J.W.; Jackson, R.W.; Harrison, R.J. Comparative genomics of *Pseudomonas syringae* reveals convergent gene gain and loss associated with specialization onto cherry (*Prunus avium*). *New Phytol.* 2018, 219, 672–696.
105. Zhang, H.L.; Ntambo, M.S.; Rott, P.C.; Chen, G.; Chen, L.L.; Huang, M.T.; Gao, S.J. Complete Genome Sequence Reveals Evolutionary and Comparative Genomic Features of *Xanthomonas albilineans* Causing Sugarcane Leaf Scald. *Microorganisms* 2020, 8, 182.
106. Labrie, S.J.; Samson, J.E.; Moineau, S. Bacteriophage resistance mechanisms. *Nat. Rev. Microbiol.* 2010, 8, 317–327.
107. Weiss, B.D.; Capage, M.A.; Kessel, M.; Benson, S.A. Isolation and characterization of a generalized transducing phage for *Xanthomonas campestris* pv. *campestris*. *J. Bacteriol.* 1994, 176, 3354–3359.
108. Newberry, E.A.; Bhandari, R.; Minsavage, G.V.; Timilsina, S.; Jibrin, M.O.; Kemble, J.; Sikora, E.J.; Jones, J.B.; Potnis, N. Independent Evolution with the Gene Flux Originating from Multiple *Xanthomonas* Species Explains Genomic Heterogeneity in *Xanthomonas perforans*. *Appl. Environ. Microbiol.* 2019, 85.
109. Bartoli, C.; Roux, F.; Lamichhane, J.R. Molecular mechanisms underlying the emergence of bacterial pathogens: An ecological perspective. *Mol. Plant Pathol.* 2016, 17, 303–310.
110. Lima, W.C.; Paquola, A.C.; Varani, A.M.; Van Sluys, M.A.; Menck, C.F. Laterally transferred genomic islands in *Xanthomonadales* related to pathogenicity and primary metabolism. *FEMS Microbiol. Lett.* 2008, 281, 87–97.
111. da Silva, A.C.; Ferro, J.A.; Reinach, F.C.; Farah, C.S.; Furlan, L.R.; Quaggio, R.B.; Monteiro-Vitorello, C.B.; Van Sluys, M.A.; Almeida, N.F.; Alves, L.M.; et al. Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature* 2002, 417, 459–463.

112. Liang, X.; Wagner, R.E.; Li, B.; Zhang, N.; Radosevich, M. Quorum Sensing Signals Alter *in vitro* Soil Virus Abundance and Bacterial Community Composition. *Front Microbiol.* 2020, 11, 1287.
113. Ghosh, D.; Roy, K.; Williamson, K.E.; Srinivasiah, S.; Wommack, K.E.; Radosevich, M. Acyl-homoserine lactones can induce virus production in lysogenic bacteria: An alternative paradigm for prophage induction. *Appl. Environ. Microbiol.* 2009, 75, 7142–7152.
114. Erez, Z.; Steinberger-Levy, I.; Shamir, M.; Doron, S.; Stokar-Avihail, A.; Peleg, Y.; Melamed, S.; Leavitt, A.; Savidor, A.; Albeck, S.; et al. Communication between viruses guides lysis-lysogeny decisions. *Nature* 2017, 541, 488–493.
115. Stokar-Avihail, A.; Tal, N.; Erez, Z.; Lopatina, A.; Sorek, R. Widespread Utilization of Peptide Communication in Phages Infecting Soil and Pathogenic Bacteria. *Cell Host Microbe* 2019, 25, 746–755.e745.
116. Wiggins, B.A.; Alexander, M. Minimum bacterial density for bacteriophage replication: Implications for significance of bacteriophages in natural ecosystems. *Appl. Environ. Microbiol.* 1985, 49, 19–23.
117. Adhikari, N.; Acharya, K.P. Effectiveness of Bacteriophage Therapy in Field Conditions and Possible Future Applications. *Curr. Pharm. Biotechnol.* 2020, 21, 364–373.
118. Hung, C.H.; Wu, H.C.; Tseng, Y.H. Mutation in the *Xanthomonas campestris* *xanA* gene required for synthesis of xanthan and lipopolysaccharide drastically reduces the efficiency of bacteriophage (phi)L7 adsorption. *Biochem. Biophys. Res. Commun.* 2002, 291, 338–343.
119. Oechslin, F. Resistance Development to Bacteriophages Occurring during Bacteriophage Therapy. *Viruses* 2018, 10.
120. Torres-Barcelo, C. The disparate effects of bacteriophages on antibiotic-resistant bacteria. *Emerg. Microbes Infect.* 2018, 7, 168.
121. Bertozzi Silva, J.; Storms, Z.; Sauvageau, D. Host receptors for bacteriophage adsorption. *FEMS Microbiol. Lett.* 2016, 363.
122. Bondy-Denomy, J.; Pawluk, A.; Maxwell, K.L.; Davidson, A.R. Bacteriophage genes that inactivate the CRISPR/Cas bacterial immune system. *Nature* 2013, 493, 429–432.
123. Seed, K.D.; Lazinski, D.W.; Calderwood, S.B.; Camilli, A. A bacteriophage encodes its own CRISPR/Cas adaptive response to evade host innate immunity. *Nature* 2013, 494, 489–491.
124. Ahmad, A.A.; Askora, A.; Kawasaki, T.; Fujie, M.; Yamada, T. The filamentous phage XacF1 causes loss of virulence in *Xanthomonas axonopodis* pv. *citri*, the causative agent of citrus canker disease. *Front Microbiol.* 2014, 5, 321.
125. Borah, P.; Jindal, J.; Verma, J. Integrated management of bacterial leaf spot of mungbean with bacteriophages of Xav and chemicals. *J. Mycol. Plant Pathol.* 2000, 30, 19–21.
126. Mallmann, W.; Hemstreest, C. Isolation of an inhibitory substance from plants. *Agric. Res.* 1924, 28, 599–602.
127. Civerolo, E.L.; Keil, H.L. Inhibition of bacterial spot of peach foliage by *Xanthomonas pruni* bacteriophage. *Phytopathology* 1969, 59, 1966–1967.
128. Civerolo, E.L. Relationship of *Xanthomonas pruni* bacteriophages to bacterial spot disease in *Prunus*. *Phytopathology* 1973, 63, 1279–1284.
129. Zaccardelli, M.; Saccardi, A.; Gambin, E.; Mazzucchi, U. *Xanthomonas campestris* pv. *pruni* bacteriophages on peach trees and their potential use for biological control. *Phytopathol. Mediterr.* 1992, 31, 133–140.
130. Saccardi, A.; Gambin, E.; Zaccardelli, M.; Barone, G.; Mazzucchi, U. *Xanthomonas campestris* pv. *pruni* control trials with phage treatments on peaches in the orchard. *Phytopathol. Mediterr.* 1993, 32, 206–210.
131. Flaherty, J.E.; Jones, J.B.; Harbaugh, B.K.; Somodi, G.C.; Jackson, L.E. Control of bacterial spot on tomato in the greenhouse and field with H-mutant bacteriophages. *HortScience* 2000, 35, 882–884.
132. Balogh, B. Strategies of Improving the Efficacy of Bacteriophages for Controlling Bacterial Spot of Tomato; University of Florida: Gainesville, FL, USA, 2002.
133. Balogh, B.; Jones, J.B.; Momol, M.T.; Olson, S.M.; Obradovic, A.; King, P.; Jackson, L.E. Improved Efficacy of Newly Formulated Bacteriophages for Management of Bacterial Spot on Tomato. *Plant Dis.* 2003, 87, 949–954.
134. Balogh, B.; Jones, J.B.; Momol, M.T.; Olson, M. S. Persistence of bacteriophages as biocontrol agents in the tomato canopy. *Acta Hortic.* 2005, 695, 299–302.
135. Obradovic, A.; Jones, J.B.; Momol, M.T.; Balogh, B.; Olson, S.M. Management of Tomato Bacterial Spot in the Field by Foliar Applications of Bacteriophages and SAR Inducers. *Plant Dis.* 2004, 88, 736–740.
136. Obradovic, A.; Jones, J.B.; Momol, M.T.; Olson, S.M.; Jackson, L.E.; Balogh, B.; Guven, K.; Iriarte, F.B. Integration of Biological Control Agents and Systemic Acquired Resistance Inducers Against Bacterial Spot on Tomato. *Plant Dis.* 2005, 89, 712–716.

137. Jones, J.B.; Momol, M.T.; Obradovic, A.; Balogh, B.; Olson, S.M. Bacterial spot management on tomatoes. *Acta Hort* 2005, 695, 119–124.
138. Jones, J.B.; Jackson, L.E.; Balogh, B.; Obradovic, A.; Iriarte, F.B.; Momol, M.T. Bacteriophages for plant disease control. *Annu. Rev. Phytopathol.* 2007, 45, 245–262.
139. Obradovic, A.; Jones, J.B.; Balogh, B.; Momol, M.T. Integrated management of tomato bacterial spot. In *Integrated Management of Plant Diseases Caused by Fungi, Phytoplasma and Bacteria*; Ciancio, A., Mukerji, K.G., Eds.; Springer Science + Business Media B. V.: Berlin/Heidelberg, Germany, 2008; pp. 211–223.
140. Abrahamian, P.; Jones, J.B.; Vallad, G.E. Efficacy of copper and copper alternatives for management of bacterial spot on tomato under transplant and field production. *Crop. Prot.* 2019, 126, 104919.
141. Gašić, K.; Kuzmanović, N.; Ivanović, M.; Prokić, A.; Šević, M.; Obradović, A. Complete Genome of the *Xanthomonas vesicatoria* Specific Bacteriophage KΦ1, Its Survival and Potential in Control of Pepper Bacterial Spot. *Front Microbiol.* 2018, 9, 2021.
142. Šević, M.; Gašić, K.; Ignjatov, M.; Mijatović, M.; Prokić, A.; Obradovic, A. Integration of biological and conventional treatments in control of pepper bacterial spot. *Crop. Prot.* 2019, 119, 46–51.
143. Lang, J.M.; Gent, D.H.; Schwartz, H.F. Management of *Xanthomonas* Leaf Blight of Onion with Bacteriophages and a Plant Activator. *Plant Dis.* 2007, 91, 871–878.
144. Balogh, B. Characterization and Use of Bacteriophages Associated with Citrus Bacterial Pathogens for Disease Control; University of Florida: Gainesville, FL, USA, 2006.
145. Balogh, B.; Canteros, B.I.; Stall, R.E.; Jones, J.B. Control of Citrus Canker and Citrus Bacterial Spot with Bacteriophages. *Plant Dis.* 2008, 92, 1048–1052.
146. Ibrahim, Y.E.; Saleh, A.A.; Al-Saleh, M.A. Management of Asiatic Citrus Canker Under Field Conditions in Saudi Arabia Using Bacteriophages and Acibenzolar-S-Methyl. *Plant Dis.* 2017, 101, 761–765.
147. Kuo, T.T.; Chang, L.C.; Yang, C.M.; Yang, S. E. Bacterial leaf blight of rice plant IV. Effect of bacteriophage on the infectivity of *Xanthomonas oryzae*. *Bot. Bull. Acad. Sin.* 1971, 12, 1–9.
148. Chae, J.C.; Hung, N.B.; Yu, S.M.; Lee, H.K.; Lee, Y.H. Diversity of bacteriophages infecting *Xanthomonas oryzae* pv. *oryzae* in paddy fields and its potential to control bacterial leaf blight of rice. *J. Microbiol. Biotechnol.* 2014, 24, 740–747.
149. Ogunyemi, S.O.; Chen, J.; Zhang, M.; Wang, L.; Masum, M.M.I.; Yan, C.; An, Q.; Li, B.; Chen, J. Identification and characterization of five new OP2-related Myoviridae bacteriophages infecting different strains of *Xanthomonas oryzae* pv. *oryzae*. *J. Plant Pathol.* 2019, 101, 263–273.
150. Flaherty, J.E.; Harbaugh, B.K.; Jones, J.B.; Somodi, G.C.; Jackson, L.E. H-mutant bacteriophages as a potential biocontrol of bacterial blight of geranium. *HortScience* 2001, 36, 98–100.
151. Nagai, H.; Miyake, N.; Kato, S.; Maekawa, D.; Inoue, Y.; Takikawa, Y. Improved control of black rot of broccoli caused by *Xanthomonas campestris* pv. *campestris* using a bacteriophage and a nonpathogenic *Xanthomonas* sp. strain. *J. Gen. Plant Pathol.* 2017, 83, 373–381.
152. Orynbayev, A.; Dzhaliilov, F.; Ignatov, A. Improved efficacy of formulated bacteriophage in control of black rot caused by *Xanthomonas campestris* pv. *campestris* on cabbage seedlings. *Arch. Phytopathol. Plant Prot.* 2020, 379–394.