

Spring Wheat

Subjects: **Plant Sciences**

Contributor: Hafeez Rehman , Matthew Reynolds

Spring wheat as durum or bread wheat is grown worldwide in mega-environments (ME; 1-6) at latitudes <40 degree during the winter or early spring season and flower in short days with relatively little vernalization requirement. Based on protein contents, kernel color and hardness of endosperm, spring wheat is classified into hard red, soft white and hard white spring with high baking quality.

megaenvironments

flowering

vernalization

heat stress

1. Introduction

Spring wheat, is a tetraploid (durum) or hexaploid (bread wheat) that is grown worldwide covering approximately 80 m ha in developing countries. Based on the growth habit, spring wheat can flower under short days with relatively little exposure to vernalization and the crop usually matures within 100-200 days. It is usually grown at latitudes <40° during the winter and early spring season [1]. Though relatively insensitive to photoperiod and vernalizing cold compared to winter wheat, flowering time and phenological pattern of spring wheat is largely a function of major genes *Ppd* and *Vrn* as well as earliness per se (*Eps*) which determines the temperature baseline for crop development [2][3].




Figure 1; CIMMYT wheat production and breeding mega-environments (MEs) adapted from (Braun et al. 2010), by Kai Sonder

2. Spring wheat mega-environments and characteristics

According to the International Maize and Wheat Improvement Center (CIMMYT) classification, among 12 mega-environments (MEs), spring wheat is produced in six meg-environments including irrigated temperate, high rainfall, acid soils, low rainfall, high temperature and high latitude showing its diverse distribution [4]. These MEs especially ME1-ME6 are classified for spring wheat, ME7-ME9 for facultative and ME10-ME12 for winter wheat (Figure 1). The concept of MEs classification in broaden is used to breed the characteristic germplasm for target environment with adaptation to major biotic and abiotic stresses (Braun et al., 2010). Later this classification has been expanded to classify these MEs based on geospatial information to address the impacts of climate change on wheat production [5][6]. For instance, ME1-ME5 include sub-tropical to tropical regions (ME5) where spring wheat is cultivated on about 9 million ha that experiences high temperature stress induced yield losses [7]. The ME5 with its subdivision (ME5A and ME5B) for humid and dry conditions, respectively include Eastern Gangetic Plains of India, Nepal and Bangladesh under 5B and Sudan and upper Egypt for example under 5A. Under changing climate conditions, with increasing temperature, much of the wheat area under ME1 is expected to change to ME5 (Hodson and White, 2007 a, b). However, increased temperature with elevated CO₂ atmospheric levels may

mitigate some of the negative impacts on wheat productivity and water use efficiency in ME1. Without significant breeding and crop management interventions, increased temperatures are expected to reduce area in ME2A, ME3, ME4 and ME5 for wheat production (Braun et al. 2010). Spring planted crops can be easily recognized by growth stages following Zadoks system [8].

3. Spring wheat classification based on kernel characteristics

Based on protein contents, color of kernel and hardness of endosperm spring wheat is also classified into hard red spring, soft white and hard white spring [9]. Spring wheat have smaller grains usually rich in protein with high baking quality [10].

References

1. Braun, H. J., Atlin, G. and Payne, T. . Multi-location testing as a tool to identify plant response to global climate change; CABI: UK , 2010; pp. 115-138 .
2. J. Le Gouis; J. Bordes; Catherine Ravel; E. Heumez; S. Faure; S. Praud; N. Galic; C. Remoué; F. Balfourier; V. Allard; et al.M. Rousset Genome-wide association analysis to identify chromosomal regions determining components of earliness in wheat. *Theoretical and Applied Genetics* **2011**, 124, 597-611, 10.1007/s00122-011-1732-3.
3. Atif Kamran; Muhammad Iqbal; Dean Spaner; Flowering time in wheat (*Triticum aestivum* L.): a key factor for global adaptability. *Euphytica* **2014**, 197, 1-26, 10.1007/s10681-014-1075-7.
4. Rajaram, S., van Ginkel, M., Fischer, R.A., 1994. In: Li, Z.S., Xin, Z.Y. (Eds.), CIMMYT's Wheat Breeding Mega-Environments (ME). Proceedings of the 8th International Wheat Genetics Symposium, Beijing, China, 20–25 July 1993. China Agriculture Scientechn Press, Beijing, pp. 1101–1106.
5. D. P. Hodson; J. W. White; PAPER PRESENTED AT INTERNATIONAL WORKSHOP ON INCREASING WHEAT YIELD POTENTIAL, CIMMYT, OBREGON, MEXICO, 20–24 MARCH 2006 Use of spatial analyses for global characterization of wheat-based production systems. *The Journal of Agricultural Science* **2007**, 145, 115, 10.1017/s0021859607006855.
6. Hodson, D. and White, J.. Climate change: what future wheat? ; CIMMYT: Mexico, 2007; pp. 41-48.
7. M. Lillemo; M. Van Ginkel; R. M. Trethowan; E. Hernandez; José Crossa; Differential Adaptation of CIMMYT Bread Wheat to Global High Temperature Environments. *Crop Science* **2005**, 45, 2443-2453, 10.2135/cropsci2004.0663.

8. J. C. Zadoks; T. T. Chang; C. F. Konzak; A decimal code for the growth stages of cereals. *Weed Research* **1974**, *14*, 415-421, 10.1111/j.1365-3180.1974.tb01084.x.
9. Mergoum M., Singh P., Pena R., Lozano-del Río A., Cooper K., Salmon D., et al.. *Triticale: a "new" crop with old challenges*; Springer: UK, 2009; pp. 267–287.
10. Dziki, D.; Cacak-Pietrzak, G.; Gawlik-Dziki, U.; Swieca, M.; Miś, A.; Rózyło, R.; Jończyk, K. Physicochemical properties and milling characteristics of spring wheat from different farming systems. *J. Agric. Sci. Technol.* **2017**, *19*, 1253–1266

Retrieved from <https://encyclopedia.pub/entry/history/show/33101>