

# Winchester Syndrome

Subjects: **Genetics & Heredity**

Contributor: Bruce Ren

Winchester syndrome is a rare inherited disease characterized by a loss of bone tissue (osteolysis), particularly in the hands and feet.

genetic conditions

## 1. Introduction

Winchester syndrome is a rare inherited disease characterized by a loss of bone tissue (osteolysis), particularly in the hands and feet. Winchester syndrome used to be considered part of a related condition now called multicentric osteolysis, nodulosis, and arthropathy (MONA). However, because Winchester syndrome and MONA are caused by mutations in different genes, they are now thought to be separate disorders.

In most cases of Winchester syndrome, bone loss begins in the hands and feet, causing pain and limiting movement. Bone abnormalities later spread to other parts of the body, with joint problems (arthropathy) occurring in the elbows, shoulders, knees, hips, and spine. Most people with Winchester syndrome develop low bone mineral density (osteopenia) and thinning of the bones (osteoporosis) throughout the skeleton. These abnormalities make bones brittle and more prone to fracture. The bone abnormalities also lead to short stature.

Some people with Winchester syndrome have skin abnormalities including patches of dark, thick, and leathery skin. Other features of the condition can include clouding of the clear front covering of the eye (corneal opacity), excess hair growth (hypertrichosis), overgrowth of the gums, heart abnormalities, and distinctive facial features that are described as "coarse."

## 2. Frequency

Winchester syndrome is a rare condition whose prevalence is unknown. It has been reported in only a few individuals worldwide.

## 3. Causes

Winchester syndrome is caused by mutations in the *MMP14* gene (also known as *MT1-MMP*). This gene provides instructions for making a protein called matrix metallopeptidase 14, which is found on the surface of cells. Matrix metallopeptidase 14 normally helps modify and break down various components of the extracellular matrix, which

is the intricate lattice of proteins and other molecules that forms in the spaces between cells. These changes influence many cell activities and functions, including promoting cell growth and stimulating cell movement (migration). Matrix metallopeptidase 14 also turns on (activates) a protein called matrix metallopeptidase 2. The activity of matrix metallopeptidase 2 appears to be important for a variety of body functions, including bone remodeling, which is a normal process in which old bone is broken down and new bone is created to replace it.

Mutations in the *MMP14* gene alter matrix metallopeptidase 14 so that less of the enzyme is able to reach the cell surface. As a result, not enough of the enzyme is available to break down components of the extracellular matrix and activate matrix metallopeptidase 2. It is unclear how a shortage of this enzyme leads to the signs and symptoms of Winchester syndrome. It is possible that a loss of matrix metallopeptidase 2 activation somehow disrupts the balance of new bone creation and the breakdown of existing bone during bone remodeling, causing a progressive loss of bone tissue. How a reduced amount of matrix metallopeptidase 14 leads to the other features of Winchester syndrome is unknown.

### 3.1 The gene associated with Winchester syndrome

- MMP14

## 4. Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

## 5. Other Names for This Condition

- Winchester disease
- WNCHRS

## References

1. Evans BR, Mosig RA, Lobl M, Martignetti CR, Camacho C, Grum-Tokars V, Glucksman MJ, Martignetti JA. Mutation of membrane type-1 metalloproteinase, MT1-MMP, causes the multicentric osteolysis and arthritis disease Winchester syndrome. *Am J Hum Genet.* 2012 Sep 7;91(3):572-6. doi:10.1016/j.ajhg.2012.07.022.
2. Itoh Y, Seiki M. MT1-MMP: a potent modifier of pericellular microenvironment. *J Cell Physiol.* 2006 Jan;206(1):1-8. Review.
3. Zankl A, Pachman L, Poznanski A, Bonafé L, Wang F, Shusterman Y, Fishman DA, Superti-Furga A. Torg syndrome is caused by inactivating mutations in MMP2 and is allelic to NAO and

Winchester syndrome. J Bone Miner Res. 2007 Feb;22(2):329-33.

---

Retrieved from <https://encyclopedia.pub/entry/history/show/13852>